

# involve

a journal of mathematics

Outer billiards and tilings of the hyperbolic plane

Filiz Dogru, Emily M. Fischer and Cristian Mihai Munteanu





# Outer billiards and tilings of the hyperbolic plane

Filiz Dogru, Emily M. Fischer and Cristian Mihai Munteanu

(Communicated by Kenneth S. Berenhaut)

We present new results regarding the periodicity of outer billiards in the hyperbolic plane around polygonal tables which are tiles in regular two-piece tilings of the hyperbolic plane.

## 1. Introduction

Outer billiards is a simple dynamical system introduced by B. H. Neumann [1959]. J. Moser [1973; 1978] popularized outer billiards as a toy model for planetary motion as a means of finding possible unbounded orbits. Since then, many mathematicians have asked and answered questions about outer billiards systems in various geometries. For example, C. Culter proved in 2004 the existence of periodic orbits for polygonal tables in the Euclidean plane (the proof is presented by S. Tabachnikov [2007]). R. Schwartz [2007; 2009] answered, in the affirmative, Moser's question about the existence of unbounded orbits for certain polygons.

The main motivation for this paper is a result of Vivaldi and Shaidenko [1987] that in the Euclidean case, outer billiards associated to quasirational polygons have all orbits bounded; see also [Kołodziej 1989; Gutkin and Simányi 1992]. As a consequence, all orbits about a lattice polygon in the Euclidean plane are periodic. We continue the work of Dogru and Tabachnikov [2003], who studied the relationship between one-tile regular tilings of the hyperbolic plane and the outer billiards system.

For a detailed account of hyperbolic geometry and the hyperbolic plane, we direct the reader to [Greenberg 1980], and for a survey of outer billiards, see [Tabachnikov and Dogru 2005; Tabachnikov 2005].


## 2. Definitions

The outer billiard map associated to a convex polygonal table  $P$  in the hyperbolic plane is defined as follows. For a point  $x \in \mathbb{H}^2 \setminus P$ , there are two lines that pass

---

*MSC2010:* 37E15.

*Keywords:* hyperbolic billiards, outer billiards, polygonal billiards, symbolic dynamics, tiling, rotation number, crochét.



**Figure 1.** Outer billiards map in the Klein model.

through  $x$  and are tangent to the table  $P$ . By convention, we consider the tangent line for which  $P$  is on the left, from the point of view of  $x$ . Then we reflect  $x$  about the tangency (support) point to get  $T(x)$  (see Figure 1). The map is well-defined whenever the tangency point is unique and so we are able to define the map  $T$  on the entire hyperbolic plane except for the clockwise continuations of the sides of  $P$  (see Figure 1) and their preimages under  $T$ . An immediate consequence of the definition is that  $T$  is a piecewise isometry.

Likewise, the inverse map  $T^{-1}$  is not defined on the counterclockwise continuations of the sides of  $P$ . We define the *web* associated to  $P$  to be the union of all preimages under  $T$  of the clockwise continuation of the sides and of all preimages under  $T^{-1}$  of the counterclockwise continuation of the sides. For each connected component of the complement of the web, the restriction of the map  $T^n$  to that component is defined by a single isometry of the hyperbolic plane for every  $n \in \mathbb{Z}$ . That means that each connected component of the complement of the web maps as a whole under the iterations of  $T$ .


Another feature of the billiards map  $T$  is that it extends continuously to a continuous circle map  $t : S^1 \rightarrow S^1$  at infinity. The map  $t$  is defined using the same reflecting procedure. In this case, the uniqueness of the support point is not needed since the distance between our initial point and the support point is infinite no matter the choice, and hence the map  $t$  is well-defined for every point at infinity. Since  $t$  is a circle map, it has a well-defined Poincaré rotation number  $\rho(t)$ , and we will prove in Section 3 that  $\rho(t)$  encodes information about the combinatorial dynamics of the outer billiards.

### 3. Outer billiards on tilings

We are studying the hyperbolic outer billiards map associated with a polygonal table that is part of a two-piece regular tiling of the hyperbolic plane. These tilings use two polygonal pieces, a regular  $M$ -gon and a regular  $N$ -gon that meet four in each vertex (see Figure 2). We describe the combinatorial dynamics for outer billiards around one of the  $M$ -gons. We note that the web associated to such a map will fall exactly on the grid lines of the tiling. This is because the reflection around a vertex of the table tile is just a rotation by  $180^\circ$  around vertices in the tiling. It follows that each tile maps as a whole under iterations of  $T$ .

**3.1. Previous results.** Previous results describing outer billiards of tiles in the hyperbolic plane are obtained in [Dogru and Tabachnikov 2003]. In this paper, the authors have proved that every orbit of the outer billiard map around a right-angled regular  $n$ -gon, for  $n \geq 5$ , is periodic. Any right-angled regular  $n$ -gon generates a tiling of the hyperbolic plane entirely consisting of  $n$ -gons. The theorems proven in the next sections have the same flavor as Theorem 4 in the above mentioned paper.

Define the *rank* of a tile as the minimum number of sides that one has to cross, when starting inside the table, to get to the given tile. This means that tiles that have one common side with the table have rank 1, and tiles that have a common side with a tile of rank 1 have rank 2, and so on.



**Figure 2.** Example of  $(M, N)$ -tiling for  $(M, N) = (6, 7)$ .

**Theorem 1** [Dogru and Tabachnikov 2003]. *For a tiling of regular  $n$ -gons meeting in four,  $n \geq 5$ , the dual billiard map  $T$  preserves the rank of a tile, and every orbit of  $T$  is periodic. The set of rank  $k$  tiles consists of*

$$q_k = n \frac{\lambda_1^k - \lambda_2^k}{\lambda_1 - \lambda_2}$$

*elements, where*

$$\lambda_{1,2} = \frac{n-2 \pm \sqrt{n(n-4)}}{2}$$

*are the roots of the equation  $\lambda^2 - (n-2)\lambda + 1 = 0$ . The action of  $T$  on the set of rank  $k$  tiles is a transitive cyclic permutation  $i \mapsto i + p_k$ , where*

$$p_k = \frac{\lambda_1^{k-1} - \lambda_2^{k-1}}{\lambda_1 - \lambda_2} + \frac{\lambda_1^k - \lambda_2^k}{\lambda_1 - \lambda_2}.$$

*The rotation number of the dual billiard map at infinity is given by the formula*

$$\rho(t) = \lim_{k \rightarrow \infty} \frac{p_k}{q_k} = \frac{n - \sqrt{n(n-4)}}{2n}.$$

The proof of this theorem uses geometric arguments for the periodicity of orbits and recurrence formulas for computing the number of tiles in each rank and the rotation number of  $t$  (see [Dogru and Tabachnikov 2003] for details). The authors make an important remark that the representation of  $\lambda_1$  (and so the rotation number of the map at infinity) as a continued fraction encodes the dynamics of the tiles under the billiard map  $T$ . We will deduce similar results for two-piece tilings.

**3.2. New results.** Our results extend Theorem 1 to two-piece regular tilings of the hyperbolic plane. We will denote a tiling of regular  $M$ -gons and regular  $N$ -gons as an  $(M, N)$ -tiling, and we will always consider the table to be an  $M$ -gon. Such an  $(M, N)$ -tiling exists if  $\frac{1}{M} + \frac{1}{N} < \frac{1}{2}$ . As mentioned earlier, these tilings have four shapes meeting at each vertex, two  $M$ -gons and two  $N$ -gons.

**3.2.1. Triangles and  $N$ -gons.** Most of the geometric arguments used here are analogous to those used by Dogru and Tabachnikov. Our counting arguments are different, although they are also based on recurrence relations.

Let us introduce a more general notation for rank in order to avoid cumbersome indexing. Observe that the layer of tiles of rank  $k$  includes tiles of the same type (all  $M$ -gons or all  $N$ -gons) and as rank changes by one, that shape changes. So triangles always have even rank and  $N$ -gons always have odd rank. We will say that a rank  $2k-1$  tile is a rank  $k$   $N$ -gon and a rank  $2k$  tile is a rank  $k$  triangle. The rest of this section is dedicated to describing the dynamics of the billiard map  $T$  in the  $(3, N)$ -tilings through the proof of the following theorem:

**Theorem 2.** *For a  $(3, N)$ -tiling,  $N \geq 7$ , the outer billiard map  $T$  preserves the rank of a tile and every orbit of  $T$  is periodic. The set of rank  $k$   $N$ -gons consists of*

$$q_k = \frac{1}{\sqrt{N-6}}(\Phi_1^{2k-3} + \Phi_2^{2k-3}) + \Phi_1^{2k-2} + \Phi_2^{2k-2}$$

*elements and the set of rank  $k$  triangles consists of*

$$l_k = \frac{N-4}{\sqrt{N-6}}(\Phi_1^{2k-3} + \Phi_2^{2k-3}) + (N-3)(\Phi_1^{2k-2} + \Phi_2^{2k-2})$$

*elements, where*

$$\Phi_{1,2} = \frac{\sqrt{N-6} \pm \sqrt{N-2}}{2}$$

*are the two roots of the equation*

$$\Phi^2 - \sqrt{N-6}\Phi - 1 = 0.$$

*The action of  $T$  on the set of rank  $k$   $N$ -gons is a cyclic permutation  $i \mapsto i + p_k$ , where*

$$p_k = \frac{q_k}{3} + \frac{\Phi_1^{2k-4} - \Phi_2^{2k-4}}{\sqrt{(N-6)(N-2)}} + \frac{\Phi_1^{2k-3} - \Phi_2^{2k-3}}{\sqrt{N-2}},$$

*and the action of  $T$  on the set of rank  $k$  triangles is also a cyclic permutation  $i \mapsto i + j_k$ , where*

$$j_k = \frac{l_k}{3} + (N-4) \frac{\Phi_1^{2k-4} - \Phi_2^{2k-4}}{\sqrt{(N-6)(N-2)}} + (N-3) \frac{\Phi_1^{2k-3} - \Phi_2^{2k-3}}{\sqrt{N-2}}.$$

*The rotation number of the outer billiard map at infinity is given by the formula*

$$\rho(t) = \lim_{k \rightarrow \infty} \frac{p_k}{q_k} = \lim_{k \rightarrow \infty} \frac{j_k}{l_k} = \frac{1}{3} + \frac{1}{3(1 + \Phi_1^2)} = \frac{1}{3} + \frac{1}{3\sqrt{N-2}\Phi_1}.$$


Theorem 2 contains many independent results and for reasons of clarity we will prove them one by one as claims.

**Claim 3.** *Every orbit of  $T$  is periodic.*

*Proof.* The proof of this result is written in much detail in [Dogru and Tabachnikov 2003]. We will present here a sketch of it and will refer the reader to the above work for detailed explanations. The statement of the claim is a consequence of the following lemma:

**Lemma 4.** *The rank of a tile is preserved under  $T$ .*

*Proof of lemma.* The proof is by induction on the rank, based on geometrical observations. Observe that rank 1 tiles are preserved by  $T$  and notice that every rank  $k$  tile is adjacent to a rank  $k-1$  tile, where these two tiles map together under a single application of  $T$ . These two facts complete the base case and the step of the induction.  $\square$



**Figure 3.** Special case for Lemma 6.

From Lemma 4, since there are finitely many tiles of rank  $k$ , every tile must eventually map back to itself after  $m$  iterations, for some natural number  $m$ . Hence the  $m$ -th iteration of  $T$  maps the entire tile to itself. This implies that  $T^{\circ m}$  (the composition of  $T$  with itself  $m$  times) is a rotation by either  $2\pi j/N$  (for  $N$ -gons) or  $2\pi j/3$  (for triangles) around some point inside the tile. Hence  $T^{\circ Nm}$  restricted to that tile is the identity if the tile is an  $N$ -gon and  $T^{\circ 3m}$  restricted to that tile is the identity if the tile is a triangle. We conclude that every orbit of  $T$  is periodic.  $\square$

**Claim 5.** For every  $k \geq 1$ ,  $T$  permutes the rank  $k$  tiles cyclically.

*Proof.* This claim is an immediate corollary to the following lemma:

**Lemma 6.** Any two consecutive rank  $k$  tiles are mapped to two consecutive rank  $k$  tiles.

*Proof of lemma.* We know by Lemma 4 that the rank of two tiles is preserved under  $T$ . If the two consecutive tiles are not separated by a clockwise continuation of one of the sides of the table then their common point is mapped, together with the two tiles, through the same vertex. Thus the tiles are mapped to two consecutive tiles.

If the two tiles are separated by such a continuation of one side of the table then the argument is more involved. A similar argument is presented in [Dogru and Tabachnikov 2003]. Figure 3 gives a pictorial representation of the situation. The first tile is reflected in  $O_1$ , while the second one is reflected in  $O_2$ . What remains to prove is that  $A'=B'$  so that the images of the two tiles still touch in one point. The following sequence of equalities completes the proof:

$$A'O_2 = A'O_1 - O_1O_2 = BO_1 + AB - O_1O_2 = BO_1 + O_1O_2 = BO_2 = B'O_2. \quad \square$$

In order to compute the formulas for  $q_k, p_k, j_k, l_k$ , we first explain why the tiling we are working with has an intrinsic self-similar geometric structure. We will refer from now on to this self-similar structure as the *crochet pattern*. To describe the crochet pattern, we consider  $N$ -gons to be of two types,  $X$ -type and  $Y$ -type (see Figure 4). Type  $X$   $N$ -gons have two parents in the sense that they touch two  $N$ -gons of the previous rank, while type  $Y$   $N$ -gons touch only one parent. The rank 1  $N$ -gons are of neither of the types, having zero parents, so we call them type 0  $N$ -gons. (This is why our counting argument begins with counting rank 2  $N$ -gons.)



The following claim gives an intuitive explanation of why we call this self-similar structure of the tiling a *crochet pattern*.

**Claim 7.** *When passing from the  $k$ -th layer of  $N$ -gons to the  $(k+1)$ -th layer of  $N$ -gons, we apply the replacement rules*


$$\begin{aligned} X &\rightarrow XY^{N-6}, \\ Y &\rightarrow XY^{N-5}, \end{aligned}$$

*i.e., when incrementing rank of the layer by 1, every  $X$  gets replaced by an  $X$  followed by  $N-6$   $Y$ s, and every  $Y$  gets replaced by an  $X$  followed by  $N-5$   $Y$ s.*

*Proof.* The methods used to prove this claim have been developed by Poincaré, and we will not dwell on the details here. The reader can find extensive explanation in *The Symmetry of Things* [Conway et al. 2008].

Instead, we will illustrate the methods used to prove the claim in the case of  $N = 7$  in order to give the geometrical intuition behind the proof. Figure 4 illustrates the local and global behavior of a  $(3, 7)$ -tiling.

In the local picture, the difference between a type  $X$  7-gon and a type  $Y$  7-gon is encoded in the different types of degenerate heptagons we associate to them. We associate to the  $Y$ -type heptagon a rectangle with three additional points on the upper side, while to the  $X$ -type heptagon we associate a rectangle with two additional points on the upper side and one on the lower side since it has two parents. Now by reducing the triangles in the global picture to points, we notice that the heptagons must meet three in each vertex. This results in the *crochet pattern* shown in Figure 4. This *crochet pattern* immediately implies the claimed replacement rules.  $\square$



**Figure 4.** The  $(3, 7)$ -tiling.

We can now use this crochet pattern to start our counting argument in order to get the exact numbers in Theorem 2.

**Claim 8.** *The formulas for  $q_k$ ,  $p_k$ ,  $j_k$ ,  $l_k$  hold as stated in Theorem 2.*

*Proof.* Denote the number of  $X$ -type and  $Y$ -type  $N$ -gons of rank  $k$  by  $x_k$  and  $y_k$  and use Claim 7 to obtain the system of linear difference equations

$$\begin{pmatrix} x_k \\ y_k \end{pmatrix} = \begin{pmatrix} 1 & 1 \\ N-6 & N-5 \end{pmatrix} \begin{pmatrix} x_{k-1} \\ y_{k-1} \end{pmatrix}.$$

The initial configuration is  $\begin{pmatrix} x_2 \\ y_2 \end{pmatrix} = \begin{pmatrix} 3 \\ 3(N-4) \end{pmatrix}$  because there must be three rank 2  $N$ -gons with two parents, and the rest of the vertices of the rank 1  $N$ -gons must serve as an anchor for a different  $Y$ -type rank 2  $N$ -gon. Solving this recurrence gives the general term formula

$$\begin{pmatrix} x_k \\ y_k \end{pmatrix} = 3 \begin{pmatrix} \frac{1}{\sqrt{N-6}}(\Phi_1^{2k-3} + \Phi_2^{2k-3}) \\ \Phi_1^{2k-2} + \Phi_2^{2k-2} \end{pmatrix},$$

where

$$\Phi_1 = \frac{\sqrt{N-2} + \sqrt{N-6}}{2} \quad \text{and} \quad \Phi_2 = \frac{-\sqrt{N-2} + \sqrt{N-6}}{2}.$$

From here the formula for  $q_k = x_k + y_k$  follows immediately.

To count the triangles of rank  $k$ , we observe that the triangles of rank  $k$  are the next layer after the  $N$ -gons of rank  $k$ , and each  $X$ -type  $N$ -gon is replaced by  $N-4$  triangles and each  $Y$ -type is replaced by  $N-3$  triangles. Hence the formula for  $l_k = (N-4)x_k + (N-3)y_k$  can be computed.

In order to count how many rank  $k$   $N$ -gons  $T$  jumps, i.e.,  $p_k$ , we need to define  $s_k$  as the number of rank  $k$   $N$ -gons in a small cone, as can be seen in Figure 4. A small cone is opposite one of the triangle's vertices and doesn't contain any side of the triangle. In the same way, a big cone (see Figure 4) is opposite one of the sides of a triangle and contains the table. The number of rank  $k$   $N$ -gons in a big cone is just  $q_k/3 - s_k$  because of the 3-fold symmetry of the tiling.

As above, we need to introduce  $x_k^s$  and  $y_k^s$ , the number of  $X$ -type and  $Y$ -type rank  $k$   $N$ -gons in a small cone. With this,  $s_k = x_k^s + y_k^s$ . The billiard map  $T$  makes any tile jump over two small cones and one big cone so in total it will jump

$$p_k = 2s_k + \left(\frac{q_k}{3} - s_k\right) = \frac{q_k}{3} + s_k.$$

By studying the structure of the small cone, we observe the crochet pattern once again. One notices that the cone that starts at the last  $X$ -type  $N$ -gon of the rank  $k$  ( $k \geq 2$ ) layer looks exactly the same as the initial small cone. That is why  $s_k$  is

equal to the total number of  $N$ -gons obtained by starting with an  $X$ -type  $N$ -gon and using the replacement rules in Claim 7. We express this as a sum,

$$\begin{pmatrix} x_k^s \\ y_k^s \end{pmatrix} = \sum_{i=0}^{k-2} \begin{pmatrix} 1 & 1 \\ N-6 & N-5 \end{pmatrix}^i \begin{pmatrix} 1 \\ 0 \end{pmatrix},$$

which, after some computation, becomes

$$\begin{pmatrix} x_k^s \\ y_k^s \end{pmatrix} = \begin{pmatrix} 1 + \frac{\Phi_1^{2k-4} - \Phi_2^{2k-4}}{\sqrt{(N-6)(N-2)}} \\ -1 + \frac{\Phi_1^{2k-3} - \Phi_2^{2k-3}}{\sqrt{N-2}} \end{pmatrix}.$$

The formula for  $p_k = q_k/3 + x_k^s + y_k^s$  follows immediately, and  $j_k$  is computed in the same manner as  $l_k$  was computed. As we have already said, every  $X$  type  $N$ -gon is replaced by  $N-4$  triangles and every  $Y$  type  $N$ -gon is replaced by  $N-3$  triangles on the next level, and this procedure leaves uncounted only one rank  $k$  triangle in the small cone, so  $j_k = (N-4)x_k^s + (N-3)y_k^s + 1$ .  $\square$

**Claim 9.** *The rotation number  $\rho(t)$  equals*

$$\frac{1}{3} + \frac{1}{3(1 + \Phi_1^2)} = \frac{1}{3} + \frac{1}{3\sqrt{N-2}\Phi_1}.$$

*Proof.* The  $k$ -th layer of  $N$ -gons gives a discrete approximation of the circle map at infinity and so  $p_k/q_k$  is an approximation of  $\rho(t)$  as  $k$  goes to  $\infty$ . By taking the limit we obtained the desired formula for the rotation number  $\rho(t)$ .  $\square$

This last claim completes the proof of all the statements in Theorem 2.

**Remark 10.** (1) One might expect the formulas in Theorem 2 to also work for  $N = 6$ , i.e., a  $(3, 6)$ -tiling of the Euclidean plane. That is not the case even though the crochet pattern works exactly the same also in the  $(3, 6)$ -tiling. The difference that appears when computing the formulas in the  $(3, 6)$ -tiling is that the matrix of the difference system is not diagonalizable and so its powers look completely different.

- (2) Note that the determinant of all the matrices given by the crochet pattern is 1. We believe this is true because the crochet pattern replacement can also be reversed, i.e., starting with the rank  $k$  layer, we can construct the rank  $k-1$  layer.
- (3) According to Theorem 2, one can express the eigenvalues  $\Phi_1$  and  $\Phi_2 = 1/\Phi_1$  via the rotation number  $\rho(t)$ . Therefore this rotation number determines the numbers  $q_k, l_k, p_k, j_k$ , and hence the whole dynamics of the map  $T$ .

**3.2.2. General  $(M, N)$ -tilings.** Next we consider the case of a general  $(M, N)$ -tiling. The theorem and subsequent proof are analogous to those in the  $(3, N)$  case in the previous subsection, but we must consider the cases separately due to a difference in the counting method. In the previous section,  $N$ -gons were classified into types  $X$  and  $Y$ , having two parents and one parent, respectively. However, due to the difference in geometry of triangles versus generic  $M$ -gons, the tilings in the  $M \geq 4$  case never produce  $N$ -gons with two parents. In this case,  $N$ -gons either have one parent or no parent, which we denote as types  $Y$  and  $Z$ . This alternate counting method will be explained in detail in the proof, but first we state the theorem:

**Theorem 11.** *For an  $(M, N)$ -tiling with  $M, N \geq 4$  and*

$$\frac{1}{M} + \frac{1}{N} < \frac{1}{2},$$

*the outer billiard map  $T$  preserves the rank of a tile and every orbit of  $T$  is periodic. The set of rank  $k$   $N$ -gons consists of*

$$q_k = \frac{M}{\sqrt{b^2 - 4}} ((b + 1)(\alpha_1^{2k-2} - \alpha_2^{2k-2}) - (\alpha_1^{2k-4} - \alpha_2^{2k-4}))$$

*elements, and the set of rank  $k$   $M$ -gons consists of*

$$l_k = \frac{M(N - 2)}{\sqrt{b^2 - 4}} (b(\alpha_1^{2k-2} - \alpha_2^{2k-2}) - (\alpha_1^{2k-4} - \alpha_2^{2k-4}))$$

*elements, where  $b = (M - 2)(N - 2) - 2$  and*

$$\alpha_{1,2} = \frac{\sqrt{b - 2} \pm \sqrt{b + 2}}{2}$$

*are the two roots of the equation  $\alpha^2 - \sqrt{b - 2}\alpha - 1 = 0$ . The action of  $T$  on the set of rank  $k$   $N$ -gons is a cyclic permutation  $i \mapsto i + p_k$ , where*

$$p_k = \frac{q_k}{M} + \frac{M - 2}{(b - 2)\sqrt{b + 2}} ((b - 1)(\alpha_1^{2k-3} - \alpha_2^{2k-3}) - (\alpha_1^{2k-5} - \alpha_2^{2k-5})),$$


*and the action of  $T$  on the set of rank  $k$   $M$ -gons is also a cyclic permutation  $i \mapsto i + j_k$ , where*

$$j_k = \frac{l_k}{M} + \frac{1}{(b - 2)\sqrt{b + 2}} ((b^2 - 2)(\alpha_1^{2k-3} - \alpha_2^{2k-3}) - b(\alpha_1^{2k-5} - \alpha_2^{2k-5})).$$

*The rotation number of the outer billiard map at infinity is given by the formula*

$$\rho(t) = \frac{1}{M} + \frac{M - 2}{M\sqrt{b - 2}\alpha_1} \frac{(b - 1)\alpha_1^2 - 1}{(b + 1)\alpha_1^2 - 1}.$$

**Remark 12.** If  $N = M$ , the statement of Theorem 11 reduces to that of Theorem 1.



**Figure 5.** A  $(4, 5)$ -tiling, with rank 1 and rank 2 pentagons labeled either as type Y (one parent) or as type Z (no parents).

The proof of Theorem 11 also consists of several steps.

**Claim 13.** *Every orbit of  $T$  is periodic.*

*Proof.* The proof of this claim is analogous to the proof in the previous section. Because the rank of each tile is preserved under the billiard map, and because there are finitely many tiles of a given rank, every tile must map back to itself after some finite number of iterations  $m$ . When the tile maps back to itself, it has rotated by  $2\pi j/M$  if it is an  $M$ -gon or by  $2\pi j/N$  if it is an  $N$ -gon. Then  $T^{omM}$  is the identity if the tile is an  $M$ -gon and  $T^{omN}$  is the identity if the tile is an  $N$ -gon.  $\square$

**Claim 14.** *For every  $k \geq 1$ ,  $T$  permutes the rank  $k$  tiles cyclically.*

*Proof.* Proof is similar to that for Claim 5.  $\square$


Recall that type Y tiles have one parent and type Z tiles have zero parents (see Figure 5). We now give a crochet pattern for general  $(M, N)$ -tilings,  $M \geq 4$ .

**Claim 15.** *The following replacement rules hold for  $(M, N)$ -tilings:*

$$Y \rightarrow (YZ^{M-3})^{N-4}YZ^{M-4}, \quad (1)$$

$$Z \rightarrow (YZ^{M-3})^{N-3}YZ^{M-4}. \quad (2)$$

*Proof.* In a similar manner to the  $(3, N)$  case, we represent type Y and Z tiles as degenerate polygons, with additional vertices. See Figure 6 for illustrations of the  $(4, 5)$  case. Type Y tiles are represented as quadrilaterals with  $N$  vertices, and type Z tiles are represented as triangles with  $N$  vertices. Because a Y tile has  $N-3$  sides available to connect with a tile of higher rank, a rank  $k$  Y tile produces  $N-3$  Y tiles of rank  $k+1$ . Then, since tiles must meet  $M$  to a vertex, there must be  $M-3$  Z tiles between every pair of Y tiles, and there must be  $M-4$  type Z



**Figure 6.** Tiling of pentagons meeting in fours. Can be extended to a  $(4, 5)$ -tiling.


tiles following the last  $Y$ . Similarly, a  $Z$  tile has  $N - 2$  edges free to connect to a tile of higher rank, so a rank  $k$   $Z$  tile produces  $N - 2$   $Y$  tiles of rank  $k + 1$ , again with  $Z$  tiles appropriately interspersed.

This crochet pattern tiles the hyperbolic plane with  $M$   $N$ -gons meeting at every vertex. From this tiling, we obtain the  $(M, N)$ -tiling by considering the points in the tiling becoming  $M$ -gons, as in Figure 7 (compare with [Conway et al. 2008]). The described crochet pattern translates to the replacement rules given above.  $\square$

We can now compute the formulas for the number of  $M$ - and  $N$ -gons of any rank, as well as for the cyclic permutation of  $M$ - and  $N$ -gons of any rank.

**Claim 16.** *The formulas for  $q_k, p_k, j_k, l_k$  hold as stated in Theorem 11.*

*Proof.* Denoting the number of  $Y$ -type and  $Z$ -type  $N$ -gons of rank  $k$  by  $y_k$  and  $z_k$ ,



**Figure 7.** Left: Tiling of the plane by hexagons meeting in fives. Right: by replacing the vertices in the previous picture with pentagons, we achieve a  $(5, 6)$ -tiling. Here two hexagons and two pentagons meet at each single vertex.

we obtain the recursion formula

$$\begin{pmatrix} y_k \\ z_k \end{pmatrix} = A \begin{pmatrix} y_{k-1} \\ z_{k-1} \end{pmatrix}, \quad (3)$$

where the matrix  $A$  is obtained from the rules given in (1) and (2), and

$$A = \begin{pmatrix} N-3 & N-2 \\ (M-3)(N-3)-1 & (M-3)(N-2)-1 \end{pmatrix}. \quad (4)$$

As mentioned above, the initial conditions are  $\begin{pmatrix} y_1 \\ z_1 \end{pmatrix} = \begin{pmatrix} 0 \\ M \end{pmatrix}$ .

Solving the recurrence, we find the general formula

$$\begin{pmatrix} y_k \\ z_k \end{pmatrix} = \begin{pmatrix} \frac{M(N-2)(\alpha_1^{2k-2} - \alpha_2^{2k-2})}{\sqrt{b^2-4}} \\ \frac{M((M-3)(N-2)-1)(\alpha_1^{2k-2} - \alpha_2^{2k-2}) + M(\alpha_2^{2k-4} - \alpha_1^{2k-4})}{\sqrt{b^2-4}} \end{pmatrix},$$

where

$$b = (M-2)(N-2)-2, \quad \alpha_1 = \frac{\sqrt{b-2} + \sqrt{b+2}}{2}, \quad \alpha_2 = \frac{\sqrt{b-2} - \sqrt{b+2}}{2}.$$

Then  $q_k = y_k + z_k$ , so

$$q_k = \frac{M}{\sqrt{b^2-4}} ((b+1)(\alpha_1^{2k-2} - \alpha_2^{2k-2}) + \alpha_2^{2k-4} - \alpha_1^{2k-4}).$$

Now that we have counted the  $N$ -gons, we count the  $M$ -gons of rank  $k$  by noticing a pattern in the tiling. We see that a type  $Y$   $N$ -gon of rank  $k$  produces  $N-3$   $M$ -gons of rank  $k$ , and a type  $Z$   $N$ -gon produces  $N-2$   $M$ -gons. Thus the number of  $M$ -gons of rank  $k$  is given by  $l_k = (N-3)y_k + (N-2)z_k$ . The formula for  $l_k$  given in Theorem 11 follows.

Next we determine  $p_k$  by counting how many tiles a rank  $k$   $N$ -gon jumps when  $T$  is applied. As in the previous section, we define  $s_k$  as the number of rank  $k$   $N$ -gons in a small cone. We call  $y_k^s$  and  $z_k^s$  the number of rank  $k$   $Y$ s and  $Z$ s in the small cone. Also, as before, applying  $T$  to any tile causes the tile to jump over two small cones and one big cone. In total, the jump is given by  $p_k = s_k + q_k/M$ .

We observe that

$$\begin{pmatrix} y_k^s \\ z_k^s \end{pmatrix} = \sum_{i=0}^{k-2} A^i \begin{pmatrix} 2 \\ M-4 \end{pmatrix}, \quad (5)$$

where  $A$  is given in (4).

This becomes

$$\begin{pmatrix} y_k^s \\ z_k^s \end{pmatrix} = \begin{pmatrix} \frac{1-\alpha_1^{2k-2}}{1-\alpha_1^2} + \frac{2}{\sqrt{b^2-4}} \frac{1-\alpha_2^{2k-2}}{1-\alpha_2^2} (\alpha_1^2 - N + 3) \\ \frac{1}{\sqrt{b^2-4}} \left( \frac{1-\alpha_1^{2k-2}}{1-\alpha_1^2} (B - \alpha_2^2(M-4)) + \frac{1-\alpha_2^{2k-2}}{1-\alpha_2^2} (-B + \alpha_1^2(M-4)) \right) \end{pmatrix},$$

where  $B = (M-3)(b-2) + (M-4)$ . Then, since  $s_k = y_k^s + z_k^s$ , we have

$$s_k = \frac{M-2}{(b-2)\sqrt{b+2}} ((b-1)(\alpha_1^{2k-3} + \alpha_2^{2k-3}) + \alpha_2^{2k-5} - \alpha_1^{2k-5}).$$

This allows us to calculate  $p_k$ , and we can compute  $j_k$  by noticing again that every  $Y$ -type  $N$ -gon will be replaced by  $N-3$   $M$ -gons and every  $Z$ -type  $(N-2)$ -gon will be replaced by  $N-3$   $M$ -gons on the next level. This procedure will leave again only one  $M$ -gon out, so  $j_k = (N-3)y_k^s + (N-2)z_k^s + 1$ .  $\square$

**Claim 17.** *The rotation number is given by*

$$\rho(t) = \frac{1}{M} + \frac{M-2}{M\sqrt{b-2}\alpha_1} \frac{(b-1)\alpha_1^2 - 1}{(b+1)\alpha_1^2 - 1}.$$

*Proof.* This results from taking the limit of  $p_k/q_k$  as  $k \rightarrow \infty$ .  $\square$

#### 4. Remarks and acknowledgments

The methods used in this paper both for geometrical and counting arguments can be used also for all other tilings with 2-fold symmetries in the vertices and so we believe that similar theorems and observations can be deduced in a more general setting.

We want to acknowledge the work of two of our colleagues during the program. Stephanie Ger classified the tilings of the hyperbolic plane in terms of symmetry groups, and Ananya Uppal provided us with a Mathematica demonstration that created images that helped to understand the patterns in the tilings. Their contributions were extremely valuable for our work.

We also want to thank our advisors during the Summer@ICERM program, Chaim Goodman-Strauss, Sergei Tabachnikov, Ryan Greene and Tarik Aougab, for all the help and the inspiring discussions. Finally, we thank ICERM for providing the opportunity for us to participate in this great program.

The tiling pictures were created using *KaleidoTile* application created by Jeff Weeks ([geometrygames.org/contact.html](http://geometrygames.org/contact.html)).

#### References

- [Conway et al. 2008] J. H. Conway, H. Burgiel, and C. Goodman-Strauss, *The symmetries of things*, A K Peters, Wellesley, MA, 2008. MR 2009c:00002 Zbl 1173.00001



- [Dogru and Tabachnikov 2003] F. Dogru and S. Tabachnikov, “On polygonal dual billiard in the hyperbolic plane”, *Regul. Chaotic Dyn.* **8**:1 (2003), 67–81. MR 2004b:37064 Zbl 1075.37524
- [Greenberg 1980] M. J. Greenberg, *Euclidean and non-Euclidean geometries: Development and history*, 2nd ed., W. H. Freeman, San Francisco, CA, 1980. MR 81f:51001 Zbl 0418.51001
- [Gutkin and Simányi 1992] E. Gutkin and N. Simányi, “Dual polygonal billiards and necklace dynamics”, *Comm. Math. Phys.* **143**:3 (1992), 431–449. MR 92k:58139 Zbl 0749.58032
- [Kołodziej 1989] R. Kołodziej, “The antibilliard outside a polygon”, *Bull. Polish Acad. Sci. Math.* **37**:1-6 (1989), 163–168. MR 92g:52002 Zbl 0762.58021
- [Moser 1973] J. Moser, *Stable and random motions in dynamical systems: With special emphasis on celestial mechanics*, Annals of Mathematics Studies **77**, Princeton University Press, 1973. MR 56 #1355 Zbl 0271.70009
- [Moser 1978] J. Moser, “Is the solar system stable?”, *Math. Intelligencer* **1**:2 (1978), 65–71. MR 58 #14029 Zbl 1288.70008
- [Neumann 1959] B. Neumann, “Sharing ham and eggs”, *Iota* (1959), 14–18.
- [Schwartz 2007] R. E. Schwartz, “Unbounded orbits for outer billiards, I”, *J. Mod. Dyn.* **1**:3 (2007), 371–424. MR 2008f:37082 Zbl 1130.37021
- [Schwartz 2009] R. E. Schwartz, *Outer billiards on kites*, Annals of Mathematics Studies **171**, Princeton University Press, 2009. MR 2011a:37081 Zbl 1205.37001
- [Tabachnikov 2005] S. Tabachnikov, *Geometry and billiards*, Student Mathematical Library **30**, Amer. Math. Soc., Providence, RI, 2005. MR 2006h:51001 Zbl 1119.37001
- [Tabachnikov 2007] S. Tabachnikov, “A proof of Culter’s theorem on the existence of periodic orbits in polygonal outer billiards”, *Geom. Dedicata* **129** (2007), 83–87. MR 2008m:37062 Zbl 1131.37039
- [Tabachnikov and Dogru 2005] S. Tabachnikov and F. Dogru, “Dual billiards”, *Math. Intelligencer* **27**:4 (2005), 18–25. MR 2006i:37121 Zbl 1088.37014
- [Vivaldi and Shaidenko 1987] F. Vivaldi and A. V. Shaidenko, “Global stability of a class of discontinuous dual billiards”, *Comm. Math. Phys.* **110**:4 (1987), 625–640. MR 89c:58067 Zbl 0653.58018

Received: 2014-05-28      Accepted: 2014-06-20

dogruf@gvsu.edu

*Department of Mathematics, Grand Valley State University,  
Allendale, MI 49401, United States*

efischer@hmc.edu

*Harvey Mudd College, Claremont, CA 91711, United States*

mihaim92@gmail.com

*Department of Mathematics, Jacobs University Bremen,  
D-28759 Bremen, Germany*



# involve

msp.org/involve

## MANAGING EDITOR

Kenneth S. Berenhaut, Wake Forest University, USA, [berenhks@wfu.edu](mailto:berenhks@wfu.edu)

## BOARD OF EDITORS

Colin Adams	Williams College, USA <a href="mailto:colin.c.adams@williams.edu">colin.c.adams@williams.edu</a>	David Larson	Texas A&M University, USA <a href="mailto:larson@math.tamu.edu">larson@math.tamu.edu</a>
John V. Baxley	Wake Forest University, NC, USA <a href="mailto:baxley@wfu.edu">baxley@wfu.edu</a>	Suzanne Lenhart	University of Tennessee, USA <a href="mailto:lenhart@math.utk.edu">lenhart@math.utk.edu</a>
Arthur T. Benjamin	Harvey Mudd College, USA <a href="mailto:benjamin@hmc.edu">benjamin@hmc.edu</a>	Chi-Kwong Li	College of William and Mary, USA <a href="mailto:ckli@math.wm.edu">ckli@math.wm.edu</a>
Martin Bohner	Missouri U of Science and Technology, USA <a href="mailto:bohner@mst.edu">bohner@mst.edu</a>	Robert B. Lund	Clemson University, USA <a href="mailto:lund@clemson.edu">lund@clemson.edu</a>
Nigel Boston	University of Wisconsin, USA <a href="mailto:boston@math.wisc.edu">boston@math.wisc.edu</a>	Gaven J. Martin	Massey University, New Zealand <a href="mailto:g.j.martin@massey.ac.nz">g.j.martin@massey.ac.nz</a>
Amarjit S. Budhiraja	U of North Carolina, Chapel Hill, USA <a href="mailto:budhiraj@email.unc.edu">budhiraj@email.unc.edu</a>	Mary Meyer	Colorado State University, USA <a href="mailto:meyer@stat.colostate.edu">meyer@stat.colostate.edu</a>
Pietro Cerone	La Trobe University, Australia <a href="mailto:P.Cerone@latrobe.edu.au">P.Cerone@latrobe.edu.au</a>	Emil Minchev	Ruse, Bulgaria <a href="mailto:eminchev@hotmail.com">eminchev@hotmail.com</a>
Scott Chapman	Sam Houston State University, USA <a href="mailto:scott.chapman@shsu.edu">scott.chapman@shsu.edu</a>	Frank Morgan	Williams College, USA <a href="mailto:frank.morgan@williams.edu">frank.morgan@williams.edu</a>
Joshua N. Cooper	University of South Carolina, USA <a href="mailto:cooper@math.sc.edu">cooper@math.sc.edu</a>	Mohammad Sal Moelehian	Ferdowsi University of Mashhad, Iran <a href="mailto:moelehian@ferdowsi.um.ac.ir">moelehian@ferdowsi.um.ac.ir</a>
Jem N. Corcoran	University of Colorado, USA <a href="mailto:corcoran@colorado.edu">corcoran@colorado.edu</a>	Zuhair Nashed	University of Central Florida, USA <a href="mailto:znashed@mail.ucf.edu">znashed@mail.ucf.edu</a>
Toka Diagana	Howard University, USA <a href="mailto:tdiagana@howard.edu">tdiagana@howard.edu</a>	Ken Ono	Emory University, USA <a href="mailto:ono@mathcs.emory.edu">ono@mathcs.emory.edu</a>
Michael Dorff	Brigham Young University, USA <a href="mailto:mdorff@math.byu.edu">mdorff@math.byu.edu</a>	Timothy E. O'Brien	Loyola University Chicago, USA <a href="mailto:tobrie1@luc.edu">tobrie1@luc.edu</a>
Sever S. Dragomir	Victoria University, Australia <a href="mailto:sever@matilda.vu.edu.au">sever@matilda.vu.edu.au</a>	Joseph O'Rourke	Smith College, USA <a href="mailto:orourke@cs.smith.edu">orourke@cs.smith.edu</a>
Behrouz Emamizadeh	The Petroleum Institute, UAE <a href="mailto:bemamizadeh@pi.ac.ae">bemamizadeh@pi.ac.ae</a>	Yuval Peres	Microsoft Research, USA <a href="mailto:peres@microsoft.com">peres@microsoft.com</a>
Joel Foisy	SUNY Potsdam <a href="mailto:foisys@potsdam.edu">foisys@potsdam.edu</a>	Y.-F. S. Pétermann	Université de Genève, Switzerland <a href="mailto:petermann@math.unige.ch">petermann@math.unige.ch</a>
Errin W. Fulp	Wake Forest University, USA <a href="mailto:fulp@wfu.edu">fulp@wfu.edu</a>	Robert J. Plemmons	Wake Forest University, USA <a href="mailto:plemmons@wfu.edu">plemmons@wfu.edu</a>
Joseph Gallian	University of Minnesota Duluth, USA <a href="mailto:jgallian@d.umn.edu">jgallian@d.umn.edu</a>	Carl B. Pomerance	Dartmouth College, USA <a href="mailto:carl.pomerance@dartmouth.edu">carl.pomerance@dartmouth.edu</a>
Stephan R. Garcia	Pomona College, USA <a href="mailto:stephan.garcia@pomona.edu">stephan.garcia@pomona.edu</a>	Vadim Ponomarenko	San Diego State University, USA <a href="mailto:vadim@sciences.sdsu.edu">vadim@sciences.sdsu.edu</a>
Anant Godbole	East Tennessee State University, USA <a href="mailto:godbole@etsu.edu">godbole@etsu.edu</a>	Bjorn Poonen	UC Berkeley, USA <a href="mailto:poonen@math.berkeley.edu">poonen@math.berkeley.edu</a>
Ron Gould	Emory University, USA <a href="mailto:rg@mathcs.emory.edu">rg@mathcs.emory.edu</a>	James Propp	U Mass Lowell, USA <a href="mailto:jpropp@cs.uml.edu">jpropp@cs.uml.edu</a>
Andrew Granville	Université Montréal, Canada <a href="mailto:andrew@dms.umontreal.ca">andrew@dms.umontreal.ca</a>	József H. Przytycki	George Washington University, USA <a href="mailto:przytyck@gwu.edu">przytyck@gwu.edu</a>
Jerrold Griggs	University of South Carolina, USA <a href="mailto:griggs@math.sc.edu">griggs@math.sc.edu</a>	Richard Rebarber	University of Nebraska, USA <a href="mailto:rrebarbe@math.unl.edu">rrebarbe@math.unl.edu</a>
Sat Gupta	U of North Carolina, Greensboro, USA <a href="mailto:sgupta@uncg.edu">sgupta@uncg.edu</a>	Robert W. Robinson	University of Georgia, USA <a href="mailto:rwr@cs.uga.edu">rwr@cs.uga.edu</a>
Jim Haglund	University of Pennsylvania, USA <a href="mailto:jhaglund@math.upenn.edu">jhaglund@math.upenn.edu</a>	Filip Saidak	U of North Carolina, Greensboro, USA <a href="mailto:f_saidak@uncg.edu">f_saidak@uncg.edu</a>
Johnny Henderson	Baylor University, USA <a href="mailto:johnny_henderson@baylor.edu">johnny_henderson@baylor.edu</a>	James A. Sellers	Penn State University, USA <a href="mailto:sellersj@math.psu.edu">sellersj@math.psu.edu</a>
Jim Hoste	Pitzer College <a href="mailto:jhoste@pitzer.edu">jhoste@pitzer.edu</a>	Andrew J. Sterge	Honorary Editor <a href="mailto:andy@ajsterge.com">andy@ajsterge.com</a>
Natalia Hritonenko	Prairie View A&M University, USA <a href="mailto:nahritonenko@pvamu.edu">nahritonenko@pvamu.edu</a>	Ann Trenk	Wellesley College, USA <a href="mailto:atrenk@wellesley.edu">atrenk@wellesley.edu</a>
Glenn H. Hurlbert	Arizona State University, USA <a href="mailto:hurlbert@asu.edu">hurlbert@asu.edu</a>	Ravi Vakil	Stanford University, USA <a href="mailto:vakil@math.stanford.edu">vakil@math.stanford.edu</a>
Charles R. Johnson	College of William and Mary, USA <a href="mailto:crjohnso@math.wm.edu">crjohnso@math.wm.edu</a>	Antonia Vecchio	Consiglio Nazionale delle Ricerche, Italy <a href="mailto:antonia.vecchio@cnr.it">antonia.vecchio@cnr.it</a>
K. B. Kulasekera	Clemson University, USA <a href="mailto:kk@ces.clemson.edu">kk@ces.clemson.edu</a>	Ram U. Verma	University of Toledo, USA <a href="mailto:verma99@msn.com">verma99@msn.com</a>
Gerry Ladas	University of Rhode Island, USA <a href="mailto:gladas@math.uri.edu">gladas@math.uri.edu</a>	John C. Wierman	Johns Hopkins University, USA <a href="mailto:wierman@jhu.edu">wierman@jhu.edu</a>
		Michael E. Zieve	University of Michigan, USA <a href="mailto:zieve@umich.edu">zieve@umich.edu</a>

## PRODUCTION

Silvio Levy, Scientific Editor


Cover: Alex Scorpan

See inside back cover or [msp.org/involve](http://msp.org/involve) for submission instructions. The subscription price for 2015 is US \$140/year for the electronic version, and \$190/year (+\$35, if shipping outside the US) for print and electronic. Subscriptions, requests for back issues from the last three years and changes of subscribers address should be sent to MSP.

Involve (ISSN 1944-4184 electronic, 1944-4176 printed) at Mathematical Sciences Publishers, 798 Evans Hall #3840, c/o University of California, Berkeley, CA 94720-3840, is published continuously online. Periodical rate postage paid at Berkeley, CA 94704, and additional mailing offices.

Involve peer review and production are managed by EditFlow® from Mathematical Sciences Publishers.

## PUBLISHED BY

 **mathematical sciences publishers**  
nonprofit scientific publishing

<http://msp.org/>

© 2015 Mathematical Sciences Publishers

# involve

2015

vol. 8

no. 4

The $\Delta^2$ conjecture holds for graphs of small order	541
COLE FRANKS	
Linear symplectomorphisms as $R$ -Lagrangian subspaces	551
CHRIS HELLMANN, BRENNAN LANGENBACH AND MICHAEL VANVALKENBURGH	
Maximization of the size of monic orthogonal polynomials on the unit circle corresponding to the measures in the Steklov class	571
JOHN HOFFMAN, MCKINLEY MEYER, MARIYA SARDARLI AND ALEX SHERMAN	
A type of multiple integral with log-gamma function	593
DUOKUI YAN, RONGCHANG LIU AND GENG-ZHE CHANG	
Knight's tours on boards with odd dimensions	615
BAOYUE BI, STEVE BUTLER, STEPHANIE DEGRAAF AND ELIZABETH DOEBEL	
Differentiation with respect to parameters of solutions of nonlocal boundary value problems for difference equations	629
JOHNNY HENDERSON AND XUEWEI JIANG	
Outer billiards and tilings of the hyperbolic plane	637
FILIZ DOGRU, EMILY M. FISCHER AND CRISTIAN MIHAI MUNTEANU	
Sophie Germain primes and involutions of $\mathbb{Z}_n^\times$	653
KARENNA GENZLINGER AND KEIR LOCKRIDGE	
On symplectic capacities of toric domains	665
MICHAEL LANDRY, MATTHEW MCMILLAN AND EMMANUEL TSUKERMAN	
When the catenary degree agrees with the tame degree in numerical semigroups of embedding dimension three	677
PEDRO A. GARCÍA-SÁNCHEZ AND CATERINA VIOLA	
Cylindrical liquid bridges	695
LAMONT COLTER AND RAY TREINEN	
Some projective distance inequalities for simplices in complex projective space	707
MARK FINCHER, HEATHER OLNEY AND WILLIAM CHERRY	



1944-4176(2015)8:4;1-3