

involve

a journal of mathematics

Cocircular relative equilibria of four vortices

Jonathan Gomez, Alexander Gutierrez,
John Little, Roberto Pelayo and Jesse Robert



Cocircular relative equilibria of four vortices

Jonathan Gomez, Alexander Gutierrez,
John Little, Roberto Pelayo and Jesse Robert

(Communicated by Martin Bohner)

We study the cocircular relative equilibria (planar central configurations) in the four-vortex problem using methods suggested by the study of cocircular central configurations in the Newtonian four-body problem in recent work of Cors and Roberts. Using mutual distance coordinates, we show that the set of four-vortex relative equilibria is a two-dimensional surface with boundary curves representing kite configurations, isosceles trapezoids, and degenerate configurations with one zero vorticity. We also show that there is a constraint on the signs of the vorticities in these configurations; either three or four of the vorticities must have the same sign, in contrast to the noncocircular cases studied by Hampton, Roberts, and Santoprete.

1. Introduction

Understanding central configurations is a problem of fundamental importance in celestial mechanics (for instance, see [Saari 2011]). Recent years have seen heightened interest in the study of central configurations, in part due to the fact that advances in computing power have made it possible to utilize tools from algebraic geometry to study such problems. These tools have led to breakthroughs such as the proof that there are only finitely many central configurations for each collection of positive masses in the four-body problem [Hampton and Moeckel 2006], and the proof of finiteness in generic cases of the five-body problem [Hampton and Jensen 2011; Albouy and Kaloshin 2012].

Similarly useful is the study of relative equilibrium configurations of collections of *Helmholtz vortices* [Hampton and Moeckel 2009; Saari 2011]. Helmholtz vortices, thought of as whirlpools lying in an infinite plane composed of a perfect fluid, were first introduced as a means of modeling the interactions of two-dimensional slices of collections of columnar vortex filaments. The study of relative equilibria of

MSC2010: primary 76B99; secondary 70F10, 13P10.

Keywords: relative equilibria, vortices, central configurations.

The PURE Math 2012 program was made possible through National Science Foundation grants DMS-1045082 and DMS-1045147 and a grant from the National Security Agency.

vortices has applications that range from basic fluid mechanics to the study of how cyclones and hurricanes interact and evolve over time.

Algebraically, the equations defining relative equilibria of vortices are very similar to those defining relative equilibria of masses. Suppose vortices of strengths Γ_i (unlike the masses in the Newtonian problem, these can have positive or negative real values) are initially located at positions $q_i \in \mathbb{R}^2$. Writing $r_{ij} = \|q_i - q_j\|$ for the mutual distance, we have a relative equilibrium if for all i ,

$$\sum_{j \neq i} \Gamma_j \frac{q_i - q_j}{r_{ij}^2} = -\lambda(q_i - c), \quad (1-1)$$

where λ is a constant and c is the center of rotation. The equations (1-1) differ from their Newtonian equivalents because of the r_{ij}^2 in the denominators (where r_{ij}^3 appears in the equations for relative equilibria of masses). The difference is caused by a logarithmic potential in the vortex case that replaces the gravitational potential in the Newtonian case.

In this paper, we study relative equilibria of collections of four point vortices whose locations lie on a circle in the plane (the cocircular configurations in the title). The inspiration for this study can be found in a recent paper in which Cors and Roberts [2012] study the corresponding problem for four cocircular masses under Newtonian gravity. Other articles devoted to the study of cocircular central configurations include [Hampton 2005; Llibre and Valls 2015]. We also use a number of general results on the vortex problem from a second recent article by Hampton, Roberts, and Santoprete [Hampton et al. 2014]. We first present a set of equations in mutual distance coordinates whose solutions correspond to these configurations in Section 2. By analyzing the set of solutions of these equations, in Section 3 we obtain a surface in \mathbb{R}^3 whose points parametrize the family of cocircular relative equilibria. Next, in Section 4, we prove a result concerning the possible signs of the vorticities for a cocircular relative equilibrium. We discuss some constraints on the positions q_i and the vorticities Γ_i in relative equilibria in Section 5. Finally, we follow [Cors and Roberts 2012], mutatis mutandis, and analyze two symmetric cases (kites and isosceles trapezoids) in Sections 6 and 7. These cases correspond to boundary points of our surface.

2. Equations for relative equilibria in mutual distance coordinates

By using results from [Hampton et al. 2014] on the general four-vortex problem and adapting results from [Cors and Roberts 2012] on the cocircular case of the four-body problem, in this section we will derive a set of equations characterizing the cocircular relative equilibria in the four-vortex problem.

By equation (10) of [Hampton et al. 2014], the following relation (a consequence of the Dziobek relations in the vortex case) is necessary and sufficient for the existence of a four-vortex relative equilibrium with mutual distances $r_{ij} > 0$,

where $1 \leq i < j \leq 4$:

$$(r_{13}^2 - r_{12}^2)(r_{23}^2 - r_{34}^2)(r_{24}^2 - r_{14}^2) - (r_{12}^2 - r_{14}^2)(r_{24}^2 - r_{34}^2)(r_{13}^2 - r_{23}^2) = 0. \quad (2-1)$$

For future reference, we note that this equation can be rearranged algebraically in many different ways. We will also need the forms

$$(r_{14}^2 - r_{24}^2)(r_{13}^2 - r_{34}^2)(r_{12}^2 - r_{23}^2) - (r_{14}^2 - r_{34}^2)(r_{13}^2 - r_{23}^2)(r_{12}^2 - r_{24}^2) = 0, \quad (2-2)$$

$$(r_{23}^2 - r_{24}^2)(r_{14}^2 - r_{34}^2)(r_{12}^2 - r_{13}^2) - (r_{24}^2 - r_{34}^2)(r_{13}^2 - r_{14}^2)(r_{12}^2 - r_{23}^2) = 0, \quad (2-3)$$

$$(r_{24}^2 - r_{23}^2)(r_{13}^2 - r_{34}^2)(r_{12}^2 - r_{14}^2) - (r_{34}^2 - r_{23}^2)(r_{13}^2 - r_{14}^2)(r_{12}^2 - r_{24}^2) = 0. \quad (2-4)$$

Now we impose the condition that the locations of the four vortices lie on a single circle in the plane. Numbering the positions sequentially around that circle, it follows that $r_{12}, r_{23}, r_{34}, r_{14}$ are the lengths of the exterior edges of a cyclic quadrilateral, and r_{13}, r_{24} are the lengths of the diagonals. Letting

$$a = r_{12}r_{34} + r_{14}r_{23}, \quad b = r_{12}r_{14} + r_{23}r_{34}, \quad c = r_{12}r_{23} + r_{14}r_{34}, \quad (2-5)$$

from the law of cosines and the fact that opposite interior angles in the quadrilateral are supplementary, it follows that

$$r_{13}^2 = \frac{ab}{c}, \quad (2-6)$$

$$r_{24}^2 = \frac{ac}{b}. \quad (2-7)$$

Multiplying the two equations above and taking square roots gives Ptolemy's theorem on cyclic quadrilaterals

$$r_{13}r_{24} = r_{12}r_{34} + r_{14}r_{23}. \quad (2-8)$$

As in [Cors and Roberts 2012], we will always fix the numbering of the vortices so that r_{12} is the largest exterior side length, and we will normalize the unit of distance so $r_{12} = 1$. Then

$$r_{23}, r_{34}, r_{14} \leq 1. \quad (2-9)$$

As noted in [Cors and Roberts 2012], we also have

$$\frac{r_{13}}{r_{24}} = \frac{b}{c},$$

so

$$r_{13} - r_{24} \geq 0 \iff b - c \geq 0 \iff (r_{14} - r_{23})(r_{12} - r_{34}) \geq 0.$$

Since $r_{12} \geq r_{34}$ by our choice of labeling,

$$r_{14} \geq r_{23} \iff r_{13} \geq r_{24}. \quad (2-10)$$

We note some additional useful consequences of the equations above relating the diagonals of the cyclic quadrilateral to the exterior sides. In words, these inequalities

will say that *the diagonals of the cyclic quadrilateral are longer than any exterior side on the opposite side of the diagonal from the longest exterior side*. For instance, from (2-6), notice that

$$r_{13}^2 - r_{14}^2 = r_{34} \left(\frac{r_{34}r_{23} + r_{23}^2r_{14} + r_{14} - r_{14}^3}{r_{23} + r_{14}r_{34}} \right) > 0 \quad (2-11)$$

(since $r_{14} - r_{14}^3 \geq 0$ by (2-9)). By similar computations, we also have

$$r_{13}^2 - r_{34}^2 > 0, \quad (2-12)$$

$$r_{24}^2 - r_{23}^2 > 0, \quad (2-13)$$

$$r_{24}^2 - r_{34}^2 > 0. \quad (2-14)$$

Let $\Gamma_i \in \mathbb{R} \setminus \{0\}$, where $i = 1, \dots, 4$, denote the strengths (vorticities) of the four vortices. The derivation of (2-1) above and a computation analogous to that giving equations (16)–(18) in [Cors and Roberts 2012] leads to the vorticity ratio formulas

$$\frac{\Gamma_2}{\Gamma_1} = \frac{r_{23}r_{24}(r_{13}^2 - r_{14}^2)}{r_{13}r_{14}(r_{24}^2 - r_{23}^2)}, \quad (2-15)$$

$$\frac{\Gamma_3}{\Gamma_1} = \frac{r_{23}r_{34}(1 - r_{14}^2)}{r_{14}(r_{23}^2 - r_{34}^2)}, \quad (2-16)$$

$$\frac{\Gamma_4}{\Gamma_1} = \frac{r_{24}r_{34}(r_{13}^2 - 1)}{r_{13}(r_{24}^2 - r_{34}^2)}. \quad (2-17)$$

We can always normalize (choose units for vorticity) to set $\Gamma_1 = 1$. By (2-3) and (2-11)–(2-14), the numerator in the formula for Γ_2 and the denominators in the formulas for Γ_2 and Γ_4 are always nonzero, so the values of Γ_2 and Γ_4 are always determined by these. Equation (2-16) gives a well-defined value for Γ_3 unless $r_{23}^2 - r_{34}^2 = 0$. Looking at (2-4), (2-12), and (2-13), we see that this implies $1 - r_{14}^2 = 0$, so the quotient is actually indeterminate. If, on the other hand, the factor $1 - r_{14}^2$ vanishes, then (2-4) and (2-11) show that $r_{23}^2 - r_{34}^2 = 0$, or $1 - r_{24}^2 = 0$. When $r_{23}^2 - r_{34}^2 = 0$, an alternate formula for Γ_3 can be derived using (2-1):

$$\Gamma_3 = \frac{(r_{13}^2 - 1)(r_{24}^2 - 1)r_{23}^2}{(r_{24}^2 - r_{23}^2)(r_{13}^2 - r_{23}^2)}. \quad (2-18)$$

There are solutions with $r_{14} = r_{12} = r_{24} = 1$ corresponding to degenerate configurations with vortices 1, 2 and 4 forming an equilateral triangle and $\Gamma_3 = 0$. Similarly, there are degenerate configurations with $r_{13} = r_{12} = r_{23} = 1$ and $\Gamma_4 = 0$. The configurations with $r_{14} = r_{12} = 1$ and $r_{23} = r_{34}$ are the symmetric kites to be studied in Section 6.

Collecting all of the results stated above, we see the following statement.

Theorem 2.1. *A cocircular configuration of four vortices with mutual distances r_{ij} , vorticities Γ_i , and with $r_{12} = 1$, $r_{14} < 1$ and $\Gamma_1 = 1$ is a relative equilibrium if and only if the r_{ij} and Γ_i give a common zero of the following set of six polynomial equations:*

$$\begin{aligned}
 F_1 &= r_{13}^2(r_{23} + r_{34}r_{14}) - (r_{34} + r_{14}r_{23})(r_{14} + r_{23}r_{34}), \\
 F_2 &= r_{24}^2(r_{14} + r_{23}r_{34}) - (r_{34} + r_{14}r_{23})(r_{23} + r_{14}r_{34}), \\
 F_3 &= (r_{13}^2 - 1)(r_{23}^2 - r_{34}^2)(r_{24}^2 - r_{14}^2) - (1 - r_{14}^2)(r_{24}^2 - r_{34}^2)(r_{13}^2 - r_{23}^2), \\
 F_4 &= r_{13}r_{14}(r_{24}^2 - r_{23}^2)\Gamma_2 - r_{23}r_{24}(r_{13}^2 - r_{14}^2), \\
 F_5 &= r_{14}(r_{23}^2 - r_{34}^2)\Gamma_3 - r_{23}r_{34}(1 - r_{14}^2), \\
 F_6 &= r_{13}(r_{24}^2 - r_{34}^2)\Gamma_4 - r_{23}r_{24}(r_{13}^2 - 1).
 \end{aligned} \tag{2-19}$$

When $r_{12} = r_{14} = 1$, the equation $F_5 = 0$ is replaced by a similar equation $F'_5 = 0$ derived from (2-18).

3. The surface of cocircular relative equilibria

As suggested by the naive count of variables and equations in the system (2-19), with our normalizations, the set of cocircular relative equilibria is two-dimensional. The equations $F_4 = F_5 = F_6 = 0$ in Theorem 2.1 express the vorticities $\Gamma_2, \Gamma_3, \Gamma_4$ in terms of the r_{ij} . Moreover, we may use the equations $F_1 = 0$ and $F_2 = 0$ to write the squared diagonals r_{13}^2 and r_{24}^2 as functions of the other mutual distances as in (2-6) and (2-7) above. Using these two relations, one can think of F_3 as a function of the three exterior side lengths r_{23}, r_{34}, r_{14} :

$$F_3(r_{23}, r_{34}, r_{14}) = (r_{13}^2 - 1)(r_{23}^2 - r_{34}^2)(r_{24}^2 - r_{14}^2) - (1 - r_{14}^2)(r_{24}^2 - r_{34}^2)(r_{13}^2 - r_{23}^2).$$

Then $F_3 = 0$ defines an algebraic surface in \mathbb{R}^3 with coordinates r_{23}, r_{34}, r_{14} .

By (2-9), we can plot the set of points on which $F_3 = 0$ implicitly in the unit cube. Figure 1 shows the view of the surface looking along the positive r_{14} -axis toward the (r_{23}, r_{34}) -plane. There is a nearly vertical portion of the surface that is obscured from this viewpoint, but visible in the rotated view on the right in Figure 1. However, the entire implicit plot is symmetric across the plane $r_{14} = r_{23}$ (this can be seen by the fact that interchanging r_{14} and r_{23} takes (2-1) to (2-2)).

Therefore we can assume without loss of generality that $r_{14} \geq r_{23}$, and so also $r_{13} \geq r_{24}$ by (2-10). We will only consider that portion of the graph in the following. Because of the shape, we will refer to it as the *bowtie surface*.

We next consider what configurations correspond to points on the boundary curves. Note that if $r_{14} = r_{23}$, then (2-5), (2-6), and (2-7) imply that $r_{13} = r_{24}$ as well, so the only cases where $r_{14} = r_{23}$ are the configurations known as isosceles trapezoids. These will be studied in more detail in Section 7. We next note that since $1 = r_{12} \geq r_{14} \geq r_{23}$, the rest of the boundary is defined by $r_{14} = 1$. Substituting




Figure 1. Two views of the surface $F_3(r_{23}, r_{34}, r_{14}) = 0$.

this into F_3 and factoring yields

$$F_3(r_{23}, r_{34}, 1) = (r_{23} - r_{34})(r_{23} + r_{34})^2(r_{23}^2 + r_{23}r_{34} + r_{34}^2 - 1).$$

The first factor vanishes on points corresponding to kite configurations where $r_{23} = r_{34}$. The kite cases will be completely characterized in Section 6.

The second factor is never zero for positive mutual distances. Hence it is left to consider cases where

$$r_{23}^2 + r_{23}r_{34} + r_{34}^2 - 1 = 0.$$

Examining (2-7), we see that when $r_{12} = r_{14} = 1$, this equation is equivalent to $r_{24}^2 = 1$. Therefore, the vortices 1, 2, and 4 are at the corners of an equilateral triangle, and it follows by (2-18) that $\Gamma_3 = 0$. Thus, the points on this curved component of the boundary shown in Figure 2 correspond to degenerate configurations.




Figure 2. Plot of $r_{23}^2 + r_{23}r_{34} + r_{34}^2 - 1 = 0$ with the graph of $F_3(r_{23}, r_{34}, r_{14}) = 0$.




Figure 3. Views of Γ_3 along r_{14} axis (left) and from the side (right).

4. The signs of the vorticities

In this section, we will analyze the possible signs of the Γ_i in solutions of the system of equations from [Theorem 2.1](#). We will see that, in fact, in any such relative equilibrium either all of the Γ_i have the same sign, or else three of the Γ_i have the same sign and the remaining vorticity has the opposite sign.

We were led to conjecture these patterns by plots showing the values for the vorticity Γ_3 obtained from the equation $F_5 = 0$ in (2-19) on the points of the bowtie surface defined by $F_3 = 0$. To generate the plots in [Figure 3](#), we solved the equation $F_3 = 0$ numerically for r_{14} as a function of r_{23} and r_{34} at a collection of points in the projection of the bowtie onto the (r_{23}, r_{34}) -plane, then plotted positive Γ_3 values in blue and negative Γ_3 values in red. [Figure 3](#) (left) shows a top view along the direction of the r_{14} -axis. [Figure 3](#) (right) shows the same plot of Γ_3 -values, but from one side.

In the remainder of this section, we will give an analytic proof that Γ_3 takes opposite signs on the two lobes of the bowtie surface. We will need the following fact; this depends only on the geometry of the cyclic quadrilateral.

Lemma 4.1 [[Cors and Roberts 2012](#), Lemma 4.6]. *Under the assumption $r_{14} \geq r_{23}$, and the consequence noted above in (2-10), it follows that*

$$\frac{r_{13}}{r_{24}} \leq \frac{r_{14}}{r_{23}}.$$

Proof. For the convenience of the reader, we reproduce the proof from [[Cors and Roberts 2012](#)]. From (2-6) and (2-7), and using the assumptions $r_{12} = 1$ and $r_{23} \leq r_{14}$, we have

$$\frac{r_{13}}{r_{24}} = \frac{b}{c} = \frac{r_{14} + r_{23}r_{34}}{r_{23} + r_{14}r_{34}} \leq \frac{r_{14}(1 + r_{34})}{r_{23}(1 + r_{34})},$$

which implies the claim. □

Lemma 4.2. *In all cocircular four-vortex relative equilibria as above, $\Gamma_2 > 0$.*

Proof. From the equation $F_4 = 0$, we have

$$\Gamma_2 = \frac{r_{24} r_{23} (r_{13}^2 - r_{14}^2)}{r_{13} r_{14} (r_{24}^2 - r_{23}^2)}. \tag{4-1}$$

The inequality $\Gamma_2 > 0$ follows from (2-11) and (2-13). □

The portion of the bowtie surface with $r_{14} \geq r_{23}$ off the boundary curves is composed of two lobes: one (on the left in Figure 1 (left)) on which $r_{23} < r_{34}$, and a second on which $r_{23} > r_{34}$. We will call these open subsets of the bowtie surface lobe I and lobe II, respectively. The closures of the two lobes of the surface intersect only at the point corresponding to a degenerate configuration that is also a kite.

We will deal with the points in the interior of lobe II first, since they follow essentially the same patterns as those found by Cors and Roberts in the cocircular four-body central configurations. We note that in [Cors and Roberts 2012, Section 2.2], the inequality $r_{23} \geq r_{34}$ was deduced from the positivity of the masses m_i . However, this inequality holds by definition on our lobe II.

Theorem 4.3. *On lobe II, we have*

$$\Gamma_2 \geq \Gamma_4 \geq \Gamma_3 > 0.$$

Hence all four of the vorticities have the same sign on lobe II.

Proof. The inequality $\Gamma_2 \geq \Gamma_4$ follows from the equations $F_4 = 0$ and $F_6 = 0$, or from (2-15) and (2-17). These say

$$\Gamma_2 = \frac{r_{23}r_{24}(r_{13}^2 - r_{14}^2)}{r_{13}r_{14}(r_{24}^2 - r_{23}^2)}, \quad \Gamma_4 = \frac{r_{34}r_{24}(r_{13}^2 - 1)}{r_{13}(r_{24}^2 - r_{34}^2)},$$

and the inequalities $r_{23} > r_{34}$, $r_{14} \leq 1$, and $r_{13} \geq r_{14}$ combine to give $\Gamma_2 \geq \Gamma_4$. Finally, $\Gamma_4 \geq \Gamma_3 > 0$ follows using Lemma 4.1 just as in the proof of Theorem 4.4 of [Cors and Roberts 2012]. □

Now we analyze the situation on lobe I:

Theorem 4.4. *On lobe I, we have*

$$\Gamma_4 > \Gamma_2 > 0 > \Gamma_3.$$

Hence three of the vorticities are positive and one is negative on lobe I.

Proof. The inequality $\Gamma_2 > 0$ follows again from Lemma 4.2. On lobe I, $r_{23} < r_{34}$ and the equation $F_5 = 0$ from (2-19) imply that $\Gamma_3 < 0$. Hence to finish the proof, we only need to show that $\Gamma_4 > \Gamma_2$ on this lobe of the bowtie.

We begin with the equations $F_4 = 0$ and $F_6 = 0$ from (2-19). Solving for Γ_2, Γ_4 and multiplying, we have

$$\Gamma_2\Gamma_4 = \frac{r_{23}r_{24}^2r_{34}}{r_{13}^2r_{14}} \frac{(r_{13}^2 - r_{14}^2)}{(r_{24}^2 - r_{23}^2)} \frac{(r_{13}^2 - 1)}{(r_{24}^2 - r_{34}^2)}.$$

We will show first that $\Gamma_2\Gamma_4 > 0$. From (2-3), we also have

$$\frac{r_{13}^2 - r_{14}^2}{r_{24}^2 - r_{23}^2} = \frac{(r_{14}^2 - r_{34}^2)(r_{13}^2 - 1)}{(1 - r_{23}^2)(r_{24}^2 - r_{34}^2)}. \quad (4-2)$$

Substituting into the previous equation, we have

$$\Gamma_2\Gamma_4 = \frac{r_{23}r_{24}^2r_{34}}{r_{13}^2r_{14}} \frac{(r_{14}^2 - r_{34}^2)}{(1 - r_{23}^2)} \left(\frac{r_{13}^2 - 1}{r_{24}^2 - r_{34}^2} \right)^2.$$

Hence the sign of $\Gamma_2\Gamma_4$ is determined by the sign of the factor $r_{14}^2 - r_{34}^2$.

By rearranging (2-2) and (2-3) (with $r_{12} = 1$), we obtain the equations

$$\frac{r_{14}^2 - r_{34}^2}{1 - r_{23}^2} = \frac{(r_{13}^2 - r_{34}^2)(r_{24}^2 - r_{14}^2)}{(r_{13}^2 - r_{23}^2)(r_{24}^2 - 1)} = \frac{(r_{13}^2 - r_{14}^2)(r_{24}^2 - r_{34}^2)}{(r_{13}^2 - 1)(r_{24}^2 - r_{23}^2)}. \quad (4-3)$$

In the rightmost expression in (4-3), all of the factors except $r_{13}^2 - 1$ are known to be positive by (2-11), (2-13), and (2-14). Similarly from (2-12) and $r_{23} < r_{34}$, the factors $r_{13}^2 - r_{34}^2$ and $r_{13}^2 - r_{23}^2$ in the middle product are also positive.

We consider the following possible cases. If $r_{24}^2 - r_{14}^2$ and $r_{24}^2 - 1$ have *the same sign*, then $\Gamma_2\Gamma_4 > 0$ and we are done.

On the other hand, we claim that the case where these factors have opposite signs, that is, $r_{24}^2 - 1 < 0$ but $r_{24}^2 - r_{14}^2 > 0$, is not possible for a four-vortex relative equilibrium (even though these relations are certainly possible for a cyclic quadrilateral). We note that in this remaining potential ‘‘bad’’ case, from (4-3), we have $r_{13}^2 - 1 < 0$, so the edge lengths are ordered as

$$r_{12} = 1 > r_{13} > r_{24} > r_{34} > r_{14} > r_{23}. \quad (4-4)$$

We will show that this is incompatible with the equation $F_3 = 0$, but in the rearranged form given in (2-4).

Denote the factors in that equation as $ABC - abc = 0$. Under the assumptions that the lengths are ordered as in (4-4), we see

$$A = r_{24}^2 - r_{23}^2 > a = r_{34}^2 - r_{23}^2 > 0.$$

We claim that it is also true that $BC > bc > 0$, so the equation $ABC - abc = 0$ cannot hold. First, $BC > 0$ and $bc > 0$ by (4-4). Expand out the products in $BC - bc$, noting one cancellation, to obtain

$$r_{13}^2r_{24}^2 + r_{14}^2 + r_{34}^2r_{14}^2 - r_{13}^2r_{14}^2 - r_{34}^2 - r_{14}^2r_{24}^2.$$

By Ptolemy’s theorem from (2-8), we can substitute for the first term and simplify to obtain

$$BC - bc = r_{14}^2(r_{23}^2 + r_{34}^2 + 1 - r_{13}^2 - r_{24}^2) + 2r_{14}r_{23}r_{34}.$$

By the law of cosines as before, we have

$$r_{23}^2 + r_{34}^2 = r_{24}^2 + 2r_{23}r_{34} \cos \theta_3,$$

where θ_3 is the interior angle of the quadrilateral at vortex 3. Hence

$$BC - bc = r_{14}^2(1 - r_{13}^2) + 2r_{14}r_{23}r_{34}(1 + r_{14} \cos \theta_3) > 0.$$

This shows that this case cannot occur. Hence $\Gamma_2\Gamma_4 > 0$ and in addition, $r_{14}^2 - r_{34}^2 > 0$.

It remains to show that $\Gamma_4 > \Gamma_2$. By (2-15) and (2-17),

$$\frac{\Gamma_4}{\Gamma_2} = \frac{r_{34}r_{14}}{r_{23}} \frac{(r_{13}^2 - 1)(r_{24}^2 - r_{23}^2)}{(r_{13}^2 - r_{14}^2)(r_{24}^2 - r_{34}^2)}.$$

As noted above, from (2-3) (with $r_{12} = 1$), we obtain

$$\frac{(r_{13}^2 - 1)(r_{24}^2 - r_{23}^2)}{(r_{13}^2 - r_{14}^2)(r_{24}^2 - r_{34}^2)} = \frac{1 - r_{23}^2}{r_{14}^2 - r_{34}^2}. \tag{4-5}$$

Hence

$$\frac{\Gamma_4}{\Gamma_2} = \frac{r_{34}r_{14}(1 - r_{23}^2)}{r_{23}(r_{14}^2 - r_{34}^2)}.$$

Note that both the numerator and the denominator are positive by the argument showing $\Gamma_2\Gamma_4 > 0$. We subtract the denominator in the last expression from the numerator and factor to obtain

$$(r_{34} - r_{14}r_{23})(r_{14} + r_{23}r_{34}).$$

The first factor is positive since $r_{34} > r_{23}$ on lobe I and $r_{14} < 1$. The second factor is automatically positive since the r_{ij} are distances. Hence $\Gamma_4 > \Gamma_2$. □

5. Further constraints on the q_i and the Γ_i

We have already seen that, as in the Newtonian case, not every cyclic quadrilateral can appear in a relative equilibrium of four vortices; there are additional geometric constraints imposed by (2-1). The following lemma is inspired by the proof of Conley’s perpendicular bisector theorem for Newtonian central configurations from [Moeckel 1990] and gives another type of constraint. To our knowledge, this sort of argument has not been used before for vortices and this sort of approach could be useful in other situations. However, the fact that the Γ_i can be positive or negative makes it somewhat difficult to foresee the circumstances where something of this sort might be used (other than for cases where it is assumed that all the Γ_i are

positive, for instance). We continue to assume that the positions of the vortices are labeled in sequential order around the circumscribed circle, $r_{12} = 1$ is the longest exterior side of the quadrilateral, $r_{23} \leq r_{14}$, and $\Gamma = 1$.

Lemma 5.1. *Let L be the perpendicular bisector of the chord of the circle connecting q_2 and q_3 . Then q_1 and q_4 lie on opposite sides of L . In particular, the arc from q_1 to q_2 along the circle not containing q_3 and q_4 is less than a semicircle.*

Proof. We begin with the observation that, by Theorems 4.3 and 4.4, $\Gamma_1 = 1$ and $\Gamma_4 > 0$ have the same sign in all of our relative equilibria. From (1-1) with $i = 2, 3$, we have the equations

$$\begin{aligned} \Gamma_1 \frac{q_2 - q_1}{r_{12}^2} + \Gamma_3 \frac{q_2 - q_3}{r_{23}^2} + \Gamma_4 \frac{q_2 - q_4}{r_{24}^2} &= -\lambda(q_2 - c), \\ \Gamma_1 \frac{q_3 - q_1}{r_{13}^2} + \Gamma_2 \frac{q_3 - q_2}{r_{23}^2} + \Gamma_4 \frac{q_3 - q_4}{r_{34}^2} &= -\lambda(q_3 - c). \end{aligned}$$

Subtracting these two equations and rearranging, we see that the vector

$$\Gamma_1 \left(\frac{q_2 - q_1}{r_{12}^2} - \frac{q_3 - q_1}{r_{13}^2} \right) + \Gamma_4 \left(\frac{q_2 - q_4}{r_{24}^2} - \frac{q_3 - q_4}{r_{34}^2} \right) \tag{5-1}$$

is a scalar multiple of $q_2 - q_3$. Let v be a unit vector orthogonal to $q_2 - q_3$. The standard inner (dot) product of v and $q_2 - q_3$ is $\langle v, q_2 - q_3 \rangle = 0$. Hence $\langle v, q_2 - q_1 \rangle = \langle v, q_3 - q_1 \rangle$ and $\langle v, q_2 - q_4 \rangle = \langle v, q_3 - q_4 \rangle$. Call the first of these scalars d_1 and the second d_4 . Then taking the inner product of (5-1) and v , we obtain

$$\Gamma_1 d_1 \left(\frac{1}{r_{12}^2} - \frac{1}{r_{13}^2} \right) + \Gamma_4 d_4 \left(\frac{1}{r_{24}^2} - \frac{1}{r_{34}^2} \right) = 0. \tag{5-2}$$

We claim that this relation can only hold when q_1 and q_4 lie on opposite sides of L . Note that $1/r_{12}^2 - 1/r_{13}^2$ (respectively, $1/r_{24}^2 - 1/r_{34}^2$) is zero only if q_1 (respectively, q_4) lies on the perpendicular bisector L . Moreover the sign is positive if q_1 (respectively, q_4) lies in the half-plane bounded by L and containing q_2 and negative on the half-plane containing q_3 . On the other hand, d_1 and d_4 both have the same sign since q_1 and q_4 lie in the same half-plane bounded by the chord through q_2 and q_3 . Hence the only way the left side of (5-2) can cancel to zero is if q_1 and q_4 lie on opposite sides of L . □

Theorem 5.2. *In all of our relative equilibria, $\Gamma_2 \leq 1$.*

Proof. In a cyclic quadrilateral, it is a standard fact that the angle between an exterior side and a diagonal is equal to the angle between the opposite side and the other diagonal. It follows that the four triangles formed by the two diagonals and the exterior sides are similar in pairs. In particular, the angle at q_4 in the triangle formed by q_1, q_2, q_4 and the angle at q_3 in the triangle formed by q_1, q_2, q_3 are

equal. Denote this angle by θ . By Lemma 5.1, $\theta < \pi/2$, so $\cos \theta > 0$. By the law of cosines in these triangles,

$$\begin{aligned} r_{13}^2 + r_{23}^2 &= r_{12}^2 + 2r_{13}r_{23} \cos \theta, \\ r_{14}^2 + r_{24}^2 &= r_{12}^2 + 2r_{14}r_{24} \cos \theta. \end{aligned}$$

By Lemma 4.1, $r_{13}r_{23} \leq r_{14}r_{24}$, and hence since $\cos \theta > 0$, it follows that $r_{13}^2 + r_{23}^2 \leq r_{14}^2 + r_{24}^2$. Thus $r_{13}^2 - r_{14}^2 \leq r_{24}^2 - r_{23}^2$ and the statement to be proved follows since each of the three factors in the product giving Γ_2 in (4-1) is at most 1. \square

It follows from this result that $\Gamma_1 = 1 \geq \Gamma_2 \geq \Gamma_4 \geq \Gamma_3 > 0$ on lobe II from the previous section. On lobe I, we have $\Gamma_4 > \Gamma_2 > 0 > \Gamma_3$, but at present we do not see how to get good bounds on Γ_4 or Γ_3 .

6. The kite configurations

We call a convex quadrilateral a kite if two opposite vertices lie on an axis of symmetry of the configuration (see Figure 4). Thus a cocircular relative equilibrium forms a kite if and only if one pair of opposite vortices lie on the diameter of the circumscribed circle. There are also kites that are not cocircular, but we will not consider them. In the following, we will assume, as in Figure 4, that the axis of symmetry passes through vortices 1 and 3.

The definition of a kite implies that adjacent sides are equal for the two vortices that lie on the diameter of the circle. Thus the conditions $r_{12} = r_{14} = 1$ and $r_{23} = r_{34}$ hold. For any kite inscribed in a circle, each side of the line of symmetry forms a right triangle. This gives us the Pythagorean relation

$$r_{13}^2 = 1 + r_{34}^2. \tag{6-1}$$

To analyze this case, we will use (2-19), but with $F_5 = 0$ replaced by the equivalent form $F'_5 = 0$ from (2-18). We will make use of Gröbner bases for the ideals generated by these polynomials. See [Cox et al. 2007] for general background




Figure 4. Kite configuration with line of symmetry through vortices 1 and 3.

on this algebraic technique. Equations for the kite configurations are obtained by substituting $r_{14} = 1$ and $r_{23} = r_{34}$. We adjoin an additional equation,

$$1 - tr_{13}r_{24}r_{34}\Gamma_2\Gamma_3\Gamma_4,$$

to force the variables appearing there to be nonzero. Using Sage [Stein et al. 2012], we compute a Gröbner basis for the substituted ideal with respect to the lexicographic order with the variables ordered as

$$t > r_{13} > r_{24} > r_{34} > \Gamma_2 > \Gamma_3 > \Gamma_4.$$

The resulting Gröbner basis contains 24 polynomials, one of which depends only on Γ_3, Γ_4 . After factoring, we see that this polynomial is

$$(4\Gamma_4^2 + \Gamma_4\Gamma_3 + \Gamma_4 - 2\Gamma_3)(-4\Gamma_4^2 + \Gamma_4\Gamma_3 + \Gamma_4 + 2\Gamma_3). \quad (6-2)$$

The next polynomial in the Gröbner basis is

$$\Gamma_2 - \Gamma_4,$$

which shows that $\Gamma_2 = \Gamma_4$ for all kite configurations, as we expect from the symmetry.

The real vanishing locus of each of the two factors in (6-2) is a hyperbola in the (Γ_3, Γ_4) -plane and each of these equations can be solved for Γ_3 in terms of Γ_4 :

$$\Gamma_3 = \frac{\mp 4\Gamma_4^2 - \Gamma_4}{\Gamma_4 \mp 2} \quad (6-3)$$

(the $-$ sign gives the solution of the equation from the left-hand factor in (6-2) and the $+$ gives the solution of the equation from the right-hand factor).

Adjoining each factor in (6-2) to the ideal individually and computing Gröbner bases again, all of the other variables can be expressed in terms of Γ_4 . From the system using the left-hand factor in (6-2), for instance, we obtain

$$r_{34}^2 = \frac{3\Gamma_4}{\Gamma_4 - 2}, \quad r_{24}^2 = \frac{6\Gamma_4}{2\Gamma_4 - 1}, \quad r_{13}^2 = \frac{4\Gamma_4 - 2}{\Gamma_4 - 2}.$$

All of the right sides must be positive since r_{ij} must be nonzero and real. In addition, $r_{34} \leq 1$ forces $-1 \leq \Gamma_4 \leq 0$. However, since $\Gamma_4 > 0$ on the interiors of lobes I and II of the bowtie surface from Theorems 4.4 and 4.3, we see that the left-hand factor from (6-2) is satisfied only for points on the surface $F_3 = 0$ with $r_{23} > r_{14}$.

With the right-hand factor in (6-2), we obtain

$$r_{34}^2 = \frac{3\Gamma_4}{\Gamma_4 + 2}, \quad r_{24}^2 = \frac{6\Gamma_4}{2\Gamma_4 + 1}, \quad r_{13}^2 = \frac{4\Gamma_4 + 2}{\Gamma_4 + 2}. \quad (6-4)$$

(The last equation also follows from (6-1).) Now the equation for r_{34}^2 shows that to get $0 < r_{34} \leq 1$, we must have $0 < \Gamma_4 \leq 1$. Using the $+$ signs in (6-3), it follows




Figure 5. An isosceles trapezoid.

that $\Gamma_3 < 0$ for $0 < \Gamma_4 < \frac{1}{4}$ and $\Gamma_3 > 0$ for $\frac{1}{4} < \Gamma_4 \leq 1$. The points with $\Gamma_3 < 0$ form one of the boundary curves of lobe I of the bowtie surface considered above, and the points with $\Gamma_3 > 0$ give one boundary curve of lobe II. When $\Gamma_4 = \frac{1}{4}$, it follows that $r_{34} = 1/\sqrt{3}$, and the corresponding configuration is the symmetric degenerate configuration mentioned before: an equilateral triangle configuration with $\Gamma_1 = 1$, $\Gamma_2 = \Gamma_4 = \frac{1}{4}$, and an additional vortex with $\Gamma_3 = 0$. When $\Gamma_4 = 1$, we have a geometric square configuration with all exterior sides equal to 1, diagonals equal to $\sqrt{2}$, and all vorticities $\Gamma_i = 1$.

We have proved the following statements.

Theorem 6.1. *There is exactly one kite configuration corresponding to each point on the intersection of the bowtie surface $F_3 = 0$ and the plane given by $r_{23} = r_{34}$. These configurations are parametrized by the value of the vorticity Γ_4 with $0 < \Gamma_4 \leq 1$ as in (6-4). The other vorticities are $\Gamma_2 = \Gamma_4$ and*

$$\Gamma_3 = \frac{4\Gamma_4^2 - \Gamma_4}{\Gamma_4 + 2}.$$

The values $0 < \Gamma_4 \leq \frac{1}{4}$ give the portion of the boundary curve in the closure of lobe I and the values $\frac{1}{4} \leq \Gamma_4 \leq 1$ give the portion of the boundary curve in the closure of lobe II.

7. The isosceles trapezoid configurations

We will call a convex quadrilateral possessing a line of symmetry passing through the midpoints of two opposite edges an isosceles trapezoid. Any such quadrilateral has a circumscribed circle. If we label the vertices as in Figure 5, then the equal pairs of distances are $r_{13} = r_{24}$ and $r_{14} = r_{23}$. The corresponding four-vortex relative equilibria have been described already in Section 7 of [Hampton et al. 2014]. Hence we will only briefly discuss how the results of Hampton, Roberts and Santoprete can be recovered with our setup.

To analyze this case, we will use (2-19). Equations for the isosceles trapezoid configurations are obtained by substituting $r_{23} = r_{14}$ and $r_{24} = r_{13}$. We adjoin an

additional equation,

$$1 - tr_{14}r_{13}r_{34}\Gamma_2\Gamma_3\Gamma_4,$$

to force the variables appearing there to be nonzero. Using Sage [Stein et al. 2012], we compute a Gröbner basis for the substituted ideal with respect to the lexicographic order with the variables ordered as

$$t > r_{14} > r_{13} > r_{34} > \Gamma_3 > \Gamma_4 > \Gamma_2.$$

The resulting Gröbner basis contains 35 polynomials. In factored form, the equations from the polynomials with the three smallest lex leading terms are

$$\begin{aligned} (\Gamma_2 - 1)(r_{34} + 1) &= 0, \\ (\Gamma_4 - 1)(\Gamma_3 - \Gamma_4)(r_{34} + 1) &= 0, \\ (r_{34} - 1)(r_{34} + 1)(\Gamma_3 - \Gamma_4) &= 0. \end{aligned}$$

The first implies that $\Gamma_2 = 1$, since $r_{34} > 0$. Similarly, the second implies either $\Gamma_4 = 1$ or $\Gamma_4 = \Gamma_3$ and the third implies $r_{34} = 1$ or $\Gamma_4 = \Gamma_3$. If $r_{34} = 1$, then the configuration must be a geometric square and $\Gamma_i = 1$ for $i = 1, \dots, 4$. Hence we see the symmetry of the vorticities directly from the form of the Gröbner basis polynomials.

From the subsequent polynomials in the basis, we can solve for the remaining distances in terms of Γ_3 with the triangular form system

$$r_{34}^2 = \frac{2\Gamma_3 + \Gamma_3^2}{2\Gamma_3 + 1}, \quad r_{13}^2 = \frac{\Gamma_3 r_{34}^2 - r_{34}}{\Gamma_3 - r_{34}}, \quad r_{14}^2 = \frac{\Gamma_3 r_{34}^2 + 2r_{13}^2 - \Gamma_3 - 2}{2r_{13}^2 - 2}. \quad (7-1)$$

From the first equation here, we see that $0 < r_{34} \leq 1$ only when $-2 < \Gamma_3 \leq -1$ or $0 < \Gamma_3 \leq 1$. The last equation then shows $r_{14}^2 > 0$ only when $0 < \Gamma_3 \leq 1$.

Theorem 7.1. *There is exactly one isosceles trapezoid configuration corresponding to each point on the intersection of the bowtie surface $F_3 = 0$ and the plane given by $r_{14} = r_{23}$. With the labeling in Figure 5, these configurations are parametrized by the value of the vorticity Γ_3 with $0 < \Gamma_3 \leq 1$ as in (7-1). The point with $\Gamma_3 = 1$ corresponds to the geometric square configuration.*

Acknowledgments

The work reported in this article was begun at the Pacific Undergraduate Research Experience (PURE Math) program at the University of Hawai'i at Hilo in summer 2012 by students Jonathan Gomez, Alexander Gutierrez, and Jesse Robert under the supervision of John Little and Roberto Pelayo. Thanks to everyone at UH Hilo for an enjoyable and productive program. Thanks also to the referees for a careful reading of the manuscript and for identifying a gap in our original proof of the inequality $\Gamma_2 \leq 1$.

References

- [Albouy and Kaloshin 2012] A. Albouy and V. Kaloshin, “Finiteness of central configurations of five bodies in the plane”, *Ann. of Math. (2)* **176**:1 (2012), 535–588. [MR 2925390](#) [Zbl 06074021](#)
- [Cors and Roberts 2012] J. M. Cors and G. E. Roberts, “Four-body co-circular central configurations”, *Nonlinearity* **25**:2 (2012), 343–370. [MR 2876872](#) [Zbl 1235.70033](#)
- [Cox et al. 2007] D. Cox, J. Little, and D. O’Shea, *Ideals, varieties, and algorithms: An introduction to computational algebraic geometry and commutative algebra*, 3rd ed., Springer, New York, 2007. [MR 2007h:13036](#) [Zbl 1118.13001](#)
- [Hampton 2005] M. Hampton, “Co-circular central configurations in the four-body problem”, pp. 993–998 in *EQUADIFF 2003*, edited by F. Dumortier et al., World Sci. Publ., Hackensack, NJ, 2005. [MR 2185162](#) [Zbl 1100.70008](#)
- [Hampton and Jensen 2011] M. Hampton and A. Jensen, “Finiteness of spatial central configurations in the five-body problem”, *Celestial Mech. Dynam. Astronom.* **109**:4 (2011), 321–332. [MR 2012d:70019](#) [Zbl 1270.70038](#)
- [Hampton and Moeckel 2006] M. Hampton and R. Moeckel, “Finiteness of relative equilibria of the four-body problem”, *Invent. Math.* **163**:2 (2006), 289–312. [MR 2008c:70019](#) [Zbl 1083.70012](#)
- [Hampton and Moeckel 2009] M. Hampton and R. Moeckel, “Finiteness of stationary configurations of the four-vortex problem”, *Trans. Amer. Math. Soc.* **361**:3 (2009), 1317–1332. [MR 2009m:76023](#) [Zbl 1161.76011](#)
- [Hampton et al. 2014] M. Hampton, G. E. Roberts, and M. Santoprete, “Relative equilibria in the four-vortex problem with two pairs of equal vorticities”, *J. Nonlinear Sci.* **24**:1 (2014), 39–92. [MR 3162500](#) [Zbl 1302.76042](#)
- [Llibre and Valls 2015] J. Llibre and C. Valls, “The co-circular central configurations of the 5-body problem”, *J. Dynam. Differential Equations* **27**:1 (2015), 55–67. [MR 3317391](#) [Zbl 06425778](#)
- [Moeckel 1990] R. Moeckel, “On central configurations”, *Math. Z.* **205**:4 (1990), 499–517. [MR 92b:70012](#) [Zbl 0684.70005](#)
- [Saari 2011] D. G. Saari, “Central configurations—a problem for the twenty-first century”, pp. 283–298 in *Expeditions in mathematics*, edited by T. Shubin et al., Mathematical Association of America, Washington, DC, 2011. [MR 2012g:00008](#) [Zbl 1214.00002](#)
- [Stein et al. 2012] W. A. Stein et al., *Sage mathematics software*, Version 5.0, Sage Development Team, 2012, available at <http://www.sagemath.org>.

Received: 2014-11-14

Revised: 2015-03-12

Accepted: 2015-05-09

gomezjon@math.hawaii.edu*Department of Mathematics, University of Hawai’i at Manoa, Honolulu, HI 96822, United States*alexg@umn.edu*Department of Mathematics, University of Minnesota, Minneapolis, MN 55455, United States*jlittle@holycross.edu*Department of Mathematics and Computer Science, College of the Holy Cross, Worcester, MA 01610, United States*robertop@hawaii.edu*Department of Mathematics, University of Hawai’i at Hilo, Hilo, HI 96720, United States*jesse20@hawaii.edu*Department of Mathematics, University of Hawai’i at Hilo, Hilo, HI 96720, United States*

INVOLVE YOUR STUDENTS IN RESEARCH

Involve showcases and encourages high-quality mathematical research involving students from all academic levels. The editorial board consists of mathematical scientists committed to nurturing student participation in research. Bridging the gap between the extremes of purely undergraduate research journals and mainstream research journals, *Involve* provides a venue to mathematicians wishing to encourage the creative involvement of students.

MANAGING EDITOR

Kenneth S. Berenhaut Wake Forest University, USA

BOARD OF EDITORS

Colin Adams	Williams College, USA	Suzanne Lenhart	University of Tennessee, USA
John V. Baxley	Wake Forest University, NC, USA	Chi-Kwong Li	College of William and Mary, USA
Arthur T. Benjamin	Harvey Mudd College, USA	Robert B. Lund	Clemson University, USA
Martin Bohner	Missouri U of Science and Technology, USA	Gaven J. Martin	Massey University, New Zealand
Nigel Boston	University of Wisconsin, USA	Mary Meyer	Colorado State University, USA
Amarjit S. Budhiraja	U of North Carolina, Chapel Hill, USA	Emil Minchev	Ruse, Bulgaria
Pietro Cerone	La Trobe University, Australia	Frank Morgan	Williams College, USA
Scott Chapman	Sam Houston State University, USA	Mohammad Sal Moslehian	Ferdowsi University of Mashhad, Iran
Joshua N. Cooper	University of South Carolina, USA	Zuhair Nashed	University of Central Florida, USA
Jem N. Corcoran	University of Colorado, USA	Ken Ono	Emory University, USA
Toka Diagana	Howard University, USA	Timothy E. O'Brien	Loyola University Chicago, USA
Michael Dorff	Brigham Young University, USA	Joseph O'Rourke	Smith College, USA
Sever S. Dragomir	Victoria University, Australia	Yuval Peres	Microsoft Research, USA
Behrouz Emamizadeh	The Petroleum Institute, UAE	Y.-F. S. Pétermann	Université de Genève, Switzerland
Joel Foisy	SUNY Potsdam, USA	Robert J. Plemmons	Wake Forest University, USA
Erin W. Fulp	Wake Forest University, USA	Carl B. Pomerance	Dartmouth College, USA
Joseph Gallian	University of Minnesota Duluth, USA	Vadim Ponomarenko	San Diego State University, USA
Stephan R. Garcia	Pomona College, USA	Bjorn Poonen	UC Berkeley, USA
Anant Godbole	East Tennessee State University, USA	James Propp	U Mass Lowell, USA
Ron Gould	Emory University, USA	József H. Przytycki	George Washington University, USA
Andrew Granville	Université Montréal, Canada	Richard Rebarber	University of Nebraska, USA
Jerrold Griggs	University of South Carolina, USA	Robert W. Robinson	University of Georgia, USA
Sat Gupta	U of North Carolina, Greensboro, USA	Filip Saidak	U of North Carolina, Greensboro, USA
Jim Haglund	University of Pennsylvania, USA	James A. Sellers	Penn State University, USA
Johnny Henderson	Baylor University, USA	Andrew J. Sterge	Honorary Editor
Jim Hoste	Pitzer College, USA	Ann Trenk	Wellesley College, USA
Natalia Hritonenko	Prairie View A&M University, USA	Ravi Vakil	Stanford University, USA
Glenn H. Hurlbert	Arizona State University, USA	Antonia Vecchio	Consiglio Nazionale delle Ricerche, Italy
Charles R. Johnson	College of William and Mary, USA	Ram U. Verma	University of Toledo, USA
K. B. Kulasekera	Clemson University, USA	John C. Wierman	Johns Hopkins University, USA
Gerry Ladas	University of Rhode Island, USA	Michael E. Zieve	University of Michigan, USA

PRODUCTION

Silvio Levy, Scientific Editor


Cover: Alex Scorpan

See inside back cover or msp.org/involve for submission instructions. The subscription price for 2016 is US \$160/year for the electronic version, and \$215/year (+\$35, if shipping outside the US) for print and electronic. Subscriptions, requests for back issues from the last three years and changes of subscribers address should be sent to MSP.

Involve (ISSN 1944-4184 electronic, 1944-4176 printed) at Mathematical Sciences Publishers, 798 Evans Hall #3840, c/o University of California, Berkeley, CA 94720-3840, is published continuously online. Periodical rate postage paid at Berkeley, CA 94704, and additional mailing offices.

Involve peer review and production are managed by EditFLOW® from Mathematical Sciences Publishers.

PUBLISHED BY

 **mathematical sciences publishers**
nonprofit scientific publishing

<http://msp.org/>

© 2016 Mathematical Sciences Publishers

involve

2016

vol. 9

no. 3

A combinatorial proof of a decomposition property of reduced residue systems	361
YOTSANAN MEEMARK AND THANAKORN PRINYASART	
Strong depth and quasigeodesics in finitely generated groups	367
BRIAN GAPINSKI, MATTHEW HORAK AND TYLER WEBER	
Generalized factorization in $\mathbb{Z}/m\mathbb{Z}$	379
AUSTIN MAHLUM AND CHRISTOPHER PARK MOONEY	
Cocircular relative equilibria of four vortices	395
JONATHAN GOMEZ, ALEXANDER GUTIERREZ, JOHN LITTLE, ROBERTO PELAYO AND JESSE ROBERT	
On weak lattice point visibility	411
NEIL R. NICHOLSON AND REBECCA RACHAN	
Connectivity of the zero-divisor graph for finite rings	415
REZA AKHTAR AND LUCAS LEE	
Enumeration of m -endomorphisms	423
LOUIS RUBIN AND BRIAN RUSHTON	
Quantum Schubert polynomials for the G_2 flag manifold	437
RACHEL E. ELLIOTT, MARK E. LEWERS AND LEONARDO C. MIHALCEA	
The irreducibility of polynomials related to a question of Schur	453
LENNY JONES AND ALICIA LAMARCHE	
Oscillation of solutions to nonlinear first-order delay differential equations	465
JAMES P. DIX AND JULIO G. DIX	
A variational approach to a generalized elastica problem	483
C. ALEX SAFSTEN AND LOGAN C. TATHAM	
When is a subgroup of a ring an ideal?	503
SUNIL K. CHEBOLU AND CHRISTINA L. HENRY	
Explicit bounds for the pseudospectra of various classes of matrices and operators	517
FEIXUE GONG, OLIVIA MEYERSON, JEREMY MEZA, MIHAI STOICIU AND ABIGAIL WARD	