

involve

a journal of mathematics

Spectrum of a composition operator with automorphic symbol

Robert F. Allen, Thong M. Le and Matthew A. Pons



Spectrum of a composition operator with automorphic symbol

Robert F. Allen, Thong M. Le and Matthew A. Pons

(Communicated by Stephan Garcia)

We give a complete characterization of the spectrum of composition operators, induced by an automorphism of the open unit disk, acting on a family of Banach spaces of analytic functions that includes the Bloch space and BMOA. We show that for parabolic and hyperbolic automorphisms the spectrum is the unit circle. For the case of elliptic automorphisms, the spectrum is either the unit circle or a finite cyclic subgroup of the unit circle.

1. Introduction

For an analytic self-map φ of the open unit disk \mathbb{D} and a Banach space X of functions analytic on \mathbb{D} , we define the composition operator with symbol φ , denoted C_φ , by the rule $C_\varphi f = f \circ \varphi$ for all $f \in X$. The study of composition operators began formally with Nordgren's paper [1968], where he explored properties of composition operators acting on the Hardy Hilbert space H^2 . Since then the study has proved to be an active area of research, most likely due to the fact that the study of such operators lies at the intersection of complex function theory and operator theory.

The spectrum of C_φ has been studied on many classical spaces of analytic functions, such as the Hardy spaces, Bergman spaces, weighted Hardy and Bergman spaces, Besov spaces, and the Dirichlet space. The interested reader is directed to [Cowen and MacCluer 1995] for general references.

The motivation for this paper was to determine the spectrum of a composition operator, induced by a disk automorphism, acting on the Bloch space. The Bloch space is the largest space of analytic functions on \mathbb{D} that is Möbius invariant. This is one reason the Bloch space is a welcoming environment to study composition operators. The techniques developed apply to a larger class of spaces that includes the Bloch space.

The purpose of this paper is to determine the spectrum of C_φ acting on a family of Banach spaces, where φ is a disk automorphism. The spectrum will depend on

MSC2010: primary 47A10, 47B33; secondary 30H05.

Keywords: composition operator, spectrum, automorphism.

the fixed point classification of the automorphisms of \mathbb{D} . This is a standard approach to the study of composition operators induced by automorphisms. We show the spectrum of C_φ , acting on a particular family of Banach spaces, induced by a disk automorphism, must be a subset of the unit circle $\partial\mathbb{D}$, and in some instances is the entire unit circle. Finally, we compare these results to particular examples of classical spaces.

2. Preliminaries

2A. Automorphisms. The automorphisms of the open unit disk $\mathbb{D} = \{z \in \mathbb{C} : |z| < 1\}$ are precisely the analytic bijections on \mathbb{D} which have the form

$$\varphi(z) = \lambda \frac{a - z}{1 - \bar{a}z},$$

where λ is a unimodular constant and a is a point in \mathbb{D} . These automorphisms form a group under composition denoted by $\text{Aut}(\mathbb{D})$. Every element of $\text{Aut}(\mathbb{D})$ has two fixed points (counting multiplicity), and thus can be classified by the location of the fixed points:

elliptic: one fixed point in \mathbb{D} and one in the complement of $\bar{\mathbb{D}}$;

parabolic: one fixed point on the unit circle $\partial\mathbb{D}$ (of multiplicity 2);

hyperbolic: two distinct fixed points on $\partial\mathbb{D}$.

Two disk automorphisms φ and ψ are conformally equivalent if there exists a disk automorphism τ for which $\psi = \tau \circ \varphi \circ \tau^{-1}$. Many properties of automorphisms are preserved under conformal equivalence. The main advantage of conformal equivalence is in the placement of the fixed points. Every elliptic disk automorphism is conformally equivalent to one whose fixed point in \mathbb{D} is the origin.

Lemma 2.1. *Let φ be an elliptic disk automorphism with fixed point a in \mathbb{D} . Then φ is conformally equivalent to $\psi(z) = \lambda z$ where $\lambda = \varphi'(a)$.*

Proof. Let τ_a be the involution automorphism which interchanges 0 and a , that is

$$\tau_a(z) = \frac{a - z}{1 - \bar{a}z}.$$

Define $\psi = \tau_a \circ \varphi \circ \tau_a^{-1}$ on \mathbb{D} . Since a is a fixed point of φ , ψ fixes the origin, and is a rotation. So there is a unimodular constant λ such that $\psi(z) = \lambda z$. To complete the proof, we will show $\lambda = \varphi'(a)$. Observe $\psi'(z) = \lambda$ for all $z \in \mathbb{D}$. In particular

$$\lambda = \psi'(0) = \tau_a'(\varphi(\tau_a(0)))\varphi'(\tau_a(0))\tau_a'(0) = \varphi'(a)\tau_a'(a)\tau_a'(0) = \varphi'(a).$$

Thus φ is conformally equivalent to the rotation $\psi(z) = \varphi'(a)z$. □

Every parabolic disk automorphism is conformally equivalent to one whose fixed point (of multiplicity 2) is 1. The following lemma is found as Exercise 2.3.5c of [Cowen and MacCluer 1995], and a complete proof can be found in [Pons 2007].

Lemma 2.2 [Pons 2007, Lemma 4.1.2]. *Let φ be a parabolic disk automorphism. Then φ is conformally equivalent to either*

$$\psi_1(z) = \frac{(1+i)z-1}{z+i-1} \quad \text{or} \quad \psi_2(z) = \frac{(1-i)z-1}{z-i-1}.$$

Every hyperbolic disk automorphism is conformally equivalent to one whose fixed points in $\partial\mathbb{D}$ are ± 1 .

Lemma 2.3 [Nordgren 1968, Theorem 6]. *Let φ be a hyperbolic disk automorphism. Then, for some $r \in (0, 1)$, φ is conformally equivalent to*

$$\psi(z) = \frac{z+r}{1+rz}.$$

2B. The space of bounded analytic functions. The set of analytic functions on \mathbb{D} is denoted by $H(\mathbb{D})$. The space of bounded analytic functions on \mathbb{D} , denoted $H^\infty = H^\infty(\mathbb{D})$, is a Banach space under the norm

$$\|f\|_\infty = \sup_{z \in \mathbb{D}} |f(z)|.$$

The bounded analytic functions on \mathbb{D} is a rich space containing many interesting types of functions, such as polynomials and Blaschke products. In addition, the disk algebra $\mathcal{A}(\mathbb{D})$, the set of analytic functions on \mathbb{D} continuous to $\partial\mathbb{D}$, is a closed subspace of H^∞ .

The following two families of functions will be used in the next section. To prove these functions are in H^∞ , we take a geometric approach using conformal mappings of the plane. To this effect let \mathbb{H}_ℓ and \mathbb{H}_r denote the open left and right half planes respectively, i.e., $\mathbb{H}_\ell = \{\operatorname{Re} z < 0\}$ and $\mathbb{H}_r = \{\operatorname{Re} z > 0\}$.

Lemma 2.4. *For $s \geq 0$, the function*

$$f_s(z) = \exp \frac{s(z+1)}{z-1}$$

is in H^∞ .

Proof. If $s = 0$, then f_s is identically 1. So, $f_s(z)$ is in H^∞ . Now suppose $s > 0$. The function f_s is comprised of the functions

- (1) $z \mapsto (z+1)/(z-1)$ (mapping \mathbb{D} onto \mathbb{H}_ℓ),
- (2) $z \mapsto sz$ (mapping \mathbb{H}_ℓ onto \mathbb{H}_ℓ),
- (3) $z \mapsto e^z$ (mapping \mathbb{H}_ℓ onto $\mathbb{D} \setminus \{0\}$).




Figure 1. Map $f_s(z) = \exp \frac{s(z+1)}{z-1}$ for $s > 0$.

So f_s maps \mathbb{D} into \mathbb{D} , as in Figure 1, and thus $f_s(z)$ is an element of H^∞ . □

Lemma 2.5. For real value t , the function

$$f_t(z) = \left(\frac{1+z}{1-z} \right)^{it}$$

is in H^∞ .

Proof. For $t = 0$, f_t is identically 1, and thus is in H^∞ . Now suppose $t > 0$. We will rewrite the function f_t as

$$f_t(z) = \exp \left(it \log \frac{1+z}{1-z} \right),$$

where \log is the principle branch of the logarithm. Then f_t is comprised of the functions

- (1) $z \mapsto (1+z)/(1-z)$ (mapping \mathbb{D} onto H_r),
- (2) $z \mapsto \log z$ (mapping H_r onto the horizontal strip $S_h = \{0 < \text{Im } z < 2\pi\}$),
- (3) $z \mapsto itz$ (mapping S_h onto the vertical strip $S_v = \{-2\pi < \text{Re } z < 0\}$),
- (4) $z \mapsto e^z$ (mapping S_v into $A(e^{-2\pi}, 1) = \{e^{-2\pi} < |z| < 1\}$).

So f_t maps \mathbb{D} into $A(e^{-2\pi}, 1) \subseteq \mathbb{D}$, as depicted in Figure 2. In the case of $t < 0$, the vertical strip S_v becomes $\{0 < \text{Re } z < 2\pi\}$. The map $z \mapsto e^z$ takes S_v into $A(1, e^{2\pi}) \subseteq e^{2\pi} \mathbb{D}$, as depicted in Figure 3. In either case, $f_t(z)$ is an element of H^∞ since $\|f_t\|_\infty < e^{2\pi}$ for all $t \in \mathbb{R}$. □




Figure 2. Map $f_t(z) = \exp \left(it \log \frac{1+z}{1-z} \right)$ for $t > 0$.




Figure 3. Map $f_t(z) = \exp\left(it \log \frac{1+z}{1-z}\right)$ for $t < 0$.

These functions above, together with the monomials, play such a pivotal role in Section 3 that we denote the union of these functions by \mathcal{F} , i.e.,

$$\mathcal{F} = \{f_s : s \geq 0\} \cup \{f_t : t \in \mathbb{R}\} \cup \{z^k : k \in \mathbb{N}\}.$$

2C. Spectrum of C_φ . In this section we collect useful results regarding the spectrum of operators on Banach spaces. For a bounded linear operator T on a Banach space X , the spectrum of T is given by

$$\sigma(T) = \{\lambda \in \mathbb{C} : T - \lambda I \text{ is not invertible}\}$$

where I denotes the identity operator on X . The spectrum is a nonempty, closed subset of \mathbb{C} . The spectral radius of T is given by

$$\rho(T) = \sup \{|\lambda| : \lambda \in \sigma(T)\}.$$

Due to the fact that the spectrum is closed, we have the spectrum of T is contained in the closed disk centered at the origin of radius $\rho(T)$.

Determining the spectrum of a particular composition operator can be difficult depending on the symbol of the operator and the space on which it is acting. However, the difficulties can be avoided if the operator is similar to a “simpler” operator. Linear operators S and T (not necessarily bounded) on a Banach space X are similar if there exists a bounded linear operator U on X , having bounded inverse, such that $T = USU^{-1}$. If S and T are both bounded operators, then similarity preserves the spectrum.

Theorem 2.6. *Let S and T be bounded operators on a Banach space X . If S and T are similar, then $\sigma(S) = \sigma(T)$.*

Proof. Suppose S and T are similar operators on X . By definition, there exists an invertible, bounded operator U such that $T = USU^{-1}$. Let $\lambda \in \mathbb{C}$ and observe that

$$\begin{aligned} T - \lambda I &= USU^{-1} - \lambda I \\ &= USU^{-1} - \lambda UU^{-1} \\ &= USU^{-1} - U(\lambda I)U^{-1} \\ &= U(S - \lambda I)U^{-1}. \end{aligned}$$

Thus, we have that $S - \lambda I$ is not invertible if and only if $T - \lambda I$ is not invertible. Therefore $\sigma(S) = \sigma(T)$. \square

3. Main results

In this section, we determine the spectrum of C_φ for φ a disk automorphism acting on a particular family of Banach spaces of analytic functions. The spaces we consider will be denoted by \mathcal{X} and have the following properties:

- (i) \mathcal{X} contains \mathcal{F} ,
- (ii) for all $\varphi \in \text{Aut}(\mathbb{D})$, C_φ is bounded on \mathcal{X} and $\rho(C_\varphi) = 1$.

The set of automorphisms of \mathbb{D} , as seen previously, is a very nice subset of the analytic self-maps of \mathbb{D} . By property (ii), every composition operator induced by a disk automorphism is bounded on \mathcal{X} . In fact, every such composition operator is invertible. This result, that we prove below, can be viewed as a consequence of Theorem 1.6 of [Cowen and MacCluer 1995].

Proposition 3.1. *Let φ be a disk automorphism and C_φ the induced composition operator on \mathcal{X} . Then C_φ is invertible with inverse $C_\varphi^{-1} = C_{\varphi^{-1}}$.*

Proof. Since $\varphi \in \text{Aut}(\mathbb{D})$, φ is invertible, and φ^{-1} is an automorphism. The composition operator $C_{\varphi^{-1}}$ is bounded by property (ii) and

$$\begin{aligned} C_\varphi(C_{\varphi^{-1}}(f)) &= C_\varphi(f \circ \varphi^{-1}) = f \circ \varphi^{-1} \circ \varphi = f, \\ C_{\varphi^{-1}}(C_\varphi(f)) &= C_{\varphi^{-1}}(f \circ \varphi) = f \circ \varphi \circ \varphi^{-1} = f. \end{aligned}$$

Therefore, C_φ is invertible with $C_\varphi^{-1} = C_{\varphi^{-1}}$. \square

Since the spectral radius of C_φ on \mathcal{X} is 1 for $\varphi \in \text{Aut}(\mathbb{D})$, we see that the search for the spectrum can be restricted to subsets of $\overline{\mathbb{D}}$. However, our search can be refined further to subsets of the unit circle.

Theorem 3.2. *Let φ be a disk automorphism and C_φ the induced composition operator on \mathcal{X} . Then $\sigma(C_\varphi) \subseteq \partial\mathbb{D}$.*

Proof. By property (ii) of \mathcal{X} , we have $\rho(C_\varphi) = 1$. So, $\sigma(C_\varphi) \subseteq \overline{\mathbb{D}}$. Since, by Proposition 3.1, C_φ is invertible with the inverse $C_\varphi^{-1} = C_{\varphi^{-1}}$, then $0 \notin \sigma(C_\varphi)$. So, the function $f(z) = z^{-1}$ is analytic in some neighborhood of $\sigma(C_\varphi)$. By the Spectral Mapping Theorem (see Theorem 5.14 of [MacCluer 2009]), we have $\sigma(f \circ C_\varphi) = f(\sigma(C_\varphi))$, and so,

$$\sigma(C_{\varphi^{-1}}) = \sigma(C_\varphi^{-1}) = \sigma(C_\varphi)^{-1} = \{\lambda^{-1} : \lambda \in \sigma(C_\varphi)\}.$$

Since $\varphi^{-1} \in \text{Aut}(\mathbb{D})$, $\sigma(C_{\varphi^{-1}}) \subseteq \overline{\mathbb{D}}$. Thus for $\lambda \in \sigma(C_\varphi)$, both λ and λ^{-1} are in $\overline{\mathbb{D}}$. This implies $\lambda \in \partial\mathbb{D}$. So $\sigma(C_\varphi) \subseteq \partial\mathbb{D}$, as desired. \square

Since the disk automorphisms are classified into three categories, according to fixed points, we will treat each type of automorphism separately. However, the strategy to determine $\sigma(C_\varphi)$ is the same. For a disk automorphism φ , we have shown φ to be conformally equivalent to a particularly “nice” disk automorphism: in the elliptic case a disk automorphism that fixes 0, in the parabolic case a disk automorphism that fixes 1, and in the hyperbolic case a disk automorphism that fixes ± 1 . In the next result, we show that conformally equivalent automorphisms induce similar composition operators on \mathcal{X} . This result is not unique to the space \mathcal{X} , but is true for any space for which automorphisms induce bounded composition operators (see p. 250 of [Cowen and MacCluer 1995]).

Proposition 3.3. *Let φ and ψ be conformally equivalent disk automorphisms. Then the induced composition operators C_φ and C_ψ on \mathcal{X} are similar.*

Proof. Suppose φ and ψ are conformally equivalent disk automorphisms. Then there exists a disk automorphism τ such that $\psi = \tau \circ \varphi \circ \tau^{-1}$. For $f \in \mathcal{X}$, observe

$$C_\psi f = f \circ (\tau \circ \varphi \circ \tau^{-1}) = ((f \circ \tau) \circ \varphi) \circ \tau^{-1} = (C_{\tau^{-1}} C_\varphi C_\tau) f.$$

Since $C_{\tau^{-1}}$ is bounded and invertible on \mathcal{X} with $C_{\tau^{-1}}^{-1} = C_\tau$, then $C_\psi = C_{\tau^{-1}} C_\varphi C_\tau^{-1}$. Therefore C_φ and C_ψ are similar. □

With Proposition 3.3 and Lemmas 2.1, 2.2, and 2.3, it suffices to determine the spectrum of composition operators induced by these “nice” disk automorphisms, since similarity of bounded operators preserves the spectrum.

Theorem 3.4. *Let φ be an elliptic disk automorphism with fixed point a in \mathbb{D} . Then the spectrum of C_φ acting on \mathcal{X} is the closure of the positive powers of $\varphi'(a)$. Moreover, this closure is a finite subgroup of the unit circle if $\varphi'(a)^n = 1$ for some natural number n , and is the unit circle otherwise.*

Proof. By Lemma 2.1, φ is conformally equivalent to $\psi(z) = \lambda z$ where $\lambda = \varphi'(a)$. By Proposition 3.3, it suffices to show that $\sigma(C_\psi)$ is the closure of the positive powers of λ . Let $G = \langle \lambda \rangle = \{\lambda^k : k \in \mathbb{N}\}$, which is a subset of $\partial\mathbb{D}$ since $|\lambda| = 1$. For each $k \in \mathbb{N}$, the function $f_k(z) = z^k$ is in \mathcal{X} by property (i), and we have $(C_\psi f_k)(z) = \lambda^k f_k(z)$. Thus λ^k is an eigenvalue of C_ψ corresponding to the eigenfunction f_k . So $G \subseteq \sigma(C_\psi)$, and since the spectrum is closed, we have $\overline{G} \subseteq \sigma(C_\psi) = \sigma(C_\varphi)$. If the order of λ is infinite, then G is dense in $\partial\mathbb{D}$, and so $\overline{G} = \partial\mathbb{D}$.

Now suppose λ has order $m < \infty$. Then $G = \{\lambda^k : k = 1, \dots, m\}$. So, $\overline{G} = G$. We now wish to show $\sigma(C_\psi) \subseteq \overline{G}$. Since $\sigma(C_\psi) \subseteq \partial\mathbb{D}$ by Theorem 3.2 it suffices to show that if $\mu \in \partial\mathbb{D} \setminus \overline{G}$ then $\mu \notin \sigma(C_\psi)$. Suppose $\mu \in \partial\mathbb{D} \setminus \overline{G}$.

Since $\mu \notin \overline{G}$, it clear that $\mu \notin G$ and $\mu^m \neq 1$. In order to show $\mu \notin \sigma(C_\psi)$, we will show that $C_\psi - \mu I$ is invertible by proving that for every $g \in \mathcal{X}$, there exists a unique $f \in \mathcal{X}$ such that $f \circ \psi - \mu f = g$.

Since the order of λ is m , we have

$$\psi^{(m)}(z) = \underbrace{(\psi \circ \dots \circ \psi)}_{m\text{-times}}(z) = \lambda^m z = z.$$

By repeated composition with ψ , we obtain the system of linear equations:

$$\begin{aligned} f \circ \psi - \mu f &= g \\ f \circ \psi^{(2)} - \mu(f \circ \psi) &= g \circ \psi \\ &\vdots \\ f - \mu(f \circ \psi^{(m-1)}) &= g \circ \psi^{(m-1)}. \end{aligned}$$

This system of linear equations can be expressed as the matrix equation $A\vec{x} = \vec{b}$ where

$$A = \begin{bmatrix} -\mu & 1 & 0 & 0 & \dots & 0 \\ 0 & -\mu & 1 & 0 & \dots & 0 \\ \vdots & 0 & \ddots & \ddots & & \vdots \\ \vdots & \vdots & \ddots & \ddots & \ddots & \vdots \\ 0 & \vdots & & \ddots & \ddots & 1 \\ 1 & 0 & \dots & \dots & 0 & -\mu \end{bmatrix}, \vec{x} = \begin{bmatrix} f \\ f \circ \psi \\ \vdots \\ \vdots \\ f \circ \psi^{(m-2)} \\ f \circ \psi^{(m-1)} \end{bmatrix}, \text{ and } \vec{b} = \begin{bmatrix} g \\ g \circ \psi \\ \vdots \\ \vdots \\ g \circ \psi^{(m-2)} \\ g \circ \psi^{(m-1)} \end{bmatrix}.$$

The determinant of A is $(-1)^m(\mu^m - 1)$, which is not zero since $\mu \notin G$. Thus there is a unique solution for \vec{x} . It gives us the unique solution f , which is a finite linear combination of function in \mathcal{X} of the form $g \circ \psi^{(j-1)}$ for $j = 1, \dots, m$, and thus f is in \mathcal{X} . It follows that $C_\psi - \mu I$ is invertible. So, $\mu \notin \sigma(C_\psi)$. Therefore, $\sigma(C_\varphi) = \sigma(C_\psi) \subseteq \overline{G}$. □

Theorem 3.5. *Let φ be a parabolic disk automorphism. Then the spectrum of C_φ acting on \mathcal{X} is the unit circle.*

Proof. From Lemma 2.2, φ is conformally equivalent to either

$$\psi_1(z) = \frac{(1+i)z - 1}{z + i - 1} \quad \text{or} \quad \psi_2(z) = \frac{(1-i)z - 1}{z - i - 1}.$$

By Theorems 3.2 and 3.3 it suffices to show that $\partial\mathbb{D}$ is a subset of $\sigma(C_{\psi_1})$ and $\sigma(C_{\psi_2})$.

First suppose φ is conformally equivalent to ψ_1 . Consider the function

$$f_s(z) = \exp \frac{s(z+1)}{z-1}$$

for $s \geq 0$. By property (i), f_s is in \mathcal{X} . Observe

$$\begin{aligned} (C_{\psi_1} f_s)(z) &= f_s(\psi_1(z)) = f_s\left(\frac{(1+i)z-1}{z+i-1}\right) \\ &= \exp \frac{s\left(\frac{(1+i)z-1}{z+i-1} + 1\right)}{\frac{(1+i)z-1}{z+i-1} - 1} = \exp \frac{s((1+i)z-1+z+i-1)}{(1+i)z-1-z-i+1} \\ &= \exp \frac{s((2+i)z+i-2)}{i(z-1)} = \exp \frac{s((1-2i)z+1+2i)}{z-1} \\ &= \exp\left(\frac{s(z+1)}{z-1} - 2is\right) = e^{i(-2s)} f_s(z). \end{aligned}$$

So, f_s is an eigenfunction of C_{ψ_1} for $s \geq 0$. Then, $\partial\mathbb{D} = \{e^{i(-2s)} : s \geq 0\}$ is a subset of $\sigma(C_{\psi_1})$. If φ is conformally equivalent to ψ_2 , then by a similar calculation, we have

$$(C_{\psi_2} f_s)(z) = e^{2is} f_s(z),$$

and so $\partial\mathbb{D} = \{e^{2is} : s \geq 0\}$ is a subset of $\sigma(C_{\psi_2})$. Therefore, $\sigma(C_\varphi) = \partial\mathbb{D}$, as desired. \square

Theorem 3.6. *Let φ be a hyperbolic disk automorphism. Then the spectrum of C_φ acting on \mathcal{X} is the unit circle.*

Proof. From Lemma 2.3, φ is conformally equivalent to $\psi(z) = (z+r)/(1+rz)$ for some $r \in (0, 1)$. By Theorems 3.2 and 3.3 it suffices to show that $\partial\mathbb{D} \subseteq \sigma(C_\psi)$. Consider the function

$$f_t(z) = \left(\frac{1+z}{1-z}\right)^{it}$$

for $t \in \mathbb{R}$. By property (i), f_t is in \mathcal{X} . Observe

$$\begin{aligned} (C_\psi f_t)(z) &= f_t(\psi(z)) = f_t\left(\frac{z+r}{1+rz}\right) \\ &= \left(\frac{1+\frac{z+r}{1+rz}}{1-\frac{z+r}{1+rz}}\right)^{it} = \left(\frac{1+rz+z+r}{1+rz-z-r}\right)^{it} \\ &= \left(\frac{(r+1)z+(r+1)}{(r-1)z-(r-1)}\right)^{it} \\ &= \left(\frac{r+1}{r-1}\right)^{it} f_t(z). \end{aligned}$$

So, f_t is an eigenfunction of C_ψ for t real. Then $\partial\mathbb{D} = \left\{\left(\frac{r+1}{r-1}\right)^{it} : 0 < r < 1, t \in \mathbb{R}\right\}$ is a subset of $\sigma(C_\psi) = \sigma(C_\varphi)$. Therefore $\sigma(C_\varphi) = \partial\mathbb{D}$, as desired. \square

4. Examples and comparisons

In this section we first consider examples of spaces that satisfy the properties of \mathcal{X} . For these spaces, our results characterize the spectrum of composition operators induced by disk automorphisms. Lastly, we consider spaces that do not satisfy the properties of \mathcal{X} but for which the spectrum of composition operators induced by automorphisms is known. We will compare the spectra for those spaces with the characterization for \mathcal{X} .

4A. Examples. First, we will discuss examples of spaces that satisfy the properties of \mathcal{X} .

4A1. Bounded analytic functions. The property (i) of \mathcal{X} is satisfied by H^∞ by Lemmas 2.4 and 2.5. In fact, on H^∞ , any analytic self-map of \mathbb{D} induces a bounded composition operator C_φ such that $\|C_\varphi\| = 1$. Equality is achieved since H^∞ contains the constant function 1. The spectral radius formula (see Theorem 5.15 of [MacCluer 2009]) then implies that $\rho(C_\varphi) = 1$. Thus, property (ii) is satisfied. Thus H^∞ belongs to the family of Banach spaces of analytic functions \mathcal{X} .

4A2. Bloch space. The Bloch space on \mathbb{D} , denoted $\mathcal{B} = \mathcal{B}(\mathbb{D})$, is the space of analytic functions on \mathbb{D} such that $\beta_f = \sup_{z \in \mathbb{D}} (1 - |z|^2)|f'(z)| < \infty$. The quantity β_f is a seminorm, called the Bloch seminorm. The Bloch space is a Banach space under the norm

$$\|f\|_{\mathcal{B}} = |f(0)| + \beta_f.$$

It is well-known that \mathcal{B} is a Banach space of analytic functions that contains H^∞ , and thus satisfies property (i) of \mathcal{X} . In fact, every analytic self-map of \mathbb{D} induces a bounded composition operator on \mathcal{B} (see [Arazy et al. 1985, p. 126]). Donaway, in his Ph.D. thesis, proved the spectral radius of every composition operator induced by an analytic function on \mathbb{D} , and in particular the disk automorphisms, is 1, by [Donaway 1999, Corollary 3.9]. So the Bloch space satisfies all the properties of \mathcal{X} .

4A3. Analytic functions of bounded mean oscillation. The space of analytic functions on \mathbb{D} with bounded mean oscillation on $\partial\mathbb{D}$, denoted BMOA, is defined to be the set of functions in $H(\mathbb{D})$ such that

$$\|f\|_* = \sup_{z \in \mathbb{D}} \|f \circ \tau_a - f(z)\|_{H^2} < \infty,$$

where H^2 is defined in Section 4B1. The space BMOA is a Banach space under the norm

$$\|f\|_{\text{BMOA}} = |f(0)| + \|f\|_*.$$

It is well-known that BMOA is a Banach space of analytic functions, a subspace of the Bloch space, and contains H^∞ as a subspace since $\|f\|_{\text{BMOA}} \leq 3\|f\|_\infty$.

Thus property (i) is satisfied by BMOA. The following result shows property (ii) is satisfied by BMOA also.

Theorem 4.1. *Let φ be an analytic self-map of \mathbb{D} . Then C_φ acting on BMOA is bounded and $\rho(C_\varphi) = 1$.*

Proof. As a result of the Littlewood subordination principle (see Theorem 1.7 of [Duren 1970]), every analytic self-map φ of \mathbb{D} induces a bounded composition operator on BMOA.

To compute the spectral radius of C_φ acting on BMOA, we first estimate the norm. By Corollary 2.2 of [Laitila 2009] there is a constant $M > 0$, independent of φ , such that

$$\|C_\varphi\| \leq M \left(\sup_{a \in \mathbb{D}} \|\tau_{\varphi(a)} \circ \varphi \circ \tau_a\|_{H^2} + \log \frac{2}{1 - |\varphi(0)|^2} \right). \tag{4-1}$$

Since the function $\tau_{\varphi(a)} \circ \varphi \circ \tau_a$ is a composition of self-maps of the disk, the first term on the right is bounded above by 1. Also,

$$\frac{1}{1 - |\varphi(0)|^2} \leq \frac{1 + |\varphi(0)|}{1 - |\varphi(0)|} \leq \frac{2}{1 - |\varphi(0)|}$$

and hence

$$\log \left(\frac{2}{1 - |\varphi(0)|^2} \right) \leq \log \left(\frac{4}{1 - |\varphi(0)|} \right) \leq 2 \log 2 - \log(1 - |\varphi(0)|).$$

Applying these estimates to Equation (4-1), we have

$$\|C_\varphi\| \leq M(1 + 2 \log 2) - M \log(1 - |\varphi(0)|).$$

This immediately implies that

$$\|C_{\varphi_n}\| \leq M(1 + 2 \log 2) - M \log(1 - |\varphi_n(0)|)$$

and it follows that $\rho(C_\varphi) = 1$ for all bounded composition operators acting on BMOA by Theorem 3.7 of [Donaway 1999]. □

Thus BMOA satisfies all the properties of \mathcal{X} .

4B. Comparisons. We now investigate spaces that do not satisfy the properties of \mathcal{X} . We compare the spectrum of induced composition operators on these spaces with those on \mathcal{X} .

4B1. Hardy spaces. For $1 \leq p < \infty$, the Hardy space, denoted $H^p = H^p(\mathbb{D})$, is the space of analytic functions on \mathbb{D} such that

$$\|f\|_{H^p}^p = \sup_{0 < r < 1} \int_{\mathbb{D}} |f(re^{i\theta})|^p \frac{d\theta}{2\pi} < \infty.$$

Under this norm, the Hardy spaces are Banach spaces and for $p = 2$ it is a Hilbert space.

It is well known that H^p is a Banach space of analytic functions that contains H^∞ as a subspace. For the cases of an elliptic or parabolic automorphism φ , it is the case that $\rho(C_\varphi) = 1$ and the spectrum of C_φ on H^p is the same as for C_φ acting on \mathcal{X} (see Theorem 3.9 of [Cowen and MacCluer 1995]). However, it is not the case that the spectral radius is 1 for every composition operator induced by an automorphism. In fact, if φ is hyperbolic, then $\rho(C_\varphi) = \varphi'(a)^{-1/p}$ where a is the Denjoy–Wolff point of φ (see Theorem 3.9 of [Cowen and MacCluer 1995]). In this situation, $\varphi'(a) < 1$ thus making $\rho(C_\varphi) > 1$. In turn, the spectrum is the annulus $\varphi'(a)^{1/p} \leq |z| \leq \varphi'(a)^{-1/p}$ (see Theorem 4.9 of [Hyvärinen et al. 2013]).

4B2. Weighted Bergman spaces. For $1 \leq p < \infty$ and $\alpha > -1$, the standard weighted Bergman space, denoted $A_\alpha^p = A_\alpha^p(\mathbb{D})$, is the space of analytic functions on \mathbb{D} such that

$$\|f\|_{A_\alpha^p}^p = \int_{\mathbb{D}} (1 - |z|^2)^\alpha |f(z)|^p dA(z) < \infty,$$

where $dA(z)$ is the normalized Lebesgue area measure on \mathbb{D} . The weighted Bergman spaces are Banach spaces under the norm $\|\cdot\|_{A_\alpha^p}$.

It is well known that A_α^p is a Banach space of analytic functions that contains H^∞ as a subspace. For the cases of an elliptic or parabolic automorphism φ , it is the case that $\rho(C_\varphi) = 1$ and the spectrum of C_φ on A_α^p is the same as for C_φ acting on \mathcal{X} (see Lemma 4.2 and Theorem 4.14 of [Hyvärinen et al. 2013]). However, as was the case for the Hardy spaces, it is not the case that the spectral radius is 1 for every composition operator induced by an automorphism. In fact, if φ is hyperbolic, then

$$\rho(C_\varphi) = \max \left\{ \frac{1}{\varphi'(a)^s}, \frac{1}{\varphi'(b)^s} \right\}$$

where $s = (\alpha + 2)/p$, a is the Denjoy–Wolff point and b is the other fixed point of φ (see Theorem 4.6 of [Hyvärinen et al. 2013]). In turn, the spectrum contains the annulus

$$\min \left\{ \frac{1}{\varphi'(a)^s}, \frac{1}{\varphi'(b)^s} \right\} \leq |z| \leq \max \left\{ \frac{1}{\varphi'(a)^s}, \frac{1}{\varphi'(b)^s} \right\}$$

(see Corollary 4.7 of [Hyvärinen et al. 2013]).

4B3. Weighted Banach spaces. For $0 < p < \infty$, the standard weighted Banach space on \mathbb{D} , denoted $H_p^\infty = H_p^\infty(\mathbb{D})$, is the space of analytic functions on \mathbb{D} such that

$$\|f\|_{H_p^\infty} = \sup_{z \in \mathbb{D}} (1 - |z|^2)^p |f(z)| < \infty,$$

The weighted Banach spaces are, not surprisingly, Banach spaces under $\|\cdot\|_{H_p^\infty}$.

It is well known that H_p^∞ is a Banach space of analytic functions that contain H^∞ as a subspace. For the cases of an elliptic or parabolic automorphism φ , it is the case that $\rho(C_\varphi) = 1$ and the spectrum of C_φ on H_p^∞ is the same as for C_φ acting on \mathcal{X} (see Lemma 4.2 and Theorem 4.14 of [Hyvärinen et al. 2013]). However, as was the case for the Hardy spaces, it is not the case that the spectral radius is 1 for every composition operator induced by an automorphism. In fact, if φ is hyperbolic, then

$$\rho(C_\varphi) = \max\left\{\frac{1}{\varphi'(a)^s}, \frac{1}{\varphi'(b)^s}\right\}$$

where a is the Denjoy–Wolff point and b is the other fixed point of φ . In turn, the spectrum contains the annulus

$$\min\left\{\frac{1}{\varphi'(a)^p}, \frac{1}{\varphi'(b)^p}\right\} \leq |z| \leq \max\left\{\frac{1}{\varphi'(a)^p}, \frac{1}{\varphi'(b)^p}\right\}$$

(see Theorem 4.6 and Corollary 4.7 of [Hyvärinen et al. 2013]).

4B4. Dirichlet space. The Dirichlet space on \mathbb{D} , denoted \mathcal{D} , is the space of analytic functions on \mathbb{D} such that

$$\int_{\mathbb{D}} |f'(z)|^2 dA(z) < \infty$$

where dA denotes the normalized Lebesgue area measure on \mathbb{D} . Under the norm

$$\|f\|_{\mathcal{D}}^2 = |f(0)|^2 + \int_{\mathbb{D}} |f'(z)|^2 dA(z)$$

the Dirichlet space has a Hilbert space structure. Although not every analytic self-map of \mathbb{D} induce bounded composition operators on \mathcal{D} , univalent maps, and thus the automorphisms, of \mathbb{D} do.

Independently, Donaway [1999, Corollary 3.11] and Martín and Vukotić [2005, Theorem 7] showed that composition operators on \mathcal{D} induced by univalent self-maps of \mathbb{D} , and thus the automorphisms, have spectral radius 1. However, by direct calculation, one can see that the functions in \mathcal{F} are not contained in the Dirichlet space; for the case of f_s this is shown in [Pons 2010] (see p. 455). Despite \mathcal{D} not satisfying all the properties of \mathcal{X} , the spectrum of automorphism induced composition operators on \mathcal{D} are precisely the same as those on \mathcal{X} .

To overcome the lack of eigenfunctions, the authors in [Higdon 1997] and [Gallardo-Gutiérrez and Montes-Rodríguez 2003] used two new approaches. In [Higdon 1997], the author produces approximate eigenfunctions and in [Gallardo-Gutiérrez and Montes-Rodríguez 2003] unitary similarity is the key tool.

Remark 4.2. For all of the spaces discussed in Sections 4A and 4B (and those discussed in the next section), the spectrum of C_φ when φ is elliptic will be the

same as that for C_φ acting on \mathcal{X} . This is due to the fact that the eigenfunctions are the monomials, which are contained in all of these spaces.

5. Open questions

We end this paper with open questions which were inspired while developing the examples and comparisons in Sections 4A and 4B.

5A. The little Bloch space. While the Bloch space contains the polynomials, they are not dense in \mathcal{B} . The closure of the polynomials with respect to $\|\cdot\|_{\mathcal{B}}$ is called the little Bloch space, denoted $\mathcal{B}_0 = \mathcal{B}_0(\mathbb{D})$. More formally, the little Bloch space consists of the functions $f \in \mathcal{B}$ such that

$$\lim_{|z| \rightarrow 1} (1 - |z|^2) |f'(z)| = 0.$$

From Theorem 12 of [Arazy et al. 1985], bounded composition operators on \mathcal{B}_0 are induced exactly by functions in \mathcal{B}_0 , which include the automorphisms. Donaway also proved the spectral radius of every bounded composition operator on \mathcal{B}_0 is 1. Thus property (ii) is satisfied by \mathcal{B}_0 . However, the following result shows that \mathcal{F} is not contained in \mathcal{B}_0 , and thus property (i) of \mathcal{X} is not satisfied.

Theorem 5.1. *The functions f_s and f_t , for $s > 0$ and $t \neq 0$, are not contained in the little Bloch space.*

Proof. Consider the function

$$f_t(z) = \exp\left(it \log \frac{1+z}{1-z}\right).$$

We show that this function is not in \mathcal{B}_0 for $t \in \mathbb{R} \setminus \{0\}$. Taking the derivative,

$$f'_t(z) = f_t(z) \left(it \frac{1-z}{1+z}\right) \frac{2}{(1-z)^2} = f_t(z) \frac{2it}{(1-z)(1+z)}.$$

For $t > 0$, $|f_t(z)| \geq e^{-2\pi}$ and, for $t < 0$, $|f_t(z)| \geq 1$. In either case, there is a constant $C > 0$ such that $|f_t(z)| \geq C$ for all $t \in \mathbb{R} \setminus \{0\}$ and all $z \in \mathbb{D}$. Hence

$$|f'_t(z)| \geq \frac{2C|t|}{|z-1||z+1|}.$$

To show that $f_t \notin \mathcal{B}_0$, we need to show that

$$\lim_{|z| \rightarrow 1} (1 - |z|^2) |f'_t(z)| \neq 0.$$

To see this, first observe that

$$\lim_{|z| \rightarrow 1} (1 - |z|^2) |f'_t(z)| \geq \lim_{|z| \rightarrow 1} (1 - |z|^2) \frac{2C|t|}{|z-1||z+1|}$$

by our estimate from above. If we now take a radial path to 1, that is, we set $z = r$ and let $r \uparrow 1$, we have

$$\lim_{r \rightarrow 1^-} (1 - r^2) \frac{2C|t|}{(1-r)(1+r)} = 2C|t| > 0$$

when $t \neq 0$. Thus

$$\lim_{|z| \rightarrow 1} (1 - |z|^2) \frac{2Ct}{|z-1||z+1|} \neq 0$$

for $t \neq 0$, and hence f_t is not in \mathcal{B}_0 .

Next consider the function

$$f_s(z) = \exp \frac{s(z+1)}{z-1}.$$

We will show that this function is not in \mathcal{B}_0 for $s > 0$. First observe that

$$f'_s(z) = \left(\exp \frac{s(z+1)}{z-1} \right) \frac{-2s}{(z-1)^2}$$

and thus we aim to show that

$$\lim_{|z| \rightarrow 1} (1 - |z|^2) |f'_s(z)| = \lim_{|z| \rightarrow 1} (1 - |z|^2) \left| \exp \left(\frac{s(z+1)}{z-1} \right) \right| \frac{2s}{|1-z|^2} \neq 0.$$

Fix $x_0 < 0$ and consider the sequence $\{z_n\}$ defined by

$$z_n = \frac{x_0 + in + 1}{x_0 + in - 1}.$$

Since $x_0 < 0$, this sequence is contained in the unit disk and $\{z_n\} \rightarrow 1$ as $n \rightarrow \infty$.

To obtain our conclusion, we show

$$\lim_{n \rightarrow \infty} (1 - |z_n|^2) \left| \exp \frac{s(z_n+1)}{z_n-1} \right| \frac{2s}{|1-z_n|^2} \neq 0.$$

First observe that the map $\psi(z) = (z+1)/(z-1)$ is its own inverse and hence $\psi(z_n) = x_0 + in$ for each $n \in \mathbb{N}$. Thus

$$\left| \exp \frac{s(z_n+1)}{z_n-1} \right| = \left| \exp(sx_0 + isn) \right| = e^{sx_0} > 0.$$

Substituting,

$$\lim_{n \rightarrow \infty} (1 - |z_n|^2) \left| \exp \frac{s(z_n+1)}{z_n-1} \right| \frac{2s}{|1-z_n|^2} = \lim_{n \rightarrow \infty} e^{sx_0} (1 - |z_n|^2) \frac{2s}{|1-z_n|^2}.$$

Next,

$$1 - |z_n|^2 = \frac{-4x_0}{(x_0 - 1)^2 + n^2}$$

and

$$|1 - z_n|^2 = \frac{4}{(x_0 - 1)^2 + n^2}.$$

Thus

$$\begin{aligned} \lim_{n \rightarrow \infty} e^{sx_0} (1 - |z_n|^2) \frac{2s}{|1 - z_n|^2} &= \lim_{n \rightarrow \infty} e^{sx_0} \left(\frac{-4x_0}{(x_0 - 1)^2 + n^2} \right) \left(\frac{s((x_0 - 1)^2 + n^2)}{2} \right) \\ &= \lim_{n \rightarrow \infty} (-2sx_0) e^{sx_0} > 0 \end{aligned}$$

and hence f_s is not in \mathcal{B}_0 for $s > 0$. \square

For the little Bloch space, we leave the reader with the following question.

Question 1. For φ a parabolic or hyperbolic automorphism, what is the spectrum of C_φ on the little Bloch space?

5B. Analytic functions of vanishing mean oscillation. Like the Bloch space, the polynomials are contained in BMOA, but they are not dense in BMOA. We denote by VMOA the closure of the polynomials in $\|\cdot\|_{\text{BMOA}}$. VMOA is the space of analytic functions with vanishing mean oscillation on $\partial\mathbb{D}$, formally defined as the functions $f \in \text{BMOA}$ such that

$$\lim_{|a| \rightarrow 1} \|f \circ \tau_a - f(a)\|_{H^2} = 0.$$

By Corollary 4.2 of [Laitila 2009], C_φ is bounded on VMOA if and only if $\varphi \in \text{VMOA}$. So every automorphism induces a bounded composition operator on VMOA. By the same argument as in Section 4A3, the spectral radius of C_φ induced by a disk automorphism is 1. Thus property (ii) of \mathcal{X} is satisfied. Since VMOA is a subspace of the little Bloch space (see [Gallardo-Gutiérrez et al. 2013]), it follows that VMOA does not satisfy property (i), a corollary of Theorem 5.1.

Corollary 5.2. *The functions f_s and f_t , for $s > 0$ and $t \neq 0$, are not contained in VMOA.*

For VMOA, we leave the reader with the following question.

Question 2. For φ parabolic or hyperbolic automorphism, what is the spectrum of C_φ on VMOA?

Acknowledgements

The authors would like to thank the referees for their careful review of the manuscript and their many suggestions, especially to generalize the results to a more general family of Banach spaces. The work of the second author was conducted while an undergraduate student at the University of Wisconsin-La Crosse and funded by the College of Science and Health Dean's Distinguished Fellowship.

References

- [Arazy et al. 1985] J. Arazy, S. D. Fisher, and J. Peetre, “Möbius invariant function spaces”, *J. Reine Angew. Math.* **363** (1985), 110–145. MR 814017 Zbl 0566.30042
- [Cowen and MacCluer 1995] C. C. Cowen and B. D. MacCluer, *Composition operators on spaces of analytic functions*, Studies in Advanced Mathematics **20**, CRC Press, Boca Raton, FL, 1995. MR 1397026 Zbl 0873.47017
- [Donaway 1999] R. L. Donaway, *Norm and essential norm estimates of composition operators on Besov-type spaces*, ProQuest LLC, Ann Arbor, MI, 1999. Thesis (Ph.D.)—University of Virginia. MR 2698630
- [Duren 1970] P. L. Duren, *Theory of H^p spaces*, Pure and Applied Mathematics **Vol. 38**, Academic Press, New York, 1970. MR 0268655 Zbl 0215.20203
- [Gallardo-Gutiérrez and Montes-Rodríguez 2003] E. A. Gallardo-Gutiérrez and A. Montes-Rodríguez, “Adjoints of linear fractional composition operators on the Dirichlet space”, *Math. Ann.* **327**:1 (2003), 117–134. MR 2005124 Zbl 1048.47016
- [Gallardo-Gutiérrez et al. 2013] E. A. Gallardo-Gutiérrez, M. J. González, F. Pérez-González, C. Pommerenke, and J. Rättyä, “Locally univalent functions, VMOA and the Dirichlet space”, *Proc. Lond. Math. Soc.* (3) **106**:3 (2013), 565–588. MR 3048550 Zbl 1276.30026
- [Higdon 1997] W. M. Higdon, *Composition operators of the Dirichlet space*, Ph.D. thesis, Michigan State University, 1997.
- [Hyvärinen et al. 2013] O. Hyvärinen, M. Lindström, I. Nieminen, and E. Saukko, “Spectra of weighted composition operators with automorphic symbols”, *J. Funct. Anal.* **265**:8 (2013), 1749–1777. MR 3079234 Zbl 1325.47054
- [Laitila 2009] J. Laitila, “Weighted composition operators on BMOA”, *Comput. Methods Funct. Theory* **9**:1 (2009), 27–46. MR 2478261 Zbl 1163.47018
- [MacCluer 2009] B. D. MacCluer, *Elementary functional analysis*, Graduate Texts in Mathematics **253**, Springer, New York, 2009. MR 2462971 Zbl 1170.46002
- [Martín and Vukotić 2005] M. J. Martín and D. Vukotić, “Norms and spectral radii of composition operators acting on the Dirichlet space”, *J. Math. Anal. Appl.* **304**:1 (2005), 22–32. MR 2124646 Zbl 1071.47028
- [Nordgren 1968] E. A. Nordgren, “Composition operators”, *Canad. J. Math.* **20** (1968), 442–449. MR 0223914 Zbl 0161.34703
- [Pons 2007] M. A. Pons, *Composition Operators on Besov and Dirichlet Type Spaces*, Ph.D. thesis, University of Virginia, 2007.
- [Pons 2010] M. A. Pons, “The spectrum of a composition operator and Calderón’s complex interpolation”, pp. 451–467 in *Topics in operator theory. Volume I. Operators, matrices and analytic functions*, Oper. Theory Adv. Appl. **202**, Birkhäuser, Basel, 2010. MR 2723292 Zbl 1217.47050

Received: 2015-06-24

Revised: 2015-08-26

Accepted: 2015-09-07

rallen@uwf.edu

Department of Mathematics and Statistics, University
of Wisconsin-La Crosse, La Crosse, WI 54601, United States

thmle@ucdavis.edu

Department of Computer Science,
University of California, Davis, Davis, CA 95616, United States

mapons@noctrl.edu

Department of Mathematics, North Central College,
Naperville, IL 60540, United States

involve

msp.org/involve

INVOLVE YOUR STUDENTS IN RESEARCH

Involve showcases and encourages high-quality mathematical research involving students from all academic levels. The editorial board consists of mathematical scientists committed to nurturing student participation in research. Bridging the gap between the extremes of purely undergraduate research journals and mainstream research journals, *Involve* provides a venue to mathematicians wishing to encourage the creative involvement of students.

MANAGING EDITOR

Kenneth S. Berenhaut Wake Forest University, USA

BOARD OF EDITORS

Colin Adams	Williams College, USA	Suzanne Lenhart	University of Tennessee, USA
John V. Baxley	Wake Forest University, NC, USA	Chi-Kwong Li	College of William and Mary, USA
Arthur T. Benjamin	Harvey Mudd College, USA	Robert B. Lund	Clemson University, USA
Martin Bohner	Missouri U of Science and Technology, USA	Gaven J. Martin	Massey University, New Zealand
Nigel Boston	University of Wisconsin, USA	Mary Meyer	Colorado State University, USA
Amarjit S. Budhiraja	U of North Carolina, Chapel Hill, USA	Emil Minchev	Ruse, Bulgaria
Pietro Cerone	La Trobe University, Australia	Frank Morgan	Williams College, USA
Scott Chapman	Sam Houston State University, USA	Mohammad Sal Moslehian	Ferdowsi University of Mashhad, Iran
Joshua N. Cooper	University of South Carolina, USA	Zuhair Nashed	University of Central Florida, USA
Jem N. Corcoran	University of Colorado, USA	Ken Ono	Emory University, USA
Toka Diagana	Howard University, USA	Timothy E. O'Brien	Loyola University Chicago, USA
Michael Dorff	Brigham Young University, USA	Joseph O'Rourke	Smith College, USA
Sever S. Dragomir	Victoria University, Australia	Yuval Peres	Microsoft Research, USA
Behrouz Emamizadeh	The Petroleum Institute, UAE	Y.-F. S. Pétermann	Université de Genève, Switzerland
Joel Foisy	SUNY Potsdam, USA	Robert J. Plemmons	Wake Forest University, USA
Errin W. Fulp	Wake Forest University, USA	Carl B. Pomerance	Dartmouth College, USA
Joseph Gallian	University of Minnesota Duluth, USA	Vadim Ponomarenko	San Diego State University, USA
Stephan R. Garcia	Pomona College, USA	Bjorn Poonen	UC Berkeley, USA
Anant Godbole	East Tennessee State University, USA	James Propp	U Mass Lowell, USA
Ron Gould	Emory University, USA	József H. Przytycki	George Washington University, USA
Andrew Granville	Université Montréal, Canada	Richard Rebarber	University of Nebraska, USA
Jerrold Griggs	University of South Carolina, USA	Robert W. Robinson	University of Georgia, USA
Sat Gupta	U of North Carolina, Greensboro, USA	Filip Saidak	U of North Carolina, Greensboro, USA
Jim Haglund	University of Pennsylvania, USA	James A. Sellers	Penn State University, USA
Johnny Henderson	Baylor University, USA	Andrew J. Sterge	Honorary Editor
Jim Hoste	Pitzer College, USA	Ann Trenk	Wellesley College, USA
Natalia Hritonenko	Prairie View A&M University, USA	Ravi Vakil	Stanford University, USA
Glenn H. Hurlbert	Arizona State University, USA	Antonia Vecchio	Consiglio Nazionale delle Ricerche, Italy
Charles R. Johnson	College of William and Mary, USA	Ram U. Verma	University of Toledo, USA
K. B. Kulasekera	Clemson University, USA	John C. Wierman	Johns Hopkins University, USA
Gerry Ladas	University of Rhode Island, USA	Michael E. Zieve	University of Michigan, USA

PRODUCTION

Silvio Levy, Scientific Editor


Cover: Alex Scorpan

See inside back cover or msp.org/involve for submission instructions. The subscription price for 2016 is US \$160/year for the electronic version, and \$215/year (+\$35, if shipping outside the US) for print and electronic. Subscriptions, requests for back issues from the last three years and changes of subscribers address should be sent to MSP.

Involve (ISSN 1944-4184 electronic, 1944-4176 printed) at Mathematical Sciences Publishers, 798 Evans Hall #3840, c/o University of California, Berkeley, CA 94720-3840, is published continuously online. Periodical rate postage paid at Berkeley, CA 94704, and additional mailing offices.

Involve peer review and production are managed by EditFLOW[®] from Mathematical Sciences Publishers.

PUBLISHED BY

 **mathematical sciences publishers**
nonprofit scientific publishing

<http://msp.org/>

© 2016 Mathematical Sciences Publishers

involve

2016

vol. 9

no. 5

An iterative strategy for Lights Out on Petersen graphs BRUCE TORRENCE AND ROBERT TORRENCE	721
A family of elliptic curves of rank ≥ 4 FARZALI IZADI AND KAMRAN NABARDI	733
Splitting techniques and Betti numbers of secant powers REZA AKHTAR, BRITTANY BURNS, HALEY DOHRMANN, HANNAH HOGANSON, OLA SOBIESKA AND ZEROTTI WOODS	737
Convergence of sequences of polygons ERIC HINTIKKA AND XINGPING SUN	751
On the Chermak–Delgado lattices of split metacyclic p -groups ERIN BRUSH, JILL DIETZ, KENDRA JOHNSON-TESCH AND BRIANNE POWER	765
The left greedy Lie algebra basis and star graphs BENJAMIN WALTER AND AMINREZA SHIRI	783
Note on superpatterns DANIEL GRAY AND HUA WANG	797
Lifting representations of finite reductive groups: a character relation JEFFREY D. ADLER, MICHAEL CASSEL, JOSHUA M. LANSKY, EMMA MORGAN AND YIFEI ZHAO	805
Spectrum of a composition operator with automorphic symbol ROBERT F. ALLEN, THONG M. LE AND MATTHEW A. PONS	813
On nonabelian representations of twist knots JAMES C. DEAN AND ANH T. TRAN	831
Envelope curves and equidistant sets MARK HUIBREGTSE AND ADAM WINCHELL	839
New examples of Brunnian theta graphs BYOUNGWOOK JANG, ANNA KRONAEUR, PRATAP LUITEL, DANIEL MEDICI, SCOTT A. TAYLOR AND ALEXANDER ZUPAN	857
Some nonsimple modules for centralizer algebras of the symmetric group CRAIG DODGE, HARALD ELLERS, YUKIHIDE NAKADA AND KELLY POHLAND	877
Acknowledgement	899



1944-4176(2016)9:5;1-0