

# involve

a journal of mathematics

Reeb dynamics of the link of the  $A_n$  singularity

Leonardo Abbrescia, I. Huq-Kuruvilla, J. Nelson and N. Sultani



# Reeb dynamics of the link of the $A_n$ singularity

Leonardo Abbrescia, I. Huq-Kuruville, J. Nelson and N. Sultani

(Communicated by Colin Adams)

The link of the  $A_n$  singularity,  $L_{A_n} \subset \mathbb{C}^3$  admits a natural contact structure  $\xi_0$  coming from the set of complex tangencies. The canonical contact form  $\alpha_0$  associated to  $\xi_0$  is degenerate and thus has no isolated Reeb orbits. We show that there is a nondegenerate contact form for a contact structure equivalent to  $\xi_0$  that has two isolated simple periodic Reeb orbits. We compute the Conley–Zehnder index of these simple orbits and their iterates. From these calculations we compute the positive  $S^1$ -equivariant symplectic homology groups for  $(L_{A_n}, \xi_0)$ . In addition, we prove that  $(L_{A_n}, \xi_0)$  is contactomorphic to the lens space  $L(n+1, n)$ , equipped with its canonical contact structure  $\xi_{\text{std}}$ .

## 1. Introduction and main results

The classical topological theory of isolated critical points of complex polynomials relates the topology of the link of the singularity to the algebraic properties of the singularity [Milnor 1968]. More generally, the link of an irreducible affine variety  $A^n \subset \mathbb{C}^N$  with an isolated singularity at  $\mathbb{0}$  is defined by  $L_A = A \cap S_\delta^{2N+1}$ . For sufficiently small  $\delta$ , the link  $L_A$  is a manifold of real dimension  $2n - 1$ , which is an invariant of the germ of  $A$  at  $\mathbb{0}$ . The links of Brieskorn varieties can sometimes be homeomorphic but not always diffeomorphic to spheres (see [Brieskorn 1966], a preliminary result which further motivated the study of such objects). Recent developments in symplectic and contact geometry have shown that the algebraic properties of a singularity are strongly connected to the contact topology of the link and symplectic topology of (the resolution of) the variety. A wide range of results demonstrating the power of investigating the symplectic and contact perspective of singularities include [Keating 2015; Kwon and van Koert 2016; McLean 2016; Ritter 2010; Seidel 2008b; Ustilovsky 1999].

In this paper we study the contact topology of the link of the  $A_n$  singularity, providing a computation of positive  $S^1$ -equivariant symplectic homology. This

---

*MSC2010:* primary 37B30, 53D35, 57R17; secondary 53D42.

*Keywords:* contact geometry, contact topology, Conley–Zehnder index,  $A_n$  singularity, Reeb dynamic, Maslov index.

is done via our construction of an explicit nondegenerate contact form and the computation of the Conley–Zehnder indices of the associated simple Reeb orbits and their iterates. Our computations show that positive  $S^1$ -equivariant symplectic homology is a free  $\mathbb{Q}[u]$  module of rank equal to the number of conjugacy classes of the finite subgroup  $A_n$  of  $\mathrm{SL}(2; \mathbb{C})$ . This provides a concrete example of the relationship between the cohomological McKay correspondence and symplectic homology, which is work in progress by McLean and Ritter [ $\geq 2017$ ]. As a result, the topological nature of the singularity is reflected by qualitative aspects of the Reeb dynamics associated to the link of the  $A_n$  singularity.

The link of the  $A_n$  singularity is defined by

$$L_{A_n} = f_{A_n}^{-1}(0) \cap S^5 \subset \mathbb{C}^3, \quad f_{A_n} = z_0^{n+1} + 2z_1z_2. \quad (1-1)$$

It admits a natural contact structure coming from the set of complex tangencies,

$$\xi_0 := TL_{A_n} \cap J_0(TL_{A_n}).$$

The contact structure can be expressed as the kernel of the canonically defined contact form,

$$\alpha_0 = \frac{i}{2} \left( \sum_{j=0}^m (z_j d\bar{z}_j - \bar{z}_j dz_j) \right) \Big|_{L_{A_n}}.$$

The contact form  $\alpha_0$  is degenerate and hence not appropriate for computing Floer-theoretic invariants as the periodic orbits of the Reeb vector field defined by

$$\alpha_0(R_{\alpha_0}) = 1, \quad \iota_{R_{\alpha_0}} d\alpha_0 = 0$$

are not isolated.

Our first result is the construction of a nondegenerate contact form  $\alpha_\epsilon$  such that  $(L_{A_n}, \ker \alpha_0)$  and  $(L_{A_n}, \ker \alpha_\epsilon)$  are contactomorphic. Define the Hamiltonian on  $\mathbb{C}^3$  by

$$H : \mathbb{C}^3 \rightarrow \mathbb{R}, \\ (z_0, z_1, z_2) \mapsto |z|^2 + \epsilon(|z_1|^2 - |z_2|^2),$$

where  $\epsilon$  is chosen so that  $H > 0$  on  $S^5$ . We will show

$$\alpha_\epsilon = \frac{1}{H} \left[ \frac{(n+1)i}{8} (z_0 d\bar{z}_0 - \bar{z}_0 dz_0) + \frac{i}{4} (z_1 d\bar{z}_1 - \bar{z}_1 dz_1 + z_2 d\bar{z}_2 - \bar{z}_2 dz_2) \right] \quad (1-2)$$

is a nondegenerate contact form. We also find the simple Reeb orbits of  $R_{\alpha_\epsilon}$  and compute the associated Conley–Zehnder index with respect to the canonical trivialization of  $\mathbb{C}^3$  of their iterates.

**Theorem 1.1.** *The 1-form  $\alpha_\epsilon$  is a nondegenerate contact form for  $L_{A_n}$  such that  $(L_{A_n}, \ker \alpha_0)$  and  $(L_{A_n}, \ker \alpha_\epsilon)$  are contactomorphic. The Reeb orbits of  $R_{\alpha_\epsilon}$  are*

defined by

$$\begin{aligned} \gamma_+(t) &= (0, e^{2i(1+\epsilon)t}, 0), & 0 \leq t \leq \pi/(1+\epsilon), \\ \gamma_-(t) &= (0, 0, e^{2i(1-\epsilon)t}), & 0 \leq t \leq \pi/(1-\epsilon). \end{aligned}$$

The Conley–Zehnder index for  $\gamma = \gamma_{\pm}^N$  in  $0 \leq t \leq N\pi/(1 \pm \epsilon)$  is

$$\mu_{CZ}(\gamma_{\pm}^N) = 2 \left( \left\lfloor \frac{2N}{(n+1)(1 \pm \epsilon)} \right\rfloor + \left\lfloor \frac{N(1 \mp \epsilon)}{1 \pm \epsilon} \right\rfloor - \left\lfloor \frac{2N}{1 \pm \epsilon} \right\rfloor \right) + 2N + 1. \quad (1-3)$$

**Remark 1.2.** If  $\epsilon$  is chosen such that  $0 < \epsilon \ll 1/N$  then (1-3) can be simplified to

$$\begin{aligned} \mu_{CZ}(\gamma_-^N) &= 2 \left\lfloor \frac{2N}{(n+1)(1-\epsilon)} \right\rfloor + 1, \\ \mu_{CZ}(\gamma_+^N) &= 2 \left\lfloor \frac{2N}{(n+1)(1+\epsilon)} \right\rfloor + 1. \end{aligned} \quad (1-4)$$

The proof of Theorem 1.1 is obtained by adapting methods of Ustilovsky [1999] to obtain  $\alpha_\epsilon$  and to compute the Conley–Zehnder indices. The Conley–Zehnder index is a Maslov index for arcs of symplectic matrices and is defined in Section 2D. These paths of matrices are obtained by linearizing the flow of the Reeb vector field along the Reeb orbit and restricting to  $\xi_0$ . To better understand the spread of the Reeb orbits and their iterates in various indices, we have the following example.

**Example 1.3.** Let  $n = 2$  and  $0 < \epsilon \ll \frac{1}{10}$ . Then

$$\begin{aligned} \mu_{CZ}(\gamma_-) &= 1, & \mu_{CZ}(\gamma_+) &= 1, \\ \mu_{CZ}(\gamma_-^2) &= 3, & \mu_{CZ}(\gamma_+^2) &= 3, \\ \mu_{CZ}(\gamma_-^3) &= 5, & \mu_{CZ}(\gamma_+^3) &= 3, \\ \mu_{CZ}(\gamma_-^4) &= 5, & \mu_{CZ}(\gamma_+^4) &= 5, \\ \mu_{CZ}(\gamma_-^5) &= 7, & \mu_{CZ}(\gamma_+^5) &= 7, \\ \mu_{CZ}(\gamma_-^6) &= 9, & \mu_{CZ}(\gamma_+^6) &= 7, \\ \mu_{CZ}(\gamma_-^7) &= 9, & \mu_{CZ}(\gamma_+^7) &= 9. \end{aligned}$$

It is interesting to note that the spread of integers is not uniform between  $\mu_{CZ}(\gamma_-^N)$  and  $\mu_{CZ}(\gamma_+^N)$ , and where these jumps in index occur. However, we see that there are  $n = 2$  Reeb orbits with Conley–Zehnder index 1 and  $n + 1 = 3$  orbits with Conley–Zehnder index  $2k + 1$  for each  $k \geq 1$ .

**Remark 1.4.** Extrapolating this to all values of  $n$  and  $N$  demonstrates that the numerology of the Conley–Zehnder index realizes the number of free homotopy classes of  $L_{A_n}$ . Recall  $[\Sigma L_{A_n}] = \pi_0(\Sigma L_{A_n}) = \pi_1(L_{A_n})/\{\text{conjugacy classes}\}$  and  $H_1(L_{A_n}, \mathbb{Z}) = \mathbb{Z}_{n+1}$ . The information that the  $(n + 1)$ -th iterate of  $\gamma_{\pm}$  is the first

contractible Reeb orbit is also encoded in the above formulas. Qualitative aspects of the Reeb dynamics reflect this topological information in the following computation of a Floer-theoretic invariant of the contact structure  $\xi_0$ .

**Theorem 1.1** allows us to easily compute positive  $S^1$ -equivariant symplectic homology  $SH_*^{+,S^1}$ . Symplectic homology is a Floer-type invariant of symplectic manifolds with contact-type boundary; see [Seidel 2008a]. Under additional assumptions, one can prove that the positive  $S^1$ -equivariant symplectic homology  $SH_*^{+,S^1}$  is in fact an invariant of the contact structure; see [Gutt 2015, Theorems 1.2 and 1.3; Bourgeois and Oancea 2012, Section 4.1.2]. Because of the behavior of the Conley–Zehnder index in **Theorem 1.1**, we can directly compute  $SH_*^{+,S^1}(L_{A_n}, \xi_0)$  and conclude that it is a contact invariant. As a result, the underlying topology of the manifold determines qualitative aspects of any Reeb vector field associated to a contact form defining  $\xi_0$ .

**Theorem 1.5.** *The positive  $S^1$ -equivariant symplectic homology of  $(L_{A_n}, \xi_0)$  is*

$$SH_*^{+,S^1}(L_{A_n}, \xi_0) = \begin{cases} \mathbb{Q}^n, & * = 1, \\ \mathbb{Q}^{n+1}, & * \geq 3 \text{ and odd}, \\ 0, & * \text{ else.} \end{cases}$$

*Proof.* To obtain a contact invariant from  $SH_*^{+,S^1}$  we need to show in dimension 3 that all contractible Reeb orbits  $\gamma$  satisfy  $\mu_{CZ}(\gamma) \geq 3$ ; see [Gutt 2015, Theorems 1.2 and 1.3; Bourgeois and Oancea 2012, Section 4.1.2]. The first iterate of  $\gamma_{\pm}$  which is contractible is the  $(n+1)$ -th iterate, and by **Theorem 1.1**, will always satisfy  $\mu_{CZ}(\gamma_{\pm}) \geq 3$ .

If  $\alpha$  is a nondegenerate contact form such that the Conley–Zehnder indices of all periodic Reeb orbits are lacunary, meaning they contain no two consecutive numbers, then we can appeal to [Gutt 2015, Theorem 1.1]. This result of Gutt allows us to conclude that over  $\mathbb{Q}$ -coefficients the differential for  $SH^{S^1,+}$  vanishes. In light of **Theorem 1.1** we obtain the above result.  $\square$

**Remark 1.4** yields the following corollary of **Theorem 1.5**, indicating a Floer-theoretic interpretation of the McKay correspondence [Ito and Reid 1996] via the Reeb dynamics of the link of the  $A_n$  singularity. The  $A_n$  singularity is the singularity of  $f_{A_n}^{-1}(0)$ , where  $f_{A_n}$  is described as (1-1). This is equivalent to its characterization as the absolutely isolated double point quotient singularity of  $\mathbb{C}^2/A_n$ , where  $A_n$  is the cyclic subgroup of  $\mathrm{SL}(2; \mathbb{C})$ ; see **Section 4A**. The cyclic group  $A_n$  acts on  $\mathbb{C}^2$  by  $(u, v) \mapsto (e^{2\pi i/(n+1)}u, e^{2\pi in/(n+1)}v)$ .

**Corollary 1.6.** *The positive  $S^1$ -equivariant symplectic homology  $SH_*^{+,S^1}(L_{A_n}, \xi_0)$  is a free  $\mathbb{Q}[u]$  module of rank equal to the number of conjugacy classes of the finite subgroup  $A_n$  of  $\mathrm{SL}(2; \mathbb{C})$ .*

**Remark 1.7.** The ongoing work of Nelson [2015; ≥ 2017] and Hutchings and Nelson [2014; ≥ 2017] is needed in order to work under the assumption that a related Floer-theoretic invariant, cylindrical contact homology is a well-defined contact invariant of  $(L_{A_n}, \xi_0)$ . Once this is complete, the index calculations provided in Theorem 1.1 show that positive  $S^1$ -equivariant symplectic homology and cylindrical contact homology agree up to a degree shift.

Bourgeois and Oancea [2012] prove that there are restricted classes of contact manifolds for which one can prove that cylindrical contact homology (with a degree shift) is isomorphic to the positive part of  $S^1$ -equivariant symplectic homology when both are defined over  $\mathbb{Q}$ -coefficients. Their isomorphism relies on having transversality for a generic choice of  $J$ , which is presently the case for unit cotangent bundles  $DT^*L$  such that  $\dim L \geq 5$  or when  $L$  is Riemannian manifold which admits no contractible closed geodesics [Bourgeois and Oancea 2015]. Our computations confirm that their results should hold for many more closed contact manifolds.

Our final result is an explicit proof that the singularity  $(L_{A_n}, \xi_0)$  and the lens space  $(L(n + 1, n), \xi_{\text{std}})$  are contactomorphic. The lens space

$$L(n + 1, n) = S^3 / ((u, v) \sim (e^{2\pi i/(n+1)}u, e^{2\pi ni/(n+1)}v))$$

admits a contact structure, which is induced by the one on  $S^3$  and can be expressed as the kernel of the contact form

$$\lambda_{\text{std}} = \frac{1}{2}i(ud\bar{u} - \bar{u}du + vd\bar{v} - \bar{v}dv).$$

**Theorem 1.8.** *The link of the  $A_n$  singularity  $(L_{A_n}, \xi_0 = \ker \alpha_0)$  and the lens space  $(L(n + 1, n), \xi_{\text{std}} = \ker \lambda_{\text{std}})$  are contactomorphic.*

Theorems 1.5 and 1.8 allow us to reprove the following result of Kwon and van Koert [2016]. Since  $(L_{A_n}, \xi_0)$  and  $(L(n + 1, n), \xi_{\text{std}})$  are contactomorphic and  $SH_*^{S^1,+}$  is a contact invariant,  $SH_*^{S^1,+}(L(n + 1, n), \xi_{\text{std}}) = SH_*^{S^1,+}(L_{A_n}, \xi_0)$ .

**Theorem 1.9** [Kwon and van Koert 2016, Appendix A]. *The positive  $S^1$ -equivariant symplectic homology of  $(L(n + 1, n), \xi_{\text{std}})$  is*

$$SH_*^{+,S^1}(L(n + 1, n), \xi_{\text{std}}) = \begin{cases} \mathbb{Q}^n, & * = 1, \\ \mathbb{Q}^{n+1}, & * \geq 3 \text{ and odd,} \\ 0, & * \text{ else.} \end{cases}$$

Their proof relies on the nondegenerate contact form on  $(L(n + 1, n), \xi_{\text{std}})$ . If  $a_1, a_2$  are any rationally independent positive real numbers then

$$\lambda_{a_1, a_2} = \frac{i}{2} \sum_{j=1}^2 a_j(z_j d\bar{z}_j - \bar{z}_j dz_j)$$

is a nondegenerate contact form for  $(L(n+1, n), \xi_{\text{std}})$ . The simple Reeb orbits on  $L(n+1, n)$  are given by

$$\begin{aligned}\gamma_1 &= (e^{it/a_1}, 0), & 0 \leq t \leq (2a_1\pi)/(n+1), \\ \gamma_2 &= (0, e^{it/a_2}), & 0 \leq t \leq (2a_2\pi)/(n+1),\end{aligned}$$

which descend from the simple isolated Reeb orbits on  $S^3$ . Again, the  $n+1$  different free homotopy classes associated to this lens space are realized by covers of the isolated Reeb orbits  $\gamma_i$  for  $i = 1$  or  $2$ . The Conley–Zehnder index for  $\gamma_1^N$  is

$$\mu_{CZ}(\gamma_1^N) = 2 \left( \left\lfloor \frac{N}{n+1} \right\rfloor + \left\lfloor \frac{Na_1}{(n+1)a_2} \right\rfloor \right) + 1, \quad (1-5)$$

with a similar formula holding for  $\gamma_2^N$ .

*Outline.* The necessary background is given in [Section 2](#). The construction of a nondegenerate contact form and the proof of [Theorem 1.1](#) is given in [Section 3](#). The proof of [Theorem 1.8](#) is given in [Section 4](#).

## 2. Background

In this section we recall all the necessary symplectic and contact background which is needed to prove [Theorems 1.1](#) and [1.8](#).

**2A. Contact structures.** First we recall some notions from contact geometry.

**Definition 2.1.** Let  $M$  be a manifold of dimension  $2n+1$ . A *contact structure* is a maximally nonintegrable hyperplane field  $\xi = \ker \alpha \subset TM$ .

**Remark 2.2.** The kernel of a 1-form  $\alpha$  on  $M^{2n+1}$ ,  $\xi = \ker \alpha$ , is a contact structure whenever

$$\alpha \wedge (d\alpha)^n \neq 0,$$

which is equivalent to the condition that  $d\alpha$  be nondegenerate on  $\xi$ .

Note that the contact structure is unaffected when we multiply the contact form  $\alpha$  by any positive or negative function on  $M$ . We say that two contact structures  $\xi_0 = \ker \alpha_0$  and  $\xi_1 = \ker \alpha_1$  on a manifold  $M$  are *contactomorphic* whenever there is a diffeomorphism  $\psi : M \rightarrow M$  such that  $\psi$  sends  $\xi_0$  to  $\xi_1$ ,

$$\psi_*(\xi_0) = \xi_1.$$

If a diffeomorphism  $\psi : M \rightarrow M$  is in fact a contactomorphism then there exists a nonzero function  $g : M \rightarrow \mathbb{R}$  such that  $\psi^*\alpha_1 = g\alpha_0$ . Finding an explicit contactomorphism often proves to be a rather difficult and messy task, but an application of Moser’s argument yields Gray’s stability theorem, which essentially states that there are no nontrivial deformations of contact structures on a fixed closed manifold.

First we give the statement of Moser's theorem, which says that one cannot vary a symplectic structure by perturbing it within its cohomology class. Recall that a *symplectic structure* on a smooth manifold  $W^{2n}$  is a nondegenerate closed 2-form  $\omega \in \Omega^2(W)$ .

**Theorem 2.3** (Moser's theorem, [McDuff and Salamon 1998, Theorem 3.17]). *Let  $W$  be a closed manifold and suppose that  $\omega_t$  is a smooth family of cohomologous symplectic forms on  $W$ . Then there is a family of diffeomorphisms  $\Psi_t$  of  $W$  such that*

$$\Psi_0 = \text{id}, \quad \psi_t^* \omega_t = \omega_0.$$

The aforementioned contact analogue of Moser's theorem is Gray's stability theorem, stated formally below.

**Theorem 2.4** (Gray's stability theorem, [Geiges 2008, Theorem 2.2.2]). *Let  $\xi_t$ ,  $t \in [0, 1]$ , be a smooth family of contact structures on a closed manifold  $V$ . Then there is an isotopy  $(\psi_t)_{t \in [0,1]}$  of  $V$  such that*

$$\psi_{t*}(\xi_0) = \xi_t \quad \text{for each } t \in [0, 1].$$

Next we give the most basic example of a contact structure.

**Example 2.5.** Consider  $\mathbb{R}^{2n+1}$  with coordinates  $(x_1, y_1, \dots, x_n, y_n, z)$  and the 1-form

$$\alpha = dz + \sum_{j=1}^n x_j dy_j.$$

Then  $\alpha$  is a contact form for  $\mathbb{R}^{2n+1}$ . The contact structure  $\xi = \ker \alpha$  is called the standard contact structure on  $\mathbb{R}^{2n+1}$ .

As in symplectic geometry, a variant of Darboux's theorem holds. This states that locally all contact structures are diffeomorphic to the standard contact structure on  $\mathbb{R}^{2n+1}$ .

A contact form gives rise to a unique Hamiltonian-like vector field as follows.

**Definition 2.6.** For any contact manifold  $(M, \xi = \ker \alpha)$  the *Reeb vector field*  $R_\alpha$  is defined to be the unique vector field determined by  $\alpha$ ,

$$\iota(R_\alpha)d\alpha = 0, \quad \alpha(R_\alpha) = 1.$$

We define the Reeb flow of  $R_\alpha$  by  $\varphi_t : M \rightarrow M$ ,  $\dot{\varphi}_t = R_\alpha(\varphi_t)$ .

The first condition says that  $R_\alpha$  points along the unique null direction of the form  $d\alpha$  and the second condition normalizes  $R_\alpha$ . Because

$$\mathcal{L}_{R_\alpha} \alpha = d\iota_{R_\alpha} \alpha + \iota_{R_\alpha} d\alpha,$$



the flow of  $R_\alpha$  preserves the form  $\alpha$  and hence the contact structure  $\xi$ . Note that if one chooses a different contact form  $f\alpha$ , the corresponding vector field  $R_{f\alpha}$  is very different from  $R_\alpha$ , and its flow may have quite different properties.

A *Reeb orbit*  $\gamma$  of period  $T$  associated to  $R_\alpha$  is defined to be a path  $\gamma : \mathbb{R}/T\mathbb{Z} \rightarrow M$  given by an integral curve of  $R_\alpha$ . That is,

$$\frac{d\gamma}{dt} = R_\alpha \circ \gamma(t), \quad \gamma(0) = \gamma(T).$$

Two Reeb orbits

$$\gamma_1, \gamma_0 : \mathbb{R}/T\mathbb{Z} \rightarrow M$$

are considered equivalent if they differ by reparametrization, i.e., precomposition with a translation of  $\mathbb{R}/T\mathbb{Z}$ .

The  $N$ -fold cover  $\gamma^N$  is defined to be the composition of  $\gamma_\pm$  with  $\mathbb{R}/NT\mathbb{Z} \rightarrow \mathbb{R}/T\mathbb{Z}$ . A *simple Reeb orbit* is one such that  $\gamma : \mathbb{R}/T\mathbb{Z} \rightarrow M$  is injective.

**Remark 2.7.** Since Reeb vector fields are autonomous, the terminology “simple Reeb orbit  $\gamma$ ” refers to the entire equivalence class of orbits, and likewise for its iterates.

A Reeb orbit  $\gamma$  is said to be *nondegenerate* whenever the linearized return map

$$d(\varphi_T)_{\gamma(0)} : \xi_{\gamma(0)} \rightarrow \xi_{\gamma(T)=\gamma(0)}$$

has no eigenvalue equal to 1. A *nondegenerate contact form* is one whose Reeb orbits are all nondegenerate and hence isolated. Note that since the Reeb flow preserves the contact structure, the linearized return map is symplectic.

Next we briefly review the canonical contact form on  $S^3$  and its Reeb dynamics.

**Example 2.8** (canonical Reeb dynamics on the 3-sphere). If we define the function  $f : \mathbb{R}^4 \rightarrow \mathbb{R}$ ,

$$f(x_1, y_1, x_2, y_2) = x_1^2 + y_1^2 + x_2^2 + y_2^2,$$

then  $S^3 = f^{-1}(1)$ . Recall that the canonical contact form on  $S^3 \subset \mathbb{R}^4$  is given to be

$$\lambda_0 := -\frac{1}{2}df \circ J = (x_1 dy_1 - y_1 dx_1 + x_2 dy_2 - y_2 dx_2)|_{S^3}. \quad (2-1)$$

The Reeb vector field is given by

$$\begin{aligned} R_{\lambda_0} &= \left( x_1 \frac{\partial}{\partial y_1} - y_1 \frac{\partial}{\partial x_1} + x_2 \frac{\partial}{\partial y_2} - y_2 \frac{\partial}{\partial x_2} \right) \\ &= (-y_1, x_1, -y_2, x_2). \end{aligned} \quad (2-2)$$

Equivalently we may reformulate these using complex coordinates by identifying  $\mathbb{R}^4$  with  $\mathbb{C}^2$  via

$$u = x_1 + iy_1, \quad v = x_2 + iy_2.$$

We obtain

$$\lambda_0 = \frac{1}{2}i(ud\bar{u} - \bar{u}du + vd\bar{v} - \bar{v}dv)|_{S^3},$$

and

$$\begin{aligned} R_{\lambda_0} &= i\left(u\frac{\partial}{\partial u} - \bar{u}\frac{\partial}{\partial \bar{u}} + v\frac{\partial}{\partial v} - \bar{v}\frac{\partial}{\partial \bar{v}}\right) \\ &= (iu, iv). \end{aligned} \tag{2-3}$$

The second expression for  $R_{\lambda_0}$  follows from (2-2) since  $iu = (-y_1, x_1)$  and  $iv = (-y_2, x_2)$ .

To see that the orbits of  $R_{\lambda_0}$  define the fibers of the Hopf fibration, recall that a fiber through a point

$$(u, v) = (x_1 + iy_1, x_2 + iy_2) \in S^3 \subset \mathbb{C}^2$$

can be parameterized as

$$\varphi(t) = (e^{it}u, e^{it}v), \quad t \in \mathbb{R}. \tag{2-4}$$

We compute the time derivative of the fiber

$$\dot{\varphi}(0) = (iu, iv) = (ix_1 - y_1, ix_2 - y_2).$$

Expressed as a real vector field on  $\mathbb{R}^4$ , which is tangent to  $S^3$ , this is the Reeb vector field  $R_{\lambda_0}$  as it appears in (2-3), so the Reeb flow does indeed define the Hopf fibration.

**2B. Hypersurfaces of contact type.** Another notion that we need from symplectic and contact geometry is that of a hypersurface of contact type in a symplectic manifold. The following notion of a Liouville vector field allows us to define hypersurfaces of contact type. Liouville vector fields will be used to understand the Reeb dynamics of the nondegenerate contact form  $\alpha_1$  as well as to construct the contactomorphism between  $(L_{A_n}, \xi_0)$  and  $(L(n+1, n), \xi_{\text{std}})$ .

**Definition 2.9.** A Liouville vector field  $Y$  on a symplectic manifold  $(W, \omega)$  is a vector field satisfying

$$\mathcal{L}_Y\omega = \omega.$$

The flow  $\psi_t$  of such a vector field is conformal symplectic, i.e.,  $\psi_t^*(\omega) = e^t\omega$ . The flow of these fields is volume expanding, so such fields may only exist locally on compact manifolds.

Whenever there exists a Liouville vector field  $Y$  defined in a neighborhood of a compact hypersurface  $Q$  of  $(W, \omega)$ , which is transverse to  $Q$ , we can define a contact 1-form on  $Q$  by

$$\alpha := \iota_Y\omega.$$

**Proposition 2.10** [McDuff and Salamon 1998, Proposition 3.58]. *Let  $(W, \omega)$  be a symplectic manifold and  $Q \subset W$  a compact hypersurface. Then the following are equivalent:*

- (i) *There exists a contact form  $\alpha$  on  $Q$  such that  $d\alpha = \omega|_Q$ .*
- (ii) *There exists a Liouville vector field  $Y : U \rightarrow TW$  defined in a neighborhood  $U$  of  $Q$ , which is transverse to  $Q$ .*

*If these conditions are satisfied then  $Q$  is said to be of contact type.*

We will need the following application of Gray’s stability theorem to hypersurfaces of contact type to prove [Theorem 1.8](#) in [Section 4](#).

**Lemma 2.11** [Geiges 2008, Lemma 2.1.5]. *Let  $Y$  be a Liouville vector field on a symplectic manifold  $(W, \omega)$ . Suppose that  $M_1$  and  $M_2$  are hypersurfaces of contact type in  $W$ . Assume that there is a smooth function*

$$h : W \rightarrow \mathbb{R} \tag{2-5}$$

*such that the time-1 map of the flow of  $hY$  is a diffeomorphism from  $M_1$  to  $M_2$ . Then this diffeomorphism is in fact a contactomorphism from  $(M_1, \ker \iota_Y \omega|_{TM_1})$  to  $(M_2, \ker \iota_Y \omega|_{TM_2})$ .*

**2C. Symplectization.** The symplectization of a contact manifold is an important notion in defining Floer-theoretic theories like symplectic and contact homology. It will also be used in our calculation of the Conley–Zehnder index. Let  $(M, \xi = \ker \alpha)$  be a contact manifold. The *symplectization* of  $(M, \xi = \ker \alpha)$  is given by the manifold  $\mathbb{R} \times M$  and symplectic form

$$\omega = e^t (d\alpha - \alpha \wedge dt) = d(e^t \alpha).$$

Here  $t$  is the coordinate on  $\mathbb{R}$ , and it should be noted that  $\alpha$  is interpreted as a 1-form on  $\mathbb{R} \times M$ , as we identify  $\alpha$  with its pullback under the projection  $\mathbb{R} \times M \rightarrow M$ .

Any contact structure  $\xi$  may be equipped with a complex structure  $J$  such that  $(\xi, J)$  is a complex vector bundle. This set is nonempty and contractible. There is a unique canonical extension of the almost complex structure  $J$  on  $\xi$  to an  $\mathbb{R}$ -invariant almost complex structure  $\tilde{J}$  on  $T(\mathbb{R} \times M)$ , whose existence is due to the splitting,

$$T(\mathbb{R} \times M) = \mathbb{R} \frac{\partial}{\partial t} \oplus \mathbb{R} R_\alpha \oplus \xi. \tag{2-6}$$

**Definition 2.12** (canonical extension of  $J$  to  $\tilde{J}$  on  $T(\mathbb{R} \times M)$ ). Let  $[a, b; v]$  be a tangent vector where  $a, b \in \mathbb{R}$  and  $v \in \xi$ . We can extend  $J : \xi \rightarrow \xi$  to  $\tilde{J} : T(\mathbb{R} \times M) \rightarrow T(\mathbb{R} \times M)$  by

$$\tilde{J}[a, b; v] = [-b, a, Jv].$$

Thus  $\tilde{J}|_{\xi} = J$  and  $\tilde{J}$  acts on  $\mathbb{R}\partial/\partial t \oplus \mathbb{R}R_\alpha$  in the same manner as multiplication by  $i$  acts on  $\mathbb{C}$ , namely  $J\partial/\partial t = R_\alpha$ .

**2D. The Conley–Zehnder index.** The Conley–Zehnder index  $\mu_{CZ}$  is a Maslov index for arcs of symplectic matrices which assigns an integer  $\mu_{CZ}(\Phi)$  to every path of symplectic matrices  $\Phi : [0, T] \rightarrow \text{Sp}(n)$ , with  $\Phi(0) = \mathbb{1}$ . In order to ensure that the Conley–Zehnder index assigns the same integer to homotopic arcs, one must also stipulate that 1 is not an eigenvalue of the endpoint of this path of matrices, i.e.,  $\det(\mathbb{1} - \Phi(T)) \neq 0$ . We define the following set of continuous paths of symplectic matrices that start at the identity and end on a symplectic matrix that does not have 1 as an eigenvalue:

$$\Sigma^*(n) = \{\Phi : [0, T] \rightarrow \text{Sp}(2n) : \Phi \text{ is continuous, } \Phi(0) = \mathbb{1}, \text{ and } \det(\mathbb{1} - \Phi(T)) \neq 0\}.$$

The Conley–Zehnder index is a functor satisfying the following properties, and is uniquely determined by the homotopy, loop, and signature properties.

**Theorem 2.13** [Robbin and Salamon 1993, Theorem 2.3, Remark 5.4; Gutt 2014, Theorem 2, Propositions 8 and 9]. *There exists a unique functor  $\mu_{CZ}$  called the Conley–Zehnder index that assigns the same integer to all homotopic paths  $\Psi$  in  $\Sigma^*(n)$ ,*

$$\mu_{CZ} : \Sigma^*(n) \rightarrow \mathbb{Z},$$

such that the following hold:

- (1) Homotopy: The Conley–Zehnder index is constant on the connected components of  $\Sigma^*(n)$ .
- (2) Naturalization: For any paths  $\Phi, \Psi : [0, 1] \rightarrow \text{Sp}(2n)$ ,
 
$$\mu_{CZ}(\Phi\Psi\Phi^{-1}) = \mu_{CZ}(\Psi).$$
- (3) Zero: If  $\Psi(t) \in \Sigma^*(n)$  has no eigenvalues on the unit circle for  $t > 0$ , then  $\mu_{CZ}(\Psi) = 0$ .
- (4) Product: If  $n = n' + n''$ , identify  $\text{Sp}(2n') \oplus \text{Sp}(2n'')$  with a subgroup of  $\text{Sp}(2n)$  in the obvious way. For  $\Psi' \in \Sigma^*(n')$  and  $\Psi'' \in \Sigma^*(n'')$ , we have  $\mu_{CZ}(\Psi' \oplus \Psi'') = \mu_{CZ}(\Psi') + \mu_{CZ}(\Psi'')$ .
- (5) Loop: If  $\Phi$  is a loop at  $\mathbb{1}$ , then  $\mu_{CZ}(\Phi\Psi) = \mu_{CZ}(\Psi) + 2\mu(\Phi)$ , where  $\mu$  is the Maslov Index.
- (6) Signature: If  $S \in M(2n)$  is a symmetric matrix with  $\|S\| < 2\pi$  and  $\Psi(t) = \exp(J_0St)$ , then  $\mu_{CZ}(\Psi) = \frac{1}{2} \text{sgn}(S)$ .

The linearized Reeb flow of  $\gamma$  yields a path of symplectic matrices

$$d(\varphi_t)_{\gamma(0)} : \xi_{\gamma(0)} \rightarrow \xi_{\gamma(t)=\gamma(0)}$$

for  $t \in [0, T]$ , where  $T$  is the period of  $\gamma$ .

Thus we can compute the Conley–Zehnder index of  $d\varphi_t$ ,  $t \in [0, T]$ . This index is typically dependent on the choice of trivialization  $\tau$  of  $\xi$  along  $\gamma$  which was used in linearizing the Reeb flow. However, if  $c_1(\xi; \mathbb{Z}) = 0$  we can use the existence of an (almost) complex volume form on the symplectization to obtain a global means of linearizing the flow of the Reeb vector field. The choice of a complex volume form is parametrized by  $H^1(\mathbb{R} \times M; \mathbb{Z})$ , so an absolute integral grading is only determined up to the choice of volume form. See [Nelson  $\geq$  2017, §1.1.1].

We define

$$\mu_{CZ}^\tau(\gamma) := \mu_{CZ}(\{d\varphi_t\}_{t \in [0, T]}).$$

In the case at hand we will be able to work in the ambient space of  $(\mathbb{C}^3, J_0)$ , and use a canonical trivialization of  $\mathbb{C}^3$ .

**2E. The canonical contact structure on Brieskorn manifolds.** The  $A_n$  link is an example of a Brieskorn manifold, which are defined generally by

$$\Sigma(\mathbf{a}) = \left\{ (z_0, \dots, z_m) \in \mathbb{C}^{m+1} \mid f := \sum_{j=0}^m z_j^{a_j} = 0, a_j \in \mathbb{Z}_{>0} \text{ and } \sum_{j=0}^m |z_j|^2 = 1 \right\}.$$

The link of the  $A_n$  singularity after a linear change of variables is  $\Sigma(n+1, 2, 2)$  for  $n > 3$ ; see (3-1). Brieskorn gave a necessary and sufficient condition on  $\mathbf{a}$  for  $\Sigma(\mathbf{a})$  to be a topological sphere, and means to show when these yield exotic differentiable structures on the topological  $(2n-1)$ -sphere in [Brieskorn 1966]. A standard calculus argument [Geiges 2008, Lemma 7.1.1] shows that  $\Sigma(\mathbf{a})$  is always a smooth manifold.

In the mid 1970s, Brieskorn manifolds were found to admit a canonical contact structure, given by their set of complex tangencies,

$$\xi_0 = T\Sigma \cap J_0(T\Sigma),$$

where  $J_0$  is the standard complex structure on  $\mathbb{C}^{m+1}$ . The contact structure  $\xi_0$  can be expressed as  $\xi_0 = \ker \alpha_0$  for the canonical 1-form

$$\alpha_0 := (-d\rho \circ J_0)|_\Sigma = \frac{i}{4} \left( \sum_{j=0}^m (z_j d\bar{z}_j - \bar{z}_j dz_j) \right) \Big|_\Sigma,$$

where  $\rho = (\|z\|^2 - 1)/4$ . A proof of this fact may be found in [Geiges 2008, Theorem 7.1.2]. The Reeb dynamics associated to  $\alpha_0$  are difficult to understand. There is a more convenient contact form  $\alpha_1$  constructed by Ustilovsky [1999, Lemma 4.1.2] via the following family.

**Proposition 2.14** [Geiges 2008, Proposition 7.1.4]. *The 1-form*

$$\alpha_t = \frac{i}{4} \sum_{j=0}^m \frac{1}{1-t+t/a_j} (z_j d\bar{z}_j - \bar{z}_j dz_j)$$

is a contact form on  $\Sigma(\mathbf{a})$  for each  $t \in [0, 1]$ .

Via Gray’s stability theorem we obtain the following corollary.

**Corollary 2.15.** *For all  $t \in (0, 1]$ , the contact manifold  $(\Sigma(\mathbf{a}), \ker \alpha_0)$  is contactomorphic to  $(\Sigma(\mathbf{a}), \ker \alpha_t)$ .*

Next, the Reeb dynamics associated to  $\alpha_1 = \frac{i}{4} \sum_{j=0}^m a_j (z_j d\bar{z}_j - \bar{z}_j dz_j)$  are computed.

**Remark 2.16.** While  $\alpha_1$  is degenerate, one can still easily check that the Reeb vector field associated to  $\alpha_1$  is given by

$$R_{\alpha_1} = 2i \sum_{j=0}^m \frac{1}{a_j} \left( z_j \frac{\partial}{\partial z_j} - \bar{z}_j \frac{\partial}{\partial \bar{z}_j} \right) = 2i \left( \frac{z_0}{a_0}, \dots, \frac{z_m}{a_m} \right).$$

Indeed, one computes

$$df(R_{\alpha_1}) = f(\mathbf{z}) \quad \text{and} \quad d\rho(R_{\alpha_1}) = 0.$$

This shows that  $R_{\alpha_1}$  is tangent to  $\Sigma(\mathbf{a})$ . The defining equations for the Reeb vector field are satisfied since

$$\alpha_1(R_{\alpha_1}) \equiv 1 \quad \text{and} \quad \iota_{R_{\alpha_1}} d\alpha_1 = -d\rho,$$

with the latter form being zero on the  $T_p \Sigma(\mathbf{a})$ . The flow of  $R_{\alpha_1}$  is given by

$$\varphi_t(z_0, \dots, z_m) = (e^{2it/a_0}, \dots, e^{2it/a_m}).$$

All the orbits of the Reeb flow are closed, and the flow defines an effective  $S^1$ -action on  $\Sigma(\mathbf{a})$ .

In the next section we perturb  $\alpha_1$  to a nondegenerate contact form.

### 3. Proof of Theorem 1.1

**3A. Constructing a nondegenerate contact form.** Here, we adapt a method used by Ustilovsky [1999, §4] to obtain a nondegenerate contact form  $\alpha_\epsilon$  on  $L_{A_n}$  whose kernel is contactomorphic to  $\xi_0$ . Ustilovsky’s methods yielded a nondegenerate contact form on Brieskorn manifolds of the form  $\Sigma(p, 2, \dots, 2)$ , which are diffeomorphic to  $S^{4m+1}$ .

We define the change of coordinates to go from  $\Sigma(n + 1, 2, 2)$  with defining function  $f = z_0^{n+1} + z_1^2 + z_2^2$  to  $L_{A_n}$  with defining function  $f_{A_n} = w_0^{n+1} + 2w_1w_2$ :

$$\Psi(w_0, w_1, w_2) = \left( \underbrace{w_0}_{:=z_0}, \underbrace{\frac{\sqrt{2}}{2}(w_1 + w_2)}_{:=z_1}, \underbrace{\frac{\sqrt{2}}{2}(-iw_1 + iw_2)}_{:=z_2} \right). \tag{3-1}$$

We obtain

$$\Psi^* f(z_0, z_1, z_2) = w_0^{n+1} + 2w_1w_2. \tag{3-2}$$

Then the pull-back of

$$\frac{\alpha_1}{2} = \frac{i}{8} \sum_{j=0}^m a_j (z_j d\bar{z}_j - \bar{z}_j dz_j)$$

is given by

$$\frac{\Psi^* \alpha_1}{2} = \frac{(n+1)i}{8} (w_0 d\bar{w}_0 - \bar{w}_0 dw_0) + \frac{i}{4} (w_1 d\bar{w}_1 - \bar{w}_1 dw_1 + w_2 d\bar{w}_2 - \bar{w}_2 dw_2).$$

We now construct the Hamiltonian function

$$H(w) = |w|^2 + \epsilon(|w_1|^2 - |w_2|^2).$$

We choose  $0 < \epsilon < 1$  such that  $H(w)$  is positive on  $S^5$ , and define the contact form

$$\alpha_\epsilon = \Psi^* \alpha_1 / (2H). \tag{3-3}$$

**Remark 3.1.** The above shows that  $(\Sigma(n + 1, 2, 2), \ker \alpha_1)$  is contactomorphic to  $(\Psi(\Sigma(n + 1, 2, 2)), \ker \alpha_\epsilon)$ . Moreover  $L_{A_n} = \Psi(\Sigma(n + 1, 2, 2))$ , where  $L_{A_n}$  was defined in (1-1).

**Proposition 3.2.** *The Reeb vector field for  $\alpha_\epsilon$  is*

$$\begin{aligned} R_{\alpha_\epsilon} &= \frac{4i}{n+1} w_0 \frac{\partial}{\partial w_0} - \frac{4i}{n+1} \bar{w}_0 \frac{\partial}{\partial \bar{w}_0} + 2i(1 + \epsilon) \left( w_1 \frac{\partial}{\partial w_1} - \bar{w}_1 \frac{\partial}{\partial \bar{w}_1} \right) \\ &\quad + 2i(1 - \epsilon) \left( w_2 \frac{\partial}{\partial w_2} - \bar{w}_2 \frac{\partial}{\partial \bar{w}_2} \right) \\ &= \left( \frac{4i}{n+1} w_0, 2i(1 + \epsilon)w_1, 2i(1 - \epsilon)w_2 \right). \end{aligned} \tag{3-4}$$

**Remark 3.3.** The second formulation of the Reeb vector field is equivalent to the first in the above proposition via the standard identification of  $\mathbb{R}^4$  with  $\mathbb{C}^2$ , as explained in [Example 2.8, \(2-3\)](#).

Before proving [Proposition 3.2](#) we need the following lemma.

**Lemma 3.4.** *On  $\mathbb{C}^3$ , the vector field*

$$X(w) = \frac{1}{2} \left( \sum_{j=0}^2 w_j \frac{\partial}{\partial w_j} + \bar{w}_j \frac{\partial}{\partial \bar{w}_j} \right) \quad (3-5)$$

*is a Liouville vector field for the symplectic form*

$$\omega_1 = \frac{d(\Psi^* \alpha_1)}{2} = \frac{i(n+1)}{4} dw_0 \wedge d\bar{w}_0 + \frac{i}{2} \sum_{j=1}^2 dw_j \wedge d\bar{w}_j.$$

*The Hamiltonian vector field  $X_H$  of  $H$  with respect to  $\omega_1$  is  $-R_{\alpha_\epsilon}$ , as in (3-4).*

*Proof.* Recall that the condition to be a Liouville vector field is  $\mathcal{L}_X \omega_1 = \omega_1$ . We show this with Cartan's formula, given as

$$\begin{aligned} \mathcal{L}_X \omega_1 &= \iota_X d\omega_1 + d(\iota_X \omega_1) \\ &= d(\iota_X \omega_1). \end{aligned}$$

We do the explicit calculation for the first term and the rest easily follows:

$$\begin{aligned} d \left( \frac{i(n+1)}{4} dw_0 \wedge d\bar{w}_0 \left( \frac{1}{2} \left( w_0 \frac{\partial}{\partial w_0} + \bar{w}_0 \frac{\partial}{\partial \bar{w}_0} \right), \cdot \right) \right) \\ &= d \left( \frac{i(n+1)}{8} w_0 d\bar{w}_0 - \bar{w}_0 dw_0 \right) \\ &= \frac{i(n+1)}{8} (dw_0 \wedge d\bar{w}_0 - d\bar{w}_0 \wedge dw_0) \\ &= \frac{i(n+1)}{4} dw_0 \wedge d\bar{w}_0, \end{aligned}$$

so  $X(w)$  is indeed a Liouville vector field for  $\omega_1$ .

Next we prove that  $\omega_1(-R_{\alpha_\epsilon}, \cdot) = dH(\cdot)$ . First we calculate  $dH$ :

$$dH = \left( \sum_{j=0}^2 w_j d\bar{w}_j + \bar{w}_j dw_j \right) + \epsilon(w_1 d\bar{w}_1 + \bar{w}_1 dw_1 - w_2 d\bar{w}_2 - \bar{w}_2 dw_2).$$

Then we compare the coefficients of  $dH$  to the coefficients of  $\omega_1(-R_{\alpha_\epsilon}, \cdot)$  associated to each term,  $(dw_i \wedge d\bar{w}_i)$ . The  $(dw_0 \wedge d\bar{w}_0)$  term is

$$\begin{aligned} \frac{i(n+1)}{4} dw_0 \wedge d\bar{w}_0 \left( -\frac{4i}{n+1} w_0 \frac{\partial}{\partial w_0} + \frac{4i}{n+1} \bar{w}_0 \frac{\partial}{\partial \bar{w}_0}, \cdot \right) \\ &= \frac{i(n+1)}{4} \left( -\frac{4i}{n+1} w_0 d\bar{w}_0 - \frac{4i}{n+1} \bar{w}_0 dw_0 \right) \\ &= w_0 d\bar{w}_0 + \bar{w}_0 dw_0. \end{aligned}$$



The  $(dw_1 \wedge d\bar{w}_1)$  term is

$$\begin{aligned} \frac{1}{2}i dw_1 \wedge d\bar{w}_1 &\left( -2i(1 + \epsilon)w_1 \frac{\partial}{\partial w_1} + 2i(1 + \epsilon)\bar{w}_1 \frac{\partial}{\partial \bar{w}_1} \right) \\ &= \frac{1}{2}i(-2i(1 + \epsilon)w_1 d\bar{w}_1 - 2i(1 + \epsilon)\bar{w}_1 dw_1) \\ &= (1 + \epsilon)w_1 d\bar{w}_1 + (1 + \epsilon)\bar{w}_1 dw_1. \end{aligned}$$

The  $(dw_2 \wedge d\bar{w}_2)$  term is obtained in a similar way. Summing the terms yields  $\omega_1(-R_{\alpha_\epsilon}, \cdot) = dH(\cdot)$ . □

*Proof of Proposition 3.2.* First we show that  $X_H = -R_{\alpha_\epsilon}$  is tangent to the link  $\Psi(\Sigma(n + 1, 2, 2))$ . We compute

$$\begin{aligned} (\Psi_*df)(R_{\alpha_\epsilon}) &= ((n + 1)w_0^n dw_0 + 2w_1 dw_2 + 2w_2 dw_1)(R_{\alpha_\epsilon}) \\ &= 4i w_0^{n+1} + 4i(1 - \epsilon)w_1 w_2 + 4i(1 + \epsilon)w_1 w_2 \\ &= 4i(\Psi^* f) \\ &= 0. \end{aligned}$$

The last equality is because  $\Psi^* f$  is constant along  $\Psi(\Sigma(n + 1, 2, 2))$ . Now we have to show that  $\frac{1}{2}\Psi^*\alpha_1(X_H) = -H$ . We have

$$\begin{aligned} \frac{1}{2}\Psi^*\alpha_1(\cdot) &= \iota_X \omega_1(\cdot) = \omega_1(X(w), \cdot) = -\omega(\cdot, X(w)), \\ \frac{1}{2}\Psi^*\alpha_1(X_H) &= -\omega(X_H, X(w)) = -dH(X(w)) \\ &= -|w|^2 - \epsilon(|w_1|^2 - |w_2|^2) \\ &= -H. \end{aligned}$$

From these, we conclude

$$\begin{aligned} \alpha_\epsilon(X_H) &= -\frac{1}{H}H = -1, \\ d\alpha_\epsilon(X_H, \cdot) &= -\frac{1}{2H^2}(dH \wedge \Psi^*\alpha_1)(X_H, \cdot) + \frac{1}{2H}d\Psi^*\alpha_1(X_H, \cdot) \\ &= -\frac{1}{2H^2}dH(X_H)\Psi^*\alpha_1(\cdot) + \frac{1}{2H^2}\Psi^*\alpha_1(X_H)dH(\cdot) + \frac{1}{H}\omega(X_H, \cdot) \\ &= -\frac{1}{2H^2}\omega_1(X_H, X_H)\Psi^*\alpha_1(\cdot) - \frac{1}{H}dH(\cdot) + \frac{1}{H}dH(\cdot) \\ &= 0. \end{aligned}$$

By Lemma 3.4, we know  $-X_H = R_{\alpha_\epsilon}$  so the result follows. □

**3B. Isolated Reeb orbits.** In this short section, we prove the following proposition.

**Proposition 3.5.** *The only simple periodic Reeb orbits of  $R_{\alpha_\epsilon}$  are nondegenerate and defined by*

$$\begin{aligned}\gamma_+(t) &= (0, e^{2i(1+\epsilon)t}, 0), & 0 \leq t \leq \pi/(1+\epsilon), \\ \gamma_-(t) &= (0, 0, e^{2i(1-\epsilon)t}), & 0 \leq t \leq \pi/(1+\epsilon).\end{aligned}$$

*Proof.* The flow of

$$R_{\alpha_\epsilon} = \left( \frac{4i}{n+1} w_0, 2i(1+\epsilon)w_1, 2i(1-\epsilon)w_2 \right)$$

is given by

$$\varphi_t(w_0, w_1, w_2) = \left( e^{4it/(n+1)} w_0, e^{2i(1+\epsilon)t} w_1, e^{2i(1-\epsilon)t} w_2 \right).$$

Since  $\epsilon$  is small and irrational, the only possible periodic trajectories are

$$\begin{aligned}\gamma_0(t) &= (e^{4i/(n+1)t}, 0, 0), \\ \gamma_+(t) &= (0, e^{2i(1+\epsilon)t}, 0), \\ \gamma_-(t) &= (0, 0, e^{2i(1-\epsilon)t}).\end{aligned}$$

It is important to note that the first trajectory does not lie in  $\Psi(\Sigma(n+1, 2, 2))$ , but rather on the total space  $\mathbb{C}^3$ . This is because the point  $\gamma_0(0) = (1, 0, 0)$  is not a zero of  $f_{A_n} = w_0^{n+1} + 2w_1w_2$ .

Next we need to check that the linearized return maps  $d\phi|_\xi$  associated to  $\gamma_+$  and  $\gamma_-$  have no eigenvalues equal to 1. We consider the first orbit  $\gamma_+$  of period  $\pi/(1+\epsilon)$ , as a similar argument applies to the return flow associated to  $\gamma_-$ . The differential of its total return map is

$$d\varphi_T = \left( \begin{array}{ccc} e^{4iT/(n+1)} & 0 & 0 \\ 0 & 1 & 0 \\ 0 & 0 & e^{2i(1-\epsilon)T} \end{array} \right) \Bigg|_{T=\pi/(1+\epsilon)}.$$

Since  $\epsilon$  is a small irrational number, the total return map only has one eigenvalue which is 1. The eigenvector associated to the eigenvalue which is 1 is in the direction of the Reeb orbit  $\gamma^+$ , but since we are restricting the return map to  $\xi$ , we can conclude that  $\gamma_+$  is nondegenerate.  $\square$

**3C. Computation of the Conley–Zehnder index.** To compute the Conley–Zehnder indices of the Reeb orbits in [Theorem 1.1](#) we use the same method as shown in [\[Ustilovsky 1999\]](#), extending the Reeb flow to give rise to a symplectomorphism of  $\mathbb{C}^3 \setminus \{0\}$ . This permits us to do the computations in  $\mathbb{C}^3$ , equipped with the

symplectic form

$$\omega_1 = \frac{d(\Psi^*\alpha_1)}{2} = \frac{i(n+1)}{4}dw_0 \wedge d\bar{w}_0 + \frac{i}{2} \sum_{j=1}^2 dw_j \wedge d\bar{w}_j.$$

We may equip the contact structure  $\xi_0$  with the symplectic form  $\omega = d\alpha_1$  instead of  $d\alpha_\epsilon$  when computing the Conley–Zehnder indices. This is because  $\ker \alpha_\epsilon = \ker \alpha_1 = \xi_0$ , as  $\alpha_\epsilon = (1/H)\alpha_1$  with  $H > 0$  and because  $\omega|_\xi = Hd\alpha_\epsilon|_\xi$  and  $H$  is constant along Reeb trajectories.

Our first proposition shows that we can construct a standard symplectic basis for the symplectic complement

$$\xi^\omega = \{v \in \mathbb{C}^3 : \omega(v, w) = 0 \text{ for all } w \in \xi\}$$

of  $\xi$  in  $\mathbb{C}^3$ . As a result,  $c_1(\xi^\omega) = 0$ . Since  $c_1(\mathbb{C}^3) = 0$ , we know  $c_1(\xi) = 0$ . Thus we may compute the Conley–Zehnder indices in the ambient space  $\mathbb{C}^3$  and use additivity of the Conley–Zehnder index under direct sums of symplectic paths to compute it in  $\xi$ .

**Proposition 3.6.** *There exists a standard symplectic basis for the symplectic complement  $\xi^\omega$  with respect to  $\omega = d\alpha_1$ .*

*Proof.* Notice that  $\xi^\omega = \text{span}(X_1, Y_1, X_2, Y_2)$ , where

$$\begin{aligned} X_1 &= (\bar{w}_0^n, \bar{w}_1, \bar{w}_2), & Y_1 &= iX_1, \\ X_2 &= R_\epsilon, & Y_2 &= w. \end{aligned}$$

We make this into a symplectic standard basis for  $\xi^\omega$  via a Gram–Schmidt process. The new basis is given by

$$\begin{aligned} \tilde{X}_1 &= \frac{X_1}{\sqrt{\omega(X_1, Y_2)}}, & \tilde{Y}_1 &= \frac{Y_1}{\sqrt{\omega(X_1, Y_1)}} = i\tilde{X}_1, \\ \tilde{X}_2 &= X_2, & \tilde{Y}_2 &= Y_2 - \frac{\omega(X_1, Y_2)Y_1 - \omega(Y_1, Y_2)X_1}{\omega(X_1, Y_1)} \\ & & &= Y_2 - \frac{n-1}{2}w_0^{n+1}w(X_1, Y_1)X_1. \end{aligned}$$

This is a standard basis for the symplectic vector space  $\xi^\omega$ ; i.e., the form  $\omega$  in this basis is given by

$$\begin{pmatrix} \begin{pmatrix} 0 & 1 \\ 1 & 0 \end{pmatrix} & \\ & \begin{pmatrix} 0 & 1 \\ 1 & 0 \end{pmatrix} \end{pmatrix}. \quad \square$$

Now we are ready to prove the Conley–Zehnder index formula in [Theorem 1.1](#).

**Proposition 3.7.** *The Conley–Zehnder index for  $\gamma = \gamma_{\pm}^N$  in  $0 \leq t \leq N\pi/(1 \pm \epsilon)$  is*

$$\mu_{CZ}(\gamma_{\pm}^N) = 2 \left( \left\lfloor \frac{2N}{(n+1)(1 \pm \epsilon)} \right\rfloor + \left\lfloor \frac{N(1 \mp \epsilon)}{1 \pm \epsilon} \right\rfloor - \left\lfloor \frac{2N}{1 \pm \epsilon} \right\rfloor \right) + 2N + 1. \quad (3-6)$$

*Proof.* The Reeb flow  $\varphi$  which we introduced in the previous section can be extended to a flow on  $\mathbb{C}^3$ , which we also denote by  $\varphi$ . The action of the extended Reeb flow on  $\mathbb{C}^3$  is given by

$$\begin{aligned} d\varphi_t(w)\tilde{X}_1 &= e^{4it}\tilde{X}_1(\varphi_t(w)), & d\varphi_t(w)\tilde{Y}_1 &= e^{4it}\tilde{Y}_1(\varphi_t(w)), \\ d\varphi_t(w)\tilde{X}_2 &= \tilde{X}_2(\varphi_t(w)), & d\varphi_t(w)\tilde{Y}_2 &= \tilde{Y}_2(\varphi_t(w)). \end{aligned}$$

Define

$$\Phi := d\varphi_t|_{\mathbb{C}^3} = \text{diag}(e^{4i/(n+1)t}, e^{2i(1+\epsilon)t}, e^{2i(1-\epsilon)t}).$$

We can now use the additivity of the Conley–Zehnder index under direct sums of symplectic paths, [Theorem 2.13\(4\)](#) to get

$$\mu_{CZ}(\gamma_{\pm}) = \mu_{CZ}(\Phi) - \mu_{CZ}(\Phi_{\xi^\omega}),$$

where

$$\Phi_{\xi^\omega} := d\varphi_t|_{\xi^\omega} = \text{diag}(e^{4it}, 1). \quad (3-7)$$

The right-hand side of (3-7) is easily computed via the crossing form; see [\[Robbin and Salamon 1993, Remark 5.4\]](#). In particular we have

$$\mu_{CZ}(\{e^{it}\}_{t \in [0, T]}) = \begin{cases} T/\pi, & T \in 2\pi\mathbb{Z}, \\ 2\lfloor T/2\pi \rfloor + 1, & \text{otherwise.} \end{cases}$$

Thus for  $\{\Phi(t)\} = \{e^{4it/(n+1)} \oplus e^{2it(1+\epsilon)} \oplus e^{2it(1-\epsilon)}\}$  with  $0 \leq t \leq T$  we obtain

$$\begin{aligned} \mu_{CZ}(\Phi) &= \begin{cases} 4T/((n+1)\pi), & T \in \frac{1}{2}(n+1)\pi\mathbb{Z}, \\ 2\lfloor 2T/((n+1)\pi) \rfloor + 1, & T \notin \frac{1}{2}(n+1)\pi\mathbb{Z}, \end{cases} \\ &+ \begin{cases} 2T(1+\epsilon)/\pi, & T \in \pi/(1+\epsilon)\mathbb{Z}, \\ 2\lfloor T(1+\epsilon)/\pi \rfloor + 1, & T \notin \pi/(1+\epsilon)\mathbb{Z}, \end{cases} \\ &+ \begin{cases} 2T(1-\epsilon)/\pi, & T \in \pi/(1-\epsilon)\mathbb{Z}, \\ 2\lfloor T(1-\epsilon)/\pi \rfloor + 1, & T \notin \pi/(1-\epsilon)\mathbb{Z}. \end{cases} \end{aligned}$$

Likewise for  $\Phi_{\xi^\omega}$  with  $0 \leq t \leq T$  we obtain

$$\mu_{CZ}(\Phi_{\xi^\omega}) = \begin{cases} 4T/\pi, & T \in \pi/2\mathbb{Z}, \\ 2\lfloor 2T/\pi \rfloor + 1, & T \notin \pi/2\mathbb{Z}. \end{cases}$$

Hence we get that the Conley–Zehnder index for  $\gamma_{\pm}^N$  in  $0 \leq t \leq N\pi/(1 \pm \epsilon)$  is given by

$$\mu_{CZ}(\gamma_{\pm}^N) = 2 \left( \left\lfloor \frac{2N}{(n+1)(1 \pm \epsilon)} \right\rfloor + \left\lfloor \frac{N(1 \mp \epsilon)}{1 \pm \epsilon} \right\rfloor - \left\lfloor \frac{2N}{1 \pm \epsilon} \right\rfloor \right) + 2N + 1. \quad (3-8)$$

This completes the proof.  $\square$

#### 4. Proof of Theorem 1.8

This section proves that  $(L_{A_n}, \xi_0)$  and  $(L(n+1, n), \xi_{\text{std}})$  are contactomorphic. This is done by constructing a 1-parameter family of contact manifolds via a canonically defined Liouville vector field and applying Gray's stability theorem.

**4A. Contact geometry of  $(L(n+1, n), \xi_{\text{std}})$ .** The lens space  $L(n+1, n)$  is obtained via the quotient of  $S^3$  by the binary cyclic subgroup  $A_n \subset \text{SL}(2, \mathbb{C})$ . The subgroup  $A_n$  is given by the action of  $\mathbb{Z}_{n+1}$  on  $\mathbb{C}^2$  defined by

$$\begin{pmatrix} u \\ v \end{pmatrix} \mapsto \begin{pmatrix} e^{2\pi i/(n+1)} & 0 \\ 0 & e^{2n\pi i/(n+1)} \end{pmatrix} \begin{pmatrix} u \\ v \end{pmatrix}.$$

The following exercise shows that  $L(n+1, n)$  is homeomorphic to  $L_{A_n}$ . This construction will be needed later in another proof, so we explain it here to set up the notation.

The origin is the only fixed point of the  $A_n$  action on  $\mathbb{C}^2$  and hence is an isolated quotient singularity of  $\mathbb{C}^2/A_n$ . We can represent  $\mathbb{C}^2/A_n$  as a hypersurface of  $\mathbb{C}^3$  as follows. Consider the monomials

$$z_0 := uv, \quad z_1 := \frac{1}{\sqrt{2}}iu^{n+1}, \quad z_2 := \frac{1}{\sqrt{2}}iv^{n+1}.$$

These are invariant under the action of  $A_n$  and satisfy the equation  $z_0^{n+1} + 2z_1z_2 = 0$ . Recall that

$$f_{A_n}(z_0, z_1, z_2) = z_0^{n+1} + 2z_1z_2 \quad \text{and} \quad L_{A_n} = S^5 \cap \{f_{A_n}^{-1}(0)\}.$$

Moreover,

$$\begin{aligned} \tilde{\varphi} : \mathbb{C}^2 &\rightarrow \mathbb{C}^3, \\ (u, v) &\mapsto (uv, \frac{1}{\sqrt{2}}iu^{n+1}, \frac{1}{\sqrt{2}}iv^{n+1}), \end{aligned} \tag{4-1}$$

descends to the map

$$\varphi : \mathbb{C}^2/A_n \rightarrow \mathbb{C}^3,$$

which sends  $\varphi(\mathbb{C}^2/A_n)$  homeomorphically onto the hypersurface  $f_{A_n}^{-1}(0)$ .

Rescaling away from the origin of  $\mathbb{C}^3$  yields a homeomorphism between  $\varphi(S^3/A_n)$  and  $L_{A_n}$ . As 3-manifolds which are homeomorphic are also diffeomorphic [Moise 1952], we obtain the following proposition.

**Proposition 4.1.**  $L(n+1, n)$  is diffeomorphic to  $L_{A_n}$ .

**Remark 4.2.** In order to prove that two manifolds are contactomorphic, one must either construct an explicit diffeomorphism or make use of Gray's stability theorem. Sadly,  $\varphi$  is not a diffeomorphism onto its image when  $u = 0$  or  $v = 0$ . As the above

diffeomorphism is only known to exist abstractly, we will need to appeal the latter method to prove that  $(L_{A_n}, \xi_0)$  and  $(L(n + 1, n), \xi_{\text{std}})$  are contactomorphic. As a result, this proof is rather involved.

Our application of Gray’s stability theorem uses the flow of a Liouville vector field to construct a 1-parameter family of contactomorphisms. First we prove that  $L(n + 1, n)$  is a contact manifold whose contact structure descends from the quotient of  $S^3$ .

Consider the standard symplectic form on  $\mathbb{C}^2$  given by

$$\begin{aligned} \omega_{\mathbb{C}^2} &= d\lambda_{\mathbb{C}^2}, \\ \lambda_{\mathbb{C}^2} &= \frac{1}{2}i(ud\bar{u} - \bar{u}du + vd\bar{v} - \bar{v}dv). \end{aligned} \tag{4-2}$$

The following proposition shows that  $\lambda_0$  restricts to a contact form on  $L(n + 1, n)$ . We define  $\ker \lambda = \xi_{\text{std}}$  on  $L(n + 1, n)$ .

**Proposition 4.3.** *The vector field*

$$Y_0 = \frac{1}{2} \left( u \frac{\partial}{\partial u} + \bar{u} \frac{\partial}{\partial \bar{u}} + v \frac{\partial}{\partial v} + \bar{v} \frac{\partial}{\partial \bar{v}} \right)$$

is a Liouville vector field on  $(\mathbb{C}^2/A_n, \omega_{\mathbb{C}^2})$  away from the origin and transverse to  $L(n + 1, n)$ .

*Proof.* We have that  $\mathbb{C}^2/A_n$  is a smooth manifold away from the origin because 0 is the only fixed point by the action of  $A_n$ . Write

$$S^3/A_n = \{(u, v) \in \mathbb{C}^2/A_n : |u|^2 + |v|^2 = 1\}.$$

Then  $L(n + 1, n) = S^3/A_n$  is a regular level set of  $g(u, v) = |u|^2 + |v|^2$ . Choose a Riemannian metric on  $\mathbb{C}^2/A_n$  and note that

$$Y_0 = \frac{1}{4} \nabla g.$$

Thus  $Y_0$  is transverse to  $L(n + 1, n)$ . Since

$$\mathcal{L}_{Y_0} \omega_{\mathbb{C}^2} = d(i_{Y_0} d\lambda_{\mathbb{C}^2}) = \omega_{\mathbb{C}^2},$$

we may conclude that  $Y_0$  is indeed a Liouville vector field on  $(\mathbb{C}^2/A_n, \omega_{\mathbb{C}^2})$  away from the origin. Thus by [Proposition 2.10](#),  $L(n + 1, n)$  is a hypersurface of contact type in  $\mathbb{C}^2/A_n$ . □

**4B.** *The proof that  $(L_{A_n}, \xi_0)$  and  $(L(n + 1, n), \xi_{\text{std}})$  are contactomorphic.* First we set up  $L_{A_n}$  and  $\varphi(L(n + 1, n))$  as hypersurfaces of contact type in  $\{f_{A_n}^{-1}(0)\} \setminus \{0\}$ . Define  $\rho : \mathbb{C}^3 \rightarrow \mathbb{R}$  by

$$\rho(z) = \frac{1}{4}|z|^2 - 1 = \frac{1}{4}z_0\bar{z}_0 + \dots + z_2\bar{z}_2 - 1.$$

The standard symplectic structure on  $\mathbb{C}^3$  is given by

$$\omega_{\mathbb{C}^3} = \frac{1}{2}i(dz_0 \wedge d\bar{z}_0 + \cdots + dz_2 \wedge d\bar{z}_2).$$

Moreover,

$$Y = \nabla \rho = \frac{1}{2} \sum_{j=0}^2 z_j \frac{\partial}{\partial z_j} + \bar{z}_j \frac{\partial}{\partial \bar{z}_j} \quad (4-3)$$

is a Liouville vector field for  $(\mathbb{C}^3, \omega_{\mathbb{C}^3})$ . We define

$$\lambda_{\mathbb{C}^3} = \iota_Y \omega_{\mathbb{C}^3}.$$

A standard calculation analogous to the proof of [Proposition 4.3](#) shows that  $Y$  is a Liouville vector field on  $(\{f_{A_n}^{-1}(0)\} \setminus \{0\}, \omega_{\mathbb{C}^3})$ .

**Remark 4.4.** Both  $\varphi(L(n+1, n))$  and  $L_{A_n}$  are hypersurfaces of contact type in  $(\{f_{A_n}^{-1}(0)\} \setminus \{0\}, \omega_{\mathbb{C}^3})$ . Note that  $\varphi(L(n+1, n))$  is in fact transverse to the Liouville vector field  $Y$  because

$$\begin{aligned} \varphi(L(n+1, n)) &= \varphi(\{|u|^2 + |v|^2 = 1\}/A_n) \\ &= \varphi(\{|u|^4 + 2|u|^2|v|^2 + |v|^4 = 1\}/A_n) \\ &= \{2|z_0|^2 + 4^{1/(n+1)}|z_1|^{4/(n+1)} + 4^{1/(n+1)}|z_2|^{4/(n+1)} = 1\} \cap f_{A_n}^{-1}(0). \end{aligned}$$

We will want  $\varphi(L(n+1, n))$  and  $L_{A_n}$  to be disjoint in  $\{f_{A_n}^{-1}(0)\}$ . This is easily accomplished by rescaling  $r$  in the definition of the link.

**Definition 4.5.** Define

$$L_{A_n}^r = f_{A_n}^{-1}(0) \cap S_r^5,$$

with the assumption that  $r$  has been chosen so that  $\varphi(L(n+1, n))$  and  $L_{A_n}^r$  are disjoint in  $\{f_{A_n}^{-1}(0)\}$  and so that the flow of the Liouville vector field  $Y$  “hits”  $\varphi(L(n+1, n))$  before  $L_{A_n}^r$ .

The first result is the following lemma, which provides a 1-parameter family of diffeomorphic manifolds starting on  $\varphi(L(n+1, n))$  and ending on  $L_{A_n}^r$ . First we set up some notation. Let

$$\psi_t : \mathbb{R} \times X \rightarrow X$$

be the flow of  $Y$  and  $\psi_t(z) = \gamma_z(t)$  the unique integral curve passing through  $z \in \varphi(L(n+1, n))$  at time  $t = 0$ . For any integral curve  $\gamma$  of  $Y$  we consider the initial value problem

$$\gamma'(t) = Y(\gamma(t)) \quad \text{and} \quad \gamma(0) = z \in \varphi(L(n+1, n)). \quad (4-4)$$

By means of the implicit function theorem and the properties of the Liouville vector field  $Y$  we can prove the following claim.

**Lemma 4.6.** *For every  $\gamma_z$ , there exists a  $\tau(z) \in \mathbb{R}_{>0}$  such that  $\gamma_z(\tau(z)) \in L_{A_n}^r$ . The choice of  $\tau(z)$  varies smoothly for each  $z \in \varphi(L(n+1, n))$ .*

*Proof.* In order to apply the implicit function theorem, we must show for all  $(t, z)$  with  $\rho \circ \gamma = 0$  that

$$\frac{\partial(\rho \circ \gamma)}{\partial t} \neq 0.$$

Note that  $\rho \circ \gamma$  is smooth. By the chain rule,

$$\left. \frac{\partial(\rho \circ \gamma)}{\partial t} \right|_{(s,p)} = \text{grad } \rho|_{\gamma(s,p)} \cdot \dot{\gamma}|_{(s,p)},$$

where  $\dot{\gamma}|_{(s,p)} = \partial\gamma/\partial t|_{(s,p)}$ .

If  $\text{grad } \rho|_{\gamma(s,p)} \cdot \dot{\gamma}|_{(s,p)} = 0$ , then  $\text{grad } \rho$  is not transverse along  $\{(\rho \circ \gamma)(s, p) = 0\}$  or  $\dot{\gamma}|_{(s,p)} = 0$ , since  $\text{grad } \rho \neq 0$ . By construction,  $\text{grad } \rho = \nabla \rho$  is a Liouville vector field transverse to  $L_{A_n}^r$ . Furthermore, the conformal symplectic nature of a Liouville vector field implies that for any integral curve  $\gamma$  satisfying the initial value problem given by (4-4),  $\dot{\gamma}|_{(s,p)} \neq 0$ . Thus we see that the conditions for the implicit function theorem are satisfied and our claim is proven.  $\square$

**Remark 4.7.** The time  $\tau(z)$  can be normalized to 1 for each  $z$ , yielding a 1-parameter family of diffeomorphic contact manifolds  $(M_t, \zeta_t)$  for  $0 \leq t \leq 1$  given by

$$M_t = \psi_t(\varphi(L(n+1, n))), \quad \zeta_t = TM_t \cap J_{\mathbb{C}^3}(TM_t),$$

where

$$M_0 = \psi_0(\varphi(L(n+1, n))) = \varphi(L(n+1, n)), \quad M_1 = \psi_1(\varphi(L(n+1, n))) = L_{A_n}.$$

Moreover, we can relate the standard contact structure on  $L(n+1, n)$  under the image of  $\varphi$ . To avoid excessive parentheses, we use  $S^3/A_n$  in place of  $L(n+1, n)$  in this lemma.

**Lemma 4.8.** *On  $\varphi(S^3/A_n)$ ,*

$$\varphi_*(\xi_{\text{std}}) = T(\varphi(S^3/A_n)) \cap J_{\mathbb{C}^3}(T(\varphi(S^3/A_n))).$$

*Proof.* Since  $A_n \subset \text{SL}(2, \mathbb{C})$ , we have

$$\tilde{\varphi}(J_{\mathbb{C}^2}TS^3) = J_{\mathbb{C}^3}(T\tilde{\varphi}(S^3)).$$

Examining  $\varphi_*(\xi_{\text{std}})$  yields

$$\begin{aligned} \varphi_*(T(S^3/A_n) \cap J_{\mathbb{C}^2}T(S^3/A_n)) &= \tilde{\varphi}_*(TS^3 \cap J_{\mathbb{C}^2}(TS^3)) = \tilde{\varphi}_*(TS^3) \cap \tilde{\varphi}_*(J_{\mathbb{C}^2}(TS^3)) \\ &= \tilde{\varphi}_*(TS^3) \cap J_{\mathbb{C}^3}\tilde{\varphi}_*(TS^3) = T\tilde{\varphi}(S^3) \cap J_{\mathbb{C}^3}(T\tilde{\varphi}(S^3)) \\ &= T(\varphi(S^3/A_n)) \cap J_{\mathbb{C}^3}(T\varphi(S^3/A_n)). \end{aligned} \quad \square$$



Lemmas 4.6 and 4.8 in conjunction with Remark 4.7 and Lemma 2.11 yield the following proposition.

**Proposition 4.9.** *The image of the lens space  $(\varphi(L(n + 1, n)), \varphi_*\xi_{\text{std}})$  is contactomorphic to  $(L_{A_n}, \xi_0)$ .*

It remains to show that  $(\varphi(L(n + 1, n)), \varphi_*\xi_{\text{std}})$  and  $(L(n + 1, n), \xi_{\text{std}})$  are contactomorphic. To accomplish this, we use Moser’s lemma to prove the following lemma.

**Lemma 4.10.** *The manifolds  $(\mathbb{C}^2 \setminus \{0\}, d\lambda_{\mathbb{C}^2})$  and  $(\mathbb{C}^2 \setminus \{0\}, d\tilde{\varphi}^*\lambda_{\mathbb{C}^3})$  are contactomorphic.*

*Proof.* Consider the family of 2-forms

$$\omega_t = (1 - t)\omega_{\mathbb{C}^2} + t\tilde{\varphi}^*\omega_{\mathbb{C}^3}$$

for  $0 \leq t \leq 1$ . Then  $\omega_t$  is exact because  $Y_0$  and  $Y$  are Liouville vector fields for  $\mathbb{C}^2 \setminus \{0\}$  equipped with the symplectic forms  $\omega_{\mathbb{C}^2}$  and  $\omega_{\mathbb{C}^3}$  respectively; thus  $d\lambda_t = \omega_t$  for

$$\lambda_t = (1 - t)\lambda_{\mathbb{C}^2} + t\tilde{\varphi}^*(\lambda_{\mathbb{C}^3})$$

for  $0 \leq t \leq 1$ . We claim that  $\lambda_t$  is a family of contact forms for each  $t \in [0, 1]$ .

We compute

$$\begin{aligned} \frac{2}{i}\tilde{\varphi}^*d\lambda_{\mathbb{C}^3} &= d(uv) \wedge d(\overline{u}\overline{v}) + d(u^{n+1}) \wedge d(\overline{u}^{n+1}) + d(v^{n+1}) \wedge d(\overline{v}^{n+1}) \\ &= ((n + 1)^2|u|^{2n} + |v|^2)du \wedge d\overline{u} + 2\Re(u\overline{v}dv \wedge d\overline{u}) \\ &\quad + ((n + 1)^2|v|^{2n} + |u|^2)dv \wedge d\overline{v}. \end{aligned}$$

Since  $\omega_t$  is exact for each  $t \in [0, 1]$ , we know  $d(\omega_t) = 0$  for each  $t \in [0, 1]$ . Moreover, a simple calculation reveals that  $\omega_t \wedge \omega_t$  is a volume form on  $\mathbb{C}^2$  for each  $t \in [0, 1]$ . Thus we may conclude that  $\omega_t$  is a symplectic form for each  $t \in [0, 1]$ . Applying Moser’s argument, Theorem 2.3, yields the desired result.  $\square$

This yields the desired corollary.

**Corollary 4.11.** *The manifolds  $(L(n + 1, n), \ker \lambda_{\mathbb{C}^2})$  and  $(L(n + 1, n), \ker \varphi^*\lambda_{\mathbb{C}^3})$  are contactomorphic.*

*Proof.* Let  $\phi : (\mathbb{C}^2 \setminus \{0\}, d\lambda_{\mathbb{C}^2})$  and  $(\mathbb{C}^2 \setminus \{0\}, d\tilde{\varphi}^*\lambda_{\mathbb{C}^3})$  be the symplectomorphism, which exists by Lemma 4.10. It induces the desired contactomorphism. On  $\mathbb{C}^2 \setminus \{0\}$ ,

$$\phi^*d(\varphi^*\lambda_{\mathbb{C}^3}) = d\lambda_{\mathbb{C}^2};$$

thus,

$$d\phi^*(\varphi^*\lambda_{\mathbb{C}^3}) = d\lambda_{\mathbb{C}^2}.$$

So on  $L(n + 1, n)$ ,

$$\phi_*(\xi_{\text{std}}) = \phi_*(\ker \lambda_{\mathbb{C}^2}) = \ker \phi_*\lambda_{\mathbb{C}^3} = \varphi_*\xi_{\text{std}}.$$

Proposition 4.9 and Corollary 4.11 complete the proof of Theorem 1.8.  $\square$

### Acknowledgements

Leonardo Abbrescia, Irit Huq-Kuruville, and Nawaz Sultani thank Dr. Jo Nelson and Robert Castellano for their patient guidance and tutelage for this project. They have expanded our mathematical knowledge and have set us on the right path for further research. We especially thank Dr. Nelson for going above and beyond the call of an REU instructor. We are very grateful to the referee for their numerous helpful suggestions on improving the exposition of this paper.

We thank Dr. Robert Lipschitz and the Columbia University math REU program for giving us the opportunity to bring this project to fruition. The REU program was partially funded by NSF Grant DMS-0739392. Since graduating from Columbia, Leonardo Abbrescia is supported in part by a NSF graduate research fellowship. Jo Nelson is supported by NSF grant DMS-1303903, the Bell Companies Fellowship, the Charles Simonyi Endowment, and the Fund for Mathematics at the Institute for Advanced Study.

### References

- [Bourgeois and Oancea 2012] F. Bourgeois and A. Oancea, “ $S^1$ -equivariant symplectic homology and linearized contact homology”, preprint, 2012. To appear in *Int. Math. Res. Not.* [arXiv](#)
- [Bourgeois and Oancea 2015] F. Bourgeois and A. Oancea, “Erratum to: An exact sequence for contact- and symplectic homology”, *Invent. Math.* **200**:3 (2015), 1065–1076. [MR](#) [Zbl](#)
- [Brieskorn 1966] E. Brieskorn, “Beispiele zur Differentialtopologie von Singularitäten”, *Invent. Math.* **2** (1966), 1–14. [MR](#) [Zbl](#)
- [Geiges 2008] H. Geiges, *An introduction to contact topology*, Cambridge Studies in Advanced Mathematics **109**, Cambridge University Press, 2008. [MR](#) [Zbl](#)
- [Gutt 2014] J. Gutt, “Generalized Conley–Zehnder index”, *Ann. Fac. Sci. Toulouse Math.* (6) **23**:4 (2014), 907–932. [MR](#) [Zbl](#)
- [Gutt 2015] J. Gutt, “The positive equivariant symplectic homology as an invariant for some contact manifolds”, preprint, 2015. [arXiv](#)
- [Hutchings and Nelson 2014] M. Hutchings and J. Nelson, “Cylindrical contact homology for dynamically convex contact forms in three dimensions”, preprint, 2014. To appear in *J. Symplectic Geom.* [arXiv](#)
- [Hutchings and Nelson  $\geq$  2017] M. Hutchings and J. Nelson, “Invariance and an integral lift of cylindrical contact homology for dynamically convex contact forms”, in preparation.
- [Ito and Reid 1996] Y. Ito and M. Reid, “The McKay correspondence for finite subgroups of  $SL(3, \mathbb{C})$ ”, pp. 221–240 in *Higher-dimensional complex varieties* (Trento, 1994), edited by M. Andreatta and T. Peternell, de Gruyter, Berlin, 1996. [MR](#) [Zbl](#)
- [Keating 2015] A. Keating, “Homological mirror symmetry for hypersurface cusp singularities”, preprint, 2015. [arXiv](#)
- [Kwon and van Koert 2016] M. Kwon and O. van Koert, “Brieskorn manifolds in contact topology”, *Bull. Lond. Math. Soc.* **48**:2 (2016), 173–241. [MR](#) [Zbl](#)

- [McDuff and Salamon 1998] D. McDuff and D. Salamon, *Introduction to symplectic topology*, 2nd ed., Oxford University Press, 1998. [MR](#) [Zbl](#)
- [McLean 2016] M. McLean, “Reeb orbits and the minimal discrepancy of an isolated singularity”, *Invent. Math.* **204**:2 (2016), 505–594. [MR](#) [Zbl](#)
- [McLean and Ritter  $\geq$  2017] M. McLean and A. Ritter, “The cohomological McKay correspondence and symplectic homology”, in preparation.
- [Milnor 1968] J. Milnor, *Singular points of complex hypersurfaces*, Annals of Mathematics Studies **61**, Princeton University Press, 1968. [MR](#) [Zbl](#)
- [Moise 1952] E. E. Moise, “Affine structures in 3-manifolds, V: The triangulation theorem and Hauptvermutung”, *Ann. Math.* **56**:2 (1952), 96–114. [MR](#) [Zbl](#)
- [Nelson 2015] J. Nelson, “Automatic transversality in contact homology, I: Regularity”, *Abh. Math. Semin. Univ. Hambg.* **85**:2 (2015), 125–179. [MR](#) [Zbl](#)
- [Nelson  $\geq$  2017] J. Nelson, “Automatic transversality in contact homology, II: Invariance and computations”, in preparation.
- [Ritter 2010] A. F. Ritter, “Deformations of symplectic cohomology and exact Lagrangians in ALE spaces”, *Geom. Funct. Anal.* **20**:3 (2010), 779–816. [MR](#) [Zbl](#)
- [Robbin and Salamon 1993] J. Robbin and D. Salamon, “The Maslov index for paths”, *Topology* **32**:4 (1993), 827–844. [MR](#) [Zbl](#)
- [Seidel 2008a] P. Seidel, “A biased view of symplectic cohomology”, pp. 211–253 in *Current developments in mathematics, 2006*, edited by B. Mazur et al., Int. Press, Somerville, MA, 2008. [MR](#) [Zbl](#)
- [Seidel 2008b] P. Seidel, *Fukaya categories and Picard–Lefschetz theory*, European Mathematical Society, Zürich, 2008. [MR](#) [Zbl](#)
- [Ustilovsky 1999] I. Ustilovsky, “Infinitely many contact structures on  $S^{4m+1}$ ”, *Internat. Math. Res. Notices* **14** (1999), 781–791. [MR](#) [Zbl](#)

Received: 2015-09-18

Revised: 2016-06-07

Accepted: 2016-06-09

[leonardo@math.msu.edu](mailto:leonardo@math.msu.edu)Department of Mathematics, Michigan State University,  
1016 Chester Rd., Apt. B14, Lansing, MI 48912, United States[ih2271@columbia.edu](mailto:ih2271@columbia.edu)Department of Mathematics, Columbia University, 2920 Broad-  
way, 6580 Lerner Hall, New York, NY 10027, United States[nelson@math.columbia.edu](mailto:nelson@math.columbia.edu)Institute for Advanced Study, Columbia University,  
2990 Broadway, Room 509, MC 4403, New York, NY 10027,  
United States[njs2155@columbia.edu](mailto:njs2155@columbia.edu)Department of Mathematics, Columbia University, 2920 Broad-  
way, 6580 Lerner Hall, New York, NY 10027, United States

## INVOLVE YOUR STUDENTS IN RESEARCH

*Involve* showcases and encourages high-quality mathematical research involving students from all academic levels. The editorial board consists of mathematical scientists committed to nurturing student participation in research. Bridging the gap between the extremes of purely undergraduate research journals and mainstream research journals, *Involve* provides a venue to mathematicians wishing to encourage the creative involvement of students.

### MANAGING EDITOR

Kenneth S. Berenhaut Wake Forest University, USA

### BOARD OF EDITORS

Colin Adams	Williams College, USA	Suzanne Lenhart	University of Tennessee, USA
John V. Baxley	Wake Forest University, NC, USA	Chi-Kwong Li	College of William and Mary, USA
Arthur T. Benjamin	Harvey Mudd College, USA	Robert B. Lund	Clemson University, USA
Martin Bohner	Missouri U of Science and Technology, USA	Gaven J. Martin	Massey University, New Zealand
Nigel Boston	University of Wisconsin, USA	Mary Meyer	Colorado State University, USA
Amarjit S. Budhiraja	U of North Carolina, Chapel Hill, USA	Emil Minchev	Ruse, Bulgaria
Pietro Cerone	La Trobe University, Australia	Frank Morgan	Williams College, USA
Scott Chapman	Sam Houston State University, USA	Mohammad Sal Moslehian	Ferdowsi University of Mashhad, Iran
Joshua N. Cooper	University of South Carolina, USA	Zuhair Nashed	University of Central Florida, USA
Jem N. Corcoran	University of Colorado, USA	Ken Ono	Emory University, USA
Toka Diagana	Howard University, USA	Timothy E. O'Brien	Loyola University Chicago, USA
Michael Dorff	Brigham Young University, USA	Joseph O'Rourke	Smith College, USA
Sever S. Dragomir	Victoria University, Australia	Yuval Peres	Microsoft Research, USA
Behrouz Emamizadeh	The Petroleum Institute, UAE	Y.-F. S. Pétermann	Université de Genève, Switzerland
Joel Foisy	SUNY Potsdam, USA	Robert J. Plemmons	Wake Forest University, USA
Erin W. Fulp	Wake Forest University, USA	Carl B. Pomerance	Dartmouth College, USA
Joseph Gallian	University of Minnesota Duluth, USA	Vadim Ponomarenko	San Diego State University, USA
Stephan R. Garcia	Pomona College, USA	Bjorn Poonen	UC Berkeley, USA
Anant Godbole	East Tennessee State University, USA	James Propp	U Mass Lowell, USA
Ron Gould	Emory University, USA	József H. Przytycki	George Washington University, USA
Andrew Granville	Université Montréal, Canada	Richard Rebarber	University of Nebraska, USA
Jerrold Griggs	University of South Carolina, USA	Robert W. Robinson	University of Georgia, USA
Sat Gupta	U of North Carolina, Greensboro, USA	Filip Saidak	U of North Carolina, Greensboro, USA
Jim Haglund	University of Pennsylvania, USA	James A. Sellers	Penn State University, USA
Johnny Henderson	Baylor University, USA	Andrew J. Sterge	Honorary Editor
Jim Hoste	Pitzer College, USA	Ann Trenk	Wellesley College, USA
Natalia Hritonenko	Prairie View A&M University, USA	Ravi Vakil	Stanford University, USA
Glenn H. Hurlbert	Arizona State University, USA	Antonia Vecchio	Consiglio Nazionale delle Ricerche, Italy
Charles R. Johnson	College of William and Mary, USA	Ram U. Verma	University of Toledo, USA
K. B. Kulasekera	Clemson University, USA	John C. Wierman	Johns Hopkins University, USA
Gerry Ladas	University of Rhode Island, USA	Michael E. Zieve	University of Michigan, USA

### PRODUCTION

Silvio Levy, Scientific Editor


Cover: Alex Scorpan

See inside back cover or [msp.org/involve](http://msp.org/involve) for submission instructions. The subscription price for 2017 is US\$175/year for the electronic version, and \$235/year (+\$35, if shipping outside the US) for print and electronic. Subscriptions, requests for back issues from the last three years and changes of subscribers address should be sent to MSP.

Involve (ISSN 1944-4184 electronic, 1944-4176 printed) at Mathematical Sciences Publishers, 798 Evans Hall #3840, c/o University of California, Berkeley, CA 94720-3840, is published continuously online. Periodical rate postage paid at Berkeley, CA 94704, and additional mailing offices.

Involve peer review and production are managed by EditFLOW<sup>®</sup> from Mathematical Sciences Publishers.

PUBLISHED BY

 **mathematical sciences publishers**  
nonprofit scientific publishing

<http://msp.org/>

© 2017 Mathematical Sciences Publishers

# involve

2017

vol. 10

no. 3

Dynamics of vertical real rhombic Weierstrass elliptic functions	361
LORELEI KOSS AND KATIE ROY	
Pattern avoidance in double lists	379
CHARLES CRATTY, SAMUEL ERICKSON, FREHIWET NEGASSI AND LARA PUDWELL	
On a randomly accelerated particle	399
MICHELLE NUNO AND JUHI JANG	
Reeb dynamics of the link of the $A_n$ singularity	417
LEONARDO ABBRESCIA, IRIT HUQ-KURUVILLA, JO NELSON AND NAWAZ SULTANI	
The vibration spectrum of two Euler–Bernoulli beams coupled via a dissipative joint	443
CHRIS ABRIOLA, MATTHEW P. COLEMAN, AGLIKA DARAKCHIEVA AND TYLER WALES	
Loxodromes on hypersurfaces of revolution	465
JACOB BLACKWOOD, ADAM DUKEHART AND MOHAMMAD JAVAHERI	
Existence of positive solutions for an approximation of stationary mean-field games	473
NOJOOD ALMAYOUF, ELENA BACHINI, ANDREA CHAPOUTO, RITA FERREIRA, DIOGO GOMES, DANIELA JORDÃO, DAVID EVANGELISTA JUNIOR, AVETIK KARAGULYAN, JUAN MONASTERIO, LEVON NURBEKYAN, GIORGIA PAGLIAR, MARCO PICCIRILLI, SAGAR PRATAPSI, MARIANA PRAZERES, JOÃO REIS, ANDRÉ RODRIGUES, ORLANDO ROMERO, MARIA SARGSYAN, TOMMASO SENECI, CHULIANG SONG, KENGO TERAJ, RYOTA TOMISAKI, HECTOR VELASCO-PEREZ, VARDAN VOSKANYAN AND XIANJIN YANG	
Discrete dynamics of contractions on graphs	495
OLENA OSTAPYUK AND MARK RONNENBERG	
Tiling annular regions with skew and T-tetrominoes	505
AMANDA BRIGHT, GREGORY J. CLARK, CHARLES DUNN, KYLE EVITTS, MICHAEL P. HITCHMAN, BRIAN KEATING AND BRIAN WHETTER	
A bijective proof of a $q$ -analogue of the sum of cubes using overpartitions	523
JACOB FORSTER, KRISTINA GARRETT, LUKE JACOBSEN AND ADAM WOOD	
Ulrich partitions for two-step flag varieties	531
IZZET COSKUN AND LUKE JASKOWIAK	