

involve

a journal of mathematics

A bijective proof of a q -analogue of
the sum of cubes using overpartitions

Jacob Forster, Kristina Garrett, Luke Jacobsen and Adam Wood



A bijective proof of a q -analogue of the sum of cubes using overpartitions

Jacob Forster, Kristina Garrett, Luke Jacobsen and Adam Wood

(Communicated by Jim Haglund)

We present a q -analogue of the sum of cubes, give an interpretation in terms of overpartitions, and provide a combinatorial proof. In addition, we note a connection between a generating function for overpartitions and the q -Delannoy numbers.

1. Introduction

The formula for the sum of the first n cubes,

$$\sum_{k=1}^n k^3 = \binom{n+1}{2}^2, \quad (1)$$

is well known and has been proven using various methods. Benjamin and Orrison [2002] gave two combinatorial proofs. More recently, Garrett and Hummel [2004] proved a q -analogue of (1) using integer partitions. (A q -analogue is an expression involving q -binomial coefficients — see Section 2.3 on the next page — and reducing to the given expression when $q \rightarrow 1^-$.) In this paper, we give an alternate q -analogue of (1) and provide a bijective proof using overpartitions. The first section is devoted to an introduction to partition theory and establishing necessary notation and facts for our work. Then we state and explain a generating function for overpartitions and relate it to the Delannoy numbers. In the last section we give our q -analogue and provide a combinatorial proof.

2. Background

In this section, we introduce aspects of partition theory that are relevant to our work. For further reading, see [Andrews 1976; Corteel and Lovejoy 2004].

MSC2010: 05A17, 05A19.

Keywords: overpartitions, combinatorial proof, Delannoy numbers, q -analogue.

2.1. Partitions.

Definition 1. A partition λ of a positive integer n is a nonincreasing sequence of positive integers $\lambda_1, \lambda_2, \dots, \lambda_k$ such that $\sum_{i=1}^k \lambda_i = n$. The λ_i are called the parts of the partition.

As an example, consider $n = 4$. The five distinct partitions of 4 are

$$4, 31, 22, 211, 1111.$$

One method of displaying partitions graphically is with Ferrers shapes. A Ferrers shape of a partition $\lambda = \lambda_1, \lambda_2, \dots, \lambda_k$, where $\lambda_i \geq \lambda_{i+1}$, is a left-justified array of cells with λ_i cells in row i of the shape and $i = 1$ defined as the top row. Below is the Ferrers shape for the partition $\lambda = 31$:



2.2. Overpartitions.

Definition 2. An overpartition λ is a partition $\lambda_1, \lambda_2, \dots, \lambda_k$ in which the first occurrence of a given part size may be overlined.

Below are the fourteen distinct overpartitions of $n = 4$:

$$4, \bar{4}, 31, \bar{3}1, 3\bar{1}, \bar{3}\bar{1}, 22, \bar{2}2, 211, \bar{2}11, 2\bar{1}1, \bar{2}\bar{1}1, 1111, \bar{1}111.$$

Overpartitions can also be graphically represented using Ferrers shapes by letting the last cell of the rows corresponding to overlined parts be shaded. For example, the Ferrers shape for the overpartition $\lambda = \bar{3}1$ is



2.3. Partitions in a $k \times (n - k)$ box. In order to discuss partitions whose Ferrers shapes fit inside of a $k \times (n - k)$ box, we must first introduce the q -binomial coefficient. The q -binomial coefficient is defined as

$$\left[\begin{matrix} n \\ k \end{matrix} \right]_q = \frac{\prod_{i=n-k+1}^n (1 - q^i)}{\prod_{i=1}^k (1 - q^i)},$$

and is a q -analogue of the binomial coefficient. It is well known that

$$g_{n,k}(q) = \left[\begin{matrix} n \\ k \end{matrix} \right]_q$$

is the generating function for the number of partitions whose Ferrers shapes fit inside of a $k \times (n - k)$ box. This generating function can be easily shown to satisfy the recurrence relation

$$g_{n,k}(q) = q^k g_{n-1,k}(q) + g_{n-1,k-1}(q),$$

which is a q -analogue of the binomial coefficient recurrence. Note that, in this case, the empty partition is included in the set of partitions that fit inside the $k \times (n - k)$ box.

3. Overpartitions in a $2 \times (n - 1)$ box

For our main theorem, we are interested in counting the number of overpartitions whose Ferrers shape fits in a $2 \times (n - 1)$ box. The generating function for the number of partitions in a $2 \times (n - 1)$ box is $\left[\begin{smallmatrix} n+1 \\ 2 \end{smallmatrix} \right]_q$. In this section, we give an analogy of this generating function for overpartitions. We will discuss the recurrence relation for overpartitions that fit in a $k \times (n - k)$ box and then use it to verify a generating function for the number of overpartitions that fit in a $2 \times (n - 1)$ box.

3.1. Recurrence relation for overpartitions. We will first discuss the general case of overpartitions in a $k \times (n - k)$ box and then consider the case of a $2 \times (n - 1)$ box. Let $\bar{p}_{n,k}$ denote the number of overpartitions that can fit in a $k \times (n - k)$ box. Then, $\bar{p}_{n,k}$ satisfies the recurrence relation

$$\bar{p}_{n,k} = \bar{p}_{n-1,k} + \bar{p}_{n-1,k-1} + \bar{p}_{n-2,k-1}. \tag{2}$$

We will explain each term in the recurrence relation. Note that, given a $k \times (n - k)$ box, this recurrence relation indicates that there are three possible disjoint ways of transforming the $k \times (n - k)$ box which, when taken together, describe all possible overpartitions that can fit inside a $k \times (n - k)$ box. These disjoint cases can be easily seen by considering the largest part of an overpartition, λ , in a $k \times (n - k)$ box and are as follows:

- (i) $lp(\lambda) < n - k$. Then the other parts of the overpartition must be less than or equal to $lp(\lambda)$. This situation describes the number of overpartitions in a $k \times (n - k - 1)$ box.
- (ii) $lp(\lambda) = n - k$ and $lp(\lambda)$ is not overlined. Then the other parts of the overpartition are less than or equal to $n - k$. Thus, this collection of overpartitions is equivalent to the number of overpartitions in a $(k - 1) \times (n - k)$ box.
- (iii) $lp(\lambda) = n - k$ and $lp(\lambda)$ is overlined. Then the other parts of the overpartition must be less than $(n - k)$. Hence, this case is equivalent to the number of overpartitions that fit inside of a $(k - 1) \times (n - k - 1)$ box.

These cases are shown in [Figure 1](#).

Hence, the three disjoint cases of the recurrence relation cover all possible cases of overpartitions that can fit in a $k \times (n - k)$ box.

To be useful when verifying the generating function in question, (2) must be written in terms of q . That is, let $G(n, k, q)$ be the generating function for the




Figure 1. Illustration of the three cases for the recurrence relation. The dimensions of all of the boxes are $k \times (n - k)$. Black denotes a fixed part, gray denotes that the portion can be filled with all possible overpartitions, and white corresponds to empty space. Left: $\text{lp}(\lambda) < n - k$. Middle: $\text{lp}(\lambda) = n - k$ and $\text{lp}(\lambda)$ is not overlined. Right: $\text{lp}(\lambda) = n - k$ and $\text{lp}(\lambda)$ is overlined.

number of overpartitions that fit in a $k \times (n - k)$ box. Then,

$$G(n, k, q) = G(n - 1, k, q) + q^{n-k}G(n - 1, k - 1, q) + q^{n-k}G(n - 2, k - 1, q).$$

In the case of overpartitions in $2 \times (n - 1)$ box, we have the recurrence relation

$$G(n + 1, 2, q) = G(n, 2, q) + q^{n-1}G(n, 1, q) + q^{n-1}G(n - 1, 1, q). \tag{3}$$

We now give the generating function.

Lemma 3. *Let n be a positive integer and $|q| < 1$. Then $f(q) = (2q + 2q^2) \begin{bmatrix} n \\ 2 \end{bmatrix}_q + 1$ is the generating function for overpartitions that fit inside of a $2 \times (n - 1)$ box.*


It can be shown that $f(q)$ satisfies (3); therefore, Lemma 3 holds.

3.2. The q -analogue of Delannoy numbers. Now that we have verified our generating function for the number of overpartitions in a $2 \times (n - 1)$ box, we will draw a connection between Lemma 3 and the Delannoy numbers.

Definition 4. Let m, n be positive integers. The Delannoy numbers $D(m, n)$ are the number of lattice paths from $(0, 0)$ to (m, n) in which only east, north, and northeast steps are allowed.

It is easy to see that when we consider the cells above the path drawn from $(0, 0)$ to (m, n) as a Ferrers shape, the Delannoy numbers are equal to the number of overpartitions that fit inside of a $m \times n$ box. Note that in this model, the northeast steps correspond to overlined, and thus shaded, cells.

Example 5. Consider a 2×4 box. The following is a lattice path in this box:




This lattice path corresponds to the Ferrers shape



and thus the overpartition $\bar{4}3$.

Example 6. Consider another lattice path in a 2×4 box:



This lattice path corresponds to the Ferrers shape



and thus the partition 31.

Lemma 7. Let n be a positive integer and $|q| < 1$. The generating function for overpartitions that fit inside a $2 \times (n - 1)$ box, $g(q) = (2q + 2q^2) \left[\begin{smallmatrix} n+1 \\ 2 \end{smallmatrix} \right]_q + 1$, is a q -analogue of the Delannoy numbers, $D(2, n - 1)$.

Proof. As per the definition of a q -analogue, we first take the limit as $q \rightarrow 1^-$ of $g(q)$ to find the expression that our generating function generalizes in terms of q . Therefore, we see

$$\lim_{q \rightarrow 1^-} (2q + 2q^2) \left[\begin{smallmatrix} n \\ 2 \end{smallmatrix} \right]_q + 1 = 4 \binom{n}{2} + 1 = 2n(n - 1) + 1.$$

Next, we must show that this result, $2n(n - 1) + 1$, is indeed the expression for the Delannoy numbers $D(2, n - 1)$. According to [Pan 2015], a formula for the Delannoy numbers is

$$D(n, k) = \sum_{d=0}^n 2^d \binom{k}{d} \binom{n}{d}.$$

Therefore, in this case, we have

$$D(2, n - 1) = \sum_{d=0}^2 \binom{n-1}{d} \binom{2}{d},$$

which readily simplifies to

$$D(2, n - 1) = 2n(n - 1) + 1.$$

Ergo, we have equality and the generating function $g(q) = (2q + 2q^2) \left[\begin{smallmatrix} n+1 \\ 2 \end{smallmatrix} \right]_q + 1$ is a q -analogue of the Delannoy numbers $D(2, n - 1)$. \square

4. A q -analogue of the sum of cubes

In [Garrett and Hummel 2004], the authors give a q -analogue of the sum of cubes and a bijective proof using partitions. We give another q -analogue of the sum of cubes and provide a bijective proof with a similar method, but using overpartitions.

Theorem 8. *Let n be a positive integer and let $|q| < 1$. Then,*

$$\sum_{i=1}^n 2q^{i-1} \left(\frac{1-q^{i-1}}{1-q} \right)^2 \left(\left(\frac{1-q^{i-2}}{1-q} \right) + \left(\frac{1-q^i}{1-q} \right) \right) = (2q + 2q^2) \left[\begin{matrix} n \\ 2 \end{matrix} \right]_q. \tag{4}$$

Note that, taking the limit as $q \rightarrow 1^-$, we obtain

$$\sum_{i=1}^n i^3 = \binom{n+1}{2}^2.$$

Thus, the above theorem is a q -analogue of the sum of cubes.

Bijective proof. We will prove Theorem 8 by interpreting the terms combinatorially and finding a weight-preserving bijection between two sets of overpartitions. Let R be a set of pairs of overpartitions, (λ, μ) , where λ is a nonempty overpartition that fits inside a $2 \times (n - 1)$ box and μ is a partition that fits inside a $2 \times (n - 2)$ box. It follows that $f(q) = \sum_{(\lambda, \mu) \in R} q^{|\lambda|+|\mu|}$ is a generating function for R and is equal to the right-hand side of (4).

Given a positive integer n , let L be a set of tuples, $(v, a, b) \cup (v, a, b')$, where the allowed values of v, a, b , and b' are:

- v is an overpartition into two parts, where the largest part is equal to at most $n - 1$ and can be overlined and the smallest part is at most $n - 2$ and cannot be overlined.
- $0 \leq a \leq n - 2$.
- $0' \leq b' \leq (n - 3)'$.
- $0 \leq b \leq n - 1$.

Let $\ell = (v, a, b) \in L$. Then $g(q) = \sum_{\ell \in L} q^{|\ell|}$, where $|\ell| = |v| + a + b$ is a generating function for L and is equal to the left-hand side.

We will now define a bijection between the finite sets R and L . Then, we can show that $f(q) = g(q)$; therefore, (4) holds. So, let $\phi : R \rightarrow L$, where $\phi(\lambda, \mu) = (v, a, b)$ and define ϕ in cases:

Case 1: $\lambda_1 > \mu_1$.

- (a) $\lambda_2 \neq 0$, and λ_2 is not overlined.
 - (i) If λ_1 is not overlined, then $\phi(\lambda, \mu) = ((\lambda_1)(\lambda_2 - 1), \mu_2, \mu_1 + 1)$.
 - (ii) If λ_1 is overlined, then $\phi(\lambda, \mu) = ((\overline{\lambda_1})(\lambda_2 - 1), \mu_2, \mu_1 + 1)$.

(b) λ_2 is overlined or $\lambda_2 = 0$.

(i) If λ_1 is not overlined, then $\phi(\lambda, \mu) = ((\lambda_1)(\lambda_2), \mu_1, \mu_2)$.

(ii) If λ_1 is overlined, then $\phi(\lambda, \mu) = ((\overline{\lambda_1})(\lambda_2), \mu_1, \mu_2)$.

Case 2: $\lambda_1 \leq \mu_1$.

(a) λ_2 is not overlined.

(i) If λ_1 is not overlined, then $\phi(\lambda, \mu) = ((\mu_1 + 1)(\mu_2), \lambda_2, (\lambda_1 - 1)')$.

(ii) If λ_1 is overlined, then $\phi(\lambda, \mu) = ((\overline{\mu_1 + 1})(\mu_2), \lambda_2, (\lambda_1 - 1)')$.

(b) λ_2 is overlined.

(i) If λ_1 is not overlined, then $\phi(\lambda, \mu) = ((\mu_1 + 1)(\mu_2), \lambda_1, (\lambda_2 - 1)')$.

(ii) If λ_1 overlined, then $\phi(\lambda, \mu) = ((\overline{\mu_1 + 1})(\mu_2), \lambda_1, (\lambda_2 - 1)')$.

To prove that ϕ is a bijection, one can show that it is one-to-one and onto. It is easier, however, to construct its inverse. We can define $\phi^{-1} : L \rightarrow R$ by the following cases, starting with the case of whether b is primed or not primed.

Case 1: b is not primed.

(a) $a \geq b$.

(i) If v_1 is not overlined, then $\phi^{-1}(v, a, b) = (v, (a)(b))$.

(ii) If v_1 is overlined, then $\phi^{-1}(v, a, b) = (v, (a)(b))$.

(b) $a < b$.

(i) If v_1 is not overlined, then $\phi^{-1}(v, a, b) = ((v_1)(v_2 + 1), (b - 1)(a))$.

(ii) If v_1 is overlined, then $\phi^{-1}(v, a, b) = ((\overline{v_1})(v_2 + 1), (b - 1)(a))$.

Case 2: b is primed.

(a) $a \geq b + 2$.

(i) If v_1 is not overlined, then $\phi^{-1}(v, a, b) = ((a)(b + 1), (v_1 - 1)(v_2))$.

(ii) If v_1 is overlined, then $\phi^{-1}(v, a, b) = ((\overline{a})(b + 1), (v_1 - 1)(v_2))$.

(b) $a < b + 2$.

(i) If v_1 is not overlined, then $\phi^{-1}(v, a, b) = ((b + 1)(a), (v_1 - 1)(v_2))$.

(ii) If v_1 is overlined, then $\phi^{-1}(v, a, b) = ((\overline{b + 1})(a), (v_1 - 1)(v_2))$.

The details of verifying that ϕ and ϕ^{-1} are inverses are not hard and are left to the reader. However, we will conclude the combinatorial proof with two examples of ϕ and ϕ^{-1} to help make the bijection clearer.

Example 9. Let $(\lambda, \mu) = (54, 22)$. First, we find i . We have $\lambda_1 > \mu_1$, so $i = 5 + 1 = 6$. For ϕ , we are in **Case 1(a)(i)**, so $\phi(\lambda, \mu) = ((\lambda_1)(\lambda_2 - 1), \mu_2, \mu_1 + 1)$. Therefore, $\phi(54, 22) = (53, 2, 3)$. Note that $|\lambda| + |\mu| = 9 + 4 = 13$ and $|v| + a + b = 8 + 2 + 3 = 13$. Next, we act on (v, a, b) with the inverse. We are in **Case 1(b)(i)**, so $\phi^{-1}(v, a, b) = ((v_1)(v_2 + 1), (b - 1)(a))$. So, $\phi^{-1}(53, 2, 3) = (54, 22)$.

Example 10. Let $(\lambda, \mu) = (\bar{2}1, 41)$. First, we find i . We have $\lambda_1 \leq \mu_1$, so $i = 4 + 2 = 6$. For ϕ , we are in [Case 2\(a\)\(ii\)](#), so $\phi(\lambda, \mu) = ((\overline{\mu_1 + 1})(\mu_2), \lambda_2, (\lambda_1 - 1)')$. Therefore, $\phi(\bar{2}1, 41) = (\bar{5}1, 1, 1')$. Note that $|\lambda| + |\mu| = 3 + 5 = 8$ and $|v| + a + b = 6 + 1 + 1 = 8$. Next, we act on (v, a, b) with the inverse. We are in [Case 2\(b\)\(ii\)](#), so $\phi^{-1}(v, a, b) = ((\overline{b + 1})(a), (v_1 - 1)(v_2))$. So, $\phi^{-1}(\bar{5}1, 1, 1') = (\bar{2}1, 41)$.

5. Conclusion

Although the specific case of overpartitions whose Ferrers shape fits in a $2 \times (n - 1)$ box is central to the proof presented here, extending this idea to the general case of a $k \times (n - k)$ box would be useful. This general work could lead to q -analogues of other expressions. In particular, investigating q -analogues for the sums of other integer powers is a natural extension of our work.

6. Acknowledgments

The student authors thank Professor Garrett for directing our research, our classmates, especially Matthew Johnson, for making us aware of the Delannoy numbers and their connection to our problem, and Professor Diveris for help with L^AT_EX.

References

- [Andrews 1976] G. E. Andrews, *The theory of partitions*, Addison-Wesley, Reading, MA, 1976. [MR](#) [Zbl](#)
- [Benjamin and Orrison 2002] A. T. Benjamin and M. E. Orrison, “Two quick combinatorial proofs”, *Coll. Math. J.* **33**:5 (2002), 406–408.
- [Corteel and Lovejoy 2004] S. Corteel and J. Lovejoy, “Overpartitions”, *Trans. Amer. Math. Soc.* **356**:4 (2004), 1623–1635. [MR](#) [Zbl](#)
- [Garrett and Hummel 2004] K. C. Garrett and K. Hummel, “A combinatorial proof of the sum of q -cubes”, *Electron. J. Combin.* **11**:1 (2004), Research Paper 9, 6 pp. [MR](#) [Zbl](#)
- [Pan 2015] H. Pan, “A Lucas-type congruence for q -Delannoy numbers”, preprint, 2015. [arXiv](#)

Received: 2016-03-14 Accepted: 2016-07-11

forsterj@stolaf.edu	<i>Department of Mathematics, Statistics and Computer Science, St. Olaf College, Northfield, MN 55057, United States</i>
garrettk@stolaf.edu	<i>Department of Mathematics, Statistics and Computer Science, St. Olaf College, Northfield, MN 55057, United States</i>
jacobsel@stolaf.edu	<i>St. Olaf College, Northfield, MN 55057, United States</i>
acw8794@gmail.com	<i>Department of Mathematics, Statistics and Computer Science, St. Olaf College, Northfield, MN 55057, United States</i>

INVOLVE YOUR STUDENTS IN RESEARCH

Involve showcases and encourages high-quality mathematical research involving students from all academic levels. The editorial board consists of mathematical scientists committed to nurturing student participation in research. Bridging the gap between the extremes of purely undergraduate research journals and mainstream research journals, *Involve* provides a venue to mathematicians wishing to encourage the creative involvement of students.

MANAGING EDITOR

Kenneth S. Berenhaut Wake Forest University, USA

BOARD OF EDITORS

Colin Adams	Williams College, USA	Suzanne Lenhart	University of Tennessee, USA
John V. Baxley	Wake Forest University, NC, USA	Chi-Kwong Li	College of William and Mary, USA
Arthur T. Benjamin	Harvey Mudd College, USA	Robert B. Lund	Clemson University, USA
Martin Bohner	Missouri U of Science and Technology, USA	Gaven J. Martin	Massey University, New Zealand
Nigel Boston	University of Wisconsin, USA	Mary Meyer	Colorado State University, USA
Amarjit S. Budhiraja	U of North Carolina, Chapel Hill, USA	Emil Minchev	Ruse, Bulgaria
Pietro Cerone	La Trobe University, Australia	Frank Morgan	Williams College, USA
Scott Chapman	Sam Houston State University, USA	Mohammad Sal Moslehian	Ferdowsi University of Mashhad, Iran
Joshua N. Cooper	University of South Carolina, USA	Zuhair Nashed	University of Central Florida, USA
Jem N. Corcoran	University of Colorado, USA	Ken Ono	Emory University, USA
Toka Diagana	Howard University, USA	Timothy E. O'Brien	Loyola University Chicago, USA
Michael Dorff	Brigham Young University, USA	Joseph O'Rourke	Smith College, USA
Sever S. Dragomir	Victoria University, Australia	Yuval Peres	Microsoft Research, USA
Behrouz Emamizadeh	The Petroleum Institute, UAE	Y.-F. S. Pétermann	Université de Genève, Switzerland
Joel Foisy	SUNY Potsdam, USA	Robert J. Plemmons	Wake Forest University, USA
Erin W. Fulp	Wake Forest University, USA	Carl B. Pomerance	Dartmouth College, USA
Joseph Gallian	University of Minnesota Duluth, USA	Vadim Ponomarenko	San Diego State University, USA
Stephan R. Garcia	Pomona College, USA	Bjorn Poonen	UC Berkeley, USA
Anant Godbole	East Tennessee State University, USA	James Propp	U Mass Lowell, USA
Ron Gould	Emory University, USA	József H. Przytycki	George Washington University, USA
Andrew Granville	Université Montréal, Canada	Richard Rebarber	University of Nebraska, USA
Jerrold Griggs	University of South Carolina, USA	Robert W. Robinson	University of Georgia, USA
Sat Gupta	U of North Carolina, Greensboro, USA	Filip Saidak	U of North Carolina, Greensboro, USA
Jim Haglund	University of Pennsylvania, USA	James A. Sellers	Penn State University, USA
Johnny Henderson	Baylor University, USA	Andrew J. Sterge	Honorary Editor
Jim Hoste	Pitzer College, USA	Ann Trenk	Wellesley College, USA
Natalia Hritonenko	Prairie View A&M University, USA	Ravi Vakil	Stanford University, USA
Glenn H. Hurlbert	Arizona State University, USA	Antonia Vecchio	Consiglio Nazionale delle Ricerche, Italy
Charles R. Johnson	College of William and Mary, USA	Ram U. Verma	University of Toledo, USA
K. B. Kulasekera	Clemson University, USA	John C. Wierman	Johns Hopkins University, USA
Gerry Ladas	University of Rhode Island, USA	Michael E. Zieve	University of Michigan, USA

PRODUCTION

Silvio Levy, Scientific Editor

Cover: Alex Scorpan

See inside back cover or msp.org/involve for submission instructions. The subscription price for 2017 is US \$175/year for the electronic version, and \$235/year (+\$35, if shipping outside the US) for print and electronic. Subscriptions, requests for back issues from the last three years and changes of subscribers address should be sent to MSP.

Involve (ISSN 1944-4184 electronic, 1944-4176 printed) at Mathematical Sciences Publishers, 798 Evans Hall #3840, c/o University of California, Berkeley, CA 94720-3840, is published continuously online. Periodical rate postage paid at Berkeley, CA 94704, and additional mailing offices.

Involve peer review and production are managed by EditFlow® from Mathematical Sciences Publishers.

PUBLISHED BY

 **mathematical sciences publishers**
nonprofit scientific publishing

<http://msp.org/>

© 2017 Mathematical Sciences Publishers

involve

2017

vol. 10

no. 3

Dynamics of vertical real rhombic Weierstrass elliptic functions	361
LORELEI KOSS AND KATIE ROY	
Pattern avoidance in double lists	379
CHARLES CRATTY, SAMUEL ERICKSON, FREHIWET NEGASSI AND LARA PUDWELL	
On a randomly accelerated particle	399
MICHELLE NUNO AND JUHI JANG	
Reeb dynamics of the link of the A_n singularity	417
LEONARDO ABBRESCIA, IRIT HUQ-KURUVILLA, JO NELSON AND NAWAZ SULTANI	
The vibration spectrum of two Euler–Bernoulli beams coupled via a dissipative joint	443
CHRIS ABRIOLA, MATTHEW P. COLEMAN, AGLIKA DARAKCHIEVA AND TYLER WALES	
Loxodromes on hypersurfaces of revolution	465
JACOB BLACKWOOD, ADAM DUKEHART AND MOHAMMAD JAVAHERI	
Existence of positive solutions for an approximation of stationary mean-field games	473
NOJOD ALMAYOUF, ELENA BACHINI, ANDREIA CHAPOUTO, RITA FERREIRA, DIOGO GOMES, DANIELA JORDÃO, DAVID EVANGELISTA JUNIOR, AVETIK KARAGULYAN, JUAN MONASTERIO, LEVON NURBEKYAN, GIORGIA PAGLIAR, MARCO PICCIRILLI, SAGAR PRATAPSI, MARIANA PRAZERES, JOÃO REIS, ANDRÉ RODRIGUES, ORLANDO ROMERO, MARIA SARGSYAN, TOMMASO SENECCI, CHULIANG SONG, KENGO TERAJ, RYOTA TOMISAKI, HECTOR VELASCO-PEREZ, VARDAN VOSKANYAN AND XIANJIN YANG	
Discrete dynamics of contractions on graphs	495
OLENA OSTAPYUK AND MARK RONNENBERG	
Tiling annular regions with skew and T-tetrominoes	505
AMANDA BRIGHT, GREGORY J. CLARK, CHARLES LUNDON, KYLE EVITTS, MICHAEL P. HITCHMAN, BRIAN KEATING AND BRIAN WHETTER	
A bijective proof of a q -analogue of the sum of cubes using overpartitions	523
JACOB FORSTER, KRISTINA GARRETT, LUKE JACOBSEN AND ADAM WOOD	
Ulrich partitions for two-step flag varieties	531
IZZET COSKUN AND LUKE JASKOWIAK	