

# involve

a journal of mathematics

Leverage centrality of knight's graphs  
and Cartesian products of regular graphs  
and path powers

Roger Vargas, Jr., Abigail Waldron, Anika Sharma,  
Rigoberto Flórez and Darren A. Narayan





# Leverage centrality of knight's graphs and Cartesian products of regular graphs and path powers

Roger Vargas, Jr., Abigail Waldron, Anika Sharma,  
Rigoberto Flórez and Darren A. Narayan

(Communicated by Kenneth S. Berenhaut)

In 2010, Joyce et al. defined the leverage centrality of a graph as a means to analyze connections within the brain. In this paper we investigate this property from a mathematical perspective and determine the leverage centrality for knight's graphs, path powers, and Cartesian products.

## 1. Introduction

We recall that the degree of a vertex  $v$  is the number of edges incident to  $v$  and is denoted  $\deg v$ . Joyce, Laurienti, Burdette, and Hayasaka [Joyce et al. 2010] defined the property of leverage centrality based on vertex degrees.

**Definition 1.** Leverage centrality is a measure of the relationship between the degree of a given node  $v$  and the degree of each of its neighbors  $v_i$ , averaged over all neighbors of  $v$ , denoted  $N_v$ , and is defined as


$$l(v) = \frac{1}{\deg v} \sum_{v_i \in N_v} \frac{\deg v - \deg v_i}{\deg v + \deg v_i}.$$

This property was used by Joyce et al. [2010] in the analysis of functional magnetic resonance imaging (fMRI) data and has been used to analyze real-world networks including airline connections, electrical power grids, and coauthorship collaborations [Li et al. 2015]. The leverage centralities of complete multipartite graphs and the Cartesian product of paths were investigated by Sharma, Vargas, Waldron, Flórez, and Narayan [Sharma et al. 2017]. Bounds on leverage centrality were determined by Li, Li, Van Mieghem, Stanley, and Wang [Li et al. 2015]. We restate one of their results as our first theorem.

---

*MSC2010:* 05C07.

*Keywords:* leverage centrality, knight's graphs.



**Figure 1.** The  $3 \times 3$ ,  $4 \times 4$ ,  $5 \times 5$  and  $6 \times 6$  knight's graphs.

**Theorem 2.** For any vertex  $v$ , we have  $|l(v)| \leq 1 - \frac{2}{n}$ . Furthermore, these bounds are tight in the cases of stars and complete graphs.

We note that the bounds are also tight for regular graphs with degree  $r > 1$ .

In this paper we investigate leverage centrality for various families of graphs including the knight's graphs, path powers, and the Cartesian products of graphs.

### 2. Leverage centrality of a knight's graph


We define an  $n \times n$  knight's graph to be the graph with  $n^2$  vertices in which every vertex represents a square in an  $n \times n$  chessboard. The vertices on the  $n \times n$  chessboard can be placed in an  $n \times n$  table where two vertices  $v_i$  and  $v_j$  are adjacent if they are exactly four entries apart (including the entries of  $v_i$  and  $v_j$ ) and they form an "L" shape. We give examples of knight's graphs of small order in Figure 1, where in each graph all of the vertices of same degree are the same color.

We next state the leverage centrality of each vertex in the  $n \times n$  knight's graph. We use  $t_j$  to denote the  $j$ -th triangular number  $\frac{1}{2}j(j + 1)$ .

**Theorem 3.** Let  $G_n$  be the  $n \times n$  knight's graph.

- (1) The leverage centrality of every vertex of  $G_3$  is zero.
- (2) If  $n = 4, 6$ , or  $8$ , then  $G_n$  has exactly  $t_{n/2}$  distinct leverage centralities.
- (3) If  $n = 5$  or  $7$ , then  $G_n$  has exactly  $t_{(n+1)/2} - 1$  distinct leverage centralities.
- (4) If  $n \geq 9$ , then  $G_n$  has exactly 15 distinct leverage centralities.

*Proof.* We first find the degree of each vertex in the knight's graph on an  $n \times n$  chessboard, where  $n \geq 3$ . To describe the degree of each vertex in the graph  $G_n$ , we will arrange the vertices of  $G_n$  in an  $n \times n$  table. The vertices corresponding to entries  $(1, 1)$ ,  $(1, n)$ ,  $(n, 1)$ , and  $(n, n)$  have degree 2. Those corresponding to entries  $(1, 2)$ ,  $(1, n - 1)$ ,  $(2, 1)$ ,  $(2, n)$ ,  $(n - 1, 1)$ ,  $(n, 2)$ ,  $(n - 1, n)$ , and  $(n, n - 1)$  have degree 3. Those corresponding to entries  $(2, 2)$ ,  $(2, n - 1)$ ,  $(n - 1, 2)$ ,  $(n - 1, n - 1)$  and  $(1, i)$ ,  $(i, 1)$ ,  $(n, i)$ , and  $(i, n)$ , where  $i = 3, 4, \dots, n - 2$ , have degree 4. Those corresponding to entries  $(2, i)$ ,  $(i, 2)$ ,  $(n - 1, i)$ , and  $(i, n - 1)$ , where  $i = 3, 4, \dots, n - 2$ , have degree 6. Vertices corresponding to entries  $(i, j)$ , where  $i = 3, 4, \dots, n - 3$  and  $j = 3, 4, \dots, n - 2$ , have degree 8; see, for example, Figure 2 (left).



**Figure 2.** The  $10 \times 10$  knight's graph (left) and the  $9 \times 9$  knight's graph (right).

If  $n$  is even we subdivide the knight's graph's vertical and horizontal axes and the two diagonals to obtain eight regions. Each region forms a right triangle where the legs have  $\frac{1}{2}n$  vertices; see, for example, Figure 2 (left). Using symmetry we can calculate the leverage centrality of all vertices by only analyzing a single triangle.

If  $n$  is odd, as in Figure 2 (right), we do the same subdivision; however, in this case two adjacent triangles will overlap—the legs of the right triangle will have  $\frac{1}{2}(n + 1)$  vertices.

We choose the triangle with vertices

$$\begin{aligned}
 v_1 &= (1, 1), & v_2 &= (1, 2), & v_3 &= (2, 2), & v_4 &= (1, 3), & v_5 &= (2, 3), \\
 v_6 &= (3, 3), & v_7 &= (1, 4), & v_8 &= (2, 4), & v_9 &= (3, 4), & v_{10} &= (4, 4), & (2-1) \\
 v_{11} &= (1, 5), & v_{12} &= (2, 5), & v_{13} &= (3, 5), & v_{14} &= (4, 5), & v_{15} &= (5, 5).
 \end{aligned}$$

Note that if  $n < 10$ , we take triangles with vertices  $v_i$  for  $i = 1, 2, \dots, k$ , where  $k = \frac{1}{2}n$  if  $n = 2(k)$  or  $k = \frac{1}{2}(n + 1)$  if  $n = 2(k) - 1$ .

**Proof of (1).** Since  $G_3$  is regular, the leverage centrality of all of its vertices is 0.

**Proof of (2).** For case  $n = 4$ , it is easy to see that  $l(v_1) = -\frac{1}{3}$ ,  $l(v_2) = -\frac{1}{21}$ , and  $l(v_3) = \frac{5}{21}$ .

Now consider the cases  $n = 6$  and  $n = 8$ . From the above analysis we only need to calculate the leverage centrality for a triangle with legs that have  $\frac{1}{2}n$  vertices (see Figure 2 (left) for an example of those triangles). Thus, to calculate the leverage centrality of these special cases, we consider the triangle with vertices  $v_1, \dots, v_{t_i}$ , where  $t_i$  is the  $i$ -th triangular number where  $i = 1, 2, \dots, \frac{1}{2}n$ , and then use Tables 1 and 3, respectively.

vertex $v_i$	deg $v_i$	AD( $v_i$ )	$l(v_i)$
$v_1$	2	6, 6	$-1/2$
$v_2$	3	4, 6, 8	$-215/693$
$v_3$	4	4, 4, 8, 8	$-1/6$
$v_4$	4	3, 6, 4, 8	$-41/420$
$v_5$	6	2, 3, 4, 6, 6, 8	$187/1260$
$v_6$	8	3, 3, 4, 4, 4, 4, 6, 6	$73/231$

**Table 1.** Leverage centrality when  $n = 6$ . Here AD( $v_i$ ) denotes the degrees of vertices adjacent to  $v_i$ .

vertex $v_i$	deg $v_i$	AD( $v_i$ )	$l(v_i)$
$v_1$	2	6, 6	$-1/2$
$v_2$	3	4, 6, 8	$-215/693$
$v_3$	4	4, 4, 8, 8	$-1/6$
$v_4$	4	3, 6, 6, 8	$-31/210$
$v_5$	6	2, 4, 4, 6, 8, 8	$43/420$
$v_6$	8	3, 3, 4, 4, 6, 6, 8, 8	$215/924$
$v_7$	4	4, 4, 8, 8	$-1/6$
$v_8$	6	3, 3, 6, 6, 8, 8	$4/63$
$v_9$	8	4, 4, 4, 4, 6, 6, 8, 8	$17/84$
$v_{10}$	8	6, 6, 6, 6, 6, 6, 6, 6	$1/7$

**Table 2.** Leverage centrality when  $n = 7$ . Here AD( $v_i$ ) denotes the degrees of vertices adjacent to  $v_i$ .

**Proof of (3).** First consider the case  $n = 5$ . From the above analysis we only need to calculate the leverage centrality for a triangle with legs that have three vertices. It is easy to see that  $l(v_1) = -\frac{1}{2}, l(v_2) = -\frac{19}{77}, l(v_3) = -\frac{1}{35}, l(v_4) = -\frac{1}{35}, l(v_5) = \frac{3}{10}$ , and  $l(v_6) = \frac{5}{11}$ . This shows that there are only five distinct leverage centralities in  $G_5$ .

Now consider the case  $n = 7$ . From the above analysis we only need to calculate the leverage centrality for a triangle with legs that have four vertices. From Table 2 we can see that  $G_7$  has only nine distinct leverage centralities.

**Proof of (4).** For  $n = 8$  and  $n = 10$ , the proof is similar to those of parts (2) and (3).

We now suppose  $n > 10$ . Consider the 15 vertices in the triangle given in (2-1) and their relevant data, given in Table 3.

The analysis for the remaining vertices in the triangle is as follows. From the definition of the knight’s graph we know that if two vertices  $v_i$  and  $v_j$  are adjacent, then they are four entries apart (including the entries of  $v_i$  and  $v_j$ ) and they form an “L” shape. This implies that if  $n \geq 11$  then the leverage centrality of every vertex located

$n$	vertex $v_i$	deg $v_i$	AD( $v_i$ )	$l(v_i)$
8, 9, 10	$v_1$	2	6, 6	$-1/2$
8, 9, 10	$v_2$	3	4, 6, 8	$-215/693$
8, 9, 10	$v_3$	4	4, 4, 8, 8	$-1/6$
8, 9, 10	$v_4$	4	3, 6, 6, 8	$-31/210$
8, 9, 10	$v_5$	6	2, 4, 4, 6, 8, 8	$43/420$
8, 9, 10	$v_6$	8	3, 3, 4, 4, 6, 6, 8, 8	$215/924$
8, 9, 10	$v_7$	4	4, 6, 8, 8	$-13/60$
8, 9, 10	$v_8$	6	3, 4, 6, 8, 8, 8	$11/630$
8, 9, 10	$v_9$	8	4, 4, 4, 6, 6, 8, 8, 8	$9/56$
8, 9, 10	$v_{10}$	8	6, 6, 6, 6, 8, 8, 8, 8	$1/14$
9, 10	$v_{11}$	4	6, 6, 8, 8	$-4/45$
9, 10	$v_{12}$	6	4, 4, 8, 8, 8, 8	$-1/35$
9, 10	$v_{13}$	8	4, 4, 6, 6, 8, 8, 8, 8	$5/42$
9, 10	$v_{14}$	8	6, 6, 8, 8, 8, 8, 8, 8	$1/28$
9, 10	$v_{15}$	8	8, 8, 8, 8, 8, 8, 8, 8	0

**Table 3.** Leverage centrality with  $n = 8, 9, 10$ . Here  $AD(v_i)$  denotes the degrees of vertices adjacent to  $v_i$ .

in entries  $(r, t)$  is zero for  $r = 5, 6, \dots, k$  and  $t = 5, 6, \dots, k$ , where  $k = \frac{1}{2}n$  if  $n = 2k$  or  $k = \frac{1}{2}(n + 1)$  if  $n = 2k - 1$ . Moreover, every vertex located in position  $(i, j)$  will have the same leverage centrality as the vertices located in entries  $(i, 6)$ , where  $i = 1, 2, \dots, k$  and  $j = 7, \dots, k$ , where  $k = \frac{1}{2}n$  if  $n = 2k$  or  $k = \frac{1}{2}(n + 1)$  if  $n = 2k - 1$ .  $\square$

### 3. Leverage centralities of $P_n^k$

Let  $P_n^k$  be the graph with vertices  $v_1, v_2, \dots, v_n$  and edges  $(v_i, v_j)$  when  $1 \leq |i - j| \leq k \leq n - 1$ . In this paper we will assume  $n > 1$ . This family contains both paths (when  $k = 1$ ) and complete graphs (when  $k = n - 1$ ). Note that  $\text{deg } v_i = \min\{i + k - 1, 2k\}$ . The neighbors of  $v_i$  are  $v_{i-1}, v_{i-2}, \dots, v_{i-s}$  and  $v_{i+1}, v_{i+2}, \dots, v_{i+t}$ , where  $s = \min\{k, i - 1\}$  and  $t = \min\{k, n - i\}$ . The above conditions can be combined in the next lemma to give the leverage centrality of any vertex in  $P_n^k$ .

**Lemma 4.** *Suppose the vertex  $v_i \in V(P_n^k)$  has neighbors  $v_{i-1}, v_{i-2}, \dots, v_{i-s}$  and  $v_{i+1}, v_{i+2}, \dots, v_{i+t}$ , where  $s = \min\{k, i - 1\}$  and  $t = \min\{k, n - i\}$ . Then*

$$l(v_i) = \frac{1}{\delta_i} \sum_{i-s \leq j \leq i+t} \frac{\delta_i - \delta_j}{\delta_i + \delta_j},$$

where  $\delta_x = \min\{x + k - 1, 2k\}$  for  $x = i, j$ .

We begin by determining the leverage centrality of vertices in a path  $P_n$ , where  $n \geq 2$ . We note that by symmetry  $l(v_i) = l(v_{n+1-i})$  for all  $1 \leq i \leq n$ . We start

with small values of  $n$ . When  $n = 2$ , both vertices have a leverage centrality of zero. When  $n = 3$ , the two vertices of degree 1 have leverage centrality  $\frac{1-2}{1+2} = -\frac{1}{3}$  and the vertex of degree 2 has leverage centrality  $\frac{2-1}{1+2} = \frac{1}{3}$ . When  $n = 4$ , the two vertices of degree 1 have leverage centrality  $\frac{1-2}{1+2} = -\frac{1}{3}$  and the two vertices of degree 2 have leverage centrality  $\frac{1}{2}(\frac{2-1}{1+2} + \frac{2-2}{2+2}) = \frac{1}{6}$ .

Next, we use the operation of edge subdivision to handle cases where  $n \geq 5$ . Recall that in an edge subdivision an edge  $u - v$  is replaced by a path on three vertices  $u - w - v$ . We note that if we extend the length of a path by subdividing the edge between vertices  $c$  and  $d$ , the new vertices will have a leverage centrality of zero. Further subdivision of an edge connecting two vertices with degree 2 will include a new vertex with leverage centrality zero. Hence, there will be exactly three distinct leverage centralities in any path with five or more vertices. The general result follows.

**Theorem 5.** *Let  $P_n$  be a path where  $n \geq 5$ . Then  $l(v_1) = l(v_n) = -\frac{1}{3}$ ,  $l(v_2) = l(v_{n-1}) = \frac{1}{6}$ , and for all  $3 \leq i \leq n - 2$ , we have  $l(v_i) = 0$ .*

**3.1. Leverage centralities of  $P_n^2$ .** We now calculate the leverage centralities for paths  $P_n^2$ . Again by symmetry, we have  $l(v_i) = l(v_{n+1-i})$  for all  $1 \leq i \leq n$ .

- $n = 3$ : For  $1 \leq i \leq 3$ , we have  $l(v_i) = 0$ .
- $n = 4$ :  $l(v_1) = -\frac{4}{15}$  and  $l(v_2) = \frac{2}{15}$ .
- $n = 5$ :  $l(v_1) = -\frac{4}{15}$ ,  $l(v_2) = \frac{2}{105}$  and  $l(v_3) = \frac{5}{21}$ .
- $n = 6$ :  $l(v_1) = -\frac{4}{15}$ ,  $l(v_2) = -\frac{1}{35}$  and  $l(v_3) = \frac{13}{84}$ .
- $n = 7$ :  $l(v_1) = -\frac{4}{15}$ ,  $l(v_2) = -\frac{1}{35}$ ,  $l(v_3) = \frac{5}{42}$  and  $l(v_4) = \frac{3}{28}$ .
- $n = 8$ :  $l(v_1) = -\frac{4}{15}$ ,  $l(v_2) = -\frac{1}{35}$ ,  $l(v_3) = \frac{5}{42}$  and  $l(v_4) = \frac{1}{28}$ .
- $n \geq 9$ :  $l(v_1) = -\frac{4}{15}$ ,  $l(v_2) = -\frac{1}{35}$ ,  $l(v_3) = \frac{5}{42}$ ,  $l(v_4) = \frac{1}{28}$  and for all  $5 \leq i \leq n - 4$ ,  $l(v_i) = 0$ .

It is clear that to calculate the leverage centralities of all vertices in  $P_n^k$  for all  $k$  in this manner would require lengthy computation. However by noticing that the leverage centralities become fixed when  $n$  becomes large enough ( $n \geq 4k + 1$ ), we can compute the leverage centralities in a more formal manner.

First we give an elementary result with the leverage centralities for the first vertex in any path power.

**Proposition 6.** *If  $v_1 \in V(P_n^k)$ , then*

$$l(v_1) = \sum_{i=1}^k \frac{-i}{2k+i}.$$



*Proof.* The vertex  $v_1$  has  $k$  neighbors, with degrees  $k + 1, k + 2, \dots, 2k$ . Then

$$l(v_1) = \frac{1}{k} \left( k \sum_{i=1}^k \frac{-i}{2k+i} \right) = \sum_{i=1}^k \frac{-i}{2k+i}. \quad \square$$

We continue with three lemmas which will help us determine the relationships between the leverage centralities of different vertices in  $P_n^k$ .

**Lemma 7.** *If  $i$  is an integer and  $1 < \frac{1}{2}a \leq i < a$  then we have*

$$\frac{1}{a} \left( \frac{a-i}{a+i} \right) > \frac{1}{a-1} \left( \frac{a-1-i}{a-1+i} \right).$$

*Proof.* Let  $\frac{1}{2}a \leq i < a$ . This implies

$$\begin{aligned} 2ia - a^2 + (a-i) > 0 &\Rightarrow -a^2 + a + 2ia - 1 - i > 0 \\ &\Rightarrow a^3 - 2a^2 + (2+i)a - 1 - i > a^3 - a^2 + (1-i)a \\ &\Rightarrow (a-i)(a-1)(a-1+i) > a(a+i)(a-1-i) \\ &\Rightarrow \frac{1}{a} \left( \frac{a-i}{a+i} \right) > \frac{1}{a-1} \left( \frac{a-1-i}{a-1+i} \right). \quad \square \end{aligned}$$

**Lemma 8.** *For all  $1 \leq a \leq 2k$ , we have*

$$\frac{1}{a} \left( \frac{-1}{2a+1} \right) > \frac{-1}{(a-1)a}.$$

*Proof.* We first note that when  $a = 2k$ ,

$$\frac{1}{a} \left( \frac{a-(a+1)}{a+a+1} \right) > \frac{1}{a-1} \left( \frac{a-1-(a+1)}{a-1+a+1} \right)$$

is clear since the left side is positive and the right side is negative.

Let  $1 \leq a$ . Then

$$\begin{aligned} 2a^2 + a > a^2 + 1 &\Rightarrow \frac{-1}{2a^2+a} > \frac{-1}{a^2+1} \\ &\Rightarrow \frac{1}{a} \left( \frac{-1}{2a+1} \right) > \frac{1}{a+1} \left( \frac{-2}{2a} \right) \\ &\Rightarrow \frac{1}{a} \left( \frac{a-(a+1)}{a+a+1} \right) > \frac{1}{a-1} \left( \frac{a-1-(a+1)}{a-1+a+1} \right) \\ &\Rightarrow \frac{1}{a} \left( \frac{-1}{a+a+1} \right) > \frac{1}{a-1} \left( \frac{-2}{a-1+a+1} \right). \quad \square \end{aligned}$$

**Lemma 9.** *Let  $2 \leq i \leq k-1$ . Then*

$$\frac{1}{k+1} \left( \frac{1-i}{2k+1+i} \right) > \frac{1}{k} \left( \frac{-i}{2k+i} \right).$$

*Proof.* Note that

$$0 > \frac{1-i}{2k+i+1} > \frac{-i}{2k+i+1} > \frac{-i}{2k+i}.$$

Hence

$$0 > \frac{k+1-(k+i)}{k+1+k+i} > \frac{k-(k+i)}{k+k+i}.$$

Since  $\frac{1}{k+1} < \frac{1}{k}$ , we have

$$0 > \frac{1}{k+1} \left( \frac{1-i}{k+1+k+i} \right) > \frac{1}{k} \left( \frac{-i}{k+k+i} \right). \quad \square$$

Proposition 6 and Lemmas 7, 8, and 9 can be combined as follows.

**Proposition 10.** *Let  $G = P_n^k$ , where  $n \geq 4k + 1$ . Then:*

- (i)  $l(v_i) = l(v_{n+1-i})$ .
- (ii) For all  $0 \leq j \leq k - 1$ ,

$$l(v_{k+j+1}) = \frac{1}{2k} \left( \sum_{i=k+j}^{2k-1} \frac{2k-i}{2k+i} \right).$$

- (iii) For all  $0 \leq j \leq k - 1$ ,

$$l(v_{k-j}) = \frac{1}{2k-j-1} \sum_{i=k}^{2k-1} \frac{2k-j-i}{2k-j+i} + \frac{k-j}{2k-1-j} \left( \frac{2k-j-1-2k}{2k-j-1+2k} \right).$$

- (iv) For all  $2k + 1 \leq j \leq n - 2k$ , we have  $l(v_j) = 0$ .

This leads to the following theorem.

**Theorem 11.** *Let  $G = P_n^k$ , where  $n \geq 4k + 1$ . Then the vertex with the largest leverage centrality in  $G$  is  $v_{k+1}$ , and furthermore  $l(v_{k+1}) > l(v_k) > \dots > l(v_1)$  and  $l(v_{k+1}) > l(v_{k+2}) > \dots > l(v_{2k+1})$ .*

*Proof.* For the first part, we recall that

$$l(v_{k+1}) = \frac{1}{2k} \sum_{i=k}^{2k-1} \left( \frac{2k-i}{2k+i} \right) + \frac{k}{2k} \left( \frac{2k-2k}{2k+2k} \right),$$

and for  $0 \leq j \leq k - 1$ ,

$$l(v_{k-j}) = \frac{1}{2k-j-1} \sum_{i=k}^{2k-1} \frac{2k-j-i}{2k-j+i} + \frac{k-j}{2k-1-j} \left( \frac{2k-j-1-2k}{2k-j-1+2k} \right).$$

We seek to show that  $l(v_{k+1}) > l(v_k) > \dots > l(v_1)$ . When comparing terms from  $l(v_r)$  with  $l(v_{r-1})$  for a fixed  $i$ , five cases are needed to show that the  $i$ -th term of  $l(v_r)$  is larger than the  $i$ -th term of  $l(v_{r-1})$ .

**Case (i):**  $2k - j - i > 1$ . Use Lemma 7.

**Case (ii):**  $2k - j - i = 1$ . In the  $i$ -th term, the numerator is positive for the  $j$ -th term and the numerator is zero for  $(j + 1)$ -th term.

**Case (iii):**  $2k - j - i = 0$ . In the  $i$ -th term, the numerator is zero for the  $j$ -th term and the numerator is negative for  $(j + 1)$ -th term.

**Case (iv):**  $2k - j - i < 0$ . Use Lemma 8.

**Case (v):**

$$\frac{k - j}{2k - 1 - j} \left( \frac{2k - j - 1 - 2k}{2k - j - 1 + 2k} \right) > \frac{k - (j + 1)}{2k - 1 - (j + 1)} \left( \frac{2k - (j + 1) - 1 - 2k}{2k - (j + 1) - 1 + 2k} \right).$$

Use Lemma 8.

The combination of these five cases yields  $l(v_{k+1}) > l(v_k) > \dots > l(v_1)$ . For the second part we note that for  $0 \leq r \leq k - 1$ , we have  $l(v_{k+r}) > l(v_{k+r+1})$  as terms with positive value are replaced by zeros in each successive case. Hence,  $l(v_{k+1}) > l(v_{k+2}) > \dots > l(v_{2k+1})$ . We note that we have not obtained a linear ordering, but two separate linear orderings both starting with the largest leverage centrality  $l(v_{k+1})$ . □

#### 4. Cartesian product of graphs

In this next section we give some general results about the leverage centrality of the Cartesian product of graphs. These build upon results by Sharma et al. [2017].

**Definition 12.** Given a graph  $F$  with vertex set  $V(F)$  and edge set  $E(F)$ , and a graph  $H$  with vertex set  $V(H)$  and edge set  $E(H)$ , we let  $G$  define the Cartesian product of  $F$  and  $H$  to be the graph  $G = F \times H$ , which is defined as

$$\begin{aligned} V(G) &= \{(u, v) \mid u \in V(F) \text{ and } v \in V(H)\}, \\ E(G) &= \{(u_1, v_1), (u_2, v_2) \mid u_1 = u_2 \text{ and } (v_1, v_2) \in E(H), \\ &\quad \text{or } v_1 = v_2 \text{ and } (u_1, u_2) \in E(F)\}. \end{aligned}$$

We next present an elementary result from graph theory.

**Lemma 13.** *If  $G = F \times H$ , then the degree of a vertex  $(u, v)$  in  $G$  is the sum of the degrees of vertices  $u$  and  $v$ , where  $u \in V(F)$  and  $v \in V(H)$ .*

**Theorem 14.** *Let  $G$  be a graph and let  $RG_r$  be a regular graph where each vertex has degree  $r$ . Let  $u \in V(RG_r)$  and let  $v_i$  and  $v_j$  be vertices in  $G$  with degrees  $k_i$  and  $k_j$  respectively. For each vertex  $(u, v_i) \in V(RG_r \times G)$  we have*

$$l(u, v_i) = \frac{1}{r + k_i} \sum_{j \neq i} \frac{k_i - k_j}{2r + k_i + k_j}.$$

*Proof.* Consider a vertex  $(u, v_i) \in V(RG_r \times G)$ . We note that

$$\deg(u, v_i) = \deg u + \deg v_i = r + k_i.$$

Then

$$l(u, v_i) = \frac{1}{r + k_i} \sum_{j \neq i} \frac{k_i - k_j}{2r + k_i + k_j}. \quad \square$$

We conclude by posing the following problem where the graphs may not be regular.

**Problem 15.** Given graphs  $F$  and  $H$  where the leverage centralities are known for all vertices in  $F$  and  $H$ , determine the leverage centralities for all vertices in  $F \times H$ .

### Acknowledgements

We would like to thank an anonymous referee for helpful comments. Research was supported by National Science Foundation Research Experience for Undergraduates Grant #1358583.

### References

- [Joyce et al. 2010] K. E. Joyce, P. J. Laurienti, J. H. Burdette, and S. Hayasaka, “A new measure of centrality for brain networks”, *PLOS ONE* 5:8 (2010), art. id. e12200.
- [Li et al. 2015] C. Li, Q. Li, P. Van Mieghem, H. E. Stanley, and H. Wang, “Correlation between centrality metrics and their application to the opinion model”, *Eur. Phys. J. B* 88:3 (2015), art. id. 65. MR
- [Sharma et al. 2017] A. Sharma, R. Vargas, A. Waldron, R. Florez, and D. A. Narayan, “Leverage centrality of graphs”, preprint, 2017. arXiv

Received: 2015-09-09    Revised: 2016-01-04    Accepted: 2016-03-25

roger_vargas@g.harvard.edu	<i>Mathematics and Statistics, Williams College, Williamstown, MA 01267, United States</i>
awaldron@presby.edu	<i>Mathematics Department, Presbyterian College, Clinton, SC 29325, United States</i>
anikasha@buffalo.edu	<i>Department of Computer Science and Department of Mathematics, University of Buffalo, Buffalo, NY 14260, United States</i>
rigoflorez@gmail.com	<i>Department of Mathematics and Computer Science, The Citadel, Charleston, SC 29409, United States</i>
dansma@rit.edu	<i>School of Mathematical Sciences, Rochester Institute of Technology, Rochester, NY 14623, United States</i>

# involve

msp.org/involve

## INVOLVE YOUR STUDENTS IN RESEARCH

*Involve* showcases and encourages high-quality mathematical research involving students from all academic levels. The editorial board consists of mathematical scientists committed to nurturing student participation in research. Bridging the gap between the extremes of purely undergraduate research journals and mainstream research journals, *Involve* provides a venue to mathematicians wishing to encourage the creative involvement of students.

### MANAGING EDITOR

Kenneth S. Berenhaut Wake Forest University, USA

### BOARD OF EDITORS

Colin Adams	Williams College, USA	Suzanne Lenhart	University of Tennessee, USA
John V. Baxley	Wake Forest University, NC, USA	Chi-Kwong Li	College of William and Mary, USA
Arthur T. Benjamin	Harvey Mudd College, USA	Robert B. Lund	Clemson University, USA
Martin Bohner	Missouri U of Science and Technology, USA	Gaven J. Martin	Massey University, New Zealand
Nigel Boston	University of Wisconsin, USA	Mary Meyer	Colorado State University, USA
Amarjit S. Budhiraja	U of North Carolina, Chapel Hill, USA	Emil Minchev	Ruse, Bulgaria
Pietro Cerone	La Trobe University, Australia	Frank Morgan	Williams College, USA
Scott Chapman	Sam Houston State University, USA	Mohammad Sal Moslehian	Ferdowsi University of Mashhad, Iran
Joshua N. Cooper	University of South Carolina, USA	Zuhair Nashed	University of Central Florida, USA
Jem N. Corcoran	University of Colorado, USA	Ken Ono	Emory University, USA
Toka Diagana	Howard University, USA	Timothy E. O'Brien	Loyola University Chicago, USA
Michael Dorff	Brigham Young University, USA	Joseph O'Rourke	Smith College, USA
Sever S. Dragomir	Victoria University, Australia	Yuval Peres	Microsoft Research, USA
Behrouz Emamizadeh	The Petroleum Institute, UAE	Y.-F. S. Pétermann	Université de Genève, Switzerland
Joel Foisy	SUNY Potsdam, USA	Robert J. Plemmons	Wake Forest University, USA
Errin W. Fulp	Wake Forest University, USA	Carl B. Pomerance	Dartmouth College, USA
Joseph Gallian	University of Minnesota Duluth, USA	Vadim Ponomarenko	San Diego State University, USA
Stephan R. Garcia	Pomona College, USA	Bjorn Poonen	UC Berkeley, USA
Anant Godbole	East Tennessee State University, USA	James Propp	U Mass Lowell, USA
Ron Gould	Emory University, USA	József H. Przytycki	George Washington University, USA
Andrew Granville	Université Montréal, Canada	Richard Rebarber	University of Nebraska, USA
Jerold Griggs	University of South Carolina, USA	Robert W. Robinson	University of Georgia, USA
Sat Gupta	U of North Carolina, Greensboro, USA	Filip Saidak	U of North Carolina, Greensboro, USA
Jim Haglund	University of Pennsylvania, USA	James A. Sellers	Penn State University, USA
Johnny Henderson	Baylor University, USA	Andrew J. Sterge	Honorary Editor
Jim Hoste	Pitzer College, USA	Ann Trenk	Wellesley College, USA
Natalia Hritonenko	Prairie View A&M University, USA	Ravi Vakil	Stanford University, USA
Glenn H. Hurlbert	Arizona State University, USA	Antonia Vecchio	Consiglio Nazionale delle Ricerche, Italy
Charles R. Johnson	College of William and Mary, USA	Ram U. Verma	University of Toledo, USA
K. B. Kulasekera	Clemson University, USA	John C. Wierman	Johns Hopkins University, USA
Gerry Ladas	University of Rhode Island, USA	Michael E. Zieve	University of Michigan, USA

### PRODUCTION

Silvio Levy, Scientific Editor


Cover: Alex Scorpan

See inside back cover or [msp.org/involve](http://msp.org/involve) for submission instructions. The subscription price for 2017 is US \$175/year for the electronic version, and \$235/year (+\$35, if shipping outside the US) for print and electronic. Subscriptions, requests for back issues and changes of subscriber address should be sent to MSP.

*Involve* (ISSN 1944-4184 electronic, 1944-4176 printed) at Mathematical Sciences Publishers, 798 Evans Hall #3840, c/o University of California, Berkeley, CA 94720-3840, is published continuously online. Periodical rate postage paid at Berkeley, CA 94704, and additional mailing offices.

Involve peer review and production are managed by EditFLOW® from Mathematical Sciences Publishers.

PUBLISHED BY

 **mathematical sciences publishers**  
nonprofit scientific publishing

<http://msp.org/>

© 2017 Mathematical Sciences Publishers

# involve

2017 vol. 10 no. 4

New algorithms for modular inversion and representation by the form $x^2 + 3xy + y^2$	541
CHRISTINA DORAN, SHEN LU AND BARRY R. SMITH	
New approximations for the area of the Mandelbrot set	555
DANIEL BITTNER, LONG CHEONG, DANTE GATES AND HIEU D. NGUYEN	
Bases for the global Weyl modules of $\mathfrak{sl}_n$ of highest weight $m\omega_1$	573
SAMUEL CHAMBERLIN AND AMANDA CROAN	
Leverage centrality of knight's graphs and Cartesian products of regular graphs and path powers	583
ROGER VARGAS, JR., ABIGAIL WALDRON, ANIKA SHARMA, RIGOBERTO FLÓREZ AND DARREN A. NARAYAN	
Equivalence classes of $GL(p, \mathbb{C}) \times GL(q, \mathbb{C})$ orbits in the flag variety of $gl(p+q, \mathbb{C})$	593
LETICIA BARCHINI AND NINA WILLIAMS	
Global sensitivity analysis in a mathematical model of the renal interstitium	625
MARIEL BEDELL, CLAIRE YILIN LIN, EMMIE ROMÁN-MELÉNDEZ AND IOANNIS SGOURALIS	
Sums of squares in quaternion rings	651
ANNA COOKE, SPENCER HAMBLÉN AND SAM WHITFIELD	
On the structure of symmetric spaces of semidihedral groups	665
JENNIFER SCHAEFER AND KATHRYN SCHLECHTWEG	
Spectrum of the Laplacian on graphs of radial functions	677
RODRIGO MATOS AND FABIO MONTENEGRO	
A generalization of Eulerian numbers via rook placements	691
ESTHER BANAIAN, STEVE BUTLER, CHRISTOPHER COX, JEFFREY DAVIS, JACOB LANDGRAF AND SCARLITTE PONCE	
The $H$ -linked degree-sum parameter for special graph families	707
LYDIA EAST KENNEY AND JEFFREY SCOTT POWELL	

