

involve

a journal of mathematics

Optimal transportation with constant constraint

Wyatt Boyer, Bryan Brown, Alyssa Loving and Sarah Tammen



Optimal transportation with constant constraint

Wyatt Boyer, Bryan Brown, Alyssa Loving and Sarah Tammen

(Communicated by Michael Dorff)

We consider optimal transportation *with constraint*, as did Korman and McCann (2013, 2015), provide simplifications and generalizations of their examples and results, and provide some new examples and results.

1. Introduction

The classical problem of optimal transportation seeks the least-cost way to move material between two locations in \mathbb{R}^n . Monge [1781] sought an optimal mapping. A more general problem, introduced in [Kantorovitch 1942], see also [Villani 2009, Theorem 3.1], seeks a cost-minimizing coupling between two measure spaces. If the coupling is absolutely continuous, it is given by a density H on the product. Recently optimal transportation has been used to better understand Riemannian manifolds and extend concepts such as Ricci curvature to more general spaces; [Cordero-Erausquin et al. 2006; Villani 2009].

Korman and McCann [2015] studied a constraint on the amount of material that can be transported between any two locations, an upper bound $h(x, y)$ on the density H , which goes back at least to [Levin 1984]. It is easy to show (Proposition 2.2) that if h is not prohibitively small, there is an optimal density H which equals 0 or h almost everywhere.

In this paper we specialize to the case of constant h . We assume $h \geq 1$, which is necessary and sufficient for existence (Proposition 2.2). Focusing on the solutions of the form 0 or h almost everywhere, for this paper we define a transportation plan as a map F from X to subsets of Y with measure $1/h$.

Our main section, Section 3, recognizes that many old and new examples of optimal transportation have the stronger “universal” property of minimizing the cost at each point separately. This leads to simplified proofs for many of the results and examples of [Korman and McCann 2013; 2015], as well as explicit examples of optimal transportation plans for all constraints $h \geq 1$. For instance, Example 3.7, due to Korman and McCann [2015, Example 1.1], provides a very short proof

MSC2010: primary 49Q20; secondary 90C08.

Keywords: optimal transportation, isoperimetric, geometry.

that optimal transportation from the unit interval to itself with cost $(x - y)^2$ with constraint $h = 2$ maps each point to whichever half of the unit interval it lies in. Proposition 3.13 proves that the intersection of two optimal transportation plans is optimal under certain conditions. Proposition 3.18 shows that in the torus or any Lie group, every admissible translation-invariant transportation plan is optimal for some continuous cost.

Proposition 4.3 presents a simplified approach to the surprising symmetries for dual cost constraints found by Korman and McCann [2013, Section 4].

Section 5 relates the case of finite spaces to some known combinatorial computations and asymptotic estimates.

2. Existence and uniqueness of optimal transportation plans

Proposition 2.2 provides existence of an optimal transportation plan F for admissible (constant) constraint h .

Definitions 2.1. Let X and Y be smooth manifolds, not necessarily compact, complete, or connected. Let f and g be nonnegative densities on X and Y , yielding probability measures on X and Y . A *transportation plan* F with constant constraint $h \geq 1$ is a measurable map from X to the power set $\mathcal{P}(Y)$ such that $F(x)$ has measure $1/h$ in Y for almost all $x \in X$ and such that $\{x \in X \mid y \in F(x)\}$ has measure $1/h$ in X for almost all $y \in Y$. (By F measurable we mean that the associated density $H(x, y)$, defined as the characteristic function of $F(x)$, is measurable.) For a cost function $c(x, y) \in L^\infty(X \times Y)$, the *total cost of transportation* is defined as

$$c[F] = \int_X \int_{F(x)} c(x, y) dy dx.$$

A transportation plan F is *optimal* if it minimizes cost.

Proposition 2.2. *Let X and Y be smooth (positive-dimensional) manifolds with non-negative densities f and g respectively and total measure 1. There exists an optimal transportation plan $F(x)$ if and only if the (constant) constraint h is at least 1.*

Proof. If $h < 1$, $F(x)$ cannot have measure $1/h$. On the other hand, if $h \geq 1$, the set of transportation densities $\Gamma(X, Y)$ is nonempty, since it includes $H(x, y) = 1$, and an optimal transportation density exists by standard compactness arguments; see [Korman and McCann 2015, Theorem 3.1].

Because $L^\infty(X, Y)$ is the dual of $L^1(X, Y)$, by Alaoglu's theorem, see [Rudin 1991, Section 3.15], the unit ball is compact in the weak-* topology. Thus the set of transportation densities $\Gamma(X, Y)$ is compact as well as convex. By the Krein–Milman theorem, every compact convex set has an extreme point, see [Wikipedia 2014a], and thus $\Gamma(X, Y)$ has an extreme point. The set of optimal transportation densities is a convex face of $\Gamma(X, Y)$ which contains an extreme point H , which is

also an extreme point of $\Gamma(X, Y)$. Such an extreme H must equal 0 or h almost everywhere, i.e., it must be a transportation plan F , see [Korman and McCann 2013, Proposition 3.2], although this can fail for finite sets of points; see Remark 2.3 below. \square

Remark 2.3. Of course there is an optimal transportation plan between finite sets (because there are only finitely many transportation plans), but the same proof does not work because there might be a better transportation density. For example the optimal transportation density from $\{0, 1\}$ to $\{0, \frac{1}{2}, 1\}$ with $h = \frac{3}{2}$ maps 0 to 1 with density $\frac{2}{3}$ and $\frac{1}{2}$ with density $\frac{1}{3}$ and maps 1 to $\frac{1}{2}$ with density $\frac{1}{3}$ and 2 with density $\frac{2}{3}$, and there is no equally good transportation plan $F(x)$; the only transportation plan maps each point to all three points. Actually Proposition 2.2 and its proof work as long as one of the two manifolds is positive-dimensional.

Although we will not need it, we provide the following uniqueness theorem of Korman and McCann.

Proposition 2.4 [Korman and McCann 2013, Theorem 3.3]. *Let X and Y be smooth manifolds with nonnegative densities f and g respectively and total measure 1. If the cost $c(x, y)$ is bounded, twice differentiable, and nondegenerate, i.e., $\det[D_{x^i y^j}^2 c(x, y)] \neq 0$ for almost all $(x, y) \in X \times Y$, then an optimal transportation plan $F(x)$ is unique (up to measure 0).*

Proof. Theorem 3.3 in [Korman and McCann 2013] gives a unique optimal density H . Since at least one optimal transportation density is an extreme point of Γ , H must be an extreme point of Γ and thus a transportation plan F . \square

Additionally, we give necessary and sufficient conditions for a map F from X to subsets of Y to be a transportation plan.

Proposition 2.5. *Let F be a measurable map from X to subsets of Y with constant constraint $h \geq 1$ such that $F(x)$ has measure $1/h$ in Y for almost all $x \in X$. Then F is a transportation plan if and only if for every $A \subset X$ of measure greater than $1/h$, $\bigcap_{x \in A} F(x)$ has measure 0.*

Proof. If F is a transportation plan, the condition holds. Suppose that F is not a transportation plan. Then it is not true that $\{x \in X \mid y \in F(x)\}$ has measure $1/h$ for almost all y . Since by Fubini's theorem the average satisfies

$$\int_Y f(\{x \in X \mid y \in F(x)\}) dy = \int_X g(F(x)) dx = 1/h$$

for some nontrivial subset of Y , we have $\{x \in X \mid y \in F(x)\}$ has measure greater than $1/h$, and the condition fails. \square

3. Universally optimal transportation

Finding optimal transportation plans for a given cost and constraint is hard. For example, the problem of optimal transportation from the unit interval $I = [0, 1]$ to itself with cost $c(x, y) = (x - y)^2$ is still open for $h \neq 2$; see [Korman and McCann 2013, Figure 1; 2015, Example 1.2]. In certain cases, however, it is possible to minimize the cost at each point separately. Further, for every optimal density, the cost function can be adjusted so that the same optimal density is also minimal at each point separately; see Remark 3.2 below.

Definition 3.1. For two smooth manifolds X and Y , a transportation plan F for the cost function c under constant constraint $h \geq 1$ is *universally optimal* if for almost every $x \in X$ it minimizes

$$\int_{F(x)} c(x, y) dy.$$

It follows immediately that F is optimal.

Remark 3.2. Korman, McCann, and Seis [Korman et al. 2015, Theorem 4.2] showed that for continuous densities f, g and $h > 1$, every optimal density is universally optimal for some equivalent cost $c(x, y) + u(x) + v(y)$ and hence for $c(x, y) + v(y)$; by Proposition 2.2 and its proof, this applies to optimal plans in the positive-dimensional case.

Morgan uses this concept of universal optimality to generalize and give shorter proofs of some of the examples of Korman and McCann.

Proposition 3.3 [Korman and McCann 2015, Example 1.3; Morgan 2013, Proposition 1]. *Let X be a Riemannian manifold of unit volume, with a transitive group of measure-preserving isometries, with cost of transportation $c(x, y)$ increasing in distance with constant constraint h . Then unique (universally) optimal transportation is that which maps each $x \in X$ to a geodesic ball about x of volume $1/h$.*

Proof. An optimal transportation plan F with constraint h must map a point $x \in X$ to a set of volume at least $1/h$, and the geodesic ball minimizes cost among such. By the symmetry assumption, all balls of the same radius have the same volume, so the set mapped to a target point $y \in Y$ is the ball about x with volume $1/h$ and the map satisfies the definition of a transportation plan and is clearly uniquely optimal (up to sets of measure 0). \square

Proposition 3.4 [Korman and McCann 2015, Example 1.1; Morgan 2013, Proposition 2]. *Let X and Y be two Riemannian manifolds of unit volume with cost of transportation $c(x, y)$ and constant constraint $h \geq 1$. Suppose that for almost all $x \in X$, $c(x, y)$ is negative for $1/h$ of the y 's in Y and nonnegative for the rest, and for almost all $y \in Y$, $c(x, y)$ is negative for $1/h$ of the x 's in X and nonnegative for*

the rest. Then unique (universally) optimal transportation maps each $x \in X$ to the subset of Y with negative cost.

Proof. By hypothesis, both $F(x)$ and $\{x \in X \mid y \in F(x)\}$ have measure $1/h$ for almost all $x \in X$ and $y \in Y$ respectively, and F is clearly universally and uniquely optimal (up to sets of measure 0). \square

Proposition 3.5. *Every transportation plan for which all images and inverse images have measure $1/h$ is optimal for some cost.*

Proof. Let $c(x, y) = -\chi_{F(x)}(y)$. Then F is optimal by Proposition 3.4. \square

Example 3.6 [Korman and McCann 2015, Example 1.1; Morgan 2013, Example 2.1]. For $h \geq 2$ an integer, let X consist of h equal-volume regions in \mathbb{R}^n such that the maximum diameter of a region is less than the minimum distance between regions. Let $c(x, y)$ be a cost function on $X \times X$ increasing in distance. Then optimal transportation from X to itself with constant constraint h maps the points of each region to itself. (To apply Proposition 3.4, subtract a constant from the cost.)

Example 3.7 [Korman and McCann 2015, Example 1.1; Morgan 2013, Example 2.2]. Let X be a centrally symmetric body in \mathbb{R}^n . For cost $c(x, y) = -2x \cdot y$, which is equivalent to $(x - y)^2$ because its integral differs by a constant, and for constraint $h = 2$, (universally) optimal transportation from X to itself is that which maps x to y with $x \cdot y$ positive (see Figure 1). In \mathbb{R}^1 central symmetry is unnecessary as long as the origin is the median. A similar result holds for any cost having the same sign at each point as $-x \cdot y$. The analysis generalizes to any centrally symmetric probability measure on \mathbb{R}^n for which hyperplanes through the origin have measure 0 and to any probability measure on \mathbb{R}^1 . Optimal transportation from X to itself with cost $-2x \cdot y$ is still open for constraint $h \neq 2$, though numerical estimates from some cases are given in [Korman and McCann 2013, Figure 1; 2015, Example 1.2].




Figure 1. Optimal transportation F from the unit ball in \mathbb{R}^2 to itself with cost $c(x, y) = (x - y)^2$ and constraint $h = 2$ maps each x to the half-ball $\{x \cdot y > 0\}$.




Figure 2. Optimal transportation F maps each x on a ray from the origin to a cone about that ray.

Example 3.8. Unique (universally) optimal transportation from the sphere \mathbb{S}^n to the ball \mathbb{B}^{n+1} with cost $c(x, y) = (x - y)^2$ and constraint $h = 2$ maps a point x to the half-ball $\{x \cdot y > 0\}$.

Proof. As in Example 3.6, the cost is equivalent to $-2x \cdot y$, which is negative precisely on the asserted half-ball, proving the asserted map uniquely universally optimal. \square

Example 3.9 [Morgan 2013, Example 2.3]. Let X be a planar region with h -fold rotational symmetry, such as a square ($h = 4$) as in Figure 2. For cost

$$c(x, y) = \cos(\pi/h)|x||y| - x \cdot y,$$

and constant constraint $h \geq 1$, (universally) optimal transportation maps all points on a ray from the origin to a cone of angle π/h about that ray.

Remark 3.10 [Morgan 2013, Example 2.4]. Such examples of universally optimal transportation plans from X_i to Y_i extend to universally optimal transportation plans from $\prod X_i$ to $\prod Y_i$ with a cost which is negative if and only if the costs of the projections are all negative: optimal transportation with constraint $h = \prod h_i$ maps to points of negative cost. In particular, Example 3.9 generalizes to a product of such actions on \mathbb{R}^{2n} with negative cost if and only if $x_i \cdot y_i \geq (\cos \pi/h_i)|x_i||y_i|$ for all i : optimal transportation with constant constraint $h = \prod h_i$ maps all points with projections on rays from the origin to a product of cones of angle π/h_i about the ray.

Remark 3.11 [Morgan 2013, Example 2.5]. Such examples of universally optimal transportation plans from X to Y with cost $c(x, y)$ extend to universally optimal transportation plans on warped products $A \times X$, $A \times Y$, as long as the cost $c'(a, x, a, y)$ has the same sign as $c(x, y)$. For example, for any $h \geq 1$, Proposition 3.4 on the sphere, with cost $c(x, y) = a|x||y| - x \cdot y$, with a chosen so that optimal transportation maps to points of negative cost, extends to the ball, with points on a ray from the origin mapped to a cone of negative cost about that ray.

Remark 3.12. Although universally optimal transportation plans are by definition optimal transportation plans, the converse is not true in general. Consider transportation from the unit interval to itself with cost of transportation increasing with distance and constant constraint $h = 2$. Minimizing cost for each x does not even give a valid transportation plan because points near 0 and 1 are mapped to by less than half of the interval.

Given two universally optimal transportation plans for two different costs, we seek ways to generate a third cost and a related universally optimal transportation plan.

Proposition 3.13. *Let F_1 and F_2 be optimal transportation plans from X to Y with costs c_1 and c_2 and constant constraints h_1 and h_2 respectively. Suppose that for almost all x , we have $F_i(x) = \{y \in Y \mid c_i(x, y) < 0\}$. If for some $1 \leq h < \infty$, for almost all $x \in X$, $F_1(x) \cap F_2(x)$ has measure $1/h$, and for almost all $y \in Y$, $\{x \in X \mid y \in F_1(x)\} \cap \{x \in X \mid y \in F_2(x)\}$ has measure $1/h$, then $F(x) = F_1(x) \cap F_2(x)$ is a universally optimal transportation plan from X to Y with cost $c(x, y) = \max(c_1, c_2)$ and constraint h .*

Proof. It suffices to show that for almost all $x \in X$, $c(x, y)$ is negative for $1/h$ of the $y \in Y$ and nonnegative for the rest and for almost all $y \in Y$, $c(x, y)$ is negative for $1/h$ of the $x \in X$ and nonnegative for the rest. By hypothesis on F , for almost all $x \in X$, $c(x, y)$ is negative for $1/h$ of the $y \in Y$. It is nonnegative for the rest because $x \notin F(x)$ implies some $c_i(x, y)$ must be nonnegative; thus $c(x, y)$ must also be nonnegative. The reverse condition holds by a similar argument. \square

Corollary 3.14. *Let X be a region with 4-fold rotational symmetry in \mathbb{R}^2 with cost of transportation from X to X given by $c(x, y) = \max((x \cdot y), \det[x \mid y])$, where $\det[x \mid y]$ is the determinant of the matrix with x and y as its column vectors. Mapping each point to the region of negative cost uniquely gives (universally) optimal transportation for $h = 4$ (see Figure 3).*




Figure 3. The intersection of optimal transportation plans yields a new optimal transportation plan under certain hypotheses.

Proof. The map $F_1(x) = \{y \in X \mid x \cdot y > 0\}$ is an optimal transportation plan from X to itself with cost $c_1(x, y) = -x \cdot y$ and constraint $h = 2$ (see Example 3.6). Similarly the cost $c_2(x, y) = \det[x \mid y]$ with constraint $h = 2$ satisfies the hypotheses of Proposition 3.4 and thus the map $F_2(x) = \{y \in X \mid \det[x \mid y] < 0\}$ is an optimal transportation plan from X to itself with cost $c_2(x, y)$ and constraint $h = 2$. By Proposition 3.13, if for almost all $x, y \in X$, $F_1(x) \cap F_2(x)$ and $\{x \in X \mid y \in F_1(x)\} \cap \{x \in X \mid y \in F_2(x)\}$ both have constant measure $1/h$ for some $h > 1$, then $F(x) = F_1(x) \cap F_2(x)$ is an optimal transportation plan from X to itself with cost $c(x, y) = \max(c_1, c_2)$ and constraint $1/h$. For almost all $x \in X$, $\partial F_1(x)$ is the line through the origin normal to the line through x and the origin and $\partial F_2(x)$ is the line through x and the origin. Because two normal lines both through the origin partition a region with 4-fold rotational symmetry centered on the origin in \mathbb{R}^2 into four congruent regions, and exactly one of these regions is equivalent to $F_1(x) \cap F_2(x)$, it follows that $F_1(x) \cap F_2(x)$ has constant measure $\frac{1}{4}$. Similarly, the boundary of the set $\{x \in X \mid y \in F_1(x)\}$ is the line through the origin normal to the line through y and the origin and the boundary of the set $\{x \in X \mid y \in F_2(x)\}$ is a line through y and the origin. Thus, by the same argument as above, $\{x \in X \mid y \in F_1(x)\} \cap \{x \in X \mid y \in F_2(x)\}$ has measure $\frac{1}{4}$. By Proposition 3.13, the asserted map is optimal. \square

Remark 3.15. If the hypotheses of Proposition 3.13 hold, then the maps $F(x) = F_1(x) \cup F_2(x)$ and $F(x) = F_1(x) \Delta F_2(x)$ are optimal transportation plans for costs $c = \min(c_1, c_2)$ and $c' = c_1 \cdot c_2$ and some constraints h and h' respectively (the symbol Δ denotes the symmetric difference of two sets).

Example 3.16. Let X be a region with 4-fold rotational symmetry in \mathbb{R}^2 . Then an optimal transportation plan F for cost

$$c(x, y) = ((\cos 3\pi/4h)|x||y| - x \cdot y)((\cos \pi/4h)|x||y| - x \cdot y)$$

and constraint $h = 2$ maps points on a ray from the origin to two cones (see Figure 4).




Figure 4. Other set operations yield even more examples of optimal transportation.




Figure 5. Optimal transportation F maps each x to an H -shaped region.

The condition in Proposition 3.13 that $F_1(x) \cap F_2(x)$ have constant measure $1/h$ for almost all $x \in X$ and some h is independent of the condition that $\{x \in X \mid y \in F_1(x)\} \cap \{x \in X \mid y \in F_2(x)\}$ have constant measure $1/h$ for almost all $y \in Y$.

Example 3.17. Let X be $\{1, 2, 3, 4\}$ or equivalently the unit interval divided into four quarters. Consider transportation F_1, F_2 from X to X such that

$$\begin{aligned} F_1(1) &= \{3, 4\}, & F_1(2) &= \{2, 3\}, & F_1(3) &= \{1, 2\}, & F_1(4) &= \{1, 4\}, \\ F_2(1) &= \{1, 4\}, & F_2(2) &= \{1, 2\}, & F_2(3) &= \{2, 3\}, & F_2(4) &= \{3, 4\}. \end{aligned}$$

Then $F(x) = F_1(x) \cap F_2(x)$ has constant measure $\frac{1}{4}$ but

$$\{x \in X \mid y \in F(x)\} = \{x \in X \mid y \in F_1(x)\} \cap \{x \in X \mid y \in F_2(x)\}$$

has measure $\frac{1}{2}$ for $\{2, 4\}$ and measure 0 for $\{1, 3\}$.

Proposition 3.18. *Let X be a Lie group. Given an open subset A of X with measure $1/h$, there exists a continuous cost function $c(x, y)$ such that the unique (universally) optimal transportation plan F from X to itself with constant constraint h maps the identity to the set A and maps each element $x \in X$ to the set $x \cdot A = xA$.*

Proof. Let the cost $c(x, y)$ equal the distance from y to the boundary of $x \cdot A$, with negative cost on the interior of $x \cdot A$ and nonnegative cost on the complement of $x \cdot A$. By Proposition 3.4, the asserted map is optimal. \square

Example 3.19. Let $X = \mathbb{S}^1 \times \mathbb{S}^1$ with unit area. Given an open subset $A \subset X$ with measure $1/h$, such as the H -shaped region in Figure 5, there exists a continuous cost function $c(x, y)$ such that the unique (universally) optimal transportation plan F from X to itself with constant constraint h maps the origin to the set A and maps almost every x to the set $\tau_x(A)$, where τ_x is the translation that takes the origin to x .

Example 3.20. Optimal transportation from a flat rectangular torus to itself with cost $c(x, y) = \min(d(x_i, y_i))$ and constraint h maps each point to a neighborhood around the coordinate axis centered at that point; see Figure 6.




Figure 6. Optimal transportation on the flat rectangular torus maps each x to a small neighborhood around the coordinate axis centered at that point.

4. Transportation and symmetry

Korman and McCann [2013, Section 4] found surprising symmetries between optimal transportation plans with dual constraints. Proposition 4.3 presents a simplified approach.

Definition 4.1. A map f from X' to X is called *measure preserving* if the measure of any $A \subset X$ equals the measure of $f^{-1}(A) \subset X'$.

Proposition 4.2. Let F be an optimal transportation plan from X to Y with cost $c(x, y)$ and constraint h . Let f and g be measure-preserving maps from X' to X and from Y' to Y respectively. Then $G(x') = g^{-1}(F(f(x')))$ provides optimal transportation from X' to Y' with cost $c \circ (f, g)$ and constraint h .

Proof. We need to show that $G(x')$ and $G^{-1}(y')$ both have measure $1/h$ and that the total cost of transportation is minimal. For $x' \in X'$, $G(x') = g^{-1}(F(f(x')))$ must have the same measure as $F(f(x'))$, which is $1/h$ by hypothesis. Similarly, for y' in Y' , $G^{-1}(y') = f^{-1}(F^{-1}(g(y')))$ must have the same measure as $F^{-1}(g(y'))$, which is also $1/h$ by hypothesis. To show that G is optimal, we will show that G and F have the same cost and that any other transportation plan G_2 from X' to Y' has the same cost as an analogous transportation plan F_2 from X to Y and therefore must be of greater total cost than G . The cost of transportation from x' to y' is equal to $c(f(x'), F(f(x')))$; thus G and F have the same total cost of transportation. Let G_2 be another transportation map from X' to Y' . Let $F_2(f(x')) = g(G_2)$. Then $G_2 = g^{-1}(F_2(f(x')))$ and the result follows. \square

Proposition 4.3 [Korman and McCann 2013, Lemma 4.1; Morgan 2013, Proposition 3]. Let M_1, M_2 be subsets of \mathbb{R}^n or Riemannian manifolds with boundary or metric measure spaces of volume V . Let T_i be a measure-preserving map from M_i to itself and let $T = T_1 \times T_2$. Let $c(x, y)$ be a cost satisfying $c \circ T = -c$. If the map F is an optimal transportation plan from M_1 to M_2 with cost $c(x, y)$ with

constraint h , then the map $T_2(F' \circ T_1)$ is an optimal transportation plan from M_1 to M_2 with cost c and constraint h' , where $1/h + 1/h' = 1$ and $F'(x) = F(x)^C$.

Proof. If F is an optimal transportation plan for cost c and constraint h , then $F(x)' = F(x)^C$ is the most expensive transportation plan for cost c with constraint h' , and hence an optimal transportation plan for cost $-c$. Therefore $T_2(F' \circ T_1)$ is an optimal transportation plan for cost $-c \circ T = c$ and constraint h' . \square

Example 4.4 [Morgan 2013, Example after Proposition 3; Korman and McCann 2013, Lemma 4.1]. Let M_1 and M_2 be subsets of \mathbb{R}^n , with M_1 centrally symmetric, and let $c(x, y) = -x \cdot y$ (which is equivalent to $(x - y)^2$). Then central inversion in x carries optimal transportation with constraint h to optimal transportation with constraint h' .

5. Transportation plans on finite sets

Consider the case where X and Y are finite sets, say $X = \{1, 2, \dots, m\}$ and $Y = \{1, 2, \dots, n\}$. In this case we may assume that the constraint h is a common divisor of m and n . A map F from X to Y is equivalent to the $n \times m$ matrix of 0's and 1's with entry $a_{ij} = 1$ if and only if $i \in F(j)$; see [Wikipedia 2014b; Weisstein]. Such a matrix gives a transportation plan if and only if the matrix has m/h 1's in each column and n/h 1's in each row. Thus the number of transportation plans is equal to the number of $n \times m$ binary matrices with constant column sums n/h and constant row sums m/h . Asymptotic estimates exist for large m, n ; see [Canfield and McKay 2005; McKay and Wang 2003].

Acknowledgements

We thank the National Science Foundation and Williams College for supporting this work as part of the Williams College SMALL Undergraduate Research Project. We thank our advisor Frank Morgan for his insight, enthusiasm, and guidance. We thank Robert McCann and Christian Seis for helpful comments. Additionally we thank the Mathematical Association of America, Williams College, and Pomona College for supporting our travel to MathFest 2014.

References

- [Canfield and McKay 2005] E. R. Canfield and B. D. McKay, "Asymptotic enumeration of dense 0-1 matrices with equal row sums and equal column sums", *Electron. J. Combin.* **12** (2005), art. id. R29. MR Zbl
- [Cordero-Erausquin et al. 2006] D. Cordero-Erausquin, R. J. McCann, and M. Schmuckenschläger, "Prékopa–Leindler type inequalities on Riemannian manifolds, Jacobi fields, and optimal transport", *Ann. Fac. Sci. Toulouse Math.* (6) **15**:4 (2006), 613–635. MR Zbl

- [Kantorovitch 1942] L. Kantorovitch, “On the translocation of masses”, *C. R. (Doklady) Acad. Sci. URSS (N.S.)* **37** (1942), 199–201. In Russian; translated in *J. Math. Sci.* **133**:4 (2006), 1381–1382. MR Zbl
- [Korman and McCann 2013] J. Korman and R. McCann, “Insights into capacity-constrained optimal transport”, *Proc. Nat. Acad. Sci. USA* **110**:25 (2013), 10064–10067.
- [Korman and McCann 2015] J. Korman and R. J. McCann, “Optimal transportation with capacity constraints”, *Trans. Amer. Math. Soc.* **367**:3 (2015), 1501–1521. MR Zbl
- [Korman et al. 2015] J. Korman, R. J. McCann, and C. Seis, “Dual potentials for capacity constrained optimal transport”, *Calc. Var. Partial Differential Equations* **54**:1 (2015), 573–584. MR Zbl
- [Levin 1984] V. L. Levin, “The problem of mass transfer in a topological space and probability measures with given marginal measures on the product of two spaces”, *Dokl. Akad. Nauk SSSR* **276**:5 (1984), 1059–1064. In Russian. MR Zbl
- [McKay and Wang 2003] B. D. McKay and X. Wang, “Asymptotic enumeration of 0-1 matrices with equal row sums and equal column sums”, *Linear Algebra Appl.* **373** (2003), 273–287. MR Zbl
- [Monge 1781] G. Monge, “Mémoire sur la théorie des déblais et des remblais”, pp. 666–704 in *Histoire de l’Académie Royale des Sciences de Paris*, De l’Imprimerie Royale, Paris, 1781.
- [Morgan 2013] F. Morgan, “Optimal transportation with constraint”, blog post, 2013, available at <http://sites.williams.edu/Morgan/2013/09/14/optimal-transportation-with-constraint/>.
- [Rudin 1991] W. Rudin, *Functional analysis*, 2nd ed., International Series in Pure and Applied Mathematics, McGraw-Hill, New York, 1991. MR Zbl
- [Villani 2009] C. Villani, *Optimal transport: old and new*, Grundlehren der Mathematischen Wissenschaften **338**, Springer, 2009. MR Zbl
- [Weisstein] E. W. Weisstein, “(0,1)-matrix”, available at <http://tinyurl.com/01matrix>. From Math-World.
- [Wikipedia 2014a] “Extreme point”, Wikipedia entry, 2014, available at http://en.wikipedia.org/wiki/Extreme_point.
- [Wikipedia 2014b] “Logical matrix”, Wikipedia entry, 2014, available at https://en.wikipedia.org/wiki/Logical_matrix.

Received: 2014-12-22

Revised: 2016-04-19

Accepted: 2016-05-02

wyatt.b.boyer@gmail.com

*Department of Mathematics and Statistics, Williams College,
Brighton, MA, United States*

bbrown@math.berkeley.edu

*Department of Mathematics, Pomona College,
Claremont, CA, United States*

aloving2@illinois.edu

*Department of Mathematics, University of Hawaii at Hilo,
Hilo, HI, United States*

setammen@uga.edu

*Department of Mathematics, University of Georgia,
Lawrenceville, GA, United States*

involve

msp.org/involve

INVOLVE YOUR STUDENTS IN RESEARCH

Involve showcases and encourages high-quality mathematical research involving students from all academic levels. The editorial board consists of mathematical scientists committed to nurturing student participation in research. Bridging the gap between the extremes of purely undergraduate research journals and mainstream research journals, *Involve* provides a venue to mathematicians wishing to encourage the creative involvement of students.

MANAGING EDITOR

Kenneth S. Berenhaut Wake Forest University, USA

BOARD OF EDITORS

Colin Adams	Williams College, USA	Suzanne Lenhart	University of Tennessee, USA
John V. Baxley	Wake Forest University, NC, USA	Chi-Kwong Li	College of William and Mary, USA
Arthur T. Benjamin	Harvey Mudd College, USA	Robert B. Lund	Clemson University, USA
Martin Bohner	Missouri U of Science and Technology, USA	Gaven J. Martin	Massey University, New Zealand
Nigel Boston	University of Wisconsin, USA	Mary Meyer	Colorado State University, USA
Amarjit S. Budhiraja	U of North Carolina, Chapel Hill, USA	Emil Minchev	Ruse, Bulgaria
Pietro Cerone	La Trobe University, Australia	Frank Morgan	Williams College, USA
Scott Chapman	Sam Houston State University, USA	Mohammad Sal Moslehian	Ferdowsi University of Mashhad, Iran
Joshua N. Cooper	University of South Carolina, USA	Zuhair Nashed	University of Central Florida, USA
Jem N. Corcoran	University of Colorado, USA	Ken Ono	Emory University, USA
Toka Diagana	Howard University, USA	Timothy E. O'Brien	Loyola University Chicago, USA
Michael Dorff	Brigham Young University, USA	Joseph O'Rourke	Smith College, USA
Sever S. Dragomir	Victoria University, Australia	Yuval Peres	Microsoft Research, USA
Behrouz Emamizadeh	The Petroleum Institute, UAE	Y.-F. S. Pétermann	Université de Genève, Switzerland
Joel Foisy	SUNY Potsdam, USA	Robert J. Plemmons	Wake Forest University, USA
Errin W. Fulp	Wake Forest University, USA	Carl B. Pomerance	Dartmouth College, USA
Joseph Gallian	University of Minnesota Duluth, USA	Vadim Ponomarenko	San Diego State University, USA
Stephan R. Garcia	Pomona College, USA	Bjorn Poonen	UC Berkeley, USA
Anant Godbole	East Tennessee State University, USA	James Propp	U Mass Lowell, USA
Ron Gould	Emory University, USA	József H. Przytycki	George Washington University, USA
Andrew Granville	Université Montréal, Canada	Richard Rebarber	University of Nebraska, USA
Jerrold Griggs	University of South Carolina, USA	Robert W. Robinson	University of Georgia, USA
Sat Gupta	U of North Carolina, Greensboro, USA	Filip Saidak	U of North Carolina, Greensboro, USA
Jim Haglund	University of Pennsylvania, USA	James A. Sellers	Penn State University, USA
Johnny Henderson	Baylor University, USA	Andrew J. Sterge	Honorary Editor
Jim Hoste	Pitzer College, USA	Ann Trenk	Wellesley College, USA
Natalia Hritonenko	Prairie View A&M University, USA	Ravi Vakil	Stanford University, USA
Glenn H. Hurlbert	Arizona State University, USA	Antonia Vecchio	Consiglio Nazionale delle Ricerche, Italy
Charles R. Johnson	College of William and Mary, USA	Ram U. Verma	University of Toledo, USA
K. B. Kulasekera	Clemson University, USA	John C. Wierman	Johns Hopkins University, USA
Gerry Ladas	University of Rhode Island, USA	Michael E. Zieve	University of Michigan, USA

PRODUCTION

Silvio Levy, Scientific Editor


Cover: Alex Scorpion

See inside back cover or msp.org/involve for submission instructions. The subscription price for 2019 is US\$/year for the electronic version, and \$/year (+\$, if shipping outside the US) for print and electronic. Subscriptions, requests for back issues and changes of subscriber address should be sent to MSP.

Involve (ISSN 1944-4184 electronic, 1944-4176 printed) at Mathematical Sciences Publishers, 798 Evans Hall #3840, c/o University of California, Berkeley, CA 94720-3840, is published continuously online. Periodical rate postage paid at Berkeley, CA 94704, and additional mailing offices.

Involve peer review and production are managed by EditFLOW® from Mathematical Sciences Publishers.

PUBLISHED BY

 **mathematical sciences publishers**
nonprofit scientific publishing

<http://msp.org/>

© 2019 Mathematical Sciences Publishers

involve

2019 vol. 12 no. 1

Optimal transportation with constant constraint	1
WYATT BOYER, BRYAN BROWN, ALYSSA LOVING AND SARAH TAMMEN	
Fair choice sequences	13
WILLIAM J. KEITH AND SEAN GRINDATTI	
Intersecting geodesics and centrality in graphs	31
EMILY CARTER, BRYAN EK, DANIELLE GONZALEZ, RIGOBERTO FLÓREZ AND DARREN A. NARAYAN	
The length spectrum of the sub-Riemannian three-sphere	45
DAVID KLAPHECK AND MICHAEL VANVALKENBURGH	
Statistics for fixed points of the self-power map	63
MATTHEW FRIEDRICHSSEN AND JOSHUA HOLDEN	
Analytical solution of a one-dimensional thermistor problem with Robin boundary condition	79
VOLODYMYR HRYNKIV AND ALICE TURCHANINOVA	
On the covering number of S_{14}	89
RYAN OPPENHEIM AND ERIC SWARTZ	
Upper and lower bounds on the speed of a one-dimensional excited random walk	97
ERIN MADDEN, BRIAN KIDD, OWEN LEVIN, JONATHON PETERSON, JACOB SMITH AND KEVIN M. STANGL	
Classifying linear operators over the octonions	117
ALEX PUTNAM AND TEVIAN DRAY	
Spectrum of the Kohn Laplacian on the Rossi sphere	125
TAWFIK ABBAS, MADELYNE M. BROWN, RAVIKUMAR RAMASAMI AND YUNUS E. ZEYTUNCU	
On the complexity of detecting positive eigenvectors of nonlinear cone maps	141
BAS LEMMENS AND LEWIS WHITE	
Antiderivatives and linear differential equations using matrices	151
YOTSANAN MEEMARK AND SONGPON SRIWONGSA	
Patterns in colored circular permutations	157
DANIEL GRAY, CHARLES LANNING AND HUA WANG	
Solutions of boundary value problems at resonance with periodic and antiperiodic boundary conditions	171
ALDO E. GARCIA AND JEFFREY T. NEUGEBAUER	

