

involve

a journal of mathematics

Analytical solution of
a one-dimensional thermistor problem
with Robin boundary condition

Volodymyr Hrynkyv and Alice Turchaninova



Analytical solution of a one-dimensional thermistor problem with Robin boundary condition

Volodymyr Hryniv and Alice Turchaninova

(Communicated by Suzanne Lenhart)

A one-dimensional nonlinear heat conduction equation of steady-state Joule heating in the presence of an electric field in a metal with temperature-dependent conductivities is considered. A technique developed by Young (1986) is adapted and used to derive an analytical solution for the problem with a Robin boundary condition.

1. Introduction

A thermal resistor, or thermistor, is a type of resistor with a highly temperature-dependent electrical conductivity. Thermistors are used as temperature-control elements in a range of equipment, such as spacecraft and air conditioning units, and have applications in the medical field, meteorology, and the chemical industry [Ng 1995; Macklen 1979]. The thermistor problem has been a source of significant mathematical interest and research but, due to the nonlinear nature of the problem, this research has been largely concerned with numerical solutions or existence proofs for a solution [Antontsev and Chipot 1994; Fowler et al. 1992; Howison et al. 1993; Shi et al. 1993; Sidi Ammi and Torres 2008; Xu 2004a; 2004b; Zhou and Westbrook 1997], rather than analytical solutions. In this paper, the thermistor problem is modeled as a nonlinear heat conduction equation of steady-state Joule heating in the presence of an electric field in a metal with temperature-dependent electrical and thermal conductivities. This paper extends the solution found in [Young 1986] to a more general case by introducing a Robin boundary condition on the temperature at an endpoint of the thermistor. This establishes the existing solution in [Young 1986] as a special case.

MSC2010: 35J60.

Keywords: thermistor problem, Robin boundary condition, Joule heating, analytical solution.

2. Formulation of the problem

Assuming that the electrical conductivity $\sigma(T)$ and the thermal conductivity $\kappa(T)$ are smooth functions, the heat conduction in an electrical conductor in the presence of Joule heating due to current can be shown [Young 1986] to satisfy the following two equations, one for the potential Φ and one for the temperature T :

$$\nabla^2 \Phi = -\frac{1}{\sigma} \nabla \sigma \cdot \nabla \Phi \quad \text{in } \Omega, \quad (1)$$

$$\frac{d^2 T}{d\Phi^2} + \left(\frac{1}{\kappa} \frac{d\kappa}{dT} - \frac{1}{\sigma} \frac{d\sigma}{dT} \right) \left(\frac{dT}{d\Phi} \right)^2 = -\frac{\sigma}{\kappa} \quad \text{in } \Omega, \quad (2)$$

with some appropriate boundary conditions. Throughout this section and Section 3 we assume that the given domain Ω lies in \mathbb{R}^n . Equations (1) and (2) respectively describe conservation of charge and the steady diffusion of heat in the presence of Joule heating due to electric current (see [Young 1986] for more details).

3. Derivation of the solution in the general case

For the sake of convenience, we recreate the technique developed in [Young 1986] for obtaining a solution to (1) and (2). Equation (2) can be simplified by introducing a new variable

$$\frac{\sigma(T)}{\kappa(T)} = e^{-\xi(T)}. \quad (3)$$

Differentiating both sides of (3), and after some manipulations, (2) can be written as

$$\frac{d^2 T}{d\Phi^2} + \frac{d\xi}{dT} \left(\frac{dT}{d\Phi} \right)^2 = -e^{-\xi}. \quad (4)$$

Next, setting

$$\theta = \frac{dT}{d\Phi}, \quad (5)$$

(4) becomes

$$\theta \frac{d\theta}{dT} + \frac{d\xi}{dT} \theta^2 = -e^{-\xi}. \quad (6)$$

Observing

$$\frac{1}{2} e^{-2\xi} \frac{d}{dT} (e^{2\xi} \theta^2) = \theta \frac{d\theta}{dT} + \frac{d\xi}{dT} \theta^2$$

allows us to rewrite (6) as

$$\frac{1}{2} e^{-2\xi} \frac{d}{dT} (e^{2\xi} \theta^2) = -e^{-\xi}. \quad (7)$$

Integrating (7), we get

$$e^{2\xi(T)} \theta^2 = C - 2 \int e^{\xi(T)} dT, \quad (8)$$

where C is a constant of integration. Solving for θ^2 in (8) and taking into account (5), we have

$$\left(\frac{dT}{d\Phi}\right)^2 = \frac{C - 2 \int e^{\xi(T)} dT}{e^{2\xi(T)}}.$$

Finally, the following equation for Φ is obtained:

$$\Phi = \int \frac{(\kappa(T)/\sigma(T)) dT}{\sqrt{C - 2 \int (\kappa(s)/\sigma(s)) ds}} + C', \quad (9)$$

where the integration constant of the integral under the square root is absorbed into C . It turns out that for many metals, the ratio of conductivities is proportional to the absolute temperature of the metal. This relationship is known as the Wiedemann–Franz–Lorenz (WFL) law [Berman 1976; Meaden 1965],

$$\frac{\kappa(T)}{\sigma(T)} = \alpha T, \quad (10)$$

where α is the Lorenz number for a given metal and may have slightly different values for different metals. Once the ratio of conductivities is specified using the WFL law, (9) can be integrated to obtain the temperature in terms of the potential, $T(\Phi)$,

$$T(\Phi) = \frac{1}{\sqrt{\alpha}} [C - (\Phi - C')^2]^{1/2}. \quad (11)$$

For (11) to be of any help, we must determine Φ that solves (1). This issue can be dealt with by introducing an auxiliary potential Ψ that satisfies Laplace's equation [Young 1986; Flynn 1969]. Namely, define this auxiliary potential Ψ as

$$\sigma_0 \nabla \Psi \equiv \sigma [T(\Phi)] \nabla \Phi,$$

where σ_0 is the electrical conductivity at some conveniently chosen reference temperature. Then clearly Ψ satisfies Laplace's equation $\nabla^2 \Psi = 0$, and it is easily seen, by isolating $\nabla \Psi$ and integrating, that

$$\Psi = \frac{1}{\sigma_0} \int \sigma [T(\Phi)] d\Phi. \quad (12)$$

Knowing how σ depends on T and using (11) will enable us to perform integration in (12). This will give us Ψ in terms of Φ , and therefore finding the inverse of this function will result in an expression for Φ in terms of a function that satisfies Laplace's equation. In the next section, we will also solve for the constants C and C' in the expression. This ends the derivation of a solution to (1) and (2) when conductivities obey the WFL law.

4. Solution in one dimension with Robin boundary condition

In this section we adapt the technique described in Section 3 for a one-dimensional problem with a Robin boundary condition at one endpoint. Namely, consider a thin rod of length L , where the potential and temperature satisfy (1) and (2), respectively. In addition, the endpoints at $z = 0$ and $z = L$ are held at potentials V_0 and 0, respectively. The boundary condition for T at the right endpoint of the rod $z = L$ is given by a Robin boundary condition, whereas the left endpoint is held at the constant temperature $T = T_0$. Note that it is reasonable to consider a Robin boundary condition for at least one end of the rod, as it models the cooling effect of that end of the thermistor through Newton's law of cooling [Howison 2005]. The boundary conditions are summarized below:

$$T = T_0, \quad \Phi = V_0 \quad \text{at } z = 0, \quad (13)$$

$$\frac{dT}{dz} + \beta(T - T_0) = 0, \quad \Phi = 0 \quad \text{at } z = L. \quad (14)$$

Observe that when β approaches infinity, the boundary condition for T at $z = L$ reduces to $T = T_0$ at $z = L$, which corresponds to that in [Young 1986]. Recall that

$$T(\Phi) = \frac{1}{\sqrt{\alpha}}[C - (\Phi - C')^2]^{1/2}.$$

Now we use the boundary conditions (13) and (14) to determine the constants C and C' . From (13), it is immediate that

$$C - C'^2 = \alpha T_0^2 + V_0^2 - 2V_0 C'. \quad (15)$$

First, we find

$$\begin{aligned} & \frac{dT}{dz} + \beta(T - T_0) \\ &= \frac{C' - \Phi}{\sqrt{\alpha}} \frac{1}{[C - (\Phi - C')^2]^{1/2}} \frac{d\Phi}{dz} + \beta \left(\frac{1}{\sqrt{\alpha}} [C - (\Phi - C')^2]^{1/2} - T_0 \right) \end{aligned} \quad (16)$$

and using (14) to evaluate (16) at $z = L$ gives us

$$\frac{C'}{\sqrt{\alpha}} \frac{\Phi_0}{[C - C'^2]^{1/2}} + \beta \left(\frac{1}{\sqrt{\alpha}} [C - C'^2]^{1/2} - T_0 \right) = 0, \quad (17)$$

where we defined $\Phi_0 := \Phi'(L)$. Note that a new parameter Φ_0 has been introduced into the problem. We will address this issue later. Rewriting (17) as

$$\frac{C'}{\beta} \frac{\Phi_0}{[C - C'^2]^{1/2}} + [C - C'^2]^{1/2} = \sqrt{\alpha} T_0 \quad (18)$$

and squaring both sides of (18), we get

$$\left(\frac{C'\Phi_0}{\beta}\right)^2 + \frac{2\Phi_0 C'}{\beta}(C - C'^2) + (C - C'^2)^2 = \alpha T_0^2(C - C'^2).$$

Now using (15) and grouping the result by C'^2 , we obtain the following quadratic equation for C' :

$$C'^2 \left[\left(\frac{\Phi_0}{\beta}\right)^2 - \frac{4V_0\Phi_0}{\beta} + 4V_0^2 \right] + C' \left[\frac{2\Phi_0}{\beta}(\alpha T_0^2 + V_0^2) - 4V_0^3 - 2\alpha V_0 T_0^2 \right] + \alpha T_0^2 V_0^2 + V_0^4 = 0.$$

Defining

$$\begin{aligned} A &= \left(\frac{\Phi_0}{\beta}\right)^2 - \frac{4V_0\Phi_0}{\beta} + 4V_0^2, \\ B &= \frac{2\Phi_0}{\beta}(\alpha T_0^2 + V_0^2) - 4V_0^3 - 2\alpha V_0 T_0^2, \\ D &= \alpha T_0^2 V_0^2 + V_0^4, \end{aligned}$$

the solution for C' is given by

$$C' = \frac{-B \pm \sqrt{B^2 - 4AD}}{2A}, \quad (19)$$

where

$$\begin{aligned} B^2 &= \frac{4\Phi_0(\alpha T_0^2 + V_0^2)^2 - 4\Phi_0\beta(\alpha T_0^2 + V_0^2)(4V_0^3 + 2\alpha V_0 T_0^3) + \beta^2(4V_0^3 + 2\alpha V_0 T_0^2)^2}{\beta^2}, \\ 4AD &= \frac{4\Phi_0^2(\alpha T_0^2 + V_0^2)V_0^2 - 16\Phi_0\beta V_0(\alpha T_0^2 + V_0^2)V_0^2 + \beta^2 16V_0^2(\alpha T_0^2 V_0 + V_0^3)V_0}{\beta^2}, \end{aligned}$$

so that

$$\begin{aligned} B^2 - 4AD &= \frac{4\Phi_0^2\alpha T_0^2(\alpha T_0^2 + V_0^2) - 8\Phi_0\beta\alpha V_0 T_0^2(\alpha T_0^2 + V_0^2) + 4\beta^2\alpha^2 V_0^2 T_0^4}{\beta^2} \\ &= \frac{4\Phi_0^2\alpha T_0^2(\alpha T_0^2 + V_0^2)}{\beta^2} - \frac{8\Phi_0\alpha V_0 T_0^2(\alpha T_0^2 + V_0^2)}{\beta} + 4\alpha^2 V_0^2 T_0^4. \end{aligned}$$

Now it can be verified that this solution to the quadratic equation will match that in [Young 1986] when $\beta \rightarrow \infty$. In this case, the first two terms above disappear and we have

$$\lim_{\beta \rightarrow \infty} A = 4V_0^2, \quad \lim_{\beta \rightarrow \infty} B = -4V_0^3 - 2\alpha V_0 T_0^2, \quad \lim_{\beta \rightarrow \infty} [B^2 - 4AD] = 4\alpha^2 V_0^2 T_0^4.$$

We are thus left with

$$\lim_{\beta \rightarrow \infty} C' = \lim_{\beta \rightarrow \infty} \left[\frac{-B \pm \sqrt{B^2 - 4AD}}{2A} \right] = \frac{4V_0^3 + 2\alpha V_0 T_0^2 \pm \sqrt{4\alpha^2 V_0^2 T_0^4}}{8V_0^2}.$$

Taking the negative square root, we get $C' = \frac{1}{2} V_0$, as in [Young 1986]. The constants C and C' in terms of β are as follows:

$$C' = \frac{-B - 2T_0 \sqrt{\alpha} \sqrt{(\Phi_0 - \beta V_0)^2 (\alpha T_0^2 + V_0^2) - \beta^2 V_0^4}}{2A}, \quad (20)$$

$$C = \alpha T_0^2 + V_0^2 - 2V_0 C' + C'^2,$$

where A and B are defined above. We take the negative root in the quadratic formula for C' because this is the root that reduces to Young's solution when $\beta \rightarrow \infty$. Now we use the auxiliary potential Ψ , given in (12),

$$\Psi = \frac{1}{\sigma_0} \int \sigma[T(\Phi)] d\Phi.$$

First, we note that it is an experimentally verified fact that the thermal conductivity κ varies very little with temperature for many metals; see [Young 1986]. Therefore, it is physically reasonable to assume that $\kappa(T) = \kappa_0$, where κ_0 is a constant. Now, taking $\sigma[T(\Phi)]$ to obey the WFL law (10), we have

$$\sigma[T(\Phi)] = \frac{\kappa[T(\Phi)]}{\alpha T(\Phi)} = \frac{\kappa_0}{\alpha T(\Phi)}. \quad (21)$$

Substituting (21) into (12), we get

$$\Psi = \frac{\kappa_0}{\alpha \sigma_0} \int \frac{d\Phi}{T(\Phi)} = \frac{\kappa_0}{\alpha \sigma_0} \int \frac{d\Phi}{(1/\sqrt{\alpha})[C - (\Phi - C')^2]^{1/2}} = \frac{\kappa_0}{\alpha \sigma_0} \sin^{-1} \left(\frac{\Phi - C'}{\sqrt{C}} \right).$$

Since $\nabla^2 \Psi = 0$, it follows that Ψ is a linear function of z . We set $\Psi(z) = a + bz$ and absorb the constant $\kappa_0/(\alpha \sigma_0)$, as well as the integration constant of $\Psi(z)$, into the coefficients of the linear function. Hence,

$$\sin^{-1} \left(\frac{\Phi - C'}{\sqrt{C}} \right) = a + bz.$$

Using the boundary conditions for Φ to determine a and b ,

$$\begin{aligned} \Phi(z=0) &= \sqrt{C} \sin(a) + C' = V_0, \\ \Phi(z=L) &= \sqrt{C} \sin(a + bL) + C' = 0, \end{aligned}$$

we obtain the expression for the general solution $\Phi(z)$,

$$\Phi(z) = \sqrt{C} \sin(a + bz) + C', \quad (22)$$

fluid 1	transmission surface	fluid 2	β	Φ_0
air	cast iron	air	5.7	-0.03864
air	mild steel	air	7.9	-0.03834
steam	cast iron	air	11.3	-0.03809
steam	mild steel	air	14.2	-0.03796
steam	copper	air	17	-0.03788

Table 1. Φ_0 for realistic values of β .

where

$$a = \sin^{-1}\left(\frac{V_0 - C'}{\sqrt{C}}\right), \quad b = \frac{1}{L}\left[\sin^{-1}\left(\frac{-C'}{\sqrt{C}}\right) - a\right],$$

with C and C' given by (20), and where Φ_0 is determined numerically from the equation

$$\Phi_0 = \Phi'(L) = b\sqrt{C} \cos(a + bL). \quad (23)$$

Equation (23) was obtained by differentiating (22) with respect to z and then evaluating the derivative at $z = L$. Note that since the right-hand side of (23) also contains Φ_0 , we view (23) as an equation where the unknown is Φ_0 . Even though (23) cannot be solved analytically for Φ_0 , as it enters the right-hand side of (23) in a complicated way, we can still solve (23) numerically by choosing physically realistic values for the parameters of the problem. Table 1 gives values of Φ_0 for realistic values of β , provided in [Engineering ToolBox 2003] for transmission surfaces between various combinations of fluids. The units of β are $\text{W}/(\text{m}^2\text{K})$ and the units of Φ_0 are V/m .

To complete the general solution, we substitute (22) back into (11) to obtain $T(z)$:

$$T(z) = \frac{1}{\sqrt{\alpha}} \sqrt{C} \cos(a + bz). \quad (24)$$

Finally, as is expected, $\Phi(z)$ in (22) tends to the one found in [Young 1986] as $\beta \rightarrow \infty$. Indeed, we have

$$\begin{aligned} \lim_{\beta \rightarrow \infty} C' &= \frac{1}{2} V_0, \\ \lim_{\beta \rightarrow \infty} C &= \alpha T_0^2 + \frac{1}{4} V_0^2, \\ \lim_{\beta \rightarrow \infty} a &= \sin^{-1}\left(\frac{\frac{1}{2} V_0}{\sqrt{\alpha T_0^2 + \frac{1}{4} V_0^2}}\right) = \Omega, \\ \lim_{\beta \rightarrow \infty} b &= \frac{1}{L} \left[\sin^{-1}\left(\frac{-\frac{1}{2} V_0}{\sqrt{\alpha T_0^2 + \frac{1}{4} V_0^2}}\right) - \Omega \right] = \frac{1}{L} [-\Omega - \Omega] = -\frac{2\Omega}{L}, \end{aligned}$$




Figure 1. B^2-4AD as a function of β for $T_0 = 273$ K, $\alpha = 2.445 \cdot 10^{-8} (\text{V/K})^2$, $V_0 = 40$ mV.




Figure 2. B^2-4AD as a function of V_0 for $T_0 = 273$ K, $\alpha = 2.445 \cdot 10^{-8} (\text{V/K})^2$, $\beta = 17$ W/(m²K).

so that

$$\lim_{\beta \rightarrow \infty} \Phi(z) = \frac{1}{2} V_0 + \sqrt{\alpha T_0^2 + \frac{1}{4} V_0^2} \sin \left[\Omega \left(1 - 2 \frac{z}{L} \right) \right],$$

which coincides with the expression derived in [Young 1986]. Similarly, it can be shown that (24) tends to the expression for $T(z)$ found in [Young 1986] as $\beta \rightarrow \infty$.

Figures 1 and 2 show the graphs of B^2-4AD as a function of β and V_0 , respectively.

These figures show that the expression $B^2 - 4AD$ under the square root in (19) is always greater than zero for a physically meaningful range of parameters $\beta > 0$ and $V_0 > 0$. This, in turn, guarantees that there is no “nonexistence” of solution to the given problem.

We can also derive a solution for the case $\beta = 0$ and verify that the general solution (22) reduces to this solution as $\beta \rightarrow 0$. Indeed, when $\beta = 0$, the boundary conditions (13) and (14) are reduced to the boundary conditions

$$\begin{aligned} T &= T_0, \quad \Phi = V_0 \quad \text{at } z = 0, \\ \frac{dT}{dz} &= 0, \quad \Phi = 0 \quad \text{at } z = L. \end{aligned}$$

With the same steps as before, the following expressions for $\Phi(z)$ and $T(z)$ can be derived:

$$\Phi(z) = \sqrt{\alpha T_0^2 + V_0^2} \sin\left[\tilde{\Omega}\left(1 - \frac{z}{L}\right)\right], \quad (25)$$

$$T(z) = \frac{1}{\sqrt{\alpha}} \sqrt{\alpha T_0^2 + V_0^2} \cos\left[\tilde{\Omega}\left(1 - \frac{z}{L}\right)\right], \quad (26)$$

where

$$\tilde{\Omega} := \sin^{-1}\left(\frac{V_0}{\sqrt{\alpha T_0^2 + V_0^2}}\right).$$

It can be easily verified that (22) and (24) approach (25) and (26), respectively, as $\beta \rightarrow 0$.

References

- [Antontsev and Chipot 1994] S. N. Antontsev and M. Chipot, “The thermistor problem: existence, smoothness, uniqueness, blowup”, *SIAM J. Math. Anal.* **25**:4 (1994), 1128–1156. MR Zbl
- [Berman 1976] R. Berman, *Thermal conduction in solids*, Oxford Univ. Press, 1976.
- [Engineering ToolBox 2003] Engineering ToolBox, “Overall heat transfer coefficients for fluids-heat exchanger surface combinations”, 2003, available at <https://tinyurl.com/etoolboxov>.
- [Flynn 1969] D. R. Flynn, “Measurement of thermal conductivity by steady state methods in which the sample is heated directly by passage of an electric current”, pp. 241–300 in *Thermal conductivity*, edited by R. P. Tye, Academic Press, London, 1969.
- [Fowler et al. 1992] A. C. Fowler, I. Frigaard, and S. D. Howison, “Temperature surges in current-limiting circuit devices”, *SIAM J. Appl. Math.* **52**:4 (1992), 998–1011. MR Zbl
- [Howison 2005] S. Howison, *Practical applied mathematics: modelling, analysis, approximation*, Cambridge Univ. Press, 2005. MR Zbl
- [Howison et al. 1993] S. D. Howison, J. F. Rodrigues, and M. Shillor, “Stationary solutions to the thermistor problem”, *J. Math. Anal. Appl.* **174**:2 (1993), 573–588. MR Zbl
- [Macklen 1979] E. D. Macklen, *Thermistors*, Electrochemical Pubs., Ayr, Scotland, 1979.
- [Meaden 1965] G. T. Meaden, *Electrical resistance of metals*, Int. Cryogenics Monograph Series **2**, Springer, 1965.

- [Ng 1995] K. K. Ng, *Complete guide to semiconductor devices*, McGraw-Hill, New York, 1995.
- [Shi et al. 1993] P. Shi, M. Shillor, and X. Xu, “Existence of a solution to the Stefan problem with Joule’s heating”, *J. Differential Equations* **105**:2 (1993), 239–263. MR Zbl
- [Sidi Ammi and Torres 2008] M. R. Sidi Ammi and D. F. M. Torres, “Numerical analysis of a nonlocal parabolic problem resulting from thermistor problem”, *Math. Comput. Simulation* **77**:2-3 (2008), 291–300. MR Zbl
- [Xu 2004a] X. Xu, “Exponential integrability of temperature in the thermistor problem”, *Differential Integral Equations* **17**:5-6 (2004), 571–582. MR Zbl
- [Xu 2004b] X. Xu, “Local regularity theorems for the stationary thermistor problem”, *Proc. Roy. Soc. Edinburgh Sect. A* **134**:4 (2004), 773–782. MR Zbl
- [Young 1986] J. H. Young, “Steady state Joule heating with temperature dependent conductivities”, *Appl. Sci. Res.* **43**:1 (1986), 55–65. Zbl
- [Zhou and Westbrook 1997] S. Zhou and D. R. Westbrook, “Numerical solutions of the thermistor equations”, *J. Comput. Appl. Math.* **79**:1 (1997), 101–118. MR Zbl

Received: 2017-04-26 Revised: 2017-10-05 Accepted: 2018-02-14

hrynkivv@uhd.edu

*Department of Mathematics and Statistics, University
of Houston - Downtown, Houston, TX, United States*

turchaninova2@gator.uhd.edu

*University of Houston - Downtown, Houston, TX,
United States*

involve

msp.org/involve

INVOLVE YOUR STUDENTS IN RESEARCH

Involve showcases and encourages high-quality mathematical research involving students from all academic levels. The editorial board consists of mathematical scientists committed to nurturing student participation in research. Bridging the gap between the extremes of purely undergraduate research journals and mainstream research journals, *Involve* provides a venue to mathematicians wishing to encourage the creative involvement of students.

MANAGING EDITOR

Kenneth S. Berenhaut Wake Forest University, USA

BOARD OF EDITORS

Colin Adams	Williams College, USA	Suzanne Lenhart	University of Tennessee, USA
John V. Baxley	Wake Forest University, NC, USA	Chi-Kwong Li	College of William and Mary, USA
Arthur T. Benjamin	Harvey Mudd College, USA	Robert B. Lund	Clemson University, USA
Martin Bohner	Missouri U of Science and Technology, USA	Gaven J. Martin	Massey University, New Zealand
Nigel Boston	University of Wisconsin, USA	Mary Meyer	Colorado State University, USA
Amarjit S. Budhiraja	U of North Carolina, Chapel Hill, USA	Emil Minchev	Ruse, Bulgaria
Pietro Cerone	La Trobe University, Australia	Frank Morgan	Williams College, USA
Scott Chapman	Sam Houston State University, USA	Mohammad Sal Moslehian	Ferdowsi University of Mashhad, Iran
Joshua N. Cooper	University of South Carolina, USA	Zuhair Nashed	University of Central Florida, USA
Jem N. Corcoran	University of Colorado, USA	Ken Ono	Emory University, USA
Toka Diagana	Howard University, USA	Timothy E. O'Brien	Loyola University Chicago, USA
Michael Dorff	Brigham Young University, USA	Joseph O'Rourke	Smith College, USA
Sever S. Dragomir	Victoria University, Australia	Yuval Peres	Microsoft Research, USA
Behrouz Emamizadeh	The Petroleum Institute, UAE	Y.-F. S. Pétermann	Université de Genève, Switzerland
Joel Foisy	SUNY Potsdam, USA	Robert J. Plemmons	Wake Forest University, USA
Errin W. Fulp	Wake Forest University, USA	Carl B. Pomerance	Dartmouth College, USA
Joseph Gallian	University of Minnesota Duluth, USA	Vadim Ponomarenko	San Diego State University, USA
Stephan R. Garcia	Pomona College, USA	Bjorn Poonen	UC Berkeley, USA
Anant Godbole	East Tennessee State University, USA	James Propp	U Mass Lowell, USA
Ron Gould	Emory University, USA	József H. Przytycki	George Washington University, USA
Andrew Granville	Université Montréal, Canada	Richard Rebarber	University of Nebraska, USA
Jerrold Griggs	University of South Carolina, USA	Robert W. Robinson	University of Georgia, USA
Sat Gupta	U of North Carolina, Greensboro, USA	Filip Saidak	U of North Carolina, Greensboro, USA
Jim Haglund	University of Pennsylvania, USA	James A. Sellers	Penn State University, USA
Johnny Henderson	Baylor University, USA	Andrew J. Sterge	Honorary Editor
Jim Hoste	Pitzer College, USA	Ann Trenk	Wellesley College, USA
Natalia Hritonenko	Prairie View A&M University, USA	Ravi Vakil	Stanford University, USA
Glenn H. Hurlbert	Arizona State University, USA	Antonia Vecchio	Consiglio Nazionale delle Ricerche, Italy
Charles R. Johnson	College of William and Mary, USA	Ram U. Verma	University of Toledo, USA
K. B. Kulasekera	Clemson University, USA	John C. Wierman	Johns Hopkins University, USA
Gerry Ladas	University of Rhode Island, USA	Michael E. Zieve	University of Michigan, USA

PRODUCTION

Silvio Levy, Scientific Editor


Cover: Alex Scorpion

See inside back cover or msp.org/involve for submission instructions. The subscription price for 2019 is US\$/year for the electronic version, and \$/year (+\$, if shipping outside the US) for print and electronic. Subscriptions, requests for back issues and changes of subscriber address should be sent to MSP.

Involve (ISSN 1944-4184 electronic, 1944-4176 printed) at Mathematical Sciences Publishers, 798 Evans Hall #3840, c/o University of California, Berkeley, CA 94720-3840, is published continuously online. Periodical rate postage paid at Berkeley, CA 94704, and additional mailing offices.

Involve peer review and production are managed by EditFLOW® from Mathematical Sciences Publishers.

PUBLISHED BY

 **mathematical sciences publishers**
nonprofit scientific publishing

<http://msp.org/>

© 2019 Mathematical Sciences Publishers

involve

2019 vol. 12 no. 1

Optimal transportation with constant constraint	1
WYATT BOYER, BRYAN BROWN, ALYSSA LOVING AND SARAH TAMMEN	
Fair choice sequences	13
WILLIAM J. KEITH AND SEAN GRINDATTI	
Intersecting geodesics and centrality in graphs	31
EMILY CARTER, BRYAN EK, DANIELLE GONZALEZ, RIGOBERTO FLÓREZ AND DARREN A. NARAYAN	
The length spectrum of the sub-Riemannian three-sphere	45
DAVID KLAPHECK AND MICHAEL VANVALKENBURGH	
Statistics for fixed points of the self-power map	63
MATTHEW FRIEDRICHSEN AND JOSHUA HOLDEN	
Analytical solution of a one-dimensional thermistor problem with Robin boundary condition	79
VOLODYMYR HRYNKIV AND ALICE TURCHANINOVA	
On the covering number of S_{14}	89
RYAN OPPENHEIM AND ERIC SWARTZ	
Upper and lower bounds on the speed of a one-dimensional excited random walk	97
ERIN MADDEN, BRIAN KIDD, OWEN LEVIN, JONATHON PETERSON, JACOB SMITH AND KEVIN M. STANGL	
Classifying linear operators over the octonions	117
ALEX PUTNAM AND TEVIAN DRAY	
Spectrum of the Kohn Laplacian on the Rossi sphere	125
TAWFIK ABBAS, MADELYNE M. BROWN, RAVIKUMAR RAMASAMI AND YUNUS E. ZEYTUNCU	
On the complexity of detecting positive eigenvectors of nonlinear cone maps	141
BAS LEMMENS AND LEWIS WHITE	
Antiderivatives and linear differential equations using matrices	151
YOTSANAN MEEMARK AND SONGPON SRIWONGSA	
Patterns in colored circular permutations	157
DANIEL GRAY, CHARLES LANNING AND HUA WANG	
Solutions of boundary value problems at resonance with periodic and antiperiodic boundary conditions	171
ALDO E. GARCIA AND JEFFREY T. NEUGEBAUER	

