

# involve

a journal of mathematics

Euler's formula for the zeta function  
at the positive even integers

Samyukta Krishnamurthy and Micah B. Milinovich





# Euler's formula for the zeta function at the positive even integers

Samyukta Krishnamurthy and Micah B. Milinovich

(Communicated by Filip Saidak)

We give a new proof of Euler's formula for the values of the Riemann zeta function at the positive even integers. The proof involves estimating a certain integral of elementary functions two different ways and using a recurrence relation for the Bernoulli polynomials evaluated at  $\frac{1}{2}$ .

## 1. Introduction

Let  $\zeta(s)$  denote the Riemann zeta function and let  $\eta(s) = (1 - 2^{1-s})\zeta(s)$ . Then the series representations

$$\zeta(s) = \sum_{n=1}^{\infty} \frac{1}{n^s} \quad \text{and} \quad \eta(s) = \sum_{n=1}^{\infty} \frac{(-1)^{n-1}}{n^s}$$

converge absolutely in the half-plane  $\text{Re}(s) > 1$ . For  $n \in \mathbb{N}$ , we define the Bernoulli polynomials  $B_n(x)$  via the generating function

$$\frac{ze^{xz}}{e^z - 1} = \sum_{n=0}^{\infty} B_n(x) \frac{z^n}{n!},$$

and (as usual) we call  $B_n := B_n(0)$  the  $n$ -th Bernoulli number. It follows that

$$B_2 = \frac{1}{6}, \quad B_4 = -\frac{1}{30}, \quad B_6 = \frac{1}{42}, \quad \dots, \quad B_{12} = -\frac{691}{2730}, \quad (1-1)$$

and that

$$B_{2n+1} = 0 \quad \text{for } n \in \mathbb{N}. \quad (1-2)$$

These and other standard properties of the Bernoulli numbers and Bernoulli polynomials can be found in [Montgomery and Vaughan 2007, Appendix B]. In this note we give an apparently new proof of Euler's well-known result which states that

$$\zeta(2k) = \sum_{n=1}^{\infty} \frac{1}{n^{2k}} = (-1)^{k+1} \frac{(2\pi)^{2k} B_{2k}}{2(2k)!} \quad \text{for } k \in \mathbb{N}. \quad (1-3)$$

---

*MSC2010:* primary 11M06; secondary 11B68, 11B37.

*Keywords:* Riemann zeta function, Euler, Basel problem, Bernoulli numbers, Bernoulli polynomials.

From (1-1) and (1-3), we see (as Euler did) that

$$\zeta(2) = \frac{1}{6} \pi^2, \quad \zeta(4) = \frac{1}{90} \pi^4, \quad \zeta(6) = \frac{1}{945} \pi^6, \quad \dots, \quad \zeta(12) = \frac{691}{638512875} \pi^{12}.$$

In 1734, before realizing the connection to the Bernoulli numbers, Euler derived the values of  $\zeta(2k)$  for  $k = 1, 2, \dots, 6$ . A few years later, in 1740, Euler discovered the formula in (1-3) relating  $\zeta(2k)$  to  $B_{2k}$  for  $k \in \mathbb{N}$ . Some historical remarks about Euler’s work on the Riemann zeta function and on other infinite series can be found in [Weil 1984, Chapter 3], see also [Ayoub 1974; Kline 1983; Varadarajan 2007], while references to numerous proofs of Euler’s formula in (1-3) can be found in [de Amo et al. 2011].

Instead of evaluating  $\zeta(2k)$  directly, our proof naturally evaluates the function  $\eta(s)$  at the positive even integers. Since

$$B_n\left(\frac{1}{2}\right) = -(1 - 2^{1-n})B_n \quad \text{for } n \geq 0, \tag{1-4}$$

we note that Euler’s result in (1-3) is equivalent to the formula

$$\eta(2k) = \sum_{n=1}^{\infty} \frac{(-1)^{n-1}}{n^{2k}} = (-1)^k \frac{(2\pi)^{2k} B_{2k}\left(\frac{1}{2}\right)}{2(2k)!} \quad \text{for } k \in \mathbb{N}. \tag{1-5}$$

We derive (1-5) in Section 3. Note that (1-1), (1-4), and (1-5) imply

$$\eta(2) = \frac{1}{12} \pi^2, \quad \eta(4) = \frac{7}{720} \pi^4, \quad \eta(6) = \frac{31}{30240} \pi^6, \quad \dots, \quad \eta(12) = \frac{1414477}{1307674368000} \pi^{12}.$$

Since our proof of (1-5) is more straightforward in the special case  $k = 1$ , we discuss this situation separately at the end of this article.

There is a striking resemblance between Euler’s formula (1-3), relating the values of  $\zeta(2k)$  to  $B_{2k}$ , and the formula (1-5), relating the values of  $\eta(2k)$  to  $B_{2k}\left(\frac{1}{2}\right)$ . We have chosen to write the expression in (1-5) in this manner for more than simply aesthetic reasons; indeed our proof of (1-5) relies naturally on a recursive formula for the sequence  $\{B_{2k}\left(\frac{1}{2}\right)\}_{k=0}^{\infty}$ .

### 2. A recursive formula for $B_{2k}\left(\frac{1}{2}\right)$

The Bernoulli polynomials satisfy the inversion formula

$$x^n = \frac{1}{n+1} \sum_{\ell=0}^n \binom{n+1}{\ell} B_{\ell}(x)$$

for every integer  $n \geq 0$ . Setting  $x = \frac{1}{2}$  and then observing that (1-2) and (1-4) imply  $B_n\left(\frac{1}{2}\right) = 0$  if  $n$  is odd, we derive the recursive formula

$$\frac{1}{2^{2k}} = \frac{1}{2k+1} \sum_{j=0}^k \binom{2k+1}{2j} B_{2j}\left(\frac{1}{2}\right) \quad \text{for } k \in \mathbb{N}. \tag{2-1}$$

### 3. Proof of (1-5)

We prove (1-5) by evaluating the integral

$$I_{2k} = \int_0^1 \frac{x(\log x)^{2k}}{(x^2 + 1)^2} dx \quad \text{for } k \in \mathbb{N}$$

in two different ways. On one hand, we show that

$$I_{2k} = \frac{(2k)!}{2^{2k+1}} \eta(2k) \tag{3-1}$$

by expressing the integrand as a series and then integrating term-by-term. The formula (3-1) actually holds for  $k = 0$  as well, since  $I_0 = \frac{1}{4}$  and it can be shown that  $\eta(0) = \frac{1}{2}$ . On the other hand, using the residue theorem in a relatively standard way, we derive the recursive formula

$$\frac{1}{2^{2k}} = \frac{1}{2k + 1} \sum_{j=0}^k \binom{2k+1}{2j} (-1)^j \frac{4I_{2j}}{\pi^{2j}}. \tag{3-2}$$

Comparing this expression to the recurrence relation for  $B_{2k}(\frac{1}{2})$  from the previous section, we can derive our desired expression for  $\eta(2k)$  from (3-1) and (3-2).

*Proof of (1-5).* Evidently, from (2-1) and (3-2), the sequences

$$\left\{ B_{2j}\left(\frac{1}{2}\right) \right\}_{j=0}^{\infty} \quad \text{and} \quad \left\{ (-1)^j \frac{4I_{2j}}{\pi^{2j}} \right\}_{j=0}^{\infty}$$

satisfy the same recursion relation. Moreover, since

$$4I_0 = 4 \int_0^1 \frac{x}{(x^2 + 1)^2} dx = 1 = B_0\left(\frac{1}{2}\right),$$

the initial terms in these sequences agree and therefore these sequences are equal.

Hence, from (3-1), we see that

$$B_{2k}\left(\frac{1}{2}\right) = (-1)^k \frac{4I_{2k}}{\pi^{2k}} = (-1)^k \frac{2(2k)!}{(2\pi)^{2k}} \eta(2k) \quad \text{for every } k \in \mathbb{N}. \quad \square$$

It remains to establish (3-1) and (3-2).

**3.1. Relating  $I_{2k}$  to  $\eta(2k)$ .** Integrating by parts  $2k$  times, we derive that

$$\int_0^1 x^{2n-1} (\log x)^{2k} dx = \frac{(2k)!}{(2n)^{2k+1}} \tag{3-3}$$

for positive integers  $k$  and  $n$ . Alternatively, we can prove this estimate by using that the gamma function,

$$\Gamma(z) = \int_0^{\infty} x^{z-1} e^{-x} dx \quad \text{for } \operatorname{Re}(z) > 0,$$

satisfies the relation  $\Gamma(n + 1) = n!$  for  $n \in \mathbb{N}$ . To see this, note that the variable change  $x \mapsto e^{-t/(2n)}$  implies

$$\int_0^1 x^{2n}(\log x)^{2k} \frac{dx}{x} = \frac{1}{(2n)^{2k+1}} \int_0^\infty e^{-t} t^{2k} dt = \frac{\Gamma(2k + 1)}{(2n)^{2k+1}} = \frac{(2k)!}{(2n)^{2k+1}}.$$

We now express the integrand of  $I_{2k}$  as a series, interchange the sum and the integral, and then use (3-3) to integrate term-by-term. Since

$$\begin{aligned} \frac{x}{(x^2 + 1)^2} &= -\frac{1}{2} \frac{d}{dx} \left\{ \frac{1}{1 + x^2} \right\} \\ &= -\frac{1}{2} \frac{d}{dx} \left\{ \sum_{n=0}^\infty (-1)^n x^{2n} \right\} = \sum_{n=1}^\infty n(-1)^{n-1} x^{2n-1} \end{aligned} \tag{3-4}$$

for  $|x| < 1$ , we have

$$\begin{aligned} I_{2k} &= \int_0^1 \frac{x(\log x)^{2k}}{(x^2 + 1)^2} dx = \int_0^1 \sum_{n=1}^\infty n(-1)^{n-1} x^{2n-1} (\log x)^{2k} dx \\ &= \sum_{n=1}^\infty n(-1)^{n-1} \int_0^1 x^{2n-1} (\log x)^{2k} dx \\ &= \sum_{n=1}^\infty n(-1)^{n-1} \frac{(2k)!}{(2n)^{2k+1}} = \frac{(2k)!}{2^{2k+1}} \eta(2k) \end{aligned}$$

for every  $k \in \mathbb{N}$ . This proves (3-1). Note that the interchange of summation and integration is justified using Fubini’s theorem since, for every  $k \in \mathbb{N}$ , (3-3) implies


$$\begin{aligned} \sum_{n=1}^\infty \int_0^1 |n(-1)^{n-1} x^{2n-1} \log^{2k} x| dx &= \sum_{n=1}^\infty n \int_0^1 x^{2n-1} (\log x)^{2k} dx \\ &= \frac{(2k)!}{2^{2k+1}} \sum_{n=1}^\infty \frac{1}{n^{2k}} < \infty. \end{aligned}$$

**3.2. A recursive formula for  $I_{2k}$ .** Making the variable change  $x \mapsto 1/x$ , it follows that

$$\begin{aligned} \int_0^1 \frac{x(\log x)^{2k}}{(x^2 + 1)^2} dx &= \int_1^\infty \frac{x(\log x)^{2k}}{(x^2 + 1)^2} dx, \\ \int_0^1 \frac{x(\log x)^{2k+1}}{(x^2 + 1)^2} dx &= -\int_1^\infty \frac{x(\log x)^{2k+1}}{(x^2 + 1)^2} dx \end{aligned}$$

for integers  $k \geq 0$ . Therefore

$$I_{2k} = \frac{1}{2} \int_0^\infty \frac{x(\log x)^{2k}}{(x^2 + 1)^2} dx \quad \text{and} \quad \int_0^\infty \frac{x(\log x)^{2k+1}}{(x^2 + 1)^2} dx = 0. \tag{3-5}$$



**Figure 1**

Now we introduce the complex-valued function

$$f(z) = \frac{z(\log z)^{2k+1}}{(1+z^2)^2},$$

where  $\log z$  denotes the branch of the logarithm in  $\mathbb{C}$  with  $|z| > 0$  and  $-\frac{\pi}{2} < \arg z < \frac{3\pi}{2}$ . Note that the power of  $\log z$  in the numerator of  $f(z)$  is one power higher than the power of  $\log x$  appearing in the integrand of  $I_{2k}$ . We integrate  $f(z)$  around the positively oriented simple closed contour (shown in Figure 1) composed of the line segment  $[\varepsilon, R]$  along the real-axis, the semicircle  $\Gamma_R$  centered at 0 of radius  $R$  starting at  $z = R$  passing through  $z = iR$  and ending at  $z = -R$ , the line segment  $[-R, -\varepsilon]$  along the real-axis, and finally the semicircle  $\Gamma_\varepsilon$  centered at 0 of radius  $\varepsilon$  starting at  $z = -\varepsilon$  passing through  $z = i\varepsilon$  and ending at  $z = \varepsilon$ . Here  $\varepsilon$  and  $R$  denote real numbers satisfying  $0 < \varepsilon < 1 < R < \infty$ . The only singularity of  $f(z)$  inside this contour is a double pole at  $z = i$ . Therefore the residue theorem implies

$$2\pi i \operatorname{Res}_{z=i} f(z) = \int_\varepsilon^R \frac{x(\log x)^{2k+1}}{(1+x^2)^2} dx + \int_{\Gamma_R} f(z) dz + \int_{-R}^{-\varepsilon} \frac{x(\log(-x) + i\pi)^{2k+1}}{(1+x^2)^2} dx + \int_{\Gamma_\varepsilon} f(z) dz, \quad (3-6)$$

where the logarithms in the first and third integrals on the right-hand side denote the natural logarithm. Estimating trivially, we have

$$\left| \int_{\Gamma_\varepsilon} f(z) dz \right| \leq \text{length}(\Gamma_\varepsilon) \cdot \max_{z \in \Gamma_\varepsilon} |f(z)| \leq (\pi \varepsilon) \left( \frac{\varepsilon(\log(-\varepsilon) + \pi)^{2k+1}}{(1-\varepsilon^2)^2} \right) \rightarrow 0$$

as  $\varepsilon \rightarrow 0^+$  and

$$\left| \int_{\Gamma_R} f(z) dz \right| \leq \text{length}(\Gamma_R) \cdot \max_{z \in \Gamma_R} |f(z)| \leq (\pi R) \left( \frac{R(\log R + \pi)^{2k+1}}{(1-R^2)^2} \right) \rightarrow 0$$

as  $R \rightarrow +\infty$ . It follows that

$$2\pi i \operatorname{Res}_{z=i} f(z) = \int_0^\infty \frac{x(\log x)^{2k+1}}{(1+x^2)^2} dx + \int_{-\infty}^0 \frac{x(\log(-x) + i\pi)^{2k+1}}{(1+x^2)^2} dx.$$

By the second expression in (3-5), the first integral on the right-hand side equals 0. Sending  $x \mapsto -x$ , the second integral on the right-hand side equals

$$-\int_0^\infty \frac{x(\log x + i\pi)^{2k+1}}{(1+x^2)^2} dx = -\sum_{\ell=0}^{2k+1} \binom{2k+1}{\ell} (i\pi)^{2k-\ell+1} \int_0^\infty \frac{x(\log x)^\ell}{(1+x^2)^2} dx.$$

Again by (3-5), the terms in the sum with  $\ell$  odd vanish. Hence, for even  $\ell$ , letting  $\ell = 2j$  and using the first expression in (3-5), we have

$$2\pi i \operatorname{Res}_{z=i} f(z) = -(i\pi)^{2k+1} \sum_{j=0}^k \binom{2k+1}{2j} (-1)^j \frac{2I_{2j}}{\pi^{2j}}. \tag{3-7}$$

On the other hand, a straightforward calculation shows that

$$\operatorname{Res}_{z=i} f(z) = \lim_{z \rightarrow i} \frac{d}{dz} \left\{ \frac{z(\log z)^{2k+1}}{(z+i)^2} \right\} = -\frac{(2k+1)(i\pi)^{2k}}{2^{2k+2}}.$$

Inserting this into (3-7) and dividing by  $-(2k+1)(i\pi)^{2k+1}/2$ , we conclude that

$$\frac{1}{2^{2k}} = \frac{1}{2k+1} \sum_{j=0}^k \binom{2k+1}{2j} (-1)^j \frac{4I_{2j}}{\pi^{2j}},$$

as claimed.

**3.3. Remarks on the case  $k = 1$ .** Historically, the *Basel problem* asked for a closed-form evaluation of the sum

$$\zeta(2) = \sum_{n=1}^\infty \frac{1}{n^2}.$$

As mentioned in the Introduction, this problem was solved by Euler in 1734. Therefore, there is perhaps special interest in a direct proof of the equivalent problem of showing that

$$\eta(2) = \sum_{n=1}^\infty \frac{(-1)^{n-1}}{n^2} = \frac{\pi^2}{12}.$$

In this special case, our proof above can be simplified since there is no need to appeal to properties of the Bernoulli polynomials, the gamma function, or recursion relations. We sketch the details of this calculation for the interested reader.

In this case, we evaluate the integral

$$I_2 = \int_0^1 \frac{x(\log x)^2}{(x^2+1)^2} dx$$

in two different ways. Integrating by parts twice, it can be shown that

$$\int_0^1 x^{2n-1} (\log x)^2 dx = \frac{1}{4n^3} \quad \text{for } n \in \mathbb{N}. \tag{3-8}$$



Therefore, using the series expansion in (3-4), it follows that

$$\begin{aligned} I_2 &= \int_0^1 \frac{x(\log x)^2}{(x^2 + 1)^2} dx = \int_0^1 \sum_{n=1}^{\infty} n(-1)^{n-1} x^{2n-1} (\log x)^2 dx \\ &= \sum_{n=1}^{\infty} n(-1)^{n-1} \int_0^1 x^{2n-1} (\log x)^2 dx \\ &= \frac{1}{4} \sum_{n=1}^{\infty} \frac{(-1)^{n-1}}{n^2} = \frac{\eta(2)}{4}. \end{aligned}$$

As in Section 3.1, the interchange of summation and integration can be justified using Fubini's theorem. On the other hand, making the variable change  $x \mapsto 1/x$ , it follows that

$$I_2 = \int_0^1 \frac{x(\log x)^2}{(x^2 + 1)^2} dx = \int_1^{\infty} \frac{x(\log x)^2}{(x^2 + 1)^2} dx.$$

Therefore

$$I_2 = \frac{1}{2} \int_0^{\infty} \frac{x(\log x)^2}{(x^2 + 1)^2} dx. \tag{3-9}$$

In order to evaluate this integral, we apply the residue theorem in a manner similar to that in the previous section. We integrate the complex-valued function

$$f(z) = \frac{z(\log z)^3}{(1 + z^2)^2}$$

around the positively oriented simple closed contour shown in Figure 1. As before,  $\log z$  denotes the branch of the logarithm in  $\mathbb{C}$  with  $|z| > 0$  and  $-\frac{\pi}{2} < \arg z < \frac{3\pi}{2}$ , while  $\varepsilon$  and  $R$  denote real numbers satisfying  $0 < \varepsilon < 1 < R < \infty$ . Then the residue theorem implies

$$\begin{aligned} 2\pi i \operatorname{Res}_{z=i} f(z) &= \int_{\varepsilon}^R \frac{x(\log x)^3}{(1 + x^2)^2} dx + \int_{\Gamma_R} f(z) dz \\ &\quad + \int_{-R}^{-\varepsilon} \frac{x(\log(-x) + i\pi)^3}{(1 + x^2)^2} dx + \int_{\Gamma_{\varepsilon}} f(z) dz, \end{aligned}$$

where the logarithms in the first and third integrals on the right-hand side denote the natural logarithm. As was shown in the previous section, the second and fourth integrals on the right-hand side tend to 0 as  $R \rightarrow +\infty$  and  $\varepsilon \rightarrow 0^+$ , respectively. Since the only singularity of  $f(z)$  inside this contour is a double pole at  $z = i$  with

$$\operatorname{Res}_{z=i} f(z) = \lim_{z \rightarrow i} \frac{d}{dz} \left\{ \frac{z(\log z)^3}{(z + i)^2} \right\} = \frac{3\pi^2}{16},$$

it follows that

$$\begin{aligned} \frac{3\pi^3 i}{8} &= \int_0^\infty \frac{x(\log x)^3}{(1+x^2)^2} dx + \int_{-\infty}^0 \frac{x(\log(-x) + i\pi)^3}{(1+x^2)^2} dx \\ &= \int_0^\infty \frac{x(\log x)^3}{(1+x^2)^2} dx - \int_0^\infty \frac{x(\log x + i\pi)^3}{(1+x^2)^2} dx. \end{aligned}$$

Here we have made the variable change  $x \mapsto -x$  in the second integral. Expanding the factor  $(\log x + i\pi)^3$ , taking imaginary parts of both sides of the equation, and then using (3-9), we deduce that

$$\frac{3\pi^3}{8} = \pi^3 \int_0^\infty \frac{x}{(1+x^2)^2} dx - 3\pi \int_0^\infty \frac{x(\log x)^2}{(1+x^2)^2} dx = \frac{\pi^3}{2} - 6\pi I_2.$$

This implies  $I_2 = \pi^2/48$ . Combining this with our previous observation that  $I_2 = \eta(2)/4$ , we conclude that  $\eta(2) = \pi^2/12$ .

### Acknowledgements

Milinovich is supported in part by the NSA Young Investigator Grant H98230-16-1-0311.

### References

- [de Amo et al. 2011] E. de Amo, M. Díaz Carrillo, and J. Fernández-Sánchez, “Another proof of Euler’s formula for  $\zeta(2k)$ ”, *Proc. Amer. Math. Soc.* **139**:4 (2011), 1441–1444. MR Zbl
- [Ayoub 1974] R. Ayoub, “Euler and the zeta function”, *Amer. Math. Monthly* **81** (1974), 1067–1086. MR Zbl
- [Kline 1983] M. Kline, “Euler and infinite series”, *Math. Mag.* **56**:5 (1983), 307–314. MR Zbl
- [Montgomery and Vaughan 2007] H. L. Montgomery and R. C. Vaughan, *Multiplicative number theory, I: Classical theory*, Cambridge Studies in Advanced Mathematics **97**, Cambridge University Press, 2007. MR Zbl
- [Varadarajan 2007] V. S. Varadarajan, “Euler and his work on infinite series”, *Bull. Amer. Math. Soc. (N.S.)* **44**:4 (2007), 515–539. MR Zbl
- [Weil 1984] A. Weil, *Number theory: an approach through history, from Hammurapi to Legendre*, Birkhäuser, Boston, MA, 1984. MR Zbl

Received: 2017-06-12    Revised: 2018-07-30    Accepted: 2018-10-28

skrishnamurt@umass.edu

*Department of Physics, University of Mississippi,  
University, MS, United States*

*Current address:*

*Department of Physics, University of Massachusetts,  
Amherst, MA, United States*

mbmilino@olemiss.edu

*Department of Mathematics, University of Mississippi,  
University, MS, United States*

# involve

msp.org/involve

## INVOLVE YOUR STUDENTS IN RESEARCH

*Involve* showcases and encourages high-quality mathematical research involving students from all academic levels. The editorial board consists of mathematical scientists committed to nurturing student participation in research. Bridging the gap between the extremes of purely undergraduate research journals and mainstream research journals, *Involve* provides a venue to mathematicians wishing to encourage the creative involvement of students.

### MANAGING EDITOR

Kenneth S. Berenhaut Wake Forest University, USA

### BOARD OF EDITORS

Colin Adams	Williams College, USA	Chi-Kwong Li	College of William and Mary, USA
Arthur T. Benjamin	Harvey Mudd College, USA	Robert B. Lund	Clemson University, USA
Martin Bohner	Missouri U of Science and Technology, USA	Gaven J. Martin	Massey University, New Zealand
Nigel Boston	University of Wisconsin, USA	Mary Meyer	Colorado State University, USA
Amarjit S. Budhiraja	U of N Carolina, Chapel Hill, USA	Frank Morgan	Williams College, USA
Pietro Cerone	La Trobe University, Australia	Mohammad Sal Moslehian	Ferdowsi University of Mashhad, Iran
Scott Chapman	Sam Houston State University, USA	Zuhair Nashed	University of Central Florida, USA
Joshua N. Cooper	University of South Carolina, USA	Ken Ono	Emory University, USA
Jem N. Corcoran	University of Colorado, USA	Yuval Peres	Microsoft Research, USA
Toka Diagana	Howard University, USA	Y.-F. S. Pétermann	Université de Genève, Switzerland
Michael Dorff	Brigham Young University, USA	Jonathon Peterson	Purdue University, USA
Sever S. Dragomir	Victoria University, Australia	Robert J. Plemmons	Wake Forest University, USA
Joel Foisy	SUNY Potsdam, USA	Carl B. Pomerance	Dartmouth College, USA
Errin W. Fulp	Wake Forest University, USA	Vadim Ponomarenko	San Diego State University, USA
Joseph Gallian	University of Minnesota Duluth, USA	Bjorn Poonen	UC Berkeley, USA
Stephan R. Garcia	Pomona College, USA	József H. Przytycki	George Washington University, USA
Anant Godbole	East Tennessee State University, USA	Richard Rebarber	University of Nebraska, USA
Ron Gould	Emory University, USA	Robert W. Robinson	University of Georgia, USA
Sat Gupta	U of North Carolina, Greensboro, USA	Javier Rojo	Oregon State University, USA
Jim Haglund	University of Pennsylvania, USA	Filip Saidak	U of North Carolina, Greensboro, USA
Johnny Henderson	Baylor University, USA	Hari Mohan Srivastava	University of Victoria, Canada
Glenn H. Hurlbert	Arizona State University, USA	Andrew J. Sterge	Honorary Editor
Charles R. Johnson	College of William and Mary, USA	Ann Trenk	Wellesley College, USA
K. B. Kulasekera	Clemson University, USA	Ravi Vakil	Stanford University, USA
Gerry Ladas	University of Rhode Island, USA	Antonia Vecchio	Consiglio Nazionale delle Ricerche, Italy
David Larson	Texas A&M University, USA	John C. Wierman	Johns Hopkins University, USA
Suzanne Lenhart	University of Tennessee, USA	Michael E. Zieve	University of Michigan, USA

### PRODUCTION

Silvio Levy, Scientific Editor


Cover: Alex Scorpan

See inside back cover or [msp.org/involve](http://msp.org/involve) for submission instructions. The subscription price for 2019 is US \$195/year for the electronic version, and \$260/year (+\$35, if shipping outside the US) for print and electronic. Subscriptions, requests for back issues and changes of subscriber address should be sent to MSP.

Involve (ISSN 1944-4184 electronic, 1944-4176 printed) at Mathematical Sciences Publishers, 798 Evans Hall #3840, c/o University of California, Berkeley, CA 94720-3840, is published continuously online. Periodical rate postage paid at Berkeley, CA 94704, and additional mailing offices.

Involve peer review and production are managed by EditFLOW® from Mathematical Sciences Publishers.

PUBLISHED BY

 **mathematical sciences publishers**  
nonprofit scientific publishing

<http://msp.org/>

© 2019 Mathematical Sciences Publishers

# involve

2019 vol. 12 no. 4

Euler's formula for the zeta function at the positive even integers	541
SAMYUKTA KRISHNAMURTHY AND MICAH B. MILINOVICH	
Descents and des-Wilf equivalence of permutations avoiding certain nonclassical patterns	549
CADEN BIELAWA, ROBERT DAVIS, DANIEL GREESON AND QINHAN ZHOU	
The classification of involutions and symmetric spaces of modular groups	565
MARC BESSON AND JENNIFER SCHAEFER	
When is $a^n + 1$ the sum of two squares?	585
GREG DRESDEN, KYLIE HESS, SAIMON ISLAM, JEREMY ROUSE, AARON SCHMITT, EMILY STAMM, TERRIN WARREN AND PAN YUE	
Irreducible character restrictions to maximal subgroups of low-rank classical groups of types $B$ and $C$	607
KEMPTON ALBEE, MIKE BARNES, AARON PARKER, ERIC ROON AND A. A. SCHAEFFER FRY	
Prime labelings of infinite graphs	633
MATTHEW KENIGSBERG AND OSCAR LEVIN	
Positional strategies in games of best choice	647
AARON FOWLKES AND BRANT JONES	
Graphs with at most two trees in a forest-building process	659
STEVE BUTLER, MISA HAMANAKA AND MARIE HARDT	
Log-concavity of Hölder means and an application to geometric inequalities	671
AUREL I. STAN AND SERGIO D. ZAPETA-TZUL	
Applying prospect theory to multiattribute problems with independence assumptions	687
JACK STANLEY AND FRANK P. A. COOLEN	
On weight-one solvable configurations of the Lights Out puzzle	713
YUKI HAYATA AND MASAKAZU YAMAGISHI	

