

involve

a journal of mathematics

Positive solutions to singular second-order boundary value
problems for dynamic equations

Curtis Kunkel and Alex Lancaster



Positive solutions to singular second-order boundary value problems for dynamic equations

Curtis Kunkel and Alex Lancaster

(Communicated by Johnny Henderson)

We study singular second-order boundary value problems with mixed boundary conditions on an infinitely discrete time scale. We prove the existence of a positive solution by means of a lower and upper solutions method and the Brouwer fixed-point theorem, in conjunction with perturbation methods used to approximate regular problems.

1. Introduction

This paper continues the work done previously by Kunkel [2008], where he studied a singular second-order boundary value problem in purely discrete time scales of nonuniform step size. Although similar throughout most of the time scale, this result is different in the fact that the time scale itself has a limit point at the right-side boundary condition, forcing a nearly continuous behavior at that end. If this limiting condition were not present, the result would be trivial using [Kunkel 2008], but as it stands, this result continues to expand the work of that paper to another type of time scale.

More specifically, [Kunkel 2008] dealt with the discrete boundary value problem

$$\begin{aligned} u^{\Delta\Delta}(t_{i-1}) + f(t_i, u(t_i), u^\Delta(t_{i-1})) &= 0, \quad t \in \mathbb{T}^\circ, \\ u^\Delta(t_0) = u(t_{n+1}) &= 0, \end{aligned}$$

where \mathbb{T}° is the discrete interval of nonuniform step size $[t_1, t_n] := \{t_1, t_2, \dots, t_n\}$ and $f(t, x, y)$ is singular in x . This work was an extension of a previous result by Rachůnková and Rachůnek [2006], where they studied a singular second-order boundary value problem for the discrete p -Laplacian, $\phi_p(x) = |x|^{p-2}x$, $p > 1$.

MSC2010: 34B16, 34B18, 34B40, 39A10.

Keywords: singular boundary value problems, time scales, mixed conditions, lower and upper solutions, Brouwer fixed-point theorem, approximate regular problems.

In particular, Rachůnková and Rachůnek dealt with the discrete boundary value problem

$$\begin{aligned}\Delta(\phi_p(\Delta u(t-1))) + f(t, u(t), \Delta u(t-1)) &= 0, \quad t \in [1, T+1], \\ \Delta u(0) = u(T+2) &= 0,\end{aligned}$$

in which $f(t, x, y)$ was singular in x .

Combine these works with [Kunkel 2006], which deals with the continuous boundary value problem

$$\begin{aligned}u''(t) + f(t, u(t)) &= 0, \quad t \in (0, 1), \\ u'(0) = u(1) &= 0,\end{aligned}$$

where $f(t, x)$ is singular in x , and you have a similar boundary value problem scenario across time scales ranging from being entirely continuous to varying degrees of discrete. This result fits between these two ends of the time scale continuum being a discrete interval with a continuous point, the ultimate goal of which would be to create a unifying theorem for this type of problem across all types of time scales (forthcoming).

The methods in this paper rely heavily on lower and upper solution methods in conjunction with an application of the Brouwer fixed-point theorem [Zeidler 1986]. We consider only the singular second-order boundary value problem, while letting our function range over an infinitely discrete interval of nonuniform step size, included in which is the limit point. We will provide definitions of appropriate lower and upper solutions. The lower and upper solutions will be applied to nonsingular perturbations of our nonlinear problem, ultimately giving rise to our boundary value problem by passing to the limit.

Lower and upper solutions have been used extensively in establishing solutions of boundary value problems for finite difference equations. Representative works include [Bao et al. 2012; Henderson and Kunkel 2006; Precup 2016].

Singular boundary value problems have also received a good deal of attention. Representative works include [Agarwal and O'Regan 1999; Precup 2016; Rachůnková and Rachůnek 2009].

2. Preliminaries

We now state some definitions used throughout the remainder of the paper, many of which can be found in [Bohner and Peterson 2001; Kelley and Peterson 1991]. Some definitions are required prior to the introduction of the problem we intend to solve.

Definition 2.1. For $i = 1, 2, 3, \dots$, let $t_i = 1 - 1/i$. Define the time scale

$$\mathbb{T} = \{t_i\}_{i=1}^{\infty} \cup \{1\}.$$

We conveniently make note of the standard notation for both forward and backward jump operators on time scales of this nature.

Definition 2.2. The forward step operator $\sigma : \mathbb{T} \rightarrow \mathbb{T}$ is defined by

$$\sigma(t) := \inf\{s \in \mathbb{T} : s > t\}.$$

The backward step operator $\rho : \mathbb{T} \rightarrow \mathbb{T}$ is defined by

$$\rho(t) := \sup\{s \in \mathbb{T} : s < t\}.$$

Definition 2.3. For the function $u : \mathbb{T} \rightarrow \mathbb{R}$, define the delta derivative $u^\Delta : \mathbb{T} \rightarrow \mathbb{R}$ by

$$u^\Delta(t) := \frac{u(\sigma(t)) - u(t)}{\mu(t)},$$

where $\mu(t) := \sigma(t) - t$. Note that μ is the graininess function.

Having introduced these definitions, we can now consider the following second-order dynamic equation, which will be our focus throughout this paper:

$$u^{\Delta\Delta}(t_{i-1}) + f(t_i, u(t_i), u^\Delta(t_{i-1})) = 0, \quad t_{i-1} \in \mathbb{T}, \tag{1}$$

satisfying the mixed boundary conditions,

$$u^\Delta(0) = u(1) = 0. \tag{2}$$

Our goal is to prove the existence of a positive solution to this problem (1), (2), where f has a specific type of singularity as explained below.

Definition 2.4. Define a solution to problem (1), (2) to mean a function $u : \mathbb{T} \rightarrow \mathbb{R}$ such that u satisfies (1) on \mathbb{T} and also satisfies the boundary conditions (2). If $u(t) > 0$ for $t \in \mathbb{T}$, except possibly at the boundary conditions, we call u a positive solution to problem (1), (2).

Definition 2.5. Let $D \subseteq \mathbb{R}^2$. We say f is continuous on $\mathbb{T} \times D$ if $f(\cdot, x, y)$ is defined on \mathbb{T} for each $(x, y) \in D$ and if $f(t, \cdot, \cdot)$ is continuous on D for each $t \in \mathbb{T}$.

Definition 2.6. Let $D \subseteq \mathbb{R}^2$. Let $f : \mathbb{T} \times D \rightarrow \mathbb{R}$. If $D = \mathbb{R}^2$, then we call (1), (2) a regular problem. If $D \subsetneq \mathbb{R}^2$ and f has singularities on the boundary of D , then we call (1), (2) a singular problem.

We assume the following throughout this paper:

- (A) $D = [0, \infty) \times \mathbb{R}$.
- (B) f is continuous on $\mathbb{T} \times D$.
- (C) $f(t, x, y)$ has a singularity at $x = 0$; i.e., $\limsup_{x \rightarrow 0^+} |f(t, x, y)| = \infty$ for $t \in \mathbb{T}$ and $y \in \mathbb{R}$.

3. Lower and upper solutions method

For the purpose of establishing a lower and upper solutions method to be used in solving our pre-existing singular problem, we first consider the following regular problem:

$$u^{\Delta\Delta}(t_{i-1}) + h(t_i, u(t_i), u^{\Delta}(t_{i-1})) = 0, \quad t_{i-1} \in \mathbb{T}, \tag{3}$$

where h is continuous on $\mathbb{T} \times \mathbb{R}^2$ and the same boundary conditions (2) are satisfied. Now, (3), (2) is clearly a regular problem and it is our current goal to establish a lower and upper solutions method as a means to establish an existence result. To this end, we first must define what is meant by a lower and an upper solution.

Definition 3.1. Let $\alpha : \mathbb{T} \rightarrow \mathbb{R}$. We call α a lower solution of problem (3), (2) if

$$\alpha^{\Delta\Delta}(t_{i-1}) + h(t_i, \alpha(t_i), \alpha^{\Delta}(t_{i-1})) \geq 0, \quad t_{i-1} \in \mathbb{T}, \tag{4}$$

satisfying

$$\alpha^{\Delta}(0) \geq 0, \quad \alpha(1) \leq 0. \tag{5}$$

Definition 3.2. Let $\beta : \mathbb{T} \rightarrow \mathbb{R}$. We call β an upper solution of problem (3), (2) if

$$\beta^{\Delta\Delta}(t_{i-1}) + h(t_i, \beta(t_i), \beta^{\Delta}(t_{i-1})) \leq 0, \quad t_{i-1} \in \mathbb{T}, \tag{6}$$

satisfying

$$\beta^{\Delta}(0) \leq 0, \quad \beta(1) \geq 0. \tag{7}$$

Theorem 3.3 (lower and upper solutions method). *Let α and β be lower and upper solutions of the regular problem (3), (2), respectively, where $\alpha \leq \beta$ on \mathbb{T} . Let $h(t, x, y)$ be continuous on $\mathbb{T} \times \mathbb{R}^2$ and nonincreasing in its y -variable. Then (3), (2) has a solution u satisfying*

$$\alpha(t) \leq u(t) \leq \beta(t), \quad t \in \mathbb{T}.$$

Proof. We proceed with this proof through a sequence of steps involving modifications of the function h .

Step 1: For $t_{i-1} \in \mathbb{T}$ and $(x, y) \in \mathbb{R}^2$, define

$$\begin{aligned} & \tilde{h}\left(t_i, x, \frac{x-y}{\mu(t_{i-1})}\right) \\ &= \begin{cases} h\left(t_i, \beta(t_i), \frac{\beta(t_i) - S(t_{i-1}, y)}{\mu(t_{i-1})}\right) - \frac{x - \beta(t_i)}{x - \beta(t_i) + 1}, & x > \beta(t_i), \\ h\left(t_i, x, \frac{x - S(t_{i-1}, y)}{\mu(t_{i-1})}\right), & \alpha(t_i) \leq x \leq \beta(t_i), \\ h\left(t_i, \alpha(t_i), \frac{\alpha(t_i) - S(t_{i-1}, y)}{\mu(t_{i-1})}\right) + \frac{\alpha(t_i) - x}{\alpha(t_i) - x + 1}, & x < \alpha(t_i), \end{cases} \tag{8} \end{aligned}$$

where,

$$S(t_{i-1}, y) = \begin{cases} \beta(t_{i-1}), & y > \beta(t_{i-1}), \\ y, & \alpha(t_{i-1}) \leq y \leq \beta(t_{i-1}), \\ \alpha(t_{i-1}), & y < \alpha(t_{i-1}). \end{cases}$$

Given this construction, \tilde{h} is continuous on $\mathbb{T} \times \mathbb{R}^2$ and there exists $M > 0$ so that

$$|\tilde{h}(t, x, y)| \leq M$$

for all $t \in \mathbb{T}$ and $(x, y) \in \mathbb{R}^2$.

We now study the auxiliary equation

$$u^{\Delta\Delta}(t_{i-1}) + \tilde{h}(t_i, u(t_i), u^{\Delta}(t_{i-1})) = 0, \quad t_{i-1} \in \mathbb{T}, \quad (9)$$

satisfying boundary conditions (2). Our immediate goal is to prove the existence of a solution to problem (9), (2).

Step 2: For this existence result, we lay the foundation to use the Brouwer fixed-point theorem. To this end, define

$$E = \{u : \mathbb{T} \rightarrow \mathbb{R} : u^{\Delta}(0) = u(1) = 0\}.$$

Also, define

$$\|u\| = \max\{|u(t)| : t \in \mathbb{T}\}.$$

Given E and $\|\cdot\|$, we say E is a Banach space. Further, we define an operator $\mathcal{T} : E \rightarrow E$ by

$$(\mathcal{T}u)(t_k) = \sum_{j=k}^{\infty} \mu(t_j) \sum_{i=2}^j \mu(t_{i-1}) \tilde{h}(t_i, u(t_i), u^{\Delta}(t_{i-1})). \quad (10)$$

\mathcal{T} is a continuous operator. Moreover, from the bounds placed on \tilde{h} in Step 1 and from (10), if $r > M$, then $T(\overline{B(r)}) \subseteq \overline{B(r)}$, where $B(r) := \{u \in E : \|u\| < r\}$. Hence, by the Brouwer fixed-point theorem [Zeidler 1986], there exists $u \in \overline{B(r)}$ such that $u = \mathcal{T}u$.

Step 3: We now show that u is a fixed point of \mathcal{T} if and only if u is a solution to the problem (9), (2).

To this end, let us first assume that u solves the problem (9), (2). Then, since the boundary conditions (2) are satisfied, $u \in E$.

It is convenient for the first part of this subproof to consider a relabeling of the points in \mathbb{T} as follows: let $\tau_{\infty} = \lim_{i \rightarrow \infty} \tau_i := t_1 = 0$, let $\tau_0 := 1$, and, for each $i > 0$, let there exist some $j > 0$ so that $t_i = \tau_j$, $t_{i+1} = \tau_{j-1}$, etc. Using this notation, we then consider

$$u^{\Delta}(\tau_1) = \frac{u(\tau_0) - u(\tau_1)}{\mu(\tau_1)} = \frac{-u(\tau_1)}{\mu(\tau_1)},$$

and we have

$$u(\tau_1) = -\mu(\tau_1)u^\Delta(\tau_1).$$

Also,

$$u^\Delta(\tau_2) = \frac{u(\tau_1) - u(\tau_2)}{\mu(\tau_2)} = \frac{-\mu(\tau_1)u^\Delta(\tau_1) - u(\tau_2)}{\mu(\tau_2)},$$

and we have

$$u(\tau_2) = -\mu(\tau_1)u^\Delta(\tau_1) - \mu(\tau_2)u^\Delta(\tau_2).$$

Continuing in this manner, we have, for $m > 0$,

$$u(\tau_m) = -\sum_{i=1}^m \mu(\tau_i)u^\Delta(\tau_i). \tag{11}$$

And, given our relabeling between the τ 's and the t 's, we can conclude that for each $m > 0$ there exists some $k > 0$ such that

$$u(\tau_m) = u(t_k) = -\sum_{i=k}^{\infty} \mu(\tau_i)u^\Delta(\tau_i).$$

We also have

$$u^{\Delta\Delta}(t_1) = \frac{u^\Delta(t_2) - u^\Delta(t_1)}{\mu(t_1)} = \frac{u^\Delta(t_2) - u^\Delta(0)}{\mu(t_1)} = \frac{u^\Delta(t_2)}{\mu(t_1)},$$

and from (9) we have $u^{\Delta\Delta}(t_1) = -\tilde{h}(t_2, u(t_2), u^\Delta(t_1))$, which yields

$$u^\Delta(t_2) = -\mu(t_1)\tilde{h}(t_2, u(t_2), u^\Delta(t_1)).$$

Similarly, we have

$$u^{\Delta\Delta}(t_2) = \frac{u^\Delta(t_3) - u^\Delta(t_2)}{\mu(t_2)} = -\tilde{h}(t_3, u(t_3), u^\Delta(t_2)),$$

and via substitution of $u^\Delta(t_2)$ and simply solving for $u^\Delta(t_3)$, we have

$$u^\Delta(t_3) = -\mu(t_1)\tilde{h}(t_2, u(t_2), u^\Delta(t_1)) - \mu(t_2)\tilde{h}(t_3, u(t_3), u^\Delta(t_2)).$$

Continuing in this manner, we conclude that

$$u^\Delta(t_j) = -\sum_{i=2}^j \mu(t_{i-1})\tilde{h}(t_i, u(t_i), u^\Delta(t_{i-1})). \tag{12}$$

By substituting (12) into (11), we see that for $k > 0$

$$\begin{aligned} u(t_k) &= -\sum_{j=k}^{\infty} \mu(t_j) \left(-\sum_{i=2}^j \mu(t_{i-1})\tilde{h}(t_i, u(t_i), u^\Delta(t_{i-1})) \right) \\ &= \sum_{j=k}^{\infty} \mu(t_j) \sum_{i=2}^j \mu(t_{i-1})\tilde{h}(t_i, u(t_i), u^\Delta(t_{i-1})) = (\mathcal{T}u)(t_k). \end{aligned}$$

We now assume that u is a fixed point of \mathcal{T} , i.e., $u = \mathcal{T}u$. Then,

$$u(t_k) = (\mathcal{T}u)(t_k) = \sum_{j=k}^{\infty} \mu(t_j) \sum_{i=2}^j \mu(t_{i-1}) \tilde{h}(t_i, u(t_i), u^\Delta(t_{i-1})).$$

Also,

$$\begin{aligned} u^\Delta(t_{k-1}) &= \frac{u(t_k) - u(t_{k-1})}{\mu(t_{k-1})} \\ &= \frac{(\sum_{j=k}^{\infty} \mu(t_j) \sum_{i=2}^j \mu(t_{i-1}) \tilde{h}(t_i, u(t_i), u^\Delta(t_{i-1})))}{\mu(t_{k-1})} \\ &\quad - \frac{(\sum_{j=k-1}^{\infty} \mu(t_j) \sum_{i=2}^j \mu(t_{i-1}) \tilde{h}(t_i, u(t_i), u^\Delta(t_{i-1})))}{\mu(t_{k-1})} \\ &= - \frac{\mu(t_{k-1}) \sum_{i=2}^{k-1} \mu(t_{i-1}) \tilde{h}(t_i, u(t_i), u^\Delta(t_{i-1}))}{\mu(t_{k-1})} \\ &= - \sum_{i=2}^{k-1} \mu(t_{i-1}) \tilde{h}(t_i, u(t_i), u^\Delta(t_{i-1})), \end{aligned}$$

and

$$\begin{aligned} u^{\Delta\Delta}(t_{k-1}) &= \frac{u^\Delta(t_k) - u^\Delta(t_{k-1})}{\mu(t_{k-1})} \\ &= \frac{(-\sum_{i=2}^k \mu(t_{i-1}) \tilde{h}(t_i, u(t_i), u^\Delta(t_{i-1})))}{\mu(t_{k-1})} - \frac{(-\sum_{i=2}^{k-1} \mu(t_{i-1}) \tilde{h}(t_i, u(t_i), u^\Delta(t_{i-1})))}{\mu(t_{k-1})} \\ &= - \frac{\mu(t_{k-1}) \tilde{h}(t_k, u(t_k), u^\Delta(t_{k-1}))}{\mu(t_{k-1})} \\ &= -\tilde{h}(t_k, u(t_k), u^\Delta(t_{k-1})). \end{aligned}$$

Thus, u solves (9).

We need now only consider the boundary conditions (2) in order to complete Step 3 of this proof. To this end, we recall the construction of the time scale \mathbb{T} and notice the following based on what was just derived as the formula for u^Δ :

$$u^\Delta(0) = u^\Delta(t_1) = - \sum_{i=2}^1 \mu(t_{i-1}) \tilde{h}(t_i, u(t_i), u^\Delta(t_{i-1})) = 0.$$

We now turn our attention over to $t = 1$ and recall from the construction of \mathbb{T} that $t_\infty = 1$. Also note that standard convention when discussing time scales of this sort is $\sigma(t) = t$ if \mathbb{T} has a maximum t , or for our purposes $\mu(t) = 0$ if \mathbb{T} has a

maximum t . As such,

$$\begin{aligned} u(1) &= u(t_\infty) = \sum_{j=\infty}^{\infty} \mu(t_j) \sum_{i=2}^j \mu(t_{i-1}) \tilde{h}(t_i, u(t_i), u^\Delta(t_{i-1})) \\ &= \mu(t_\infty) \sum_{i=2}^{\infty} \mu(t_{i-1}) \tilde{h}(t_i, u(t_i), u^\Delta(t_{i-1})) \\ &= \mu(1) \sum_{i=2}^{\infty} \mu(t_{i-1}) \tilde{h}(t_i, u(t_i), u^\Delta(t_{i-1})) \\ &= 0 \cdot \sum_{i=2}^{\infty} \mu(t_{i-1}) \tilde{h}(t_i, u(t_i), u^\Delta(t_{i-1})) = 0. \end{aligned}$$

Therefore, we get that u solves (9), (2) and Step 3 is complete.

Step 4: The remaining piece we need to show is that solutions of (9), (2) satisfy

$$\alpha(t) \leq u(t) \leq \beta(t), \quad t \in \mathbb{T}.$$

To this end, without loss of generality consider the case of obtaining $u(t) \leq \beta(t)$, and let $v(t) = u(t) - \beta(t)$. For the purpose of establishing a contradiction, assume that $\max\{v(t) : t \in \mathbb{T}\} := v(l) > 0$. From (2) and (7), we see that l must be an interior point in \mathbb{T} ; i.e., $l := t_j \in \mathbb{T} \setminus \{0, 1\}$. With t_j necessarily being an interior point, t_{j-1} and t_{j+1} are well-defined, and we have

$$v(t_{j-1}) \leq v(t_j) \quad \text{and} \quad v(t_{j+1}) \leq v(t_j).$$

Consequently,

$$v^\Delta(t_{j-1}) \geq 0 \quad \text{and} \quad v^\Delta(t_j) \leq 0.$$

Further, we now know also that

$$v^{\Delta\Delta}(t_{j-1}) = \frac{v^\Delta(t_j) - v^\Delta(t_{j-1})}{\mu(t_{j-1})} \leq 0.$$

Therefore,

$$u^{\Delta\Delta}(t_{j-1}) - \beta^{\Delta\Delta}(t_{j-1}) \leq 0. \tag{13}$$

On the other hand, since h is nonincreasing in its third variable, we have from (9) and (8) that

$$\begin{aligned} u^{\Delta\Delta}(t_{j-1}) - \beta^{\Delta\Delta}(t_{j-1}) &= -\tilde{h}(t_j, u(t_j), u^\Delta(t_{j-1})) - \beta^{\Delta\Delta}(t_{j-1}) \\ &= -\left(\tilde{h}(t_j, \beta(t_j), \beta^\Delta(t_{j-1})) - \frac{u(t_j) - \beta(t_j)}{u(t_j) - \beta(t_j) + 1} \right) - \beta^{\Delta\Delta}(t_{j-1}) \\ &= -\tilde{h}(t_j, \beta(t_j), \beta^\Delta(t_{j-1})) + \frac{v(t_j)}{v(t_j) + 1} - \beta^{\Delta\Delta}(t_{j-1}) \\ &\geq \beta^{\Delta\Delta}(t_{j-1}) + \frac{v(l)}{v(l) + 1} - \beta^{\Delta\Delta}(t_{j-1}) = \frac{v(l)}{v(l) + 1} > 0. \end{aligned}$$

Hence we have a contradiction to (13) and we conclude that $\max\{v(t) : t \in \mathbb{T}\} \leq 0$. Thus, $v(t) \leq 0$ for all $t \in \mathbb{T}$, or rather

$$u(t) \leq \beta(t) \quad \text{for all } t \in \mathbb{T}.$$

A similar argument shows that $\alpha(t) \leq u(t)$ for all $t \in \mathbb{T}$.

Thus, our conclusion holds and the proof is complete. □

4. Main result

In this section, we make use of Theorem 3.3 to obtain positive solutions to the singular problem (1), (2). In particular, in applying Theorem 3.3, we deal with a sequence of regular perturbations of (1), (2). Ultimately, we obtain a desired solution by passing to the limit on a sequence of solutions for the perturbations.

Theorem 4.1. *Assume conditions (A), (B), and (C) hold, along with the following:*

(D) *There exists $c \in (0, \infty)$ so that $f(t, c, y) \leq 0$ for all $t \in \mathbb{T}$ and $y \in \mathbb{R}$.*

(E) *$f(t, x, y)$ is nonincreasing in its y -variable for all $t \in \mathbb{T}$ and $x \in (0, c)$.*

(F) *$\lim_{x \rightarrow 0^+} f(t, x, y) = \infty$ for $t \in \mathbb{T}$ and $y \in (-c, c)$.*

Then, (1), (2) has a solution u satisfying

$$0 < u(t) \leq c, \quad t \in \mathbb{T} \setminus \{1\}.$$

Proof. We proceed through this proof via a sequence of steps.

Step 1: For $k > 0$, $t \in \mathbb{T}$, and $y \in \mathbb{R}$, define

$$f_k(t, x, y) = \begin{cases} f(t, |x|, y) & \text{if } |x| \geq 1/k, \\ f(t, 1/k, y) & \text{if } |x| < 1/k. \end{cases}$$

Then, f_k is continuous on $\mathbb{T} \times \mathbb{R}^2$.

Assumption (F) implies that there exists k_0 such that, for all $k \geq k_0$,

$$f_k(t, 0, y) = f\left(t, \frac{1}{k}, y\right) > 0 \quad \text{for all } t \in \mathbb{T}, y \in \mathbb{R}.$$

We now consider

$$u^{\Delta\Delta}(t_{i-1}) + f_k(t_i, u(t_i), u^\Delta(t_{i-1})) = 0, \quad t \in \mathbb{T}. \tag{14}$$

Now, let $\alpha(t) = 0$ and $\beta(t) = c$. Then, for each $k \geq k_0$, α and β are lower and upper solutions of (14), (2), respectively. Also, $\alpha(t) \leq \beta(t)$ for $t \in \mathbb{T}$. Thus, by Theorem 3.3, for each $k \geq k_0$, there exists a solution u_k to each problem (14), (2) that satisfies $0 \leq u_k(t) \leq c$ for $t \in \mathbb{T}$.

Consequently, for all $t_i \in \mathbb{T}$,

$$|u^\Delta(t_i)| \leq c \cdot \mu(t_{i-1}). \tag{15}$$

Step 2: For $k \geq k_0$, let $\delta \in (0, 1)$ and consider the time scale $\mathbb{T}_1 := \mathbb{T} \cap [0, \delta]$. Since u_k solves (14), we get from work similar to that in the proof of [Theorem 3.3](#) that

$$u_k^\Delta(t_j) = - \sum_{i=2}^j \mu(t_{i-1}) f_k(t_i, u(t_i), u^\Delta(t_{i-1})). \quad (16)$$

We use this version of u_k^Δ as follows:

By assumption (F), there exists $\varepsilon_1 \in (0, 1/k_0)$ such that for all $k \geq 1/\varepsilon_1$

$$f_k(t_2, x, y) > c, \quad x \in (0, \varepsilon_1), \quad y \in (-c, c). \quad (17)$$

For the sake of establishing a contradiction, assume that for $k \geq 1/\varepsilon_1$ we have $u_k(t_2) < \varepsilon_1$. Then, by (16) and (17),

$$\begin{aligned} u_k^\Delta(t_2) &= - \sum_{i=2}^2 \mu(t_{i-1}) f_k(t_i, u(t_i), u^\Delta(t_{i-1})) \\ &= -\mu(t_1) f_k(t_2, u(t_2), u^\Delta(t_1)) < -\mu(t_1) \cdot c. \end{aligned}$$

However, this contradicts (15). Thus, $u_k(t_2) \geq \varepsilon_1$ for all $k \geq 1/\varepsilon_1$.

Continuing, also by assumption (F), there now exists $\varepsilon_2 \in (0, \varepsilon_1)$ such that for all $k \geq 1/\varepsilon_2$

$$f_k(t_3, x, y) > c + m_1, \quad x \in (0, \varepsilon_2), \quad y \in (-c, c), \quad (18)$$

where $m_1 = \max\{|f_k(t_2, x, y)| : x \in [\varepsilon_1, c], y \in (-c, c)\}$. For the sake of establishing a contradiction, assume that for $k \geq 1/\varepsilon_2$ we have $u_k(t_3) < \varepsilon_2$. Then, by (16) and (18),

$$\begin{aligned} u_k^\Delta(t_3) &= - \sum_{i=2}^3 \mu(t_{i-1}) f_k(t_i, u(t_i), u^\Delta(t_{i-1})) \\ &= -\mu(t_1) f_k(t_2, u(t_2), u^\Delta(t_1)) - \mu(t_2) f_k(t_3, u(t_3), u^\Delta(t_2)) \\ &\leq \mu(t_1) \cdot m_1 - \mu(t_2)(c + m_1) < \mu(t_2) \cdot c. \end{aligned}$$

However, this contradicts (15). Thus, $u_k(t_3) \geq \varepsilon_2$ for all $k \geq 1/\varepsilon_2$.

We continue in this manner, proceeding across the interval \mathbb{T}_1 for $j = 3, 4, 5, \dots, l-1$ and we create a nested sequence of epsilons, $0 < \varepsilon_{l-1} < \dots < \varepsilon_2 < \varepsilon_1$, where $u_k(t_j) \geq \varepsilon_{j-1}$ when $k \geq 1/\varepsilon_{j-1}$.

Continuing, by assumption (F), there exists $\varepsilon_l \in (0, \varepsilon_{l-1})$ such that for all $k \geq 1/\varepsilon_l$

$$f_k(t_{l+1}, x, y) > c + \sum_{i=1}^{l-1} m_i, \quad x \in [\varepsilon_l, c], \quad y \in (-c, c), \quad (19)$$

where $m_i = \max\{|f_k(t_{i+1}, x, y)| : x \in [\varepsilon_i, c], y \in (-c, c)\}$. For the sake of establishing a contradiction, assume that for $k \geq 1/\varepsilon_l$ we have $u_k(t_{l+1}) < \varepsilon_l$. Then, by

(16) and (19),

$$\begin{aligned}
 u_k^\Delta(t_{l+1}) &= - \sum_{i=2}^{l+1} \mu(t_{i-1}) f_k(t_i, u(t_i), u^\Delta(t_{i-1})) \\
 &= -\mu(t_1) f_k(t_2, u(t_2), u^\Delta(t_1)) - \mu(t_2) f_k(t_3, u(t_3), u^\Delta(t_2)) \\
 &\quad - \cdots - \mu(t_l) f_k(t_{l+1}, u(t_{l+1}), u^\Delta(t_l)) \\
 &\leq \mu(t_1) m_1 - \cdots - \mu(t_{l-1}) m_{l-1} - \mu(t_l) \left(c + \sum_{i=1}^{l-1} m_i \right) \\
 &< \mu(t_l) \cdot c.
 \end{aligned}$$

However, this contradicts (15). Thus, $u_k(t_{l+1}) \geq \varepsilon_l$ for all $k \geq 1/\varepsilon_l$.

Now, recall that we are on the interval \mathbb{T}_1 , which just so happens to be an interval with a finite number of points included in its time scale. Call the largest of these points t_M , and based on the previous argument, there exists $\varepsilon_{m-1} > 0$ so that $u_k(t_M) \geq \varepsilon_{m-1}$ for $k \geq 1/\varepsilon_{m-1}$. Choose $\varepsilon = \varepsilon_{m-1}/2$ and note that

$$0 < \varepsilon \leq u_k(t) \leq c \quad \text{for all } t \in \mathbb{T}_1, \quad k \geq \frac{1}{\varepsilon}.$$

We now need only discuss what happens when

$$t \in \mathbb{T}_2 := \mathbb{T} \setminus \mathbb{T}_1 = \mathbb{T} \cap [\delta, 1].$$

Note that for each δ as $\delta \rightarrow 1$, via previous arguments, we have for sufficiently large k that $u_k(t) > 0$, $t \in \mathbb{T}_2$. Also note that for sufficiently large k , as $\delta \rightarrow 1$, we have $u_k(t) \geq 0$. This leads to the fact that for sufficiently large k , we get $u_k(t) > 0$ for $t \in \mathbb{T} \setminus \{1\}$ and, as our boundary condition states, $u_k(1) = 0$.

We now choose a subsequence $\{u_{k_n}(t)\} \subseteq \{u_k(t)\}$ so that

$$\lim_{n \rightarrow \infty} u_{k_n}(t) = u(t), \quad t \in \mathbb{T},$$

and note that $u(t) \in E$, where E is defined as in the proof of [Theorem 3.3](#).

Moreover, (16) yields, for sufficiently large n ,

$$u_{k_n}^\Delta(t_j) = - \sum_{i=2}^j \mu(t_{i-1}) f(t_i, u_{k_n}(t_i), u_{k_n}^\Delta(t_{i-1})),$$

and so letting $n \rightarrow \infty$ and from the continuity of f we get

$$u^\Delta(t_j) = - \sum_{i=2}^j \mu(t_{i-1}) f(t_i, u(t_i), u^\Delta(t_{i-1})).$$

Consequently,

$$u^{\Delta\Delta}(t_{i-1}) = -f(t_i, u(t_i), u^\Delta(t_{i-1})). \quad \square$$

5. Example

Let \mathbb{T} be as given in [Definition 2.1](#). Let $\alpha \in [0, \infty)$, $c, \beta \in (0, \infty)$, and $a : \mathbb{T} \rightarrow \mathbb{R}$. Then, by [Theorem 4.1](#), the problem

$$u^{\Delta\Delta}(t_{i-1}) + (a(t_i) + (u(t_i))^\alpha + (u(t_i))^{-\beta})(c - u(t_i)) - (u^\Delta(t_{i-1}))^3 = 0, \quad t_{i-1} \in \mathbb{T},$$

along with the boundary conditions [\(2\)](#), has a solution u satisfying the desired inequality.

References

- [Agarwal and O'Regan 1999] R. P. Agarwal and D. O'Regan, "Singular discrete boundary value problems", *Appl. Math. Lett.* **12**:4 (1999), 127–131. [MR](#) [Zbl](#)
- [Bao et al. 2012] G. Bao, X. Xu, and Y. Song, "Positive solutions for three-point boundary value problems with a non-well-ordered upper and lower solution condition", *Appl. Math. Lett.* **25**:4 (2012), 767–770. [MR](#) [Zbl](#)
- [Bohner and Peterson 2001] M. Bohner and A. Peterson, *Dynamic equations on time scales*, Birkhäuser, Boston, 2001. [MR](#) [Zbl](#)
- [Henderson and Kunkel 2006] J. Henderson and C. J. Kunkel, "Singular discrete higher order boundary value problems", *Int. J. Difference Equ.* **1**:1 (2006), 119–133. [MR](#) [Zbl](#)
- [Kelley and Peterson 1991] W. G. Kelley and A. C. Peterson, *Difference equations: an introduction with applications*, Academic Press, Boston, 1991. [MR](#) [Zbl](#)
- [Kunkel 2006] C. J. Kunkel, "Singular second order boundary value problems for differential equations", pp. 119–124 in *Proceedings of neural, parallel, and scientific computations, III* (Atlanta, 2006), edited by G. S. Ladde et al., Dynamic, Atlanta, 2006. [MR](#) [Zbl](#)
- [Kunkel 2008] C. J. Kunkel, "Singular second order boundary value problems on purely discrete time scales", *J. Difference Equ. Appl.* **14**:4 (2008), 411–420. [MR](#) [Zbl](#)
- [Precup 2016] R. Precup, "Abstract method of upper and lower solutions and application to singular boundary value problems", *Stud. Univ. Babeş-Bolyai Math.* **61**:4 (2016), 443–451. [MR](#) [Zbl](#)
- [Rachůnková and Rachůnek 2006] I. Rachůnková and L. Rachůnek, "Singular discrete second order BVPs with p -Laplacian", *J. Difference Equ. Appl.* **12**:8 (2006), 811–819. [MR](#) [Zbl](#)
- [Rachůnková and Rachůnek 2009] I. Rachůnková and L. Rachůnek, "Singular discrete and continuous mixed boundary value problems", *Math. Comput. Modelling* **49**:3-4 (2009), 413–422. [MR](#) [Zbl](#)
- [Zeidler 1986] E. Zeidler, *Nonlinear functional analysis and its applications, I: Fixed-point theorems*, Springer, 1986. [MR](#) [Zbl](#)

Received: 2019-02-11

Revised: 2019-03-25

Accepted: 2019-03-30

ckunkel@utm.edu

*Department of Mathematics and Statistics,
University of Tennessee at Martin, Martin, TN, United States*

alej lanc@ut.utm.edu

*Department of Mathematics and Statistics,
University of Tennessee at Martin, Martin, TN, United States*

INVOLVE YOUR STUDENTS IN RESEARCH

Involve showcases and encourages high-quality mathematical research involving students from all academic levels. The editorial board consists of mathematical scientists committed to nurturing student participation in research. Bridging the gap between the extremes of purely undergraduate research journals and mainstream research journals, *Involve* provides a venue to mathematicians wishing to encourage the creative involvement of students.

MANAGING EDITOR

Kenneth S. Berenhaut Wake Forest University, USA

BOARD OF EDITORS

Colin Adams	Williams College, USA	Chi-Kwong Li	College of William and Mary, USA
Arthur T. Benjamin	Harvey Mudd College, USA	Robert B. Lund	Clemson University, USA
Martin Bohner	Missouri U of Science and Technology, USA	Gaven J. Martin	Massey University, New Zealand
Nigel Boston	University of Wisconsin, USA	Mary Meyer	Colorado State University, USA
Amarjit S. Budhiraja	U of N Carolina, Chapel Hill, USA	Frank Morgan	Williams College, USA
Pietro Cerone	La Trobe University, Australia	Mohammad Sal Moslehian	Ferdowsi University of Mashhad, Iran
Scott Chapman	Sam Houston State University, USA	Zuhair Nashed	University of Central Florida, USA
Joshua N. Cooper	University of South Carolina, USA	Ken Ono	Univ. of Virginia, Charlottesville
Jem N. Corcoran	University of Colorado, USA	Yuval Peres	Microsoft Research, USA
Toka Diagana	Howard University, USA	Y.-F. S. Pétermann	Université de Genève, Switzerland
Michael Dorff	Brigham Young University, USA	Jonathon Peterson	Purdue University, USA
Sever S. Dragomir	Victoria University, Australia	Robert J. Plemmons	Wake Forest University, USA
Joel Foisy	SUNY Potsdam, USA	Carl B. Pomerance	Dartmouth College, USA
Errin W. Fulp	Wake Forest University, USA	Vadim Ponomarenko	San Diego State University, USA
Joseph Gallian	University of Minnesota Duluth, USA	Bjorn Poonen	UC Berkeley, USA
Stephan R. Garcia	Pomona College, USA	József H. Przytycki	George Washington University, USA
Anant Godbole	East Tennessee State University, USA	Richard Rebarber	University of Nebraska, USA
Ron Gould	Emory University, USA	Robert W. Robinson	University of Georgia, USA
Sat Gupta	U of North Carolina, Greensboro, USA	Javier Rojo	Oregon State University, USA
Jim Haglund	University of Pennsylvania, USA	Filip Saidak	U of North Carolina, Greensboro, USA
Johnny Henderson	Baylor University, USA	Hari Mohan Srivastava	University of Victoria, Canada
Glenn H. Hurlbert	Virginia Commonwealth University, USA	Andrew J. Sterge	Honorary Editor
Charles R. Johnson	College of William and Mary, USA	Ann Trenk	Wellesley College, USA
K. B. Kulasekera	Clemson University, USA	Ravi Vakil	Stanford University, USA
Gerry Ladas	University of Rhode Island, USA	Antonia Vecchio	Consiglio Nazionale delle Ricerche, Italy
David Larson	Texas A&M University, USA	John C. Wierman	Johns Hopkins University, USA
Suzanne Lenhart	University of Tennessee, USA	Michael E. Zieve	University of Michigan, USA

PRODUCTION

Silvio Levy, Scientific Editor


Cover: Alex Scorpan

See inside back cover or msp.org/involve for submission instructions. The subscription price for 2019 is US \$195/year for the electronic version, and \$260/year (+\$35, if shipping outside the US) for print and electronic. Subscriptions, requests for back issues and changes of subscriber address should be sent to MSP.

Involve (ISSN 1944-4184 electronic, 1944-4176 printed) at Mathematical Sciences Publishers, 798 Evans Hall #3840, c/o University of California, Berkeley, CA 94720-3840, is published continuously online. Periodical rate postage paid at Berkeley, CA 94704, and additional mailing offices.

Involve peer review and production are managed by EditFlow[®] from Mathematical Sciences Publishers.

PUBLISHED BY

 **mathematical sciences publishers**

nonprofit scientific publishing

<http://msp.org/>

© 2019 Mathematical Sciences Publishers

involve

2019

vol. 12

no. 6

Occurrence graphs of patterns in permutations	901
BJARNI JENS KRISTINSSON AND HENNING ULFARSSON	
Truncated path algebras and Betti numbers of polynomial growth	919
RYAN COOPERGARD AND MARJU PURIN	
Orbit spaces of linear circle actions	941
SUZANNE CRAIG, NAICHE DOWNEY, LUCAS GOAD, MICHAEL J. MAHONEY AND JORDAN WATTS	
On a theorem of Besicovitch and a problem in ergodic theory	961
ETHAN GWALTNEY, PAUL HAGELSTEIN, DANIEL HERDEN AND BRIAN KING	
Algorithms for classifying points in a 2-adic Mandelbrot set	969
BRANDON BATE, KYLE CRAFT AND JONATHON YULY	
Sidon sets and 2-caps in \mathbb{F}_3^n	995
YIXUAN HUANG, MICHAEL TAIT AND ROBERT WON	
Covering numbers of upper triangular matrix rings over finite fields	1005
MERRICK CAI AND NICHOLAS J. WERNER	
Nonstandard existence proofs for reaction diffusion equations	1015
CONNOR OLSON, MARSHALL MUELLER AND SIGURD B. ANGENENT	
Improving multilabel classification via heterogeneous ensemble methods	1035
YUJUE WU AND QING WANG	
The number of fixed points of AND-OR networks with chain topology	1051
ALAN VELIZ-CUBA AND LAUREN GEISER	
Positive solutions to singular second-order boundary value problems for dynamic equations	1069
CURTIS KUNKEL AND ALEX LANCASTER	