

involve

a journal of mathematics

A systematic development of Jeans' criterion
with rotation for gravitational instabilities

Kohl Gill, David J. Wollkind and Bonni J. Dichone



A systematic development of Jeans' criterion with rotation for gravitational instabilities

Kohl Gill, David J. Wollkind and Bonni J. Dichone

(Communicated by Martin J. Bohner)

An inviscid fluid model of a self-gravitating infinite expanse of a uniformly rotating adiabatic gas cloud consisting of the continuity, Euler's, and Poisson's equations for that situation is considered. There exists a static homogeneous density solution to this model relating that equilibrium density to the uniform rotation. A systematic linear stability analysis of this exact solution then yields a gravitational instability criterion equivalent to that developed by Sir James Jeans in the absence of rotation instead of the slightly more complicated stability behavior deduced by Subrahmanyan Chandrasekhar for this model with rotation, both of which suffered from the same deficiency in that neither of them actually examined whether their perturbation analysis was of an exact solution. For the former case, it was not and, for the latter, the equilibrium density and uniform rotation were erroneously assumed to be independent instead of related to each other. Then this gravitational instability criterion is employed in the form of Jeans' length to show that there is very good agreement between this theoretical prediction and the actual mean distance of separation of stars formed in the outer arms of the spiral galaxy Andromeda M31. Further, the uniform rotation determined from the exact solution relation to equilibrium density and the corresponding rotational velocity for a reference radial distance are consistent with the spectroscopic measurements of Andromeda and the observational data of the spiral Milky Way galaxy.

1. Introduction and formulation of the problem

Consider the governing equations for a self-gravitational adiabatic inviscid fluid of infinite extent undergoing uniform rotation [Chandrasekhar 1961]:

$$\text{continuity equation: } \frac{D\rho}{Dt} + \rho \nabla \cdot \mathbf{v} = 0, \quad (1-1a)$$

MSC2010: 35B36, 35Q85, 76E07, 76E99.

Keywords: Andromeda and Milky Way star formation, Jeans' self-gravitational instabilities, rotating adiabatic inviscid gas dynamics, astrophysics.

$$\text{Euler's equation: } \frac{D\mathbf{v}}{Dt} + 2\boldsymbol{\Omega} \times \mathbf{v} + \boldsymbol{\Omega} \times (\boldsymbol{\Omega} \times \mathbf{r}) = -\frac{1}{\rho} \mathcal{P}'(\rho) \nabla \rho + \mathbf{g}, \quad (1-1b)$$

$$\text{Poisson's equation: } \nabla \cdot \mathbf{g} = -4\pi G_0 \rho. \quad (1-1c)$$

Here $t \equiv$ time, $\mathbf{r} = (x, y, z) \equiv$ position vector, $\boldsymbol{\Omega} = (0, 0, \Omega_0) \equiv$ uniform rotation vector, $\rho \equiv$ density (mass/[unit volume]), $\mathbf{v} = (u, v, w) \equiv$ velocity vector with respect to the rotating frame, $\nabla = (\partial/\partial x, \partial/\partial y, \partial/\partial z) \equiv$ gradient operator, $D/Dt = \partial/\partial t + \mathbf{v} \cdot \nabla \equiv$ material derivative, $\mathcal{P}(\rho) = p_0(\rho/\rho_0)^{\gamma_0} \equiv$ adiabatic pressure, $\mathbf{g} = -\nabla \varphi \equiv$ gravitational acceleration vector with $\varphi \equiv$ self-gravitating potential, and $G_0 \equiv$ universal gravitational constant. The continuity and Euler's equations follow from the conservation of mass and momentum for an inviscid fluid [Lin and Segel 1974] with the addition of the extra second and third terms on the left-hand side of (1-1b), which represent the Coriolis effect and centrifugal force, respectively, due to the rotation [Greenspan 1968]. Poisson's equation follows from the divergence theorem and Newton's law of universal gravitation [Binney and Tremaine 1987; Lin and Segel 1974]. Since

$$\begin{aligned} \boldsymbol{\Omega} \times (\boldsymbol{\Omega} \times \mathbf{r}) &= (\boldsymbol{\Omega} \cdot \mathbf{r}) \boldsymbol{\Omega} - (\boldsymbol{\Omega} \cdot \boldsymbol{\Omega}) \mathbf{r} = -\Omega_0^2(x, y, 0), \\ \boldsymbol{\Omega} \times \mathbf{v} &= \Omega_0(-v, u, 0), \quad \nabla \cdot \mathbf{g} = -\nabla^2 \varphi \end{aligned} \quad (1-1d)$$

[Segel 1977], the Euler's and Poisson's equations become

$$\frac{D\mathbf{v}}{Dt} + 2\Omega_0(-v, u, 0) - \Omega_0^2(x, y, 0) = -\frac{1}{\rho} \mathcal{P}'(\rho) \nabla \rho - \nabla \varphi, \quad (1-1e)$$

$$\nabla^2 \varphi = 4\pi G_0 \rho, \quad \text{where } \nabla^2 \equiv \nabla \cdot \nabla. \quad (1-1f)$$

Sir James Jeans [1902; 1928] proposed that a gravitational instability mechanism occurring in the spiral arms of protogalactic nebulae could result in the formation of chains of condensations, which eventually developed into those stars visible in the outer regions of fully evolved galaxies. He suggested that a nonrotating self-gravitating unbounded interstellar cloud of adiabatic gas, which is initially uniform in density and quiescent, should undergo an instability mechanism of this sort when acted on by random infinitesimal perturbations. Jeans [1902; 1928] deduced a criterion for which such an interstellar cloud would exhibit a gravitational instability by performing a linear stability analysis on what he assumed to be an exact solution to his governing inviscid gas dynamical model system that was equivalent to equations (1-1) in the absence of rotation, arriving at the following secular equation satisfied by σ and λ , the growth rate and wavelength, respectively, of his small density fluctuations:

$$\sigma^2 = 4\pi \left(G_0 \rho_0 - \pi \frac{c_0^2}{\lambda^2} \right), \quad (1-2a)$$

where c_0 is the speed of sound in an adiabatic medium of uniform density ρ_0 . This relation differed from that for the propagation of sound in a homogeneous medium

only due to the presence of the gravity term in (1-2a). Then, from (1-2a), Jeans concluded that there would be instability corresponding to $\sigma^2 > 0$ provided

$$\lambda > \lambda_J = c_0 \sqrt{\frac{\pi}{G_0 \rho_0}} \equiv \text{Jeans' length}, \quad (1-2b)$$

which is known as Jeans' criterion for gravitational instabilities.

The only problem with this derivation is that Jeans represented his exact static solution to those governing equations symbolically as $\mathbf{v} \equiv \mathbf{0} = (0, 0, 0)$, $\rho = \rho_0$, $\varphi = \varphi_0$. Since this analysis was for a nonrotating system with $\Omega_0 = 0$, when he assumed in addition that ρ_0 was uniform to make his perturbation equations constant coefficient this implicitly required $\nabla \varphi_0 = \mathbf{0}$, which implied $\nabla^2 \varphi_0 = 0 = 4\pi G_0 \rho_0$ or $\rho_0 = 0$ and hence is termed Jeans' swindle by Binney and Tremaine [1987]. Kiessling [2003] refutes their claim that Jeans' derivation represents a swindle because it can be justified by taking the proper limit of the appropriate cosmological model.

Since spectroscopic evidence (reviewed by Rubin and Ford [1970]) ultimately showed these nebulae to be rotating, Subrahmanyan Chandrasekhar [1961] considered the effect of adding rotation to Jeans' governing system of perturbation equations and repeated that analysis, demonstrating in the process that its stability behavior was slightly more complicated in that it involved an extra instability condition as well as Jeans' criterion. Chandrasekhar's perturbation analysis suffered from the same deficiency as Jeans' in that he did not develop a parameter relationship for his implicit exact solution and thus treated ρ_0 and Ω_0 as independent. We shall demonstrate that the proper relationship between these parameters eliminates this extra condition and only yields Jeans' instability criterion. Many subsequent linear stability analyses of similar problems influenced by the methodology of these works have treated their associated perturbation systems independently of the actual exact solution of the governing equations and thus replicate this deficiency including recent studies and reviews of gravitational instabilities [Stahler and Palla 2004]. Hence, we believe there is some merit in performing a systematic linear stability analysis of the relevant exact solution for Chandrasekhar's problem and toward that end present an investigation of this sort in the next section.

2. The exact static homogeneous density solution and its linear stability

There exists an exact static homogeneous density solution of our basic equations of the form

$$\mathbf{v} \equiv \mathbf{0} = (0, 0, 0), \quad \rho \equiv \rho_0, \quad \varphi = \varphi_0, \quad (2-1a)$$

where φ_0 satisfies

$$\nabla \varphi_0 = \Omega_0^2(x, y, 0), \quad \nabla^2 \varphi_0 = 4\pi G_0 \rho_0 \quad (2-1b)$$

or

$$\varphi_0(x, y) = \frac{1}{2}\Omega_0^2(x^2 + y^2), \quad \text{with } \Omega_0^2 = 2\pi G_0\rho_0 > 0. \quad (2-1c)$$

Now seeking a linear perturbation solution of these basic equations of the form

$$\begin{aligned} \mathbf{v} &= \varepsilon \mathbf{v}_1 + \mathbf{O}(\varepsilon^2), \quad \text{where } \mathbf{v}_1 = (u_1, v_1, w_1), \\ \rho &= \rho_0[1 + \varepsilon s + \mathbf{O}(\varepsilon^2)], \quad \varphi = \varphi_0 + \varepsilon\varphi_1 + \mathbf{O}(\varepsilon^2), \end{aligned} \quad (2-2)$$

with $|\varepsilon| \ll 1$, substituting (2-2) into those equations, neglecting terms of $\mathbf{O}(\varepsilon^2)$, and canceling the resulting ε common factor; we deduce that the perturbation quantities to this exact solution satisfy

$$\frac{\partial s}{\partial t} + \frac{\partial u_1}{\partial x} + \frac{\partial v_1}{\partial y} + \frac{\partial w_1}{\partial z} = 0, \quad (2-3a)$$

$$\frac{\partial u_1}{\partial t} - 2\Omega_0 v_1 + c_0^2 \frac{\partial s}{\partial x} + \frac{\partial \varphi_1}{\partial x} = 0, \quad \text{where } c_0^2 = \mathcal{P}'(\rho_0) = \gamma_0 \frac{p_0}{\rho_0} > 0, \quad (2-3b)$$

$$\frac{\partial v_1}{\partial t} + 2\Omega_0 u_1 + c_0^2 \frac{\partial s}{\partial y} + \frac{\partial \varphi_1}{\partial y} = 0, \quad (2-3c)$$

$$\frac{\partial w_1}{\partial t} + c_0^2 \frac{\partial s}{\partial z} + \frac{\partial \varphi_1}{\partial z} = 0, \quad (2-3d)$$

$$2\Omega_0^2 s - \nabla^2 \varphi_1 = 0. \quad (2-3e)$$

Then assuming a normal mode solution for these perturbation quantities of the form

$$[u_1, v_1, w_1, s, \varphi_1](x, y, z, t) = [A, B, C, E, F]e^{i(k_1x + k_2y + k_3z) + \sigma t}, \quad (2-4)$$

where $|A|^2 + |B|^2 + |C|^2 + |E|^2 + |F|^2 \neq 0$, $i = \sqrt{-1}$, and $k_{1,2,3} \in \mathbb{R}$ satisfy the implicit far-field boundedness property for those quantities, and substituting (2-4) into (2-3), we obtain the following equations for $[A, B, C, E, F]$ upon cancellation of the exponential common factor:

$$ik_1A + ik_2B + ik_3C + \sigma E = 0, \quad (2-5a)$$

$$\sigma A - 2\Omega_0 B + ic_0^2 k_1 E + ik_1 F = 0, \quad (2-5b)$$

$$2\Omega_0 A + \sigma B + ic_0^2 k_2 E + ik_2 F = 0, \quad (2-5c)$$

$$\sigma C + ic_0^2 k_3 E + ik_3 F = 0, \quad (2-5d)$$

$$2\Omega_0^2 E + k^2 F = 0, \quad \text{where } k^2 = k_1^2 + k_2^2 + k_3^2. \quad (2-5e)$$

Setting the determinant of the 5×5 coefficient matrix for the linear homogeneous system (2-5) of constants equal to zero to satisfy their nontriviality property, we obtain

$$k^2[\sigma^4 + (c_0^2 k^2 + 2\Omega_0^2)\sigma^2] + 4\Omega_0^2(c_0^2 k^2 - 2\Omega_0^2)k_3^2 = 0. \quad (2-6)$$

Defining the wavenumber vector $\mathbf{k} = (k_1, k_2, k_3)$, its dot product with $\boldsymbol{\Omega}$ satisfies

$$\mathbf{k} \cdot \boldsymbol{\Omega} = k_3 \Omega_0 = |\mathbf{k}| |\boldsymbol{\Omega}| \cos(\theta) = k \Omega_0 \cos(\theta), \quad (2-7a)$$

θ being the azimuthal angle between \mathbf{k} and $\boldsymbol{\Omega}$, which implies

$$k_3 = k \cos(\theta). \quad (2-7b)$$

Then, substitution of (2-7b) into (2-6) and cancellation of k^2 yields the secular equation

$$\sigma^4 + (c_0^2 k^2 + 2\Omega_0^2) \sigma^2 + 4\Omega_0^2 (c_0^2 k^2 - 2\Omega_0^2) \cos^2(\theta) = 0. \quad (2-8)$$

Since this secular equation is a quadratic in σ^2 , we first demonstrate that $\sigma^2 \in \mathbb{R}$ by showing that its discriminant \mathcal{D} satisfies

$$\mathcal{D} = (c_0^2 k^2 + 2\Omega_0^2)^2 - 16\Omega_0^2 (c_0^2 k^2 - 2\Omega_0^2) \cos^2(\theta) \geq 0. \quad (2-9a)$$

Consider the two cases of $c_0^2 k^2 - 2\Omega_0^2 \leq 0$ and $c_0^2 k^2 - 2\Omega_0^2 > 0$ separately. For the former case it is obvious, while for the latter one it can be deduced by noting that

$$\mathcal{D} \geq (c_0^2 k^2 + 2\Omega_0^2)^2 - 16\Omega_0^2 (c_0^2 k^2 - 2\Omega_0^2) = (c_0^2 k^2 - 6\Omega_0^2)^2. \quad (2-9b)$$

For $\theta = \frac{\pi}{2}$, we can conclude from (2-8) that

$$\sigma^2 = 0 \quad \text{or} \quad \sigma^2 = -(c_0^2 k^2 + 2\Omega_0^2) < 0, \quad (2-10a)$$

while for $\theta \neq \frac{\pi}{2}$, the stability criteria governing such quadratics, namely,

$$\text{given } \omega^2 + a\omega + b = 0 \text{ with } \mathcal{D} = a^2 - 4b \geq 0, \quad \omega < 0 \iff a, b > 0 \quad (2-10b)$$

[Uspensky 1948], implies that

$$\sigma^2 < 0 \iff c_0^2 k^2 - 2\Omega_0^2 > 0. \quad (2-10c)$$

Making an interpretation of these results, we can deduce from (2-10) and (2-1c) that there will only be $\sigma^2 > 0$ and hence unstable behavior provided

$$c_0^2 k^2 - 4\pi G_0 \rho_0 < 0, \quad (2-11a)$$

which is equivalent to Jeans' gravitational instability criterion (1-2b)

$$\lambda > \lambda_J = c_0 \sqrt{\frac{\pi}{G_0 \rho_0}} \equiv \text{Jeans' length} \quad (2-11b)$$

since

$$\lambda = \frac{2\pi}{k}. \quad (2-11c)$$




Figure 1. A Galaxy Evolution Explorer image of the Andromeda galaxy M31, courtesy NASA/JPL-Caltech.

3. Comparisons

Let us return to Jeans' analysis. In writing (1-2), one must implicitly assume that $\rho_0 > 0$, which seems plausible in that $\rho_0 = 0$ corresponds to a completely empty space [Scheffler and Elsässer 1988]. When gravity is taken into account in the absence of rotation, however, such an assumption is not strictly compatible with the equations of hydrostatic equilibrium, as we have seen. Thus, Jeans' uniform density solution, as mentioned above, was not exact. The problem under examination demonstrates that adding rotation to the system as Chandrasekhar did and again performing a standard linear stability analysis of its exact static solution yields Jeans' instability criterion but in a systematic manner and such a model also has the added advantage of being more astrophysically realistic. Jeans got the right answer for the wrong reason, as was shown in [Kiesling 2003] by taking the proper limit of the appropriate cosmological model to fix that analysis. In his review of hydrodynamic stability theory, the renowned comprehensive applied mathematical modeler Lee Segel [1966] stated that "Anyone can get the right answer for the right reason. It takes a genius or a physicist to get the right answer for the wrong reason." In this context, Sir James Jeans was both.

The formula for λ_J in (2-11b) is of fundamental importance in astrophysics and cosmology where many significant deductions concerning the formation of galaxies and stars have been based upon it. In particular, Jeans' interpretation of the criterion, now bearing his name, was that a gas cloud of characteristic dimension much greater than λ_J would tend to form condensations with mean distance of separation comparable to λ_J that then developed into those protostars observable in the outer arms of spiral galaxies such as Andromeda M31 (see Figure 1). Sekimura




Figure 2. Schematic plots in the k - σ plane depicting the methodology employed by [Sekimura et al. 1999] applied to the Jeans' secular equation $\sigma = \sigma(k; c) = \sqrt{2\Omega_0^2 - c^2k^2}$. That curve is plotted for both a general speed of sound c and our specific speed $c_0 > c$ in this figure where $k_J = 2\pi/\lambda_J$ is such that $\sigma(k_J; c_0) = 0$. In a weakly nonlinear stability analysis one takes the disturbance wavenumber $k \equiv k_J$ and its growth rate to be equal to $\sigma_J(c) = \sigma(k_J; c) = \delta^2 > 0$ where c is close enough to c_0 so that δ is a small parameter. Then in the $\lim_{c \rightarrow c_0} \sigma_J(c) = 0$ which is a requirement for the application of weakly nonlinear stability theory and any re-equilibrated pattern will exhibit a wavelength of λ_J . Here $c^2 = \gamma(p_0/\rho_0)$ with $\gamma < \gamma_0$ and hence the operation $\lim_{c \rightarrow c_0}$ is equivalent to $\lim_{\gamma \rightarrow \gamma_0}$.

et al. [1999] have demonstrated that, for a secular equation similar in form to (1-2a), λ_J actually corresponds to the so-called critical wavelength λ_c of linear stability theory associated with $\sigma = 0$ (see Figure 2), while nonlinear stability analyses of physical phenomena involving related secular equations have shown that the observed wavelengths are determined to a close approximation by that λ_c rather than by the dominant wavelength λ_d at which σ achieves its maximum value from linear theory (see, e.g., [Tian and Wollkind 2003]). Hence, Jeans' interpretation, although unusual for linear stability theory (where it is often presumed that such a disturbance associated with the largest growth rate predominates), both anticipated and is consistent with these nonlinear results, since, by the time perturbations have grown enough for the effect of the maximum growth rate to be observed, the neglected nonlinearities may have rendered that linear analysis inaccurate [Segel and Stoeckly 1972]. In this context, note that for a typical value of $\theta \neq \frac{\pi}{2}$, namely $\theta = 0$, we can factor our secular equation (2-8) to obtain the roots $\sigma^2 = -4\Omega_0^2$ and

$\sigma^2 = 2\Omega_0^2 - c_0^2 k^2$. Observe that this last condition, which yields our instability, is equivalent to the Jeans' secular equation (1-2a).

Thus using the formula for Jeans' length λ_J with the parameters c_0 and ρ_0 assigned the values

$$c_0 = \frac{2}{3} \times 10^4 \frac{\text{cm}}{\text{sec}} \quad \text{and} \quad \rho_0 = 10^{-22} \frac{\text{gm}}{\text{cm}^3} \quad (3-1a)$$

employed by [Jeans 1928] for this purpose but when the polytropic index γ_0 is $\frac{4}{3}$ [Bonnor 1957], while taking

$$G_0 = 6.67 \times 10^{-8} \frac{\text{cm}^3}{\text{gmsec}^2} \quad (3-1b)$$

in cgs units yields

$$\lambda_J = 4.58 \times 10^{18} \text{ cm} = 1.48 \text{ pc}, \quad (3-1c)$$

where $1 \text{ pc} \equiv 3.09 \times 10^{18} \text{ cm}$, which compares quite favorably with the mean distance between actual adjacent condensations originally formed in the outer arms of Andromeda since, in those parts of M31, the averaged observed distance between protostars in such chains is about 1.4 pc or somewhat more if allowances are made for foreshortening [Jeans 1928].

Given the small size of ρ_0 in (3-1a), Chandrasekhar [1961] was one of those individuals who regarded Jeans' analysis as a close approximation to reality [Schefler and Elsässer 1988]. Although he oriented his axes so that $\boldsymbol{\Omega} = (0, \Omega_y, \Omega_z)$ with $|\boldsymbol{\Omega}| = \Omega_0$ and $\mathbf{k} = (0, 0, k)$, using our more general orientation Chandrasekhar, in effect, considered uniform rotation Ω_0 in his perturbation equations through the Coriolis force terms of (2-3b) and (2-3c) in order to make the model more realistic while retaining the coefficient $4\pi\rho_0 G_0$ for s in (2-3e). In so doing, he implicitly assumed that Ω_0 and ρ_0 were independent rather than related parameters. Chandrasekhar plotted σ^2 versus k for $\theta = 0, \frac{\pi}{4}, \frac{\pi}{2}$ and $\Lambda^2 \equiv \Omega_0^2 / (\pi G_0 \rho_0) = 0.5, 1.0, 2.0$. Besides Jeans' criterion for $\theta \neq \frac{\pi}{2}$, this yielded an extra extraneous instability criterion for the case of $\theta = \frac{\pi}{2}$, namely,

$$c_0^2 k^2 < 4(\pi G_0 \rho_0 - \Omega_0^2) \quad \text{should} \quad \Omega_0^2 < \pi G_0 \rho_0. \quad (3-2)$$

In point of fact, $\Lambda^2 = 0.5$ is a representative value of that quantity for this instability condition of (3-2), while $\Lambda^2 = 2.0$, his upper bound, actually corresponds to its value as per our formula of (2-1c) relating these parameters, which implies

$$\Omega_0 = \sqrt{2\pi\rho_0 G_0}. \quad (3-3a)$$

Let us examine the plausibility of (3-3a), which violates (3-2) identically. In conjunction with the values for ρ_0 and G_0 of (3-1), (3-3a) yields the uniform

rotation

$$\Omega_0 = 6.47 \times 10^{-15} / \text{sec} \quad (3-3b)$$

and the corresponding rotational velocity

$$V_0 = r_0 \Omega_0 = 200 \frac{\text{km}}{\text{sec}} \quad (3-3c)$$

for the reference radial distance of

$$r_0 = 1 \text{ kpc} = 10^3 \text{ pc} = 3.09 \times 10^{21} \text{ cm} = 3.09 \times 10^{16} \text{ km}, \quad (3-3d)$$

both of which are consistent with the spectroscopic measurements of the Andromeda nebula and the observational data of the spiral Milky Way galaxy [Rubin and Ford 1970].

In conclusion our development presents a systematic linear stability analysis of Chandrasekhar's [1961] gravitational instability model in the presence of uniform rotation. We close by noting that Binney and Tremaine [1987] considered this gravitational instability model in a cylindrical rotating system as a problem in Chapter 5 of their book *Galactic Dynamics*. They observed that rotation allowed the Jeans' instability to be analyzed exactly. Since the first part of their problem was to find the condition on Ω_0 so that the homogeneous quiescent gas would be in equilibrium, Binney and Tremaine did not examine the plausibility of this condition. Further, the last part of their problem was to show, upon finding the resulting secular equation from its linear stability analysis, that waves propagating perpendicular to the rotation vector were always stable, while those propagating parallel to it were unstable if and only if the usual Jeans' criterion without rotation was satisfied. Although the latter conclusion for $\theta = 0$ agrees with our predictions, the former does not since, when $\theta = \frac{\pi}{2}$, we predicted $\sigma^2 = 0$, as well as those $\sigma^2 < 0$ which only implies a condition of neutral stability. Our results demonstrate that the best way to test the validity of a model for a natural science phenomenon is to compare its theoretical predictions with observable data of this phenomenon. Sir Arthur Conan Doyle characterized that philosophy probably as well as anyone by a Sherlock Holmes quote from "A scandal in Bohemia" in his 1891 collection entitled *The adventures of Sherlock Holmes*:

It is a capital mistake to theorize before one has data. Insensibly one begins to twist facts to suit theories, instead of theories to suit facts.

References

- [Binney and Tremaine 1987] J. Binney and S. Tremaine, *Galactic dynamics*, Princeton Univ. Press, 1987. [Zbl](#)
- [Bonnor 1957] W. B. Bonnor, "Jeans' formula for gravitational instability", *Monthly Not. Roy. Astr. Soc.* **117** (1957), 104–117. [MR](#) [Zbl](#)

- [Chandrasekhar 1961] S. Chandrasekhar, *Hydrodynamic and hydromagnetic stability*, Int. Series Monogr. Phys. **19**, Clarendon, Oxford, 1961. [MR](#) [Zbl](#)
- [Greenspan 1968] H. P. Greenspan, *The theory of rotating fluids*, Cambridge Monogr. Mech. Appl. Math. **12**, Cambridge Univ. Press, 1968. [Zbl](#)
- [Jeans 1902] J. H. Jeans, “The stability of a spherical nebula”, *Philos. Trans. R. Soc. Lond. Ser. A* **199** (1902), 1–53. [Zbl](#)
- [Jeans 1928] J. H. Jeans, *Astronomy and cosmogony*, Cambridge Univ. Press, 1928. [Zbl](#)
- [Kiessling 2003] M. K.-H. Kiessling, “The ‘Jeans swindle’: a true story — mathematically speaking”, *Adv. Appl. Math.* **31**:1 (2003), 132–149. [MR](#) [Zbl](#)
- [Lin and Segel 1974] C. C. Lin and L. A. Segel, *Mathematics applied to deterministic problems in the natural sciences*, Macmillan, New York, 1974. [MR](#) [Zbl](#)
- [Rubin and Ford 1970] V. C. Rubin and W. K. Ford, Jr., “Rotation of the Andromeda Nebula from a spectroscopic survey of emission”, *Astrophys. J.* **159** (1970), 379–403.
- [Scheffler and Elsässer 1988] H. Scheffler and H. Elsässer, *Physics of the galaxy and interstellar matter*, Springer, 1988.
- [Segel 1966] L. A. Segel, “Non-linear hydrodynamic stability theory and its applications to thermal convection and curved flows”, pp. 165–197 in *Non-equilibrium thermodynamics, variational techniques and stability* (Chicago, 1965), edited by R. J. Donnelly et al., Univ. Chicago Press, 1966. [MR](#)
- [Segel 1977] L. A. Segel, *Mathematics applied to continuum mechanics*, Macmillan, New York, 1977. [MR](#) [Zbl](#)
- [Segel and Stoeckly 1972] L. A. Segel and B. Stoeckly, “Instability of a layer of chemotactic cells, attractant and degrading enzyme”, *J. Theoret. Biol.* **37**:3 (1972), 561–585.
- [Sekimura et al. 1999] T. Sekimura, M. Zhu, J. Cook, P. K. Maini, and J. D. Murray, “Pattern formation of scale cells in Lepidoptera by differential origin-dependent cell adhesion”, *Bull. Math. Biol.* **61**:5 (1999), 807–828. [Zbl](#)
- [Stahler and Palla 2004] S. W. Stahler and F. Palla, *The formation of stars*, Wiley-VCH, Weinheim, Germany, 2004.
- [Tian and Wollkind 2003] E. M. Tian and D. J. Wollkind, “A nonlinear stability analysis of pattern formation in thin liquid films”, *Interfaces Free Bound.* **5**:1 (2003), 1–25. [MR](#) [Zbl](#)
- [Uspensky 1948] J. V. Uspensky, *Theory of equations*, McGraw-Hill, New York, 1948.

Received: 2018-01-03

Revised: 2019-05-22

Accepted: 2019-05-22

gillkohl@gmail.com*Department of Mathematics, Washington State University, Pullman, WA, United States*dwoollkind@wsu.edu*Department of Mathematics, Washington State University, Pullman, WA, United States*dichone@gonzaga.edu*Department of Mathematics, Gonzaga University, Spokane, WA, United States*

INVOLVE YOUR STUDENTS IN RESEARCH

Involve showcases and encourages high-quality mathematical research involving students from all academic levels. The editorial board consists of mathematical scientists committed to nurturing student participation in research. Bridging the gap between the extremes of purely undergraduate research journals and mainstream research journals, *Involve* provides a venue to mathematicians wishing to encourage the creative involvement of students.

MANAGING EDITOR

Kenneth S. Berenhaut Wake Forest University, USA

BOARD OF EDITORS

Colin Adams	Williams College, USA	Robert B. Lund	Clemson University, USA
Arthur T. Benjamin	Harvey Mudd College, USA	Gaven J. Martin	Massey University, New Zealand
Martin Bohner	Missouri U of Science and Technology, USA	Mary Meyer	Colorado State University, USA
Amarjit S. Budhiraja	U of N Carolina, Chapel Hill, USA	Frank Morgan	Williams College, USA
Pietro Cerone	La Trobe University, Australia	Mohammad Sal Moslehian	Ferdowsi University of Mashhad, Iran
Scott Chapman	Sam Houston State University, USA	Zuhair Nashed	University of Central Florida, USA
Joshua N. Cooper	University of South Carolina, USA	Ken Ono	Univ. of Virginia, Charlottesville
Jem N. Corcoran	University of Colorado, USA	Yuval Peres	Microsoft Research, USA
Toka Diagana	Howard University, USA	Y.-F. S. Pétermann	Université de Genève, Switzerland
Michael Dorff	Brigham Young University, USA	Jonathon Peterson	Purdue University, USA
Sever S. Dragomir	Victoria University, Australia	Robert J. Plemmons	Wake Forest University, USA
Joel Foisy	SUNY Potsdam, USA	Carl B. Pomerance	Dartmouth College, USA
Erin W. Fulp	Wake Forest University, USA	Vadim Ponomarenko	San Diego State University, USA
Joseph Gallian	University of Minnesota Duluth, USA	Bjorn Poonen	UC Berkeley, USA
Stephan R. Garcia	Pomona College, USA	József H. Przytycki	George Washington University, USA
Anant Godbole	East Tennessee State University, USA	Richard Rebarber	University of Nebraska, USA
Ron Gould	Emory University, USA	Robert W. Robinson	University of Georgia, USA
Sat Gupta	U of North Carolina, Greensboro, USA	Javier Rojo	Oregon State University, USA
Jim Haglund	University of Pennsylvania, USA	Filip Saidak	U of North Carolina, Greensboro, USA
Johnny Henderson	Baylor University, USA	Hari Mohan Srivastava	University of Victoria, Canada
Glenn H. Hurlbert	Virginia Commonwealth University, USA	Andrew J. Sterge	Honorary Editor
Charles R. Johnson	College of William and Mary, USA	Ann Trenk	Wellesley College, USA
K. B. Kulasekera	Clemson University, USA	Ravi Vakil	Stanford University, USA
Gerry Ladas	University of Rhode Island, USA	Antonia Vecchio	Consiglio Nazionale delle Ricerche, Italy
David Larson	Texas A&M University, USA	John C. Wierman	Johns Hopkins University, USA
Suzanne Lenhart	University of Tennessee, USA	Michael E. Zieve	University of Michigan, USA
Chi-Kwong Li	College of William and Mary, USA		

PRODUCTION

Silvio Levy, Scientific Editor

Cover: Alex Scorpan

See inside back cover or msp.org/involve for submission instructions. The subscription price for 2019 is US \$195/year for the electronic version, and \$260/year (+\$35, if shipping outside the US) for print and electronic. Subscriptions, requests for back issues and changes of subscriber address should be sent to MSP.

Involve (ISSN 1944-4184 electronic, 1944-4176 printed) at Mathematical Sciences Publishers, 798 Evans Hall #3840, c/o University of California, Berkeley, CA 94720-3840, is published continuously online. Periodical rate postage paid at Berkeley, CA 94704, and additional mailing offices.

Involve peer review and production are managed by EditFlow® from Mathematical Sciences Publishers.

PUBLISHED BY

 **mathematical sciences publishers**

nonprofit scientific publishing

<http://msp.org/>

© 2019 Mathematical Sciences Publishers

involve

2019

vol. 12

no. 7

Asymptotic expansion of Warlimont functions on Wright semigroups MARCO ALDI AND HANQIU TAN	1081
A systematic development of Jeans' criterion with rotation for gravitational instabilities KOHL GILL, DAVID J. WOLLKIND AND BONNI J. DICHONE	1099
The linking-unlinking game ADAM GIAMBRONE AND JAKE MURPHY	1109
On generalizing happy numbers to fractional-base number systems ENRIQUE TREVIÑO AND MIKITA ZHYLINSKI	1143
On the Hadwiger number of Kneser graphs and their random subgraphs ARRAN HAMM AND KRISTEN MELTON	1153
A binary unrelated-question RRT model accounting for untruthful responding AMBER YOUNG, SAT GUPTA AND RYAN PARKS	1163
Toward a Nordhaus–Gaddum inequality for the number of dominating sets LAUREN KEOUGH AND DAVID SHANE	1175
On some obstructions of flag vector pairs (f_1, f_{04}) of 5-polytopes HYE BIN CHO AND JIN HONG KIM	1183
Benford's law beyond independence: tracking Benford behavior in copula models REBECCA F. DURST AND STEVEN J. MILLER	1193
Closed geodesics on doubled polygons IAN M. ADELSTEIN AND ADAM Y. W. FONG	1219
Sign pattern matrices that allow inertia \mathcal{S}_n ADAM H. BERLINER, DEREK DEBLIECK AND DEEPAK SHAH	1229
Some combinatorics from Zeckendorf representations TYLER BALL, RACHEL CHAISER, DEAN DUSTIN, TOM EDGAR AND PAUL LAGARDE	1241