

involve

a journal of mathematics

On the zero-sum group-magicness
of cartesian products

Adam Fong, John Georges, David Mauro, Dylan Spagnuolo,
John Wallace, Shufan Wang and Kirsti Wash



On the zero-sum group-magicalness of cartesian products

Adam Fong, John Georges, David Mauro, Dylan Spagnuolo,
John Wallace, Shufan Wang and Kirsti Wash

(Communicated by Kenneth S. Berenhaut)

Let $G = (V(G), E(G))$ be a graph and let $\mathbb{A} = (A, +)$ be an abelian group with identity 0. Then an \mathbb{A} -magic labeling of G is a function ϕ from $E(G)$ into $A \setminus \{0\}$ such that for some $a \in A$, $\sum_{e \in E(v)} \phi(e) = a$ for every $v \in V(G)$, where $E(v)$ is the set of edges incident to v . If ϕ exists such that $a = 0$, then G is zero-sum \mathbb{A} -magic. Let G be the cartesian product of two or more graphs. We establish that G is zero-sum \mathbb{Z} -magic and we introduce a graph invariant $j^*(G)$ to explore the zero-sum integer-magic spectrum (or null space) of G . For certain G , we establish $\mathcal{A}(G)$, the set of nontrivial abelian groups for which G is zero-sum group-magic. Particular attention is given to $\mathcal{A}(G)$ for regular G , odd/even G , and G isomorphic to a product of paths.

1. Introduction

Let $G = (V(G), E(G))$ be a graph. Let \mathcal{A} be the set all nontrivial abelian groups and let $\mathbb{A} = (A, +) \in \mathcal{A}$, where 0 denotes the identity of \mathbb{A} . Then an \mathbb{A} -labeling of G is a function ϕ from $E(G)$ into $A \setminus \{0\}$. For fixed $e \in E(G)$, $\phi(e)$ is called the label of e under ϕ , and for fixed $v \in V(G)$, the sum of the labels of the edges incident to v is called the weight of v under ϕ . The graph G is \mathbb{A} -magic if and only if there exists an \mathbb{A} -labeling ϕ of G such that for some $a \in A$, the weight of every vertex in $V(G)$ under ϕ is a . In such a case, ϕ is called an \mathbb{A} -magic labeling of G . Additionally, G is zero-sum \mathbb{A} -magic if and only if there is an \mathbb{A} -labeling ϕ of G such that the weight of every vertex in $V(G)$ under ϕ is 0. In this case, ϕ is called a zero-sum \mathbb{A} -magic labeling of G . Letting $\mathcal{A}(G)$ denote the set of all $\mathbb{A} \in \mathcal{A}$ such that G is zero-sum \mathbb{A} -magic, we observe that if $\mathbb{H}_0 \in \mathcal{A}(G)$ and \mathbb{H}_0 is isomorphic to a subgroup of an abelian group \mathbb{H} , then $\mathbb{H} \in \mathcal{A}(G)$. Particularly, if $\mathbb{A} \in \mathcal{A}(G)$, then for a positive integer k , we have $\mathbb{A}^k \in \mathcal{A}(G)$. In Figure 1, we

MSC2010: 05C78.

Keywords: cartesian product of graphs, grid graph, magic labeling, group-magic labeling, zero-sum group-magic labeling, zero-sum integer-magic spectrum.




Figure 1. Zero-sum \mathbb{Z}_4 and \mathbb{Z}_4^2 -magic labelings of G .

illustrate a zero-sum \mathbb{Z}_4 -magic labeling and a zero-sum \mathbb{Z}_4^2 -magic labeling of a graph G . (The group \mathbb{Z}_4 is the cyclic group of integers under addition mod 4. In the specification of elements of \mathbb{Z}_4^2 , parentheses are omitted.) Note that an alternative zero-sum \mathbb{Z}_4^2 -magic labeling can be formed by changing each label $(a, 0)$ to (a, a) . Other obvious alternatives exist.

The *zero-sum integer-magic spectrum* of a graph G , denoted by $\text{zim}(G)$, is the set of positive integers k such that G is zero-sum \mathbb{Z}_k -magic, where \mathbb{Z}_1 is the group \mathbb{Z} of integers under addition and, for $k \geq 2$, \mathbb{Z}_k is the cyclic group of integers under addition mod k . Let \mathcal{N} denote the set of positive integers. By the fundamental theorem of finite abelian groups, $\mathcal{N} \setminus \{1\}$ is a subset of $\text{zim}(G)$ if and only if G is zero-sum \mathbb{A} -magic for all finite \mathbb{A} in \mathcal{A} . Moreover, for any infinite abelian group $\mathbb{A} = (A, +)$ with nonzero $a \in A$, a generates a subgroup of \mathbb{A} that is isomorphic to either \mathbb{Z} or \mathbb{Z}_k for some $k \geq 2$. Thus $\text{zim}(G) = \mathcal{N}$ if and only if $\mathcal{A}(G) = \mathcal{A}$. We note especially that if $\text{zim}(G) = \mathcal{N} \setminus \{2\}$, then $\mathcal{A} \setminus \{\mathbb{Z}_2^k \mid k \in \mathcal{N}\} \subseteq \mathcal{A}(G)$.

Sedláček [1964] first introduced magic labelings, motivated by magic squares in number theory. Stanley [1973] later showed that the study of magic labelings is related to the study of linear homogeneous diophantine equations. It was shown independently in [Low and Lee 2006] and [Shiu et al. 2004] that if G and H are \mathbb{A} -magic graphs, then the cartesian product of G and H is also \mathbb{A} -magic. More recently, Akbari et al. [2014] proved that every r -regular graph G with $r \geq 3$, $r \neq 5$ has $\text{zim}(G) = \mathcal{N}$ if r is even; otherwise $\text{zim}(G) \supseteq \mathcal{N} \setminus \{2, 4\}$. And, Shiu and Low [2018] have determined the zero-sum integer-magic spectrum of the cartesian product of two trees. For a dynamic survey of results on magic labelings, see [Gallian 2018].

In this paper, we consider the zero-sum \mathbb{A} -magicness of cartesian products. In Section 2, we give definitions and preliminary results. In Section 3, we develop our main results, with particular attention given to graph parity and regularity. And in Section 4, we consider the cartesian products of paths, also known as grid graphs.

2. Definitions and preliminary results

Throughout this paper, graphs will be finite, nontrivial, simple, loopless, and connected unless specified otherwise. An *even graph* (resp. *odd graph*) shall refer




Figure 2. The cartesian product $G_1 \square G_2$.

to a graph G of which each vertex v has even (resp. odd) degree, denoted by $d_G(v)$. Abelian groups shall have identity element 0 and binary operator $+$.

For $i \in \{1, \dots, n\}$, let $G_i = (V(G_i), E(G_i))$ be a graph. The *cartesian product* of G_1, G_2, \dots, G_n , denoted by $\square_{i=1}^n G_i$ or $G_1 \square G_2 \square \dots \square G_n$, is the graph G such that

- (1) the vertex set of G is $\prod_{i=1}^n V(G_i)$, and
- (2) vertices (u_1, u_2, \dots, u_n) and (w_1, w_2, \dots, w_n) of G are adjacent if and only if (u_1, u_2, \dots, u_n) and (w_1, w_2, \dots, w_n) differ in precisely one component i_0 , and u_{i_0} is adjacent to w_{i_0} in G_{i_0} .

The following are well known. Note that the results in (b), (c), and (d) extend to cartesian products of arbitrary finite length by part (a):

- (a) As a binary operator on the set of graphs, \square is associative and commutative with respect to isomorphism.
- (b) If (u, w) is a vertex of $G_1 \square G_2$, then $d_{G_1 \square G_2}((u, w)) = d_{G_1}(u) + d_{G_2}(w)$.
- (c) $G_1 \square G_2$ is regular if and only if each of G_1 and G_2 is regular.
- (d) If G_1 and G_2 are graphs with no isolated vertices, then $G_1 \square G_2$ is bridgeless.

In Figure 2, we demonstrate the cartesian product $G_1 \square G_2$ where G_1 is isomorphic to the claw on four vertices and G_2 is isomorphic to the cycle C_3 .

Let G be a graph and let h be a positive integer. Then a *factor* of G is a spanning subgraph of G , and an h -factor of G is an h -regular factor of G . (The term *factor* may also refer to a factor of a cartesian product. Its usage will clarify its intended meaning.)

For $n \geq 2$, let G be a cartesian product $\square_{i=1}^n G_i$. For fixed $i_0, 1 \leq i_0 \leq n$, suppose that H_{i_0} is an h_{i_0} -factor of G_{i_0} . Then there exists a natural h_{i_0} -factor of G , comprising the union of C disjoint subgraphs of G , each isomorphic to H_{i_0} , where C is the product of the orders of the graphs $G_i, 1 \leq i \leq n$, except for $i = i_0$. If, for




Figure 3. An illustration of regular factors.

each i , $E(G_i)$ is partitioned by the edge sets of regular factors $H_{i,1}, H_{i,2}, \dots, H_{i,m_i}$ of G_i , then these n partitions of $E(G_1), E(G_2), \dots$, and $E(G_n)$ induce a natural partition of $E(G)$ by the edge sets of regular factors $F_{i,j}$, $1 \leq i \leq n$, $1 \leq j \leq m_i$, where the regularity of $F_{i,j}$ equals the regularity of $H_{i,j}$. Note that each vertex of G is a vertex of each $F_{i,j}$ as illustrated in Figure 3. Therein, graph $G_1 \cong P_2$ has a 1-factor $H_{1,1} = G_1$ whose edge set trivially partitions $E(G_1)$. And, G_2 has a 1-factor $H_{2,1}$ isomorphic to four vertex-disjoint copies of P_2 and a 2-factor $H_{2,2}$ isomorphic to two vertex-disjoint copies of C_4 , where $E(H_{2,1})$ and $E(H_{2,2})$ together partition $E(G_2)$. These regular factors respectively induce $F_{1,1}, F_{2,1}$, and $F_{2,2}$, which together partition $E(G_1 \square G_2)$. (Note that G_2 is itself a cartesian product isomorphic to $P_2 \square C_4$.)

Let G be a graph that has a zero-sum \mathbb{Z} -magic labeling and let $\mathcal{Z}(G)$ represent the (nonempty) set of all zero-sum \mathbb{Z} -magic labelings of G . For each $\phi \in \mathcal{Z}(G)$, let $j(\phi)$ equal $\max_{e \in E(G)} |\phi(e)|$. Then $j^*(G)$ shall denote $\min_{\phi \in \mathcal{Z}(G)} j(\phi)$. To illustrate, we note that there exists a zero-sum \mathbb{Z} -magic labeling ϕ of K_4 such that $\max_{e \in E(K_4)} |\phi(e)| = 2$. Yet, since K_4 is an odd graph, there is no such labeling ϕ such that $\max_{e \in E(K_4)} |\phi(e)| = 1$. Thus $j^*(K_4) = 2$. On the other hand, some graphs, including C_{2n-1} and P_n for $n \geq 2$, are not zero-sum \mathbb{Z} -magic. For such graphs G , $j^*(G)$ does not exist.

Theorem 1. *If G is zero-sum \mathbb{Z} -magic, then G is zero-sum \mathbb{Z}_k -magic for all $k > j^*(G)$. Additionally, if $j^*(G) = 1$, then $\mathcal{A}(G) = \mathcal{A}$.*

Proof. Let ϕ be any zero-sum \mathbb{Z} -magic labeling of G such that $j(\phi) = j^*(G)$, and fix $k > j^*(G)$. We form a zero-sum \mathbb{Z}_k -magic labeling of G by assigning $\phi(e) \pmod k$

to $e \in E(G)$. Consequently, if $j^*(G) = 1$, then $\text{zim}(G) = \mathcal{N}$, implying that G is zero-sum \mathbb{A} -magic for all $\mathbb{A} \in \mathcal{A}$. \square

We observe that Georges, Mauro, and Wash [Georges et al. 2017] established necessary and sufficient conditions under which a graph G is zero-sum \mathbb{Z}_{2j}^k -magic. In particular, they showed that G has a bridge whose removal results in an isolate or bipartite component if and only if G is not zero-sum \mathbb{Z}_{2j}^k -magic for any positive integers j, k . Theorem 1 immediately implies that such G is not zero-sum \mathbb{Z} -magic for otherwise G would be zero-sum \mathbb{Z}_{2j} -magic for $2j > j^*(G)$.

Theorems 2 through 8, useful in the sequel, can be found in the existing literature.

Theorem 2 [Petersen 1891]. *Let G be a $2t$ -regular graph. Then there exist t 2-factors of G that partition $E(G)$.*

Theorem 3 [Ore 1957]. *Let G be a bridgeless regular graph of odd degree k and let h be an even integer, $2 \leq h \leq \frac{2}{3}k$. Then there exists an h -factor of G .*

Theorem 4 [Georges et al. 2010]. *Let G be a 3-regular graph. Then G is zero-sum \mathbb{Z}_2^2 -magic if and only if the chromatic index of G is 3.*

Theorem 5 [Georges et al. 2010]. *Let G be a graph with a bridge. Then for each positive integer k , G is not zero-sum \mathbb{Z}_2^k -magic.*

Theorem 6 [Choi et al. 2012]. *Let G be a bridgeless graph. Then for each positive integer $k \geq 3$, G is zero-sum \mathbb{Z}_2^k -magic.*

Theorem 7 [Low and Lee 2006; Shiu et al. 2004]. *If G_1, G_2, \dots, G_n are zero-sum \mathbb{A} -magic graphs, then $\square_{i=1}^n G_i$ is zero-sum \mathbb{A} -magic.*

Theorem 8 [Akbari et al. 2014]. *Let G be an r -regular graph, $r \geq 3$, $r \neq 5$. If r is even, then $\text{zim}(G) = \mathcal{N}$. Otherwise, $\mathcal{N} \setminus \{2, 4\} \subseteq \text{zim}(G)$.*

Corollary 9. *Suppose that G is a graph with $j^*(G) \leq 2$. Then the following hold:*

- (a) *If G is bridgeless, then $\mathcal{A} \setminus \{\mathbb{Z}_2, \mathbb{Z}_2^2\} \subseteq \mathcal{A}(G)$.*
- (b) *If G is bridgeless and even, then $\mathcal{A}(G) = \mathcal{A}$.*
- (c) *If G is bridgeless with a vertex of odd degree, then*

$$\mathcal{A} \setminus \{\mathbb{Z}_2, \mathbb{Z}_2^2\} \subseteq \mathcal{A}(G) \subseteq \mathcal{A} \setminus \{\mathbb{Z}_2\}.$$

- (d) *If G has a bridge, then $\mathcal{A}(G) = \mathcal{A} \setminus \{\mathbb{Z}_2^k \mid k \in \mathcal{N}\}$.*

Proof. (a) We note that $1 \in \text{zim}(G)$ since $j^*(G)$ is assumed to exist. The result follows by Theorems 1 and 6.

(b) Since G is even, G is zero-sum \mathbb{Z}_2 -magic (assign 1 to each edge of G), and hence zero-sum \mathbb{Z}_2^2 -magic. The result follows by part (a).

(c) Part (c) follows from part (a) and the fact that any \mathbb{Z}_2 -labeling of G must assign 1 to each edge; hence, no graph with a vertex of odd degree can be zero-sum \mathbb{Z}_2 -magic.

(d) We again observe that $1 \in \text{zim}(G)$ and, by Theorem 1, $k \in \text{zim}(G)$ for $k \geq 3$. Thus $\mathcal{N} \setminus \{2\} \subseteq \text{zim}(G)$. The result now follows from Theorem 5. \square

We note that $j^*(K_4) = 2$ and, by the upcoming Corollary 10 and Theorem 11, $j^*(K_7) = 2$ as well. Yet $\text{zim}(K_4) = \mathcal{N} \setminus \{2\}$ and, by Theorem 8, $\text{zim}(K_7) = \mathcal{N}$. Moreover, the three graphs K_4 , the Petersen graph PG , and the (unique) 3-regular graph G on 10 vertices with one bridge (see Figure 1) are each easily verified to have zero-sum integer-magic spectrum $\mathcal{N} \setminus \{2\}$. Yet, by Theorem 4 and Corollary 9, $\mathcal{A}(K_4) = \mathcal{A} \setminus \{\mathbb{Z}_2\}$, $\mathcal{A}(PG) = \mathcal{A} \setminus \{\mathbb{Z}_2, \mathbb{Z}_2^2\}$, and $\mathcal{A}(G) = \mathcal{A} \setminus \{\mathbb{Z}_2^k \mid k \in \mathcal{N}\}$.

In the following corollary, we utilize $j^*(G)$ to give an alternative proof of Theorem 8 in the case r is even, $r \geq 4$.

Corollary 10. *Let G be an even-regular graph with degree $r = 2t$ such that $t \geq 2$. Then the following hold:*

- (a) *If t is even, then $j^*(G) = 1$ and $\mathcal{A}(G) = \mathcal{A}$.*
- (b) *If t is odd, then $j^*(G) \leq 2$ and $\mathcal{A}(G) = \mathcal{A}$.*

Proof. By Theorem 2, $E(G)$ partitions into t 2-factors. If t is even, we construct a zero-sum \mathbb{Z} -magic labeling ϕ of G by assigning the label 1 to each edge of precisely $\frac{t}{2}$ 2-factors, and -1 to each edge of the remaining $\frac{t}{2}$ 2-factors. Since $j(\phi) = 1$, we have $j^*(G) = 1$, implying $\mathcal{A}(G) = \mathcal{A}$ by Theorem 1 or (since even graphs are bridgeless) Corollary 9(b). On the other hand, if t is odd, we construct a zero-sum \mathbb{Z} -magic labeling ϕ of G by assigning the label 1 to each edge of precisely $\frac{1}{2}(t + 1)$ 2-factors, -1 to each edge of precisely $\frac{1}{2}(t - 3)$ 2-factors, and -2 to each edge of the one remaining 2-factor. Since $j(\phi) = 2$, we have $j^*(G) \leq 2$, implying $\mathcal{A}(G) = \mathcal{A}$ by Corollary 9(b). \square

For illustration, Figure 4 displays vertex α of a 14-regular graph G whose edge set is partitioned by the 2-factors $H_1, H_2, H_3, H_4, H_5, H_6, H_7$. Suppose that for each i , the edge set of H_i includes the edges αu_i and αw_i . Since $t = 7$, we assign the label 1 to the edges of each of four 2-factors H_1, H_2, H_3 , and H_4 , we assign the label -1 to the edges of each of two 2-factors H_5 and H_6 , and we assign the label -2 to the edges of the remaining 2-factor H_7 . Since these labels bear upon every vertex, the weight of every vertex is 0.

We observe that the graphs $K_6 \square P_2$ and K_7 illustrate the case t odd, yet $j^*(K_6 \square P_2) = 1$ (see Theorem 11), while $j^*(K_7) = 2$.

Theorem 11. *Let G be a (connected) graph. Then $j^*(G) = 1$ if and only if G is an even graph with even size.*




Figure 4. An illustration of Corollary 10.

Proof. Suppose $j^*(G) = 1$. Then it is clear that G is even. To show that G has even size, let ϕ be a zero-sum \mathbb{Z} -magic labeling of G with $j(\phi) = 1$. Also let x denote the number of edges of G which receive the label 1 under ϕ , and let y denote the number of edges of G which receive the label -1 under ϕ . Then

$$0 = \sum_{v \in V(G)} w_v(\phi) = 2 \sum_{e \in E(G)} \phi(e) = \sum_{e \in E(G)} \phi(e),$$

implying $x = y$. Thus G has even size.

Now suppose that G is even with even size. Since G is even (and connected), there exists an Eulerian circuit in G with even length. We produce a zero-sum \mathbb{Z} -magic labeling ϕ of G with $j(\phi) = 1$ by assigning alternating labels of 1 and -1 to the edges of the circuit. □




Figure 5. Zero-sum \mathbb{Z} -magic labelings of $G_1 \square G_2$ and $G_1 \square G_3$.

Because this paper emphasizes cartesian products, Theorem 11 is illustrated in Figure 5 with an even graph $G_1 \square G_2$ with even size, and a noneven graph $G_1 \square G_3$. We exhibit a zero-sum \mathbb{Z} -magic labeling ϕ of $G_1 \square G_2$ with $j(\phi) = 1$ along with indication (via subscripts) of a corresponding Eulerian circuit. And, we show $j^*(G_1 \square G_3) = 2$ by exhibiting a zero-sum \mathbb{Z} -magic labeling of $G_1 \square G_3$ with maximum absolute label 2.

If M and N are connected odd graphs, then each has even order and $M \square N$ is an even graph with even size $|V(M)||E(N)| + |V(N)||E(M)|$. As well, if M or N is not connected, then each component of $M \square N$ is even with even size. Therefore we have the following.

Corollary 12. *Let M and N be odd graphs (not necessarily connected). Then $j^*(M \square N) = 1$.*

3. Main results

As noted in the preceding section, not all graphs are zero-sum \mathbb{Z} -magic. On the other hand, if G is the cartesian product of nontrivial graphs, we have the following result.

Theorem 13. *Let M and N be graphs with $\delta(M), \delta(N) \geq 1$. Then $M \square N$ is zero-sum \mathbb{Z} -magic with $j^*(M \square N) \leq \max\{\Delta(M), \Delta(N)\}$.*

Proof. Let $V(M) = \{u_1, \dots, u_m\}$ and $V(N) = \{w_1, \dots, w_n\}$. For fixed i , $1 \leq i \leq n$, let $M(i)$ be the subgraph of $M \square N$ induced by the vertices in $\{(u_j, w_i) \mid 1 \leq j \leq m\}$. Similarly, for fixed i , $1 \leq i \leq m$, let $N(i)$ be the subgraph of $M \square N$ induced by the vertices in $\{(u_i, w_j) \mid 1 \leq j \leq n\}$. Form a \mathbb{Z} -labeling of $M \square N$ as follows: to each edge of $M(\beta)$, assign the label $d_N(w_\beta)$ and to each edge of $N(\alpha)$, assign the label $-d_M(u_\alpha)$. Then the weight of any vertex $(u_\alpha, w_\beta) \in V(M \square N)$ is $d_M(u_\alpha)d_N(w_\beta) - d_M(u_\alpha)d_N(w_\beta) = 0$. Since the maximum absolute label is $\max\{\Delta(M), \Delta(N)\}$, the result follows. □

It is easy to show the following.

Theorem 14. *Let M and N be zero-sum \mathbb{Z} -magic graphs. Then $M \square N$ is zero-sum \mathbb{Z} -magic with $j^*(M \square N) \leq \max\{j^*(M), j^*(N)\}$.*

Theorem 15. *Let $G = \square_{i=1}^n G_i$ where $n \geq 2$ and $\delta(G_i) \geq 1$ for each i . Then G is zero-sum \mathbb{Z} -magic. Moreover, for $\zeta = \max\{\Delta(G_i)\}$ over i , we have $j^*(G) \leq \zeta$ if n is even, and $j^*(G) \leq 2\zeta$ if n is odd.*

Proof. By Theorem 13, G is zero-sum \mathbb{Z} -magic.

If $n = 2t$, let $H_i = G_{2i-1} \square G_{2i}$ for $1 \leq i \leq t$. Then by Theorem 13, each H_i is zero-sum \mathbb{Z} -magic with $j^*(H_i) \leq \max\{\Delta(G_{2i-1}), \Delta(G_{2i})\} \leq \zeta$. Since G is isomorphic to $\square_{i=1}^t H_i$, it follows from Theorem 14 that $j^*(G) \leq \max\{j^*(H_i)\} \leq \zeta$.

If $n = 2t + 1$, let $H_1 = G_1 \square (G_2 \square G_3)$ and let $H_i = G_{2i} \square G_{2i+1}$ for $2 \leq i \leq t$. By Theorem 13, each H_i is zero-sum \mathbb{Z} -magic. Observing that $\Delta(G_2 \square G_3) = \Delta(G_2) + \Delta(G_3)$, we have from Theorem 13 that $j^*(H_1) \leq \max\{\zeta, 2\zeta\} = 2\zeta$. Since $j^*(H_i) \leq \zeta$ for $i \geq 2$, we have $j^*(G) \leq 2\zeta$ by Theorem 14. \square

Lemma 16. *Let G_1, G_2 and G_3 be odd graphs, and let $M = \square_{i=1}^3 G_i$. Then $j^*(M) = 2$ and $\mathcal{A}(M) = \mathcal{A} \setminus \{\mathbb{Z}_2\}$.*

Proof. Since M is odd, we show $j^*(M) = 2$ by developing a zero-sum \mathbb{Z} -magic labeling λ of M with $j(\lambda) = 2$.

Let G_1, G_2 , and G_3 have respective vertex sets $\{u_i \mid 1 \leq i \leq n_1\}$, $\{v_i \mid 1 \leq i \leq n_2\}$, and $\{w_i \mid 1 \leq i \leq n_3\}$. Letting $V(P_2) = \{1, 2\}$, we have $j^*(G_i \square P_2) = 1$ by Corollary 12. We thus let ϕ''_i denote a zero-sum \mathbb{Z} -magic labeling of $G_i \square P_2$ such that $j(\phi''_i) = 1$. Let ϕ'_i be the \mathbb{Z} -labeling of G_i such that for each edge $\alpha\beta$ of G_i , $\phi'_i(\alpha\beta) = \phi''_i((\alpha, 1)(\beta, 1))$. Note that the weight of each vertex under ϕ'_i is 1 or -1 .

Now consider $G_1 \square G_2$. We construct a \mathbb{Z} -labeling ψ of $G_1 \square G_2$ as follows. To each edge $(u_x, v)(u_y, v)$ of $G_1 \square G_2$, assign the label $\phi'_1(u_x u_y)w_{\phi'_2(v)}$. And, to each edge $(u, v_x)(u, v_y)$ of $G_1 \square G_2$, assign the label $\phi'_2(v_x v_y)w_{\phi'_1(u)}$. We observe that under ψ , each edge of $G_1 \square G_2$ has label 1 or -1 and each vertex of $G_1 \square G_2$ has weight 2 or -2 .

Each edge e in $E(M)$ has one of the following two forms:

- Type I: $e = (z_x, w)(z_y, w)$, where $z_x z_y \in E(G_1 \square G_2)$ and $w \in V(G_3)$.
- Type II: $e = (z, w_x)(z, w_y)$, where $z \in V(G_1 \square G_2)$ and $w_x w_y \in E(G_3)$.

Let λ be a \mathbb{Z} -labeling of M such that

$$\lambda(e) = \begin{cases} \psi(z_x z_y)w_{\phi'_3(w)} & \text{if } e \text{ is of type I,} \\ -w_\psi(z)\phi'_3(w_x w_y) & \text{if } e \text{ is of type II.} \end{cases}$$

Since $\psi(z_x z_y) \in \{-1, 1\}$, $w_{\phi'_3(w)} \in \{-1, 1\}$, $w_\psi(z) \in \{-2, 2\}$, and $\phi'_3(w_x w_y) \in \{-1, 1\}$, the edges of type 1 receive ± 1 under λ and edges of type II receive labels of ± 2 under λ . It is easily checked that λ is a zero-sum \mathbb{Z} -magic labeling of M and that $j(\lambda) = 2$. Hence $j^*(M) = 2$. (In Figure 6, the evolution of λ is illustrated via edges incident to $(u, v, w) \in V(G_1 \square G_2 \square G_3)$, where each of u, v , and w is assumed to have degree 3 in G_1, G_2 , and G_3 respectively. Labels assigned to edges under ϕ'_1, ϕ'_2 , and ϕ'_3 are notional, from which labels under ψ and λ follow.)

By Corollary 9(c), it suffices to produce a zero-sum \mathbb{Z}_2^2 -magic labeling of M . For each edge $e \in E(M)$ of the form $(u_i, v, w)(u_j, v, w)$, let $\rho(e) = (0, 1)$. For each edge e of the form $(u, v_i, w)(u, v_j, w)$, let $\rho(e) = (1, 0)$. And for each edge e of the form $(u, v, w_i)(u, v, w_j)$, let $\rho(e) = (1, 1)$. It follows from the oddness of each G_i that the weight of each vertex under ρ is 0. \square




Figure 6. An illustration of the labeling λ of Lemma 16.

Theorem 17. For $1 \leq i \leq n$, let G_i be an odd graph, and let $G = \square_{i=1}^n G_i$. Then

- (a) If n is even, $j^*(G) = 1$ and $\mathcal{A}(G) = \mathcal{A}$.
- (b) If n is odd, $j^*(G) = 2$ and $\mathcal{A}(G) = \mathcal{A} \setminus \{\mathbb{Z}_2\}$.

Proof. (a) Let $n = 2t$. Then G is isomorphic to $\square_{i=1}^t H_i$, where $H_i = G_{2i-1} \square G_{2i}$. By Corollary 12, $j^*(H_i) = 1$. The result now follows from Theorems 14 and 1.

(b) Let $n = 2t + 1$, where $t \geq 1$. Let $H_1 = G_1 \square G_2 \square G_3$ and for $2 \leq i \leq t$, let $H_i = G_{2i} \square G_{2i+1}$. By Lemma 16, $j^*(H_1) = 2$ and $\mathcal{A}(H_1) = \mathcal{A} \setminus \{\mathbb{Z}_2\}$. And, by Corollary 12, $j^*(H_i) = 1$ for $2 \leq i \leq t$, implying $\mathcal{A}(H_i) = \mathcal{A}$. Thus, by Theorem 7, Theorem 14 and the fact that G is odd, $j^*(G) = 2$ and $\mathcal{A}(G) = \mathcal{A} \setminus \{\mathbb{Z}_2\}$. \square

We now consider graphs $G = \square_{i=1}^n G_i$ such that $n \geq 2$, G_i is nontrivial, and G is r -regular. Since the regularity of G coimplies the regularity of each G_i , we assume G_i is r_i -regular, $r_i \geq 1$. Then the degree of G is $r = \sum_{i=1}^n r_i$.

The next theorem follows from Theorem 8 or alternatively Corollary 10 with the 4-cycle handled as a trivial special case.

Theorem 18. Let $n \geq 2$ and let $G = \square_{i=1}^n G_i$ be an even-regular graph. Then $\mathcal{A}(G) = \mathcal{A}$.

We turn to the case r is odd. For odd-regular graphs M except 5-regular, Theorem 8 indicates that M is zero-sum \mathbb{A} -magic for all \mathbb{A} except possibly $\mathbb{Z}_2^k, \mathbb{Z}_4^k, k \geq 1$. (Certainly M is not zero-sum \mathbb{Z}_2 -magic.) However, if M is a cartesian product of at least two factors, then more definitive results can be given for any odd regularity.

Lemma 19. *Let $G = G_1 \square G_2$ be an odd-regular graph, where $\delta(G_i) \geq 1$. Then $\mathcal{A} \setminus \{\mathbb{Z}_2, \mathbb{Z}_2^2\} \subseteq \mathcal{A}(G) \subseteq \mathcal{A} \setminus \{\mathbb{Z}_2\}$. Moreover, if either G_1 or G_2 has degree 1, then $\mathcal{A}(G) = \mathcal{A} \setminus \{\mathbb{Z}_2\}$.*

Proof. Since G is regular if and only if each G_i is regular, we assume without loss of generality G_i is r_i -regular for odd r_1 and even r_2 . Let $r_2 = 2t$.

If $r_1 \geq 3$, then by Theorem 3, $E(G_1)$ partitions into a $\frac{1}{2}(r_1 + 1)$ -factor and a $\frac{1}{2}(r_1 - 1)$ -factor. By Petersen’s theorem, $E(G_2)$ partitions into t 2-factors. Thus $E(G)$ naturally partitions into t 2-factors F_1, F_2, \dots, F_t , a $\frac{1}{2}(r_1 + 1)$ -factor F' , and a $\frac{1}{2}(r_1 - 1)$ -factor F'' . We form a zero-sum \mathbb{Z} -magic labeling ϕ of G with $j(\phi) = 2$ based on the parity of t .

If $t = 2k + 1$,

$$\phi(e) = \begin{cases} 1 & \text{if } e \in F_i, 1 \leq i \leq k, \\ -1 & \text{if } e \in F_i, k + 1 \leq i \leq 2k + 1, \\ 2 & \text{if } e \in F', \\ -2 & \text{if } e \in F''. \end{cases}$$

If $t = 2k$,

$$\phi(e) = \begin{cases} 1 & \text{if } e \in F_i, 1 \leq i \leq k, \\ -1 & \text{if } e \in F_i, k + 1 \leq i \leq 2k - 1, \\ -2 & \text{if } e \in F_{2k}, \\ 2 & \text{if } e \in F', \\ -2 & \text{if } e \in F''. \end{cases}$$

Since $j(\phi) = 2$ in each case, we have $j^*(G) \leq 2$. The result follows by Corollary 9(c).

Suppose $r_1 = 1$. Then the edge set $E(G)$ partitions naturally into a 1-factor F' (of which there are $|V(G_2)|$ components each isomorphic to P_2) and, by Petersen’s theorem, t 2-factors F_1, F_2, \dots, F_t . We form a zero-sum \mathbb{Z} -magic labeling ϕ of G with $j(\phi) = 2$ as above, accounting for the vacuous $\frac{1}{2}(r_1 - 1)$ -factor.

If $t = 2k + 1$ is odd,

$$\phi(e) = \begin{cases} 1 & \text{if } e \in F_i, 1 \leq i \leq k, \\ -1 & \text{if } e \in F_i, k + 1 \leq i \leq 2k + 1, \\ 2 & \text{if } e \in F'. \end{cases}$$

If $t = 2k$ is even,

$$\phi(e) = \begin{cases} 1 & \text{if } e \in F_i, 1 \leq i \leq k, \\ -1 & \text{if } e \in F_i, k + 1 \leq i \leq 2k - 1, \\ -2 & \text{if } e \in F_{2k}, \\ 2 & \text{if } e \in F'. \end{cases}$$

Thus $j^*(G) \leq 2$, implying $\mathcal{A} \setminus \{\mathbb{Z}_2, \mathbb{Z}_2^2\} \subseteq \mathcal{A}(G) \subseteq \mathcal{A} \setminus \{\mathbb{Z}_2\}$ by Corollary 9(c). It remains to show that G is zero-sum \mathbb{Z}_2^2 -magic.

Let H denote a 2-factor of G_2 . Then $P_2 \square H$ is a 3-factor of G (not necessarily connected) in which each component is a prism. Since each prism is hamiltonian and hence 3-edge colorable, $P_2 \square H$ is zero-sum \mathbb{Z}_2^2 -magic by Theorem 4. Letting ϕ' be a zero-sum \mathbb{Z}_2^2 -magic labeling of $P_2 \square H$, we form a zero-sum \mathbb{Z}_2^2 -magic labeling ϕ of G as follows:

$$\phi(e) = \begin{cases} \phi'(e) & \text{if } e \in E(P_2 \square H), \\ (1, 1) & \text{if } e \in E(G - (P_2 \square H)). \end{cases} \quad \square$$

Lemma 20. *Suppose $G = G_1 \square G_2 \square G_3$ is odd-regular, where G_i is r_i -regular, $r_i \geq 1$. Suppose also that either $r_i = 1$ for some i or r_i is odd for all i . Then $\mathcal{A}(G) = \mathcal{A} \setminus \{\mathbb{Z}_2\}$.*

Proof. If r_i is odd for all i , then the result follows from Theorem 17. So, with no loss of generality, suppose $r_1 = 1$ and r_2, r_3 are even. Since G is isomorphic to $G_1 \square H$, where $H = G_2 \square G_3$, the result follows from Lemma 19. \square

Theorem 21. *Suppose $G = \square_{i=1}^n G_i$ is odd-regular, where $n \geq 3$ and G_i is r_i -regular, $r_i \geq 1$. Let ω denote the number of odd-regular G_i . Then the following hold:*

- (a) *If $\omega \geq 3$, then $\mathcal{A}(G) = \mathcal{A} \setminus \{\mathbb{Z}_2\}$.*
- (b) *If $\omega = 1$, then $\mathcal{A} \setminus \{\mathbb{Z}_2, \mathbb{Z}_2^2\} \subseteq \mathcal{A}(G) \subseteq \mathcal{A} \setminus \{\mathbb{Z}_2\}$. Moreover, if the sole odd-regular G_i has degree 1, then $\mathcal{A}(G) = \mathcal{A} \setminus \{\mathbb{Z}_2\}$.*

Proof. We observe that ω must be odd since G is odd-regular with degree equal to $\sum_{i=1}^n r_i$.

(a) With no loss of generality, let G_1 and G_2 be odd-regular. Then $\square_{i=3}^n G_i$ is odd regular as well, which implies that G is isomorphic to the cartesian product of three odd-regular graphs G_1, G_2 , and $\square_{i=3}^n G_i$. The result follows by Lemma 20.

(b) With no loss of generality, let G_1 be odd-regular. Then $\square_{i=2}^n G_i$ is even-regular, which implies that G is isomorphic to the cartesian product of one odd-regular graph G_1 and one even-regular graph $\square_{i=2}^n G_i$. The result follows by Lemma 19. \square

4. Grid graphs

In this section we consider the cartesian product of paths $G = \square_{i=1}^n P_{a_i}$, where $n \geq 2$ and $a_i \geq 2$. We denote the vertex set of P_m by $\{1, 2, 3, 4, \dots, m\}$, where x is adjacent to y if and only if $|x - y| = 1$. We note that if α is the number of a_i 's equal to 2, then $\Delta(G) = 2n - \alpha$, $\delta(G) = n$, and for all integers j , $n \leq j \leq 2n - \alpha$, there exists a vertex of G with degree j . We also note that G is regular (in particular, n -regular) if and only if $a_i = 2$ for all i .

Lemma 22. *Let M be a graph such that $j^*(M) \leq 2$ and $j^*(M \square P_2) \leq 2$. Then $j^*(M \square P_n) \leq 2$ for $n \geq 3$.*

Proof. Let H' denote the subgraph of $M \square P_n$ induced by the vertices in $\{(v, 1) \mid v \in V(M)\}$ and let H'' denote the subgraph of $M \square P_n$ induced by the vertices in $\{(v, i) \mid v \in V(M), i = 1, 2\}$. Since H' and H'' are respectively isomorphic to M and $M \square P_2$, we can find zero-sum \mathbb{Z} -magic labelings ϕ' of H' and ϕ'' of H'' such that $j(\phi')$ and $j(\phi'')$ are each at most 2. As follows, we construct a zero-sum \mathbb{Z} -magic labeling ϕ of $M \square P_n$ that draws its labels from the images of ϕ' and ϕ'' (this labeling is illustrated in Figure 7 with $n = 5$ and a graph M that is seen by inspection to satisfy the hypotheses of the lemma):

$$\phi(e) = \begin{cases} \phi''(e) & \text{if } e = (u, 1)(w, 1), \\ \phi'((u, 1)(w, 1)) & \text{if } e = (u, i)(w, i), 2 \leq i \leq n - 1, \\ \phi''((u, 1)(u, 2)) & \text{if } e = (u, i)(u, i + 1), 1 \leq i \leq n - 1, i \text{ odd}, \\ -\phi''((u, 1)(u, 2)) & \text{if } e = (u, i)(u, i + 1), 1 \leq i \leq n - 1, i \text{ even}, \\ -\phi''((u, 1)(w, 1)) & \text{if } e = (u, n)(w, n), n \text{ odd}, \\ \phi''((u, 1)(w, 1)) & \text{if } e = (u, n)(w, n), n \text{ even}. \end{cases}$$

Thus $j(\phi) \leq 2$, giving the result. □

Lemma 23. *Let M be a graph and let $\mathbb{A} \in \mathcal{A}$ such that both M and $M \square P_2$ are zero-sum \mathbb{A} -magic. Then $M \square P_n$ is zero-sum \mathbb{A} -magic for $n \geq 3$.*

Proof. Let H' and H'' denote the subgraphs of $M \square P_n$ given in the proof of Lemma 22, and let ϕ' and ϕ'' be zero-sum \mathbb{A} -magic labelings of H' and H'' , respectively. Then the labeling ϕ of that proof is a zero-sum \mathbb{A} -magic labeling of $M \square P_n$. □

Theorem 24. *Suppose $G = \square_{i=1}^n P_{a_i}$, where $n \geq 2$. If n is even and $a_i = 2$ for all i , then $\mathcal{A}(G) = \mathcal{A}$. Otherwise, $\mathcal{A}(G) = \mathcal{A} \setminus \{\mathbb{Z}_2\}$.*

Proof. If n is even and $a_i = 2$ for all i , the result follows from Theorem 18. If n is odd and $a_i = 2$ for all i , the result follows from Lemma 19 and the observation that G is isomorphic to $H_1 \square H_2$, where $H_1 = P_2$ and $H_2 = \square_{i=2}^n P_2$. We thus assume $a_i \geq 3$ for some i , which implies that G has a vertex of odd degree.




Figure 7. An illustration of the labeling ϕ of Lemma 22.

Express G as follows:

- If $n = 2k$, then $G = \square_{i=1}^k H_i$, where $H_i = P_{a_{2i-1}} \square P_{a_{2i}}$.
- If $n = 2k + 1$, then $G = \square_{i=1}^k H_i$, where $H_1 = P_{a_1} \square P_{a_2} \square P_{a_3}$ and $H_i = P_{a_{2i}} \square P_{a_{2i+1}}$ for $2 \leq i \leq k$.

Each H_i must be isomorphic to one of the following:

- Class 1: $P_2 \square P_2$.
- Class 2: $P_r \square P_s$, $r \geq 2$, $s > 2$.
- Class 3: $P_2 \square P_2 \square P_2$.
- Class 4: $P_2 \square P_2 \square P_s$, $s \geq 3$.
- Class 5: $P_2 \square P_s \square P_t$, $s, t \geq 3$.
- Class 6: $P_s \square P_t \square P_r$, $s, t, r \geq 3$.

We first show that for each H_i , we have $j^*(H_i) \leq 2$.

By Theorem 15, the graphs H_i of classes 1, 2, and 3 have $j^*(H_i) \leq 2$.

Consider graphs of class 4. Since $j^*(P_2 \square P_2) \leq 2$ and $j^*((P_2 \square P_2) \square P_2) \leq 2$, graphs H_i of class 4 have $j^*(H_i) \leq 2$ by Lemma 22.

Consider graphs of class 5. Since $j^*(P_2 \square P_s) \leq 2$ and $j^*((P_2 \square P_s) \square P_2) \leq 2$, graphs H_i of class 5 have $j^*(H_i) \leq 2$ by Lemma 22.




Figure 8. A zero-sum \mathbb{Z}_2^2 -magic labeling of $P_4 \square P_5$.

Consider graphs of class 6. Since $j^*(P_s \square P_t) \leq 2$ and $j^*((P_s \square P_t) \square P_2) \leq 2$, graphs H_i of class 6 have $j^*(H_i) \leq 2$ by Lemma 22.

Thus, by Theorem 14, $j^*(G) \leq 2$, which implies by Corollary 9(c) that

$$\mathcal{A} \setminus \{\mathbb{Z}_2, \mathbb{Z}_2^2\} \subseteq \mathcal{A}(G) \subseteq \mathcal{A} \setminus \{\mathbb{Z}_2\}.$$

It now suffices to show that G is zero-sum \mathbb{Z}_2^2 -magic. To that end, we next show that each H_i is zero-sum \mathbb{Z}_2^2 -magic.

By inspection, the graph H_i of class 1 has $\mathcal{A}(H_i) = \mathcal{A}$.

By Corollary 9(c), the graphs H_i of class 2 are zero-sum \mathbb{A} -magic for all \mathbb{A} except \mathbb{Z}_2 and possibly \mathbb{Z}_2^2 . But it is an easy matter to construct a zero-sum \mathbb{Z}_2^2 -magic labeling of H_i , thereby establishing that $\mathcal{A}(H_i) = \mathcal{A} \setminus \{\mathbb{Z}_2\}$. Particularly, consider the following sequence of vertices that specifies a cycle:

$$(1, 1), \dots, (1, s), (2, s), \dots, (r, s), (r, s - 1), \dots, (r, 1), (r - 1, 1), \dots, (1, 1).$$

Moving around the cycle, assign each edge a label from $\{(0, 1), (1, 0)\}$ according to the following algorithm and as illustrated in Figure 8 (where parentheses are omitted): assign $(0, 1)$ to the edge $(1, 1)(1, 2)$. For every two distinct edges e' and e'' of the cycle, assign distinct labels if e' and e'' are incident to a common vertex of degree 3 in G ; otherwise, assign equal labels if e' and e'' are incident to a common vertex of degree 2 in G . To each other edge of G , assign $(1, 1)$.

By Theorem 21(a), the graph H_i of class 3 has $\mathcal{A}(H_i) = \mathcal{A} \setminus \{\mathbb{Z}_2\}$.

Consider graphs of class 4. Since $P_2 \square P_2$ is of class 1, it is zero-sum \mathbb{A} -magic for all \mathbb{A} . And since $(P_2 \square P_2) \square P_2$ is of class 3, it is zero-sum \mathbb{A} -magic for all \mathbb{A} except \mathbb{Z}_2 . Thus, by Lemma 23 and the fact that graphs H_i of class 4 have vertices of odd degree, $\mathcal{A}(H_i) = \mathcal{A} \setminus \{\mathbb{Z}_2\}$.

Consider graphs of class 5. Since $P_2 \square P_s$ is of class 2, it is zero-sum \mathbb{A} -magic for all \mathbb{A} except \mathbb{Z}_2 . And since $(P_2 \square P_s) \square P_2$ is of class 4, it is zero-sum \mathbb{A} -magic for all \mathbb{A} except \mathbb{Z}_2 . Thus, by Lemma 23 and the fact that graphs H_i of class 5 have vertices of odd degree, $\mathcal{A}(H_i) = \mathcal{A} \setminus \{\mathbb{Z}_2\}$.

Consider graphs of class 6. Since $P_s \square P_t$ is of class 2, it is zero-sum \mathbb{A} -magic for all \mathbb{A} except \mathbb{Z}_2 . And since $(P_s \square P_t) \square P_2$ is of class 5, it is zero-sum \mathbb{A} -magic

for all \mathbb{A} except \mathbb{Z}_2 . Thus, by Lemma 23 and the fact that graphs H_i of class 6 have vertices of odd degree, $\mathcal{A}(H_i) = \mathcal{A} \setminus \{\mathbb{Z}_2\}$.

We therefore see by Theorem 7 that $\mathcal{A} \setminus \{\mathbb{Z}_2\} \subseteq \mathcal{A}(G)$. But since G has a vertex of odd degree, G is not zero-sum \mathbb{Z}_2 -magic. Thus $\mathcal{A}(G) = \mathcal{A} \setminus \{\mathbb{Z}_2\}$. \square

We close this section with a theorem that utilizes Corollary 12.

Theorem 25. *Let $n \geq 3$ and let G be an odd graph. Then $\mathcal{A}(G \square P_n) = \mathcal{A} \setminus \{\mathbb{Z}_2\}$.*

Proof. Let H' be the subgraph of $G \square P_n$ induced by $\{(v, i) \mid v \in V(G), i = 1, 2\}$. Since H' is isomorphic to $G \square P_2$, Corollary 12 implies the existence of a zero-sum \mathbb{Z} -magic labeling ϕ' of H' such that $j(\phi') = 1$. We establish a zero-sum \mathbb{Z} -magic labeling ϕ of $G \square P_n$ with $j(\phi) = 2$, thereby establishing $j^*(G \square P_n) \leq 2$:

$$\phi(e) = \begin{cases} \phi'((u, 1)(w, 1)) & \text{if } e = (u, 1)(w, 1) \text{ or } (u, n)(w, n), \\ 2\phi'((u, 1)(w, 1)) & \text{if } e = (u, i)(w, i), 2 \leq i \leq n - 1, \\ \phi'((u, 1)(u, 2)) & \text{if } e = (u, i)(u, i + 1), 1 \leq i \leq n - 1. \end{cases}$$

By Corollary 9(c), it now suffices to establish a zero-sum \mathbb{Z}_2^2 -magic labeling ϕ of $G \square P_n$:

$$\phi(e) = \begin{cases} (0, 1) & \text{if } e = (u, 1)(w, 1), \\ (1, 1) & \text{if } e = (u, i)(w, i), 2 \leq i \leq n - 1, \\ (0, 1) & \text{if } e = (u, n)(w, n), n \text{ even}, \\ (1, 0) & \text{if } e = (u, n)(w, n), n \text{ odd}, \\ (0, 1) & \text{if } e = (u, i)(u, i + 1), i \text{ odd}, \\ (1, 0) & \text{if } e = (u, i)(u, i + 1), i \text{ even}. \end{cases} \quad \square$$

5. Closing remarks

We close this paper with suggestions for further study.

If G is the cartesian product of two odd graphs, then $j^*(G) = 1$. What can be said of $j^*(G)$ if one or each factor is even?

If T_1, T_2, \dots, T_n is a collection of nontrivial trees and $G = \square_{i=1}^n T_i$, we are able to show that $j^*(G) \leq 4$. Is this a sharp upper bound?

What can be said of $j^*(G)$ if G is an odd cartesian product?

We have seen that the graphs G under study have $j^*(G) \leq 2$, with $j^*(G) = 1$ if and only if G is an even graph with even size. We have also seen that $j^*(G) = 1$ is a sufficient but not necessary condition for $\mathcal{A}(G) = \mathcal{A}$. Are there cartesian products G such that $j^*(G) \geq 3$? Are there necessary and sufficient conditions for $j^*(G) = 2$?

Acknowledgments

This paper reflects the efforts of the participants of an undergraduate research seminar in graph theory at Trinity College. The authors, including students A. Fong,

D. Spagnuolo, J. Wallace, and S. Wang, gratefully acknowledge the support and sponsorship of Dr. T. Savage. We also express our gratitude to the referee for thoughtful and constructive direction.

References

- [Akbari et al. 2014] S. Akbari, F. Rahmati, and S. Zare, “Zero-sum magic labelings and null sets of regular graphs”, *Electron. J. Combin.* **21**:2 (2014), Paper 2.17. MR Zbl
- [Choi et al. 2012] J. O. Choi, J. P. Georges, and D. Mauro, “Relating edge-coverings to the classification of \mathbb{Z}_2^k -magic graphs”, *Discrete Math.* **312**:19 (2012), 2938–2945. MR Zbl
- [Gallian 2018] J. A. Gallian, “A dynamic survey of graph labeling”, *Electron. J. Combin.* **5** (2018), Dynamic Survey 6. Zbl
- [Georges et al. 2010] J. P. Georges, D. W. Mauro, and Y. Wang, “On the structures of V_4 -magic and \mathbb{Z}_4 -magic graphs”, *J. Combin. Math. Combin. Comput.* **75** (2010), 137–152. MR Zbl
- [Georges et al. 2017] J. P. Georges, D. Mauro, and K. Wash, “On zero-sum \mathbb{Z}_{2j}^k -magic graphs”, *J. Comb. Optim.* **34**:1 (2017), 94–113. MR Zbl
- [Low and Lee 2006] R. M. Low and S.-M. Lee, “On the products of group-magic graphs”, *Australas. J. Combin.* **34** (2006), 41–48. MR Zbl
- [Ore 1957] O. Ore, “Graphs and subgraphs”, *Trans. Amer. Math. Soc.* **84** (1957), 109–136. MR Zbl
- [Petersen 1891] J. Petersen, “Die Theorie der regulären graphs”, *Acta Math.* **15**:1 (1891), 193–220. MR Zbl
- [Sedláček 1964] J. Sedláček, “Problem 27”, pp. 163–167 in *Theory of graphs and its applications* (Smolenice, 1963), edited by M. Fiedler, Publ. House Czechoslovak Acad. Sciences, Prague, 1964.
- [Shiu and Low 2018] W. C. Shiu and R. M. Low, “The integer-magic spectra and null sets of the Cartesian product of trees”, *Australas. J. Combin.* **70** (2018), 157–167. MR Zbl
- [Shiu et al. 2004] W. C. Shiu, P. C. B. Lam, and P. K. Sun, “Construction of group-magic graphs and some A -magic graphs with A of even order”, pp. 97–107 in *Proceedings of the Thirty-Fifth Southeastern International Conference on Combinatorics, Graph Theory and Computing* (Boca Raton, FL), Congressus Numerantium **167**, Utilitas Math., Winnipeg, MB, 2004. MR Zbl
- [Stanley 1973] R. P. Stanley, “Linear homogeneous Diophantine equations and magic labelings of graphs”, *Duke Math. J.* **40** (1973), 607–632. MR Zbl

Received: 2018-03-14 Revised: 2019-08-27 Accepted: 2019-09-20

adam.y.w.fong@gmail.com *Department of Mathematics, Trinity College, Hartford, CT, United States*

john.georges@trincoll.edu *Department of Mathematics, Trinity College, Hartford, CT, United States*

david.mauro@trincoll.edu *Department of Mathematics, Trinity College, Hartford, CT, United States*

dylanspag@gmail.com *Department of Mathematics, Trinity College, Hartford, CT, United States*

jackwallacemv@gmail.com *Department of Mathematics, Trinity College, Hartford, CT, United States*

1278 FONG, GEORGES, MAURO, SPAGNUOLO, WALLACE, WANG AND WASH

shufan.wang.2018@trincoll.edu *Department of Mathematics, Trinity College, Hartford, CT,
United States*

kwashmath@gmail.com *Department of Mathematics, Trinity College, Hartford, CT,
United States*

involve

msp.org/involve

INVOLVE YOUR STUDENTS IN RESEARCH

Involve showcases and encourages high-quality mathematical research involving students from all academic levels. The editorial board consists of mathematical scientists committed to nurturing student participation in research. Bridging the gap between the extremes of purely undergraduate research journals and mainstream research journals, *Involve* provides a venue to mathematicians wishing to encourage the creative involvement of students.

MANAGING EDITOR

Kenneth S. Berenhaut Wake Forest University, USA

BOARD OF EDITORS

Colin Adams	Williams College, USA	Robert B. Lund	Clemson University, USA
Arthur T. Benjamin	Harvey Mudd College, USA	Gaven J. Martin	Massey University, New Zealand
Martin Bohner	Missouri U of Science and Technology, USA	Mary Meyer	Colorado State University, USA
Amarjit S. Budhiraja	U of N Carolina, Chapel Hill, USA	Frank Morgan	Williams College, USA
Pietro Cerone	La Trobe University, Australia	Mohammad Sal Moslehian	Ferdowsi University of Mashhad, Iran
Scott Chapman	Sam Houston State University, USA	Zuhair Nashed	University of Central Florida, USA
Joshua N. Cooper	University of South Carolina, USA	Ken Ono	Univ. of Virginia, Charlottesville
Jem N. Corcoran	University of Colorado, USA	Yuval Peres	Microsoft Research, USA
Toka Diagana	Howard University, USA	Y.-F. S. Pétermann	Université de Genève, Switzerland
Michael Dorff	Brigham Young University, USA	Jonathon Peterson	Purdue University, USA
Sever S. Dragomir	Victoria University, Australia	Robert J. Plemmons	Wake Forest University, USA
Joel Foisy	SUNY Potsdam, USA	Carl B. Pomerance	Dartmouth College, USA
Errin W. Fulp	Wake Forest University, USA	Vadim Ponomarenko	San Diego State University, USA
Joseph Gallian	University of Minnesota Duluth, USA	Bjorn Poonen	UC Berkeley, USA
Stephan R. Garcia	Pomona College, USA	József H. Przytycki	George Washington University, USA
Anant Godbole	East Tennessee State University, USA	Richard Rebarber	University of Nebraska, USA
Ron Gould	Emory University, USA	Robert W. Robinson	University of Georgia, USA
Sat Gupta	U of North Carolina, Greensboro, USA	Javier Rojo	Oregon State University, USA
Jim Haglund	University of Pennsylvania, USA	Filip Saidak	U of North Carolina, Greensboro, USA
Johnny Henderson	Baylor University, USA	Hari Mohan Srivastava	University of Victoria, Canada
Glenn H. Hurlbert	Virginia Commonwealth University, USA	Andrew J. Sterge	Honorary Editor
Charles R. Johnson	College of William and Mary, USA	Ann Trenk	Wellesley College, USA
K. B. Kulasekera	Clemson University, USA	Ravi Vakil	Stanford University, USA
Gerry Ladas	University of Rhode Island, USA	Antonia Vecchio	Consiglio Nazionale delle Ricerche, Italy
David Larson	Texas A&M University, USA	John C. Wierman	Johns Hopkins University, USA
Suzanne Lenhart	University of Tennessee, USA	Michael E. Zieve	University of Michigan, USA
Chi-Kwong Li	College of William and Mary, USA		

PRODUCTION

Silvio Levy, Scientific Editor


Cover: Alex Scorpan

See inside back cover or msp.org/involve for submission instructions. The subscription price for 2019 is US \$195/year for the electronic version, and \$260/year (+\$35, if shipping outside the US) for print and electronic. Subscriptions, requests for back issues and changes of subscriber address should be sent to MSP.

Involve (ISSN 1944-4184 electronic, 1944-4176 printed) at Mathematical Sciences Publishers, 798 Evans Hall #3840, c/o University of California, Berkeley, CA 94720-3840, is published continuously online. Periodical rate postage paid at Berkeley, CA 94704, and additional mailing offices.

Involve peer review and production are managed by EditFLOW[®] from Mathematical Sciences Publishers.

PUBLISHED BY

 **mathematical sciences publishers**
nonprofit scientific publishing

<http://msp.org/>

© 2019 Mathematical Sciences Publishers

involve

2019 vol. 12 no. 8

On the zero-sum group-magickness of cartesian products	1261
ADAM FONG, JOHN GEORGES, DAVID MAURO, DYLAN SPAGNUOLO, JOHN WALLACE, SHUFAN WANG AND KIRSTI WASH	
The variable exponent Bernoulli differential equation	1279
KAREN R. RÍOS-SOTO, CARLOS E. SEDA-DAMIANI AND ALEJANDRO VÉLEZ-SANTIAGO	
The supersingularity of Hurwitz curves	1293
ERIN DAWSON, HENRY FRAUENHOFF, MICHAEL LYNCH, AMETHYST PRICE, SEAMUS SOMERSTEP, ERIC WORK, DEAN BISOGNO AND RACHEL PRIES	
Multicast triangular semilattice network	1307
ANGELINA GROSSO, FELICE MANGANIELLO, SHIWANI VARAL AND EMILY ZHU	
Edge-transitive graphs and combinatorial designs	1329
HEATHER A. NEWMAN, HECTOR MIRANDA, ADAM GREGORY AND DARREN A. NARAYAN	
A logistic two-sex model with mate-finding Allee effect	1343
ELIZABETH ANDERSON, DANIEL MAXIN, JARED OTT AND GWYNETH TERRETT	
Unoriented links and the Jones polynomial	1357
SANDY GANZELL, JANET HUFFMAN, LESLIE MAVRAKIS, KAITLIN TADEMY AND GRIFFIN WALKER	
Nonsplit module extensions over the one-sided inverse of $k[x]$	1369
ZHEPING LU, LINHONG WANG AND XINGTING WANG	
Split Grothendieck rings of rooted trees and skew shapes via monoid representations	1379
DAVID BEERS AND MATT SZCZESNY	
On the classification of Specht modules with one-dimensional summands	1399
AUBREY PIPER COLLINS AND CRAIG J. DODGE	
The monochromatic column problem with a prime number of colors	1415
LORAN CROWELL AND STEVE SZABO	
Total Roman domination edge-critical graphs	1423
CHLOE LAMPMAN, KIEKA (C. M.) MYNHARDT AND SHANNON OGDEN	



1944-4176(2019)12:8;1-Y