

involve

a journal of mathematics

Total Roman domination edge-critical graphs

Chloe Lampman, Kieka (C. M.) Mynhardt and Shannon Ogden



Total Roman domination edge-critical graphs

Chloe Lampman, Kieka (C. M.) Mynhardt and Shannon Ogden

(Communicated by Anant Godbole)

A total Roman dominating function on a graph G is a function $f : V(G) \rightarrow \{0, 1, 2\}$ such that every vertex v with $f(v) = 0$ is adjacent to some vertex u with $f(u) = 2$, and the subgraph of G induced by the set of all vertices w such that $f(w) > 0$ has no isolated vertices. The weight of f is $\sum_{v \in V(G)} f(v)$. The total Roman domination number $\gamma_{tR}(G)$ is the minimum weight of a total Roman dominating function on G . A graph G is k - γ_{tR} -edge-critical if $\gamma_{tR}(G + e) < \gamma_{tR}(G) = k$ for every edge $e \in E(\bar{G}) \neq \emptyset$, and k - γ_{tR} -edge-supercritical if it is k - γ_{tR} -edge-critical and $\gamma_{tR}(G + e) = \gamma_{tR}(G) - 2$ for every edge $e \in E(\bar{G}) \neq \emptyset$. We present some basic results on γ_{tR} -edge-critical graphs and characterize certain classes of γ_{tR} -edge-critical graphs. In addition, we show that, when k is small, there is a connection between k - γ_{tR} -edge-critical graphs and graphs which are critical with respect to the domination and total domination numbers.

1. Introduction

We consider the behaviour of the total Roman domination number of a graph G upon the addition of edges to G . A *dominating set* S in a graph G is a set of vertices such that every vertex in $V(G) - S$ is adjacent to at least one vertex in S . The *domination number* $\gamma(G)$ is the cardinality of a minimum dominating set in G . A *total dominating set* S (abbreviated by *TD-set*) in a graph G with no isolated vertices is a set of vertices such that every vertex in $V(G)$ is adjacent to at least one vertex in S . The *total domination number* $\gamma_t(G)$ (abbreviated by *TD-number*) is the cardinality of a minimum total dominating set in G . For $S \subseteq V(G)$ and a function $f : S \rightarrow \mathbb{R}$, define $f(S) = \sum_{s \in S} f(s)$. A *Roman dominating function* (abbreviated by *RD-function*) on a graph G is a function $f : V(G) \rightarrow \{0, 1, 2\}$ such that every

MSC2010: 05C69.

Keywords: Roman domination, total Roman domination, total Roman domination edge-critical graphs.

Lampman was supported by an Undergraduate Student Research Award from the Natural Sciences and Engineering Research Council of Canada. Mynhardt was supported by a Discovery Grant from the Natural Sciences and Engineering Research Council of Canada. Ogden was supported by a Science Undergraduate Research Award from the University of Victoria.

vertex v with $f(v) = 0$ is adjacent to some vertex u with $f(u) = 2$. The *weight* of f , denoted by $\omega(f)$, is defined as $f(V(G))$. The *Roman domination number* $\gamma_R(G)$ (abbreviated by *RD-number*) is defined as $\min\{\omega(f) : f \text{ is an RD-function on } G\}$. For an RD-function f , let $V_f^i = \{v \in V(G) : f(v) = i\}$ and $V_f^+ = V_f^1 \cup V_f^2$. Thus, we can uniquely express an RD-function f as $f = (V_f^0, V_f^1, V_f^2)$.

As defined by Ahangar, Henning, Samodivkin and Yero [2016], a *total Roman dominating function* (abbreviated by *TRD-function*) on a graph G with no isolated vertices is a Roman dominating function with the additional condition that $G[V_f^+]$ has no isolated vertices. The *total Roman domination number* $\gamma_{tR}(G)$ (abbreviated by *TRD-number*) is the minimum weight of a TRD-function on G ; that is, $\gamma_{tR}(G) = \min\{\omega(f) : f \text{ is a TRD-function on } G\}$. A TRD-function f such that $\omega(f) = \gamma_{tR}(G)$ is called a $\gamma_{tR}(G)$ -*function*, or a γ_{tR} -*function* if the graph G is clear from the context; γ_R -*functions* are defined analogously.

The addition of an edge to a graph has the potential to change its total domination or Roman domination number. Van der Merwe, Mynhardt and Haynes [1998b] studied γ_t -*edge-critical graphs*, that is, graphs G for which $\gamma_t(G + e) < \gamma_t(G)$ for each $e \in E(\bar{G})$ and $E(\bar{G}) \neq \emptyset$. We consider the same concept for total Roman domination. A graph G is *total Roman domination edge-critical*, or simply γ_{tR} -*edge-critical*, if $\gamma_{tR}(G + e) < \gamma_{tR}(G)$ for every edge $e \in E(\bar{G})$ and $E(\bar{G}) \neq \emptyset$. We say that G is k - γ_{tR} -*edge-critical* if $\gamma_{tR}(G) = k$ and G is γ_{tR} -edge-critical. If $\gamma_{tR}(G + e) \leq \gamma_{tR}(G) - 2$ for every edge $e \in E(\bar{G})$ and $E(\bar{G}) \neq \emptyset$, we say that G is γ_{tR} -*edge-supercritical*. If $\gamma_{tR}(G + e) = \gamma_{tR}(G)$ for all $e \in E(\bar{G})$, or $E(\bar{G}) = \emptyset$, we say that G is *stable*.

Pushpam and Padmapriya [2017] established bounds on the total Roman domination number of a graph in terms of its order and girth. Total Roman domination in trees was studied by Amjadi, Nazari-Moghaddam, Sheikholeslami and Volkman [2017], as well as by Amjadi, Sheikholeslami and Soroudi [2019]. Amjadi, Sheikholeslami, and Soroudi [2018] also studied Nordhaus–Gaddum bounds for total Roman domination. Campanelli and Kuziak [2019] considered total Roman domination in the lexicographic product of graphs. We refer the reader to the well-known books [Chartrand and Lesniak 2016; Haynes, Hedetniemi, and Slater 1998] for graph theory concepts not defined here. Frequently used or lesser known concepts are defined where needed.

We begin with some general results regarding the addition of an edge $e \in E(\bar{G})$ to a graph G in Section 2. In Section 3, we characterize n - γ_{tR} -edge-critical graphs of order n . We characterize 4- γ_{tR} -edge-critical graphs in Section 4, and, after investigating γ_{tR} -edge-supercritical graphs in Section 5, we present a necessary condition for 5- γ_{tR} -edge-critical graphs in Section 6. In Section 7, we determine the total Roman domination number of spiders and characterize γ_{tR} -edge-critical spiders. As can be expected, every graph G with $\gamma_{tR}(G) = k \geq 4$ is a spanning

subgraph of a $k\text{-}\gamma_{IR}(G)$ -edge-critical graph; a short proof is given in Section 8, where we also show that for any $k \geq 4$, there exists a $k\text{-}\gamma_{IR}$ -edge-critical graph of diameter 2. We conclude in Section 9 with ideas for future research.

2. Adding an edge

We begin with a result from [Van der Merwe, Mynhardt, and Haynes 1998a] which bounds the effect the addition of an edge can have on the total domination number of a graph and show that the same bounds hold with respect to the total Roman domination number.

Proposition 2.1 [Van der Merwe, Mynhardt, and Haynes 1998a]. *For a graph G with no isolated vertices, if $uv \in E(\bar{G})$, then $\gamma_t(G) - 2 \leq \gamma_t(G + uv) \leq \gamma_t(G)$.*

An edge $uv \in E(\bar{G})$ is *critical* if $\gamma_{IR}(G + uv) < \gamma_{IR}(G)$. The following proposition restricts the possible values assigned to the vertices of a critical edge uv by a $\gamma_{IR}(G + uv)$ -function f , which will be useful in proving subsequent results. For a graph G and a vertex $v \in V(G)$, the *open neighbourhood* of v in G is $N_G(v) = \{u \in V(G) : uv \in E(G)\}$, and the *closed neighbourhood* of v in G is $N_G[v] = N_G(v) \cup \{v\}$. When $G \neq K_2$, the unique neighbour of an end-vertex of G is called a *support vertex*.

Proposition 2.2. *Given a graph G with no isolated vertices, if $uv \in E(\bar{G})$ is a critical edge and f is a $\gamma_{IR}(G + uv)$ -function, then*

$$\{f(u), f(v)\} \in \{\{2, 2\}, \{2, 1\}, \{2, 0\}, \{1, 1\}\}.$$

If, in addition, $\deg(u) = \deg(v) = 1$, then there exists a $\gamma_{IR}(G + uv)$ -function f such that $f(u) = f(v) = 1$.

Proof. Let G be a graph with no isolated vertices, $uv \in E(\bar{G})$ such that $\gamma_{IR}(G + uv) < \gamma_{IR}(G)$, and f a $\gamma_{IR}(G + uv)$ -function. Suppose for a contradiction that $\{f(u), f(v)\} \notin \{\{2, 2\}, \{2, 1\}, \{2, 0\}, \{1, 1\}\}$. Then $\{f(u), f(v)\} \in \{\{0, 0\}, \{0, 1\}\}$. Note that, in either case, the edge uv cannot affect whether u and v are dominated or whether, in the case where (say) $f(v) = 1$, v is isolated. Hence f is a TRD-function of G , contradicting $\gamma_{IR}(G + uv) < \gamma_{IR}(G)$. Therefore $\{f(u), f(v)\} \in \{\{2, 2\}, \{2, 1\}, \{2, 0\}, \{1, 1\}\}$.

Now, suppose in addition that $\deg(u) = \deg(v) = 1$, and let f be a $\gamma_{IR}(G + uv)$ -function such that $|V_f^2|$ is as small as possible. Let w and x be the unique neighbours of u and v , respectively, noting that possibly $w = x$. Suppose for a contradiction that $f(u) = 2$ (without loss of generality). If $f(v) = 0$, then $f(w) > 0$, otherwise u would be isolated in $G[V_f^+]$. Thus, regardless of whether $w = x$ or not, consider the function $f' : V(G) \rightarrow \{0, 1, 2\}$ defined by $f'(u) = f'(v) = 1$ and $f'(y) = f(y)$ for all other $y \in V(G)$. Otherwise, if $f(v) \geq 1$, then clearly $f(w) = 0$. Thus,

regardless of whether $w = x$ or not, consider the function $f' : V(G) \rightarrow \{0, 1, 2\}$ defined by $f'(u) = f'(w) = 1$ and $f'(y) = f(y)$ for all other $y \in V(G)$. In either case, f' is a γ_{IR} -function on $G + uv$. However, $|V_{f'}^2| < |V_f^2|$, contradicting $|V_f^2|$ being as small as possible. Hence $f(u) \neq 2$, and thus $f(u) = f(v) = 1$. $\square$

Proposition 2.3. *Given a graph G with no isolated vertices, if $uv \in E(\bar{G})$, then $\gamma_{IR}(G) - 2 \leq \gamma_{IR}(G + uv) \leq \gamma_{IR}(G)$.*

Proof. Let G be a graph with no isolated vertices. Clearly, adding an edge cannot increase the total Roman domination number; hence the upper bound holds. Now, let $uv \in E(\bar{G})$. Note that when $\gamma_{IR}(G + uv) = \gamma_{IR}(G)$, the lower bound clearly holds. So assume $\gamma_{IR}(G + uv) < \gamma_{IR}(G)$ and let f be a $\gamma_{IR}(G + uv)$ -function. By Proposition 2.2, $\{f(u), f(v)\} \in \{\{2, 2\}, \{2, 1\}, \{2, 0\}, \{1, 1\}\}$.

First assume $\{f(u), f(v)\} \in \{\{2, 2\}, \{2, 1\}, \{1, 1\}\}$. Then f is an RD-function of G , and the only possible isolated vertices in $G[V_f^+]$ are u and v . Consider the function $f' : V(G) \rightarrow \{0, 1, 2\}$ defined as follows: If u is isolated in $G[V_f^+]$, choose $u' \in N_G(u)$ and let $f'(u') = 1$. Similarly, if v is isolated in $G[V_f^+]$, choose $v' \in N_G(v)$ and let $f'(v') = 1$. Let $f'(x) = f(x)$ for all other $x \in V(G)$. Now, assume instead that $f(u) = 2$ and $f(v) = 0$ (without loss of generality). Since u is not isolated in $G[V_f^+]$, f is a TRD-function of $G - v$. Consider the function $f' : V(G) \rightarrow \{0, 1, 2\}$ defined as follows: Let $f'(v) = 1$. Then, if v is isolated in $G[V_{f'}^+]$, choose $v' \in N_G(v)$ and let $f'(v') = 1$. Let $f'(x) = f(x)$ for all other $x \in V(G)$. In either case, f' is a TRD-function of G and $\omega(f') \leq \gamma_{IR}(G + uv) + 2$. Thus $\gamma_{IR}(G) \leq \gamma_{IR}(G + uv) + 2$, and hence the lower bound holds. $\square$

3. γ_{IR} -edge-critical graphs with large TRD-numbers

We now investigate the γ_{IR} -edge-critical graphs G which have the largest TRD-number, namely $|V(G)|$. A *subdivided star* is a tree obtained from a star on at least three vertices by subdividing each edge exactly once. A *double star* is a tree obtained from two disjoint nontrivial stars by joining the two central vertices (choosing either central vertex in the case of K_2). The *corona* $\text{cor}(G)$ (sometimes denoted by $G \circ K_1$) of G is obtained by joining each vertex of G to a new end-vertex.

Connected graphs G for which $\gamma_{IR}(G) = |V(G)|$ were characterized in [Ahangar, Henning, Samodivkin, and Yero 2016]. There $\mathcal{G}$ was defined as the family of connected graphs obtained from a 4-cycle v_1, v_2, v_3, v_4, v_1 by adding $k_1 + k_2 \geq 1$ vertex-disjoint paths P_2 , and joining v_i to the end of k_i such paths for $i \in \{1, 2\}$. Note that possibly $k_1 = 0$ or $k_2 = 0$. Furthermore, they defined $\mathcal{H}$ to be the family of graphs obtained from a double star by subdividing each pendant edge once and the nonpendant edge $r \geq 0$ times. For $r \geq 0$, we define $\mathcal{H}_r \subseteq \mathcal{H}$ as the family of graphs in $\mathcal{H}$ where the nonpendant edge was subdivided r times.

Proposition 3.1 [Ahangar, Henning, Samodivkin, and Yero 2016]. *If G is a connected graph of order $n \geq 2$, then $\gamma_{tR}(G) = n$ if and only if one of the following holds:*

- (i) G is a path or a cycle.
- (ii) G is the corona of a graph.
- (iii) G is a subdivided star.
- (iv) $G \in \mathcal{G} \cup \mathcal{H}$.

Using Proposition 3.1, we characterize connected n - γ_{tR} -edge-critical graphs as follows.

Theorem 3.2. *A connected graph G of order $n \geq 4$ is n - γ_{tR} -edge-critical if and only if G is one of the following graphs:*

- (i) C_n , $n \geq 4$.
- (ii) $\text{cor}(K_r)$, $r \geq 3$.
- (iii) a subdivided star of order $n \geq 7$.
- (iv) $G \in \mathcal{G}$.
- (v) $G \in \mathcal{H} - \mathcal{H}_0 - \mathcal{H}_2$.

Proof. Let G be a connected graph of order $n \geq 4$ with $\gamma_{tR}(G) = n$. First, suppose G is any of the graphs listed in (i)–(v) above. Then, for any $e \in E(\bar{G})$, $G + e$ is not one of the graphs listed in Proposition 3.1. Therefore $\gamma_{tR}(G + e) < n$ for all $e \in E(\bar{G})$, and thus G is γ_{tR} -edge-critical.

Otherwise, suppose G is not one of the graphs listed in (i)–(v) above. Note that since $\gamma_{tR}(G) = n$, G is still listed in Proposition 3.1(i)–(iv). If $G \cong P_n : v_1, \dots, v_n$, $n \geq 4$, then $G + v_1 v_n \cong C_n$ and $\gamma_{tR}(G) = \gamma_{tR}(C_n) = n$. If $G \cong \text{cor}(F)$, where F is not a complete graph of order at least 3, then $\gamma_{tR}(G) = \gamma_{tR}(G + uv)$ for any $uv \in E(\bar{F})$. If G is a subdivided star of order less than 7, then $G = P_5$. In each of these cases, G is clearly not γ_{tR} -edge-critical.

Now consider $G \in \mathcal{H}$. Let $w_1, \dots, w_k$ be the leaves of G , $u_1, \dots, u_k$ be their respective support vertices, and $v_1, \dots, v_m$ be the path such that v_1 and v_m are the two support vertices in the original double star S , labelled so that w_1 is adjacent, in S , to v_1 . Note that $m = r + 2$, and therefore $m \geq 2$. If $G \in \mathcal{H}_0$, consider the graph $G + v_2 w_1$, and note that $G + v_2 w_1 \in \mathcal{G}$. Therefore, by Proposition 3.1, $\gamma_{tR}(G + v_2 w_1) = n$, and thus G is not γ_{tR} -edge-critical. Similarly, if $G \in \mathcal{H}_2$, consider the graph $G + v_1 v_4$, and note that $G + v_1 v_4 \in \mathcal{G}$. Therefore, by Proposition 3.1, $\gamma_{tR}(G + v_1 v_4) = n$, and again G is not γ_{tR} -edge-critical. $\square$

4. $4\text{-}\gamma_{tR}$ -edge-critical graphs

Before we characterize the graphs G such that $\gamma_{tR}(G) = 4$ and $\gamma_{tR}(G + e) = 3$ for any $e \in E(\overline{G})$ (that is, the graphs which are $4\text{-}\gamma_{tR}$ -edge-critical), we present the following result from [Pushpam and Padmapriya 2017] which characterizes the graphs with a total Roman domination number of 3, the smallest possible TRD-number. Note that while the authors required that G has girth 3, the result actually holds in general for any graph G on at least three vertices, as we now show. A *universal vertex* of G is a vertex that is adjacent to all other vertices of G .

Proposition 4.1. *For a graph G of order $n \geq 3$ with no isolated vertices, $\gamma_{tR}(G) = 3$ if and only if $\Delta(G) = n - 1$, that is, G has a universal vertex.*

Proof. Suppose $\gamma_{tR}(G) = 3$ and let $f = (V_f^0, V_f^1, V_f^2)$ be a $\gamma_{tR}(G)$ -function. If $V_f^2 = \emptyset$, then $|V_f^1| = 3$, and thus $n = 3$. Since G has no isolated vertices, this implies that $G = K_3$ or P_3 , both of which have a universal vertex. Otherwise, assume $|V_f^2| = 1$ and $|V_f^1| = 1$. Pick $u, v \in V(G)$ so that $f(u) = 1$ and $f(v) = 2$. Since $G[V_f^+]$ has no isolated vertices, $uv \in E(G)$. Furthermore, since $\gamma_{tR}(G) = 3$, $f(x) = 0$ for all other $x \in V(G)$. Therefore $N_G[v] = V(G)$, and thus v is a universal vertex.

Conversely, suppose G has a universal vertex v , and take any $u \in N_G(v)$. Consider the TRD-function $f : V(G) \rightarrow \{0, 1, 2\}$ defined by $f(v) = 2$, $f(u) = 1$, and $f(x) = 0$ for all other $x \in V(G)$. Since G has at least three vertices, $\gamma_{tR}(G) > 2$. Therefore, since $\omega(f) = 3$, we conclude that $\gamma_{tR}(G) = 3$. $\square$

A *galaxy* is defined as the disjoint union of two or more nontrivial stars. The characterization of $4\text{-}\gamma_{tR}$ -edge-critical graphs follows; note that this class of graphs is exactly the class of $2\text{-}\gamma$ -edge-critical graphs, as characterized in [Sumner and Blich 1983].

Theorem 4.2. *A graph G with no isolated vertices is $4\text{-}\gamma_{tR}$ -edge-critical if and only if $\overline{G}$ is a galaxy.*

Proof. Let G be a graph of order n with no isolated vertices. Suppose first that G is $4\text{-}\gamma_{tR}$ -edge-critical. Then for any $e \in E(\overline{G})$, we have $\gamma_{tR}(G + e) = 3$, and thus Proposition 4.1 implies that the addition of any edge to G creates a universal vertex. Therefore, for each edge $uv \in E(\overline{G})$, one of u and v has degree $n - 2$ in G ; that is, one of u and v is a leaf in $\overline{G}$. Since each edge of $\overline{G}$ connects a leaf to either a support vertex or another leaf, the components of $\overline{G}$ are nontrivial stars. Moreover, $\overline{G}$ has at least two components, otherwise G has an isolated vertex.

Conversely, suppose $\overline{G}$ is a galaxy. Since $\overline{G}$ has no isolated vertices, G has no universal vertices, and thus, by Proposition 4.1, $\gamma_{tR}(G) > 3$. Let u and v be vertices in different components of $\overline{G}$, and define $f : V(G) \rightarrow \{0, 1, 2\}$ by $f(u) = f(v) = 2$ and $f(x) = 0$ for all other $x \in V(G)$. Clearly f is a TRD-function on G , and hence

$\gamma_{tR}(G) = 4$. Since the deletion of any edge in $\bar{G}$ produces an isolated vertex, the addition of any edge to G creates a universal vertex. Therefore, by Proposition 4.1, $\gamma_{tR}(G + e) = 3$ for all $e \in E(\bar{G})$, and hence G is 4- γ_{tR} -edge-critical. $\square$

Corollary 4.3. *If G is a connected $(n-2)$ -regular graph, then G is 4- γ_{tR} -edge-critical.*

Having characterized 4- γ_{tR} -edge-critical graphs, our next result demonstrates the existence of stable graphs with total Roman domination number 4.

Proposition 4.4. *If G is an $(n-3)$ -regular graph of order $n \geq 6$, then $\gamma_{tR}(G) = 4$. Moreover, G is stable.*

Proof. We prove that $\gamma(G) = 2$. Since G is $(n-3)$ -regular, its complement $\bar{G}$ is 2-regular. If $\bar{G}$ is disconnected, let u and v be vertices in different components of $\bar{G}$. Otherwise, if $\bar{G}$ is connected, then $\bar{G} \cong C_n$, $n \geq 6$, and thus we can choose $u, v \in V(\bar{G})$ such that $d_{\bar{G}}(u, v) \geq 3$. In either case, $N_{\bar{G}}[u] \cap N_{\bar{G}}[v] = \emptyset$. In G , u dominates all vertices in $G - N_{\bar{G}}(u)$ and v dominates all vertices in $G - N_{\bar{G}}(v)$. Therefore $\{u, v\}$ dominates G , and thus, since G has no universal vertex, $\gamma(G) = 2$.

Now, define $f : V(G) \rightarrow \{0, 1, 2\}$ by $f(u) = f(v) = 2$ and $f(y) = 0$ for all other $y \in V(G)$. Since $uv \in E(G)$, f is a TRD-function on G and $\omega(f) = 4$, so $\gamma_{tR}(G) \leq 4$. Since G has no universal vertex, $\gamma_{tR}(G) > 3$ by Proposition 4.1, and thus $\gamma_{tR}(G) = 4$, as required. Furthermore, since the addition of any edge to G does not create a universal vertex, it follows from Proposition 4.1 that $\gamma_{tR}(G + e) = \gamma_{tR}(G)$ for all $e \in E(\bar{G})$. Therefore G is stable. $\square$

5. γ_{tR} -edge-supercritical graphs

We now consider the graphs G which attain the lower bound in Proposition 2.3 for all $e \in E(\bar{G})$, that is, γ_{tR} -edge-supercritical graphs. An edge $uv \in E(\bar{G})$ is *supercritical* if $\gamma_{tR}(G + uv) = \gamma_{tR}(G) - 2$. Van der Merwe, Mynhardt, and Haynes [1998a] defined a graph G to be γ_t -edge-supercritical if $\gamma_t(G + e) = \gamma_t(G) - 2$ for all $e \in E(\bar{G})$. We begin with their characterization of γ_t -edge-supercritical graphs.

Proposition 5.1 [Van der Merwe, Mynhardt, and Haynes 1998a]. *A graph G is γ_t -edge-supercritical if and only if G is the union of two or more nontrivial complete graphs.*

The proof of the previous result relies on the fact that, if u and v are vertices of a graph G with $d(u, v) = 2$, then $\gamma_t(G) - 1 \leq \gamma_t(G + uv)$. However, the analogous result does not hold with respect to the total Roman domination number, as we now show. Consider the graph $G = \text{cor}(K_3)$. By Proposition 3.1, $\gamma_{tR}(G) = 6$. Consider any two nonadjacent vertices u and v in G such that $\deg(u) = 1$ and $\deg(v) = 3$. Clearly uv is a supercritical edge with $d(u, v) = 2$, and thus $d(u, v) = 2$ does not always imply that $\gamma_{tR}(G) - 1 \leq \gamma_{tR}(G + uv)$.

As a result, the classification of γ_{tR} -edge-supercritical graphs will be less straightforward than that of γ_t -edge-supercritical graphs. However, it is easy to see that there are no $5\text{-}\gamma_{tR}$ -edge-supercritical graphs, where 5 is the smallest possible TRD-number of a γ_{tR} -edge-supercritical graph, and that the disjoint union of two or more complete graphs of order at least 3 is γ_{tR} -edge-supercritical.

Proposition 5.2. (i) *There are no $5\text{-}\gamma_{tR}$ -edge-supercritical graphs.*

(ii) *If G is the disjoint union of $k \geq 2$ complete graphs, each of order at least 3, then G is $3k\text{-}\gamma_{tR}$ -edge-supercritical.*

Proof. (i) Suppose for a contradiction that G is a $5\text{-}\gamma_{tR}$ -edge-supercritical graph. Then $\gamma_{tR}(G + uv) = 3$ for any edge $uv \in E(\overline{G})$. However, as in the proof of Theorem 4.2, this implies that $\overline{G}$ is a galaxy, that is, G is $4\text{-}\gamma_{tR}$ -edge-critical, a contradiction.

(ii) It follows from Proposition 4.1 that $\gamma_{tR}(G) = 3k$. Moreover, joining any two vertices in different components of G results in a graph with TRD-number $3k - 2$. $\square$

6. $5\text{-}\gamma_{tR}$ -edge-critical graphs

We now investigate the graphs which are $5\text{-}\gamma_{tR}$ -edge-critical. We begin with the following results, which bound $\gamma_{tR}(G)$ in terms of $\gamma_t(G)$.

Proposition 6.1 [Ahangar, Henning, Samodivkin, and Yero 2016]. *If G is a graph with no isolated vertices, then $\gamma_t(G) \leq \gamma_{tR}(G) \leq 2\gamma_t(G)$. Furthermore, $\gamma_{tR}(G) = \gamma_t(G)$ if and only if G is the disjoint union of copies of K_2 .*

Note that Amjadi, Nazari-Moghaddam, Sheikholeslami, and Volkmann [2017] characterized the trees which attain the upper bound in Proposition 6.1.

Proposition 6.2 [Ahangar, Henning, Samodivkin, and Yero 2016]. *Let G be a connected graph of order $n \geq 3$. Then $\gamma_{tR}(G) = \gamma_t(G) + 1$ if and only if $\Delta(G) = n - 1$, that is, G has a universal vertex.*

By Proposition 4.1, Proposition 6.2 implies that, if G is a connected graph of order $n \geq 3$, then $\gamma_{tR}(G) = \gamma_t(G) + 1$ if and only if $\gamma_{tR}(G) = 3$. These results lead to the following observation.

Observation 6.3. *If G is a connected graph of order $n \geq 3$ such that $\Delta(G) \leq n - 2$, then $\gamma_t(G) + 2 \leq \gamma_{tR}(G) \leq 2\gamma_t(G)$.*

We now provide a result characterizing graphs with $\gamma_{tR} \in \{3, 4\}$ in terms of their domination and total domination numbers that will be useful in describing $5\text{-}\gamma_{tR}$ -edge-critical graphs.

Proposition 6.4. *If G is a connected graph of order $n \geq 3$, then $\gamma_{tR}(G) \in \{3, 4\}$ if and only if $\gamma_t(G) = 2$. Moreover, $\gamma(G) = 1$ when $\gamma_{tR}(G) = 3$, and $\gamma(G) = 2$ when $\gamma_{tR}(G) = 4$.*

Proof. Suppose first that $\gamma_t(G) = 2$. By Proposition 6.1, $2 \leq \gamma_{tR}(G) \leq 4$. Clearly $\gamma_{tR}(G) \neq 2$, since $n \geq 3$. Therefore $\gamma_{tR}(G) \in \{3, 4\}$.

Conversely, suppose $\gamma_{tR}(G) \in \{3, 4\}$. First, if $\gamma_{tR}(G) = 3$, then Proposition 4.1 implies that G has a universal vertex. Therefore $\gamma_t(G) = 2$ and $\gamma(G) = 1$. Otherwise, if $\gamma_{tR}(G) = 4$, then Proposition 4.1 implies that G has no universal vertex. Therefore, by Observation 6.3, $\gamma_t(G) + 2 \leq 4$, and thus $\gamma_t(G) = 2$. Furthermore, since $\gamma(G) \leq \gamma_t(G)$ and G has no universal vertex, $\gamma(G) = 2$. $\square$

Proposition 6.5. *For any graph G , if G is $5\text{-}\gamma_{tR}$ -edge-critical, then G is either $3\text{-}\gamma_t$ -edge-critical or $G = K_2 \cup K_n$ for $n \geq 3$, in which case G is $4\text{-}\gamma_t$ -edge-supercritical.*

Proof. Suppose G is $5\text{-}\gamma_{tR}$ -edge-critical. By Proposition 6.4, $\gamma_t(G) > 2$ and $\gamma_t(G + e) = 2$ for any $e \in E(\bar{G})$. Therefore, by Proposition 2.1, G is either $3\text{-}\gamma_t$ -edge-critical or $4\text{-}\gamma_t$ -edge-supercritical. If G is $4\text{-}\gamma_t$ -edge-supercritical, then by Proposition 5.1, G is the union of two or more nontrivial complete graphs. Since $\gamma_{tR}(G) = 5$, this implies that $G = K_2 \cup K_n$ for $n \geq 3$. $\square$

Note that if we add the condition that G is not $6\text{-}\gamma_{tR}$ -edge-supercritical, then the above becomes a necessary and sufficient condition. Clearly $G = K_2 \cup K_n$ is $5\text{-}\gamma_{tR}$ -edge-critical for any $n \geq 3$. Otherwise, if G is $3\text{-}\gamma_t$ -edge-critical, then by Proposition 6.4, $\gamma_{tR}(G) > 4$ and $\gamma_{tR}(G + e) \in \{3, 4\}$ for any $e \in E(\bar{G})$. By Proposition 6.1, $\gamma_{tR}(G) \leq 6$, and thus, since G is not $6\text{-}\gamma_{tR}$ -edge-supercritical, $\gamma_{tR}(G) = 5$. Hence G is $5\text{-}\gamma_{tR}$ -edge-critical, as required.

7. γ_{tR} -edge-critical spiders

A (generalized) spider $\text{Sp}(l_1, \dots, l_k)$, $l_i \geq 1$, $k \geq 2$, is a tree obtained from the star $K_{1,k}$ with centre u and leaves $v_1, \dots, v_k$ by subdividing the edge uv_i exactly $l_i - 1$ times, $i = 1, \dots, k$. Thus, a spider $\text{Sp}(2, \dots, 2)$ is a subdivided star. The $u - v_i$ paths (of length l_i) are called the *legs* of the spider, while u is its *head*. We now investigate the spiders which are γ_{tR} -edge-critical. Note that when $k = 2$, $\text{Sp}(l_1, \dots, l_k) \cong P_n$ for $n \geq 3$, which, by Theorem 3.2, is not γ_{tR} -edge-critical. We begin with two propositions restricting γ_{tR} -edge-criticality in general graphs, which will be useful in classifying γ_{tR} -edge-critical spiders.

Proposition 7.1. *For a graph G with no isolated vertices, if G has an end-vertex w with support vertex x such that $G[N(x) - \{w\}]$ is not complete, then G is not γ_{tR} -edge-critical.*

Proof. Suppose $u, v \in N_G(x) - \{w\}$ such that $uv \in E(\bar{G})$. We claim $\gamma_{tR}(G + uv) = \gamma_{tR}(G)$. Suppose for a contradiction that $\gamma_{tR}(G + uv) < \gamma_{tR}(G)$, and consider a γ_{tR} -function $f = (V_f^0, V_f^1, V_f^2)$ on $G + uv$. Note that, since w is an end-vertex, $f(x) > 0$. By Proposition 2.2, $\{f(u), f(v)\} \in \{\{2, 2\}, \{2, 1\}, \{2, 0\}, \{1, 1\}\}$. Since $ux, vx \in E(G)$ and at least one of $f(u)$ and $f(v)$ is positive, we can assume

without loss of generality that $f(x) = 2$. In any case, f is also a TRD-function on G , contradicting $\gamma_{tR}(G + uv) < \gamma_{tR}(G)$. Therefore $\gamma_{tR}(G + uv) = \gamma_{tR}(G)$ and G is not γ_{tR} -edge-critical. $\square$

In a tree, the support vertex of a leaf is called a *stem*. A stem is called *weak* if it is adjacent to exactly one leaf, and *strong* if it is adjacent to two or more leaves. A vertex b of a tree such that $\deg(b) \geq 3$ is called a *branch vertex*. An *endpath* in a tree is a path from a branch vertex to a leaf, where all of the internal vertices of the path have degree 2. The next result follows immediately from Proposition 7.1.

Corollary 7.2. *If T is a γ_{tR} -edge-critical tree, then T contains no stems of degree at least 3, and hence no strong stems.*

Proposition 7.3. *For a graph G with no isolated vertices, if G has two endpaths $v_0, v_1, \dots, v_k$ and $u_0, u_1, \dots, u_m$, where $k, m \geq 3$ and v_k and u_m are leaves, then G is not γ_{tR} -edge-critical.*

Proof. We claim that $\gamma_{tR}(G + v_k u_m) = \gamma_{tR}(G)$. Suppose for a contradiction that $\gamma_{tR}(G + v_k u_m) < \gamma_{tR}(G)$, and let f be a γ_{tR} -function on $G + v_k u_m$. Then, by Proposition 2.2, we may assume $f(u_m) = f(v_k) = 1$. Define $f' : V(G) \rightarrow \{0, 1, 2\}$ as follows: If $f(v_{k-1}) = 0$, then clearly $f(v_{k-2}) = 2$ and $f(v_{k-3}) \geq 1$, so let $f'(v_{k-1}) = f'(v_{k-2}) = 1$. Otherwise, let $f'(v_{k-1}) = f(v_{k-1})$ and $f'(v_{k-2}) = f(v_{k-2})$. Similarly, if $f(u_{m-1}) = 0$, then clearly $f(u_{m-2}) = 2$ and $f(u_{m-3}) \geq 1$, so let $f'(u_{m-1}) = f'(u_{m-2}) = 1$. Otherwise, let $f'(u_{m-1}) = f(u_{m-1})$ and $f'(u_{m-2}) = f(u_{m-2})$. Finally, let $f'(y) = f(y)$ for all other $y \in V(G)$. Clearly f' is a TRD-function on G and $\omega(f') = \omega(f)$, contradicting $\gamma_{tR}(G + v_k u_m) < \gamma_{tR}(G)$. Therefore $\gamma_{tR}(G + v_k u_m) = \gamma_{tR}(G)$, and thus G is not γ_{tR} -edge-critical. $\square$

Let S be a spider with $k \geq 3$ legs. In what follows, let c be the head of S , and let the k legs be labelled $c, v_{i1}, v_{i2}, \dots, v_{im_i}$, where $i \in \{1, 2, \dots, k\}$, in order of increasing length. Let $m = m_k$; that is, m is the length of a longest leg of S . We begin by determining the TRD-number of spiders.

Proposition 7.4. *If S is a spider of order n with $k \geq 3$ legs such that S has y legs of length 2, then*

$$\gamma_{tR}(S) = \begin{cases} n & \text{if } y \geq k - 1, \\ n - k + y + 1 & \text{if } 1 \leq y < k - 1, \\ n - k + 2 & \text{if } y = 0. \end{cases}$$

Proof. Suppose S has x legs of length 1, and consider a γ_{tR} -function f on S such that $|V_f^2|$ is as small as possible. First, suppose $y \geq k - 1$. If $y = k$, then S is a subdivided star. Otherwise, if $y = k - 1$, then S has exactly one leg that is not of length 2, and thus either $x = 1$ or $x = 0$. If $x = 1$, then S is the corona of a graph

(specifically, $S = \text{cor}(K_{1,k-1})$). Otherwise, if $x = 0$, then $m = m_k \geq 3$, and $S \in \mathcal{H}_r$, where $r = m - 3$. In any case, by Proposition 3.1, $\gamma_{tR}(S) = n$.

Assume therefore that $y < k - 1$. Then S has at least two legs that are not of length 2. Therefore S is not one of the graphs listed in Proposition 3.1, and thus $\gamma_{tR}(S) < n$. Hence there is some vertex $u \in V(S)$ such that $f(u) = 2$ and $f(w) = 0$ for at least two vertices w adjacent to u . Furthermore, since f is a TRD-function, such a vertex u is not isolated in $S[V_f^+]$, and thus $\deg(u) \geq 3$. Since c is the only vertex in S with degree at least 3, $f(c) = 2$. Therefore c Roman dominates each v_{i1} , and we need $f(v_{i1})$ to be positive for at least one i to ensure that $S[V_f^+]$ has no isolated vertices.

Consider an arbitrary leg $c, v_{i1}, v_{i2}, \dots, v_{im_i}$ of S . If $m_i = 1$, then $f(v_{i1}) \in \{0, 1\}$ in order for f to totally Roman dominate c and v_{i1} . If $m_i = 2$, a total weight of 2 on v_{i1} and v_{i2} is required in order for f to total Roman dominate $\{v_{i1}, v_{i2}\}$. Since $|V_f^2|$ is as small as possible, $f(v_{i1}) = f(v_{i2}) = 1$. Finally, if $m_i > 2$, by Proposition 3.1 and since $f(c) = 2$, a total weight of at least $m_i - 1$ on $v_{i1}, \dots, v_{im_i}$ is required in order for f to totally Roman dominate c and $\{v_{i1}, \dots, v_{im_i}\}$. Moreover, by the choice of f , $f(v_{i1}) \in \{0, 1\}$ and $f(v_{i2}) = \dots = f(v_{im_i}) = 1$. Therefore $\omega(f) \geq n - k + y + 1$.

Now, if $y > 0$, where (say) $m_j = 2$, then $f(v_{j1}) = 1$. By minimality and since c is adjacent to v_{j1} , $f(v_{i1}) = 0$ for each i such that $m_i \neq 2$. Then $\gamma_{tR}(S) = \omega(f) = n - k + y + 1$, as required. Otherwise, if $y = 0$, then $f(v_{i1}) = 1$ for some i to ensure that c is not isolated in $S[V_f^+]$. By minimality, $f(v_{j1}) = 0$ for each $j \neq i$. Therefore $\gamma_{tR}(S) = \omega(f) = n - k + 2$. $\square$

The characterization of γ_{tR} -edge-critical spiders follows. Our result also shows that a spider of order n is γ_{tR} -edge-critical if and only if it is n - γ_{tR} -edge-critical.

Theorem 7.5. *A spider $S = \text{Sp}(l_1, \dots, l_k)$, $k \geq 3$, is γ_{tR} -edge-critical if and only if $l_i = 2$ for each i , $1 \leq i \leq k - 1$, and $l_k \in \{2, m\}$, where $m = 4$ or $m \geq 6$.*

Proof. Suppose S has order n . If $l_i = 2$ for each $i = 1, \dots, k$, then S is a subdivided star and, by Theorem 3.2, S is n - γ_{tR} -edge-critical. Now, suppose S has exactly one leg of length $m \neq 2$. If $m = 1$, then by Proposition 7.1, S is not γ_{tR} -edge-critical. If $m = 3$ or $m = 5$, then $S \in \mathcal{H}_r$ with $r = 0$ or 2 , respectively, and thus, by Theorem 3.2, S is not γ_{tR} -edge-critical. If $m = 4$ or $m \geq 6$, then $S \in \mathcal{H}_r$ with $r = m - 3$, and therefore, by Theorem 3.2, S is n - γ_{tR} -edge-critical. Finally, suppose S has at least two legs that are not of length 2. Again, by Proposition 7.1, if S has a leg of length 1, S is not γ_{tR} -edge-critical. Otherwise, assume S has at least two legs of length at least 3. Then, by Proposition 7.3, S is not γ_{tR} -edge-critical. $\square$

8. k - γ_{tR} -edge-critical graphs with minimum diameter

We now consider the minimum diameter possible in a k - γ_{tR} -edge-critical graph for $k \geq 4$. There are no γ_{tR} -edge-critical graphs with diameter 1, as the only graphs with

diameter 1 are nontrivial complete graphs, which are clearly not γ_{tR} -edge-critical since $E(\overline{G}) = \emptyset$. Therefore, the minimum possible diameter for a γ_{tR} -edge-critical graph is 2. Asplund, Loizeaux and Van der Merwe [2018] constructed families of 3- γ_t -edge-critical graphs with diameter 2. We will show that, for any $k \geq 4$, there exists a k - γ_{tR} -edge-critical graph of diameter 2. We first present the following proposition which shows that every graph G without a dominating vertex is a spanning subgraph of a $\gamma_{tR}(G)$ -edge-critical graph with the same total Roman domination number, which will be useful in proving our result.

Proposition 8.1. *For a graph G with no isolated vertices, if $\gamma_{tR}(G) = k \geq 4$, then G is a spanning subgraph of a k - $\gamma_{tR}(G)$ -edge-critical graph.*

Proof. Suppose $\gamma_{tR}(G) = k \geq 4$. If G is k - $\gamma_{tR}(G)$ -edge-critical, then we are done. Otherwise, there is, by definition, some edge $e_1 \in E(\overline{G})$ such that $\gamma_{tR}(G + e_1) = \gamma_{tR}(G)$. Let $G_1 = G + e_1$. If G_1 is k - $\gamma_{tR}(G)$ -edge-critical, then we are done. Otherwise, there is some edge $e_2 \in E(\overline{G}_1)$ such that $\gamma_{tR}(G_1 + e_2) = \gamma_{tR}(G_1)$. Let $G_2 = G_1 + e_2$. Continuing in this way, we eventually obtain a graph G_i such that for all $e \in E(\overline{G}_i)$, $\gamma_{tR}(G_i + e) < \gamma_{tR}(G_i)$ and $\gamma_{tR}(G_i) = \gamma_{tR}(G_{i-1}) = \cdots = \gamma_{tR}(G_1) = \gamma_{tR}(G)$. Since $k \geq 4$, G_i is not complete and thus $E(G_i) \neq \emptyset$. Therefore, G_i is a k - $\gamma_{tR}(G)$ -edge-critical graph, of which G is a spanning subgraph. $\square$

Before demonstrating the existence of k - γ_{tR} -edge-critical graphs of diameter 2 for any $k \geq 4$, we determine the TRD-number of $K_n \square K_m$, where $n, m \geq 2$. Consider the vertices of $K_n \square K_m$ as an $n \times m$ matrix, where vertices v_{ij} and v_{st} are adjacent if and only if $i = s$ or $j = t$. The rows and columns of the matrix form disjoint copies of K_m and K_n , respectively. If v_{ij} and v_{st} are nonadjacent, then v_{sj} is adjacent to both v_{ij} and v_{st} , and hence $\text{diam}(K_n \square K_m) = 2$.

Proposition 8.2. *If m and n are integers such that $m \geq n \geq 2$, then $\gamma_{tR}(K_n \square K_m) = 2n$.*

Proof. Let $G = K_n \square K_m$. To see that $\gamma_{tR}(G) \leq 2n$, consider the TRD-function $g = (V_g^0, V_g^1, V_g^2)$ on G where $V_g^1 = \emptyset$ and $V_g^2 = \{v_{i1} : 1 \leq i \leq n\}$.

Now, suppose for a contradiction that $\gamma_{tR}(G) \leq 2n - 1$ and consider a TRD-function $f = (V_f^0, V_f^1, V_f^2)$ on G with $\omega(f) = 2n - 1$. Each vertex v dominates one row and one column of G , so if $|V_f^2| = x$ (note that $x \leq n - 1$), then at most x rows and at most x columns are dominated by elements of V_f^2 . Let S be the set of vertices undominated by V_f^2 . Then $|S| \geq (n - x)(m - x) \geq (n - x)^2$. Moreover, $|V_f^1| = (2n - 1) - 2x$ since $\omega(f) = 2n - 1$ and $|V_f^2| = x$.

If $x = n - 1$, then $|V_f^1| = 1$. Since f is a TRD-function and $|S| \geq (n - x)^2$, we have $|S| = 1$; say $S = \{w\}$. Hence $V_f^1 = \{w\}$. However, V_f^2 does not dominate w , and thus w is isolated in $G[V_f^+]$, which is a contradiction. Therefore, there is no TRD-function on G with weight $2n - 1$ when $x = n - 1$.

Otherwise, if $x < n - 1$, then

$$\begin{aligned} |S| - |V_f^1| &\geq (n - x)^2 - (2n - 1 - 2x) \\ &= x^2 - 2(n - 1)x + (n - 1)^2 \\ &= (n - 1 - x)^2 > 0. \end{aligned}$$

Therefore, the number of vertices undominated by V_f^2 is greater than $|V_f^1|$, contradicting f being a TRD-function. Thus there is no TRD-function on G with weight $2n - 1$ when $x < n - 1$. We conclude that $\gamma_{tR}(G) = 2n$. $\square$

Theorem 8.3. *If $k \geq 4$, then there exists a k - γ_{tR} -edge-critical graph of diameter 2.*

Proof. First, assume that k is even; say $k = 2l$ for some $l \geq 2$. Let $G_l = K_l \square K_l$. By Proposition 8.2, $\gamma_{tR}(G_l) = 2l$, and, by Proposition 8.1, G_l is a spanning subgraph of a k - γ_{tR} -edge-critical graph G'_l . Since $k > 3$, Proposition 4.1 implies that G'_l has no dominating vertex, and hence $2 \leq \text{diam}(G'_l) \leq \text{diam}(G_l) = 2$.

Now, consider the case where k is odd; say $k = 2l + 1$ for some $l \geq 2$. Let G_l^d be the graph formed by taking $K_{l+1} \square K_{l+1}$ and deleting the vertices in the set $\{v_{j1} : \lfloor \frac{l}{2} \rfloor + 2 \leq j \leq l + 1\}$. Similarly to G_l , $\text{diam}(G_l^d) = 2$. See Figure 1.

We claim that $\gamma_{tR}(G_l^d) = 2l + 1$. To see that $\gamma_{tR}(G_l^d) \leq 2l + 1$, consider the following TRD-function on G_l^d : If l is even, place two 2's in each of the first $\frac{l}{2} - 1$ rows, and one 2 in each of rows $\frac{l}{2}$ and $\frac{l}{2} + 1$, such that they span columns 2 through $l + 1$. At this point, every vertex in G_l^d is dominated. However, the 2's in rows $\frac{l}{2}$ and $\frac{l}{2} + 1$ are isolated, so place a 1 in row $\frac{l}{2}$ such that it shares a column with the 2 in row $\frac{l}{2} + 1$. Otherwise, if l is odd, place two 2's in each of the first $\frac{l-1}{2}$ rows, and one 2 in row $\frac{l+1}{2}$, such that they span columns 2 through $l + 1$. Similarly to the even case, every vertex in G_l^d is now dominated. However, the 2 in row $\frac{l+1}{2}$ is isolated, so place a 1 in row $\frac{l-1}{2}$ such that it shares a column with that 2. In either case, we have a TRD-function on G_l^d with weight $2l + 1$; hence $\gamma_{tR}(G_l^d) \leq 2l + 1$.

Now, suppose for a contradiction that $\gamma_{tR}(G_l^d) < 2l + 1$, and consider a TRD-function $f = (V_f^0, V_f^1, V_f^2)$ on G_l^d with $\omega(f) = 2l$. We claim that $f(v_{j1}) = 0$ for all $1 \leq j \leq \lfloor \frac{l}{2} \rfloor + 1$. If $f(v_{j1}) = 2$ for $x \geq 1$ vertices in column 1, the undominated vertices in columns 2 through $l + 1$ form the graph $K_l \square K_{l+1-x}$. By Proposition 8.2, a TRD-function on $K_l \square K_{l+1-x}$ requires a weight of $2 \min\{l, l + 1 - x\} = 2(l + 1 - x)$. However, since $2x + 2(l + 1 - x) > 2l$, this is impossible. Therefore $f(v_{j1}) \neq 2$ for all $1 \leq j \leq \lfloor \frac{l}{2} \rfloor + 1$. If $f(v_{j1}) = 1$ for $x \geq 1$ vertices in column 1, the undominated vertices in columns 2 through $l + 1$ (that is, those for which f could be assigned a 2) form the graph $K_l \square K_{l+1}$. Again by Proposition 8.2, a TRD-function on $K_l \square K_{l+1}$ requires a weight of $2 \min\{l, l + 1\} = 2l$. However, $x + 2l > 2l$ for $x \geq 1$, so this is also not possible. Therefore, $f(v_{j1}) = 0$ for all $1 \leq j \leq \lfloor \frac{l}{2} \rfloor + 1$.

As a result, in order to totally Roman dominate the first column, there must be a 2 in each of the first $\lfloor \frac{l}{2} \rfloor + 1$ rows, none of which can be in the first column. That




Figure 1. The graphs G_3 and G_3^d with a γ_{tR} -function.

is, for each $1 \leq s \leq \lfloor \frac{l}{2} \rfloor + 1$, $f(v_{st}) = 2$ for some $2 \leq t \leq l + 1$. Let S be the set of these vertices. Note that, thus far, we have accounted for a total weight of

$$2(\lfloor \frac{l}{2} \rfloor + 1) = \begin{cases} l + 2 & \text{if } l \text{ is even,} \\ l + 1 & \text{if } l \text{ is odd,} \end{cases}$$

which leaves a weight of $l - 2$ if l is even and $l - 1$ if l is odd to be assigned. That is, a weight of $2(\lceil \frac{l}{2} \rceil - 1)$ remains to be accounted for. We now claim that no two vertices in S can be in the same column. If the vertices in S span fewer than $\lfloor \frac{l}{2} \rfloor + 1$ columns, then the vertices which are undominated by S induce a graph containing $K_{\lceil l/2 \rceil} \square K_{\lceil l/2 \rceil}$ as subgraph. If $l = 2$, then no weight remains to dominate this vertex, as $2(\lceil \frac{l}{2} \rceil - 1) = 0$. Otherwise, if $l > 2$, Proposition 8.2 implies that $\gamma_{tR}(K_{\lceil l/2 \rceil} \square K_{\lceil l/2 \rceil}) = 2(\lceil \frac{l}{2} \rceil)$. However, $2(\lceil \frac{l}{2} \rceil) > 2(\lceil \frac{l}{2} \rceil - 1)$. In either case, this contradicts f being a TRD-function, and thus no vertices of S share a column.

Therefore, the vertices left undominated by S induce a graph $T \cong K_{\lceil l/2 \rceil} \square K_{\lceil l/2 \rceil - 1}$, with $\lceil \frac{l}{2} \rceil$ rows and $\lceil \frac{l}{2} \rceil - 1$ columns. Moreover, the vertices in S are all isolated, as none share a row or column. By Proposition 8.2, $\gamma_{tR}(T) = 2(\lceil \frac{l}{2} \rceil - 1)$. Thus the entire remaining weight is required in order to dominate T ; necessarily, the vertices in $V_f^+ - S$ belong to rows and columns that do not contain vertices in S . However, this still leaves the vertices in S isolated, which contradicts f being a TRD-function on G_l^d . Therefore $\gamma_{tR}(G_l^d) \geq 2l + 1$ and we conclude that $\gamma_{tR}(G_l^d) = 2l + 1$. As in the case where k is even, G_l^d is a spanning subgraph of a k - γ_{tR} -edge-critical graph with diameter 2. $\square$

9. Future work

We showed in Section 5 that the disjoint union of two or more complete graphs, each having order at least 3, is γ_{tR} -edge-supercritical. We also explained that a proof similar to that of Proposition 5.1 does not work for total Roman domination. Hence we pose the following question.

Question 1. *Are the disjoint unions of two or more complete graphs, each having order at least 3, the only γ_{tR} -edge-supercritical graphs?*

Note that if this is the case, Proposition 6.5 automatically becomes a necessary and sufficient condition for a graph to be 5- γ_{tR} -edge-critical.

Now consider, for a moment, Roman dominating functions, and suppose a graph G has nonadjacent vertices u and v such that $f(u) = f(v) = 0$ for every γ_R -function f on G . We claim that $\gamma_R(G + uv) = \gamma_R(G)$. Suppose $\gamma_R(G + uv) < \gamma_R(G)$ and let f be a γ_R -function on $G + uv$. Similar to Proposition 2.2, we may assume without loss of generality that $f(u) = 2$ and $f(v) = 0$, otherwise f is an RD-function on G such that $\omega(f) < \gamma_R(G)$. However, the function f' defined by $f'(v) = 1$ and $f'(y) = f(y)$ for all other $y \in V(G)$ is a γ_R -function on G such that $f'(v) > 0$, contrary to our assumption. The situation for total Roman domination is different.

For a graph G , we define $u \in V(G)$ to be a *dead vertex* if every γ_{tR} -function f on G has $f(u) = 0$. Not only do there exist graphs G containing nonadjacent dead vertices u and v such that $\gamma_{tR}(G + uv) < \gamma_{tR}(G)$, but it is possible to find such a graph G with $\gamma_{tR}(G + uw) < \gamma_{tR}(G)$ for every edge $uw \in E(\bar{G})$; that is, every edge in $E(\bar{G})$ incident with the dead vertex u is critical. We define the graph D_n below and show that D_n is such a graph.

Let D_n be the graph composed of $n \geq 2$ copies of $K_4 - e$ sharing a single central vertex as follows: let c be the central vertex, $w_1, \dots, w_n$ be the degree-2 vertices, and $u_1, \dots, u_n$ and $v_1, \dots, v_n$ be the remaining vertices (where u_i and v_i are adjacent for each i) such that c, u_i, w_i, v_i, c is a 4-cycle in D_n for each $1 \leq i \leq n$. See Figure 2.

Proposition 9.1. *If $n \geq 2$, then $\gamma_{tR}(D_n) = 2n + 1$. Moreover, w_i is a dead vertex for each $1 \leq i \leq n$.*

Proof. To see that $\gamma_{tR}(D_n) \leq 2n + 1$, consider the TRD-function $g : V(D_n) \rightarrow \{0, 1, 2\}$ on D_n defined by $g(c) = 1$, $g(u_i) = 2$ for $1 \leq i \leq n$, and $g(y) = 0$ for all other $y \in V(D_n)$.

We claim that, if f is a TRD-function on D_n with $\omega(f) \leq 2n + 1$, then $f(c) = 1$. If $f(c) = 2$, then the only vertices that remain undominated in D_n are w_i for $1 \leq i \leq n$. However, since $d(w_i, w_j) = 4$ for all $i \neq j$, a weight of $2n$ is required in order to totally Roman dominate these vertices, contradicting $\omega(f) \leq 2n + 1$. If $f(c) = 0$, then since $D_n - c$ is the disjoint union of n triangles, Proposition 3.1 implies that a weight of $3n$ is required to totally Roman dominate the remaining vertices, contradicting $\omega(f) \leq 2n + 1$. Therefore $f(c) = 1$. Since a weight of at least $2n$ is required to totally Roman dominate the remaining disjoint union of n triangles, we conclude that $\gamma_{tR}(D_n) = 2n + 1$.

Now, let f be any γ_{tR} -function on D_n . Then $\omega(f) = 2n + 1$ and $f(c) = 1$. To dominate each triangle of $D_n - c$ with a weight of 2, $\{f(u_i), f(v_i)\} = \{0, 2\}$ and $f(w_i) = 0$ for each $1 \leq i \leq n$. Hence each w_i is a dead vertex. $\square$




Figure 2. The graphs D_3 and D_4 .

The following result shows that, for $n \geq 3$, every edge in $E(\bar{D}_n)$ incident with w_i is critical.

Proposition 9.2. *If $n \geq 3$, $i \in \{1, \dots, n\}$, and $w_i v \in E(\bar{D}_n)$, then $\gamma_{tR}(D_n + w_i v) < \gamma_{tR}(D_n)$.*

Proof. Without loss of generality, consider an edge $w_1 v \in E(\bar{D}_n)$. Then (without loss of generality) $v \in \{w_2, u_2, c\}$. If $v = w_2$, define $f : V(D_n + w_1 v) \rightarrow \{0, 1, 2\}$ by $f(w_1) = f(w_2) = 1$, $f(c) = f(u_3) = \dots = f(u_n) = 2$, and $f(y) = 0$ for all other $y \in V(D_n)$. Otherwise, if $v \in \{u_2, c\}$, define $f : V(D_n + w_1 v) \rightarrow \{0, 1, 2\}$ by $f(c) = f(u_2) = f(u_3) = \dots = f(u_n) = 2$ and $f(y) = 0$ for all other $y \in V(D_n)$. In either case, f is a TRD-function on $D_n + w_1 v$ and $\omega(f) = 2n$. Therefore, by Proposition 9.1, every edge $w_i v \in E(\bar{D}_n)$ is critical. $\square$

However, for $n \geq 3$, the graph D_n is not γ_{tR} -edge-critical since (for example) $\gamma_{tR}(D_n + u_1 u_2) = 2n + 1$. Furthermore, the graph D_2 is not γ_{tR} -edge-critical since (for example) $\gamma_{tR}(D_2 + w_1 w_2) = 5$. However, adding edges to D_n until a $(2n+1)$ - γ_{tR} -edge-critical graph D'_n is obtained results in D'_n having no dead vertices. Hence we pose the following question.

Question 2. *Do there exist γ_{tR} -edge-critical graphs containing dead vertices?*

We characterized γ_{tR} -edge-critical spiders in Theorem 7.5. Finding other classes of γ_{tR} -edge-critical trees and, indeed, characterizing γ_{tR} -edge-critical trees, remain open problems.

References

- [Ahangar, Henning, Samodivkin, and Yero 2016] H. A. Ahangar, M. A. Henning, V. Samodivkin, and I. G. Yero, “Total Roman domination in graphs”, *Appl. Anal. Discrete Math.* **10**:2 (2016), 501–517. MR Zbl

- [Amjadi, Nazari-Moghaddam, Sheikholeslami, and Volkmann 2017] J. Amjadi, S. Nazari-Moghaddam, S. M. Sheikholeslami, and L. Volkmann, “Total Roman domination number of trees”, *Australas. J. Combin.* **69** (2017), 271–285. MR Zbl
- [Amjadi, Sheikholeslami, and Soroudi 2018] J. Amjadi, S. M. Sheikholeslami, and M. Soroudi, “Nordhaus–Gaddum bounds for total Roman domination”, *J. Comb. Optim.* **35**:1 (2018), 126–133. MR Zbl
- [Amjadi, Sheikholeslami, and Soroudi 2019] J. Amjadi, S. M. Sheikholeslami, and M. Soroudi, “On the total Roman domination in trees”, *Discuss. Math. Graph Theory* **39**:2 (2019), 519–532. MR Zbl
- [Asplund, Loizeaux, and Van der Merwe 2018] J. Asplund, M. Loizeaux, and L. van der Merwe, “A note on total domination edge critical graphs”, *J. Combin. Math. Combin. Comput.* **104** (2018), 245–260. MR Zbl
- [Campanelli and Kuziak 2019] N. Campanelli and D. Kuziak, “Total Roman domination in the lexicographic product of graphs”, *Discrete Appl. Math.* **263** (2019), 88–95. MR Zbl
- [Chartrand and Lesniak 2016] G. Chartrand and L. Lesniak, *Graphs & digraphs*, 6th ed., Chapman & Hall/CRC, Boca Raton, FL, 2016.
- [Haynes, Hedetniemi, and Slater 1998] T. W. Haynes, S. T. Hedetniemi, and P. J. Slater, *Fundamentals of domination in graphs*, Monographs and Textbooks in Pure and Applied Mathematics **208**, Marcel Dekker, New York, 1998. MR Zbl
- [Pushpam and Padmapricea 2017] P. R. L. Pushpam and S. Padmapricea, “On total Roman domination in graphs”, pp. 326–331 in *Theoretical computer science and discrete mathematics*, edited by S. Arumugam et al., Lecture Notes in Comput. Sci. **10398**, Springer, 2017. MR Zbl
- [Sumner and Blitch 1983] D. P. Sumner and P. Blitch, “Domination critical graphs”, *J. Combin. Theory Ser. B* **34**:1 (1983), 65–76. MR Zbl
- [Van der Merwe, Mynhardt, and Haynes 1998a] L. C. van der Merwe, C. M. Mynhardt, and T. W. Haynes, “Criticality index of total domination”, pp. 67–73 in *Proceedings of the Twenty-ninth Southeastern International Conference on Combinatorics, Graph Theory and Computing* (Boca Raton, FL, 1998), Congressus Numerantium **131**, Utilitas Math., Winnipeg, MB, 1998. MR Zbl
- [Van der Merwe, Mynhardt, and Haynes 1998b] L. C. van der Merwe, C. M. Mynhardt, and T. W. Haynes, “Total domination edge critical graphs”, *Util. Math.* **54** (1998), 229–240. MR Zbl

Received: 2019-07-23 Accepted: 2019-09-26

chloe.lampman3@gmail.com Department of Mathematics and Statistics,
University of Victoria, Victoria, BC, Canada

kieka@uvic.ca Department of Mathematics and Statistics,
University of Victoria, Victoria, BC, Canada

sogden@uvic.ca Department of Mathematics and Statistics,
University of Victoria, Victoria, BC, Canada

involve

msp.org/involve

INVOLVE YOUR STUDENTS IN RESEARCH

Involve showcases and encourages high-quality mathematical research involving students from all academic levels. The editorial board consists of mathematical scientists committed to nurturing student participation in research. Bridging the gap between the extremes of purely undergraduate research journals and mainstream research journals, *Involve* provides a venue to mathematicians wishing to encourage the creative involvement of students.

MANAGING EDITOR

Kenneth S. Berenhaut Wake Forest University, USA

BOARD OF EDITORS

Colin Adams	Williams College, USA	Robert B. Lund	Clemson University, USA
Arthur T. Benjamin	Harvey Mudd College, USA	Gaven J. Martin	Massey University, New Zealand
Martin Bohner	Missouri U of Science and Technology, USA	Mary Meyer	Colorado State University, USA
Amarjit S. Budhiraja	U of N Carolina, Chapel Hill, USA	Frank Morgan	Williams College, USA
Pietro Cerone	La Trobe University, Australia	Mohammad Sal Moslehian	Ferdowsi University of Mashhad, Iran
Scott Chapman	Sam Houston State University, USA	Zuhair Nashed	University of Central Florida, USA
Joshua N. Cooper	University of South Carolina, USA	Ken Ono	Univ. of Virginia, Charlottesville
Jem N. Corcoran	University of Colorado, USA	Yuval Peres	Microsoft Research, USA
Toka Diagana	Howard University, USA	Y.-F. S. Pétermann	Université de Genève, Switzerland
Michael Dorff	Brigham Young University, USA	Jonathon Peterson	Purdue University, USA
Sever S. Dragomir	Victoria University, Australia	Robert J. Plemmons	Wake Forest University, USA
Joel Foisy	SUNY Potsdam, USA	Carl B. Pomerance	Dartmouth College, USA
Errin W. Fulp	Wake Forest University, USA	Vadim Ponomarenko	San Diego State University, USA
Joseph Gallian	University of Minnesota Duluth, USA	Bjorn Poonen	UC Berkeley, USA
Stephan R. Garcia	Pomona College, USA	József H. Przytycki	George Washington University, USA
Anant Godbole	East Tennessee State University, USA	Richard Rebarber	University of Nebraska, USA
Ron Gould	Emory University, USA	Robert W. Robinson	University of Georgia, USA
Sat Gupta	U of North Carolina, Greensboro, USA	Javier Rojo	Oregon State University, USA
Jim Haglund	University of Pennsylvania, USA	Filip Saidak	U of North Carolina, Greensboro, USA
Johnny Henderson	Baylor University, USA	Hari Mohan Srivastava	University of Victoria, Canada
Glenn H. Hurlbert	Virginia Commonwealth University, USA	Andrew J. Sterge	Honorary Editor
Charles R. Johnson	College of William and Mary, USA	Ann Trenk	Wellesley College, USA
K. B. Kulasekera	Clemson University, USA	Ravi Vakil	Stanford University, USA
Gerry Ladas	University of Rhode Island, USA	Antonia Vecchio	Consiglio Nazionale delle Ricerche, Italy
David Larson	Texas A&M University, USA	John C. Wierman	Johns Hopkins University, USA
Suzanne Lenhart	University of Tennessee, USA	Michael E. Zieve	University of Michigan, USA
Chi-Kwong Li	College of William and Mary, USA		

PRODUCTION

Silvio Levy, Scientific Editor

Cover: Alex Scorpan

See inside back cover or msp.org/involve for submission instructions. The subscription price for 2019 is US \$195/year for the electronic version, and \$260/year (+\$35, if shipping outside the US) for print and electronic. Subscriptions, requests for back issues and changes of subscriber address should be sent to MSP.

Involve (ISSN 1444-4184 electronic, 1444-4176 printed) at Mathematical Sciences Publishers, 798 Evans Hall #3840, c/o University of California, Berkeley, CA 94720-3840, is published continuously online. Periodical rate postage paid at Berkeley, CA 94704, and additional mailing offices.

Involve peer review and production are managed by EditFlow® from Mathematical Sciences Publishers.

PUBLISHED BY

 **mathematical sciences publishers**
nonprofit scientific publishing

<http://msp.org/>

© 2019 Mathematical Sciences Publishers

involve

2019 vol. 12 no. 8

On the zero-sum group-magicness of cartesian products	1261
ADAM FONG, JOHN GEORGES, DAVID MAURO, DYLAN SPAGNUOLO, JOHN WALLACE, SHUFAN WANG AND KIRSTI WASH	
The variable exponent Bernoulli differential equation	1279
KAREN R. RÍOS-SOTO, CARLOS E. SEDA-DAMIANI AND ALEJANDRO VÉLEZ-SANTIAGO	
The supersingularity of Hurwitz curves	1293
ERIN DAWSON, HENRY FRAUENHOFF, MICHAEL LYNCH, AMETHYST PRICE, SEAMUS SOMERSTEP, ERIC WORK, DEAN BISOGNO AND RACHEL PRIES	
Multicast triangular semilattice network	1307
ANGELINA GROSSO, FELICE MANGANIELLO, SHIWANI VARAL AND EMILY ZHU	
Edge-transitive graphs and combinatorial designs	1329
HEATHER A. NEWMAN, HECTOR MIRANDA, ADAM GREGORY AND DARREN A. NARAYAN	
A logistic two-sex model with mate-finding Allee effect	1343
ELIZABETH ANDERSON, DANIEL MAXIN, JARED OTT AND GWYNETH TERRETT	
Unoriented links and the Jones polynomial	1357
SANDY GANZELL, JANET HUFFMAN, LESLIE MAVRAKIS, KAITLIN TADAMY AND GRIFFIN WALKER	
Nonsplit module extensions over the one-sided inverse of $k[x]$	1369
ZHEPING LU, LINHONG WANG AND XINGTING WANG	
Split Grothendieck rings of rooted trees and skew shapes via monoid representations	1379
DAVID BEERS AND MATT SZCZESNY	
On the classification of Specht modules with one-dimensional summands	1399
AUBREY PIPER COLLINS AND CRAIG J. DODGE	
The monochromatic column problem with a prime number of colors	1415
LORAN CROWELL AND STEVE SZABO	
Total Roman domination edge-critical graphs	1423
CHLOE LAMPMAN, KIEKA (C. M.) MYNHARDT AND SHANNON OGDEN	



1944-4176(2019)12:8;1-Y