

involve

a journal of mathematics

Uniqueness of a three-dimensional
stochastic differential equation

Carl Mueller and Giang Truong



Uniqueness of a three-dimensional stochastic differential equation

Carl Mueller and Giang Truong

(Communicated by Amarjit Singh Budhiraja)

In order to extend the study of the uniqueness property of multidimensional systems of stochastic differential equations, we look at the following three-dimensional system of equations, of which the two-dimensional case has been well studied: $dX_t = Y_t dt$, $dY_t = Z_t dt$, $dZ_t = |X_t|^\alpha dB_t$. We prove that if $(X_0, Y_0, Z_0) \neq (0, 0, 0)$ and $\frac{3}{4} < \alpha < 1$, then the system of equations has a unique solution in the strong sense.

1. Introduction and main results

The uniqueness of ordinary differential equations (ODEs) has been extensively studied; see for example [Hartman 1964]. In particular, if $F(u)$ is Lipschitz continuous, then

$$u'(t) = F(u(t)), \quad u(0) = u_0,$$

has a unique solution for all $t \geq 0$. In the case above, F , $u(t)$, and u_0 take values in \mathbb{R}^d , $d \geq 1$. The stochastic differential equation (SDE) realm, on the contrary, has different criteria for uniqueness of solutions; see for example [Protter 1990]. One of the most well-known results regarding strong uniqueness of SDEs is due to [Watanabe and Yamada 1971]. The result states that if $f(x)$ is locally Hölder continuous with index $\alpha \in [\frac{1}{2}, 1]$ and with linear growth, then

$$dX = f(X) dW, \quad X_0 = x_0,$$

has a unique strong solution for all times $t \geq 0$. Yamada and Watanabe's theory essentially focuses on one-dimensional SDEs.

MSC2010: primary 60H10; secondary 34F05.

Keywords: white noise, stochastic differential equations, uniqueness.

Mueller was supported by a Simons grant.

One motivation for studying higher-dimensional SDEs comes from the wave equation

$$\begin{aligned}\partial_t^2 u &= \Delta u, \\ u(0, x) &= u_0(x), \\ \partial_t u(0, x) &= u_1(x).\end{aligned}$$

In this equation, we have

$$\partial_t^2 u = \partial_x^2 u = \Delta u.$$

If we let

$$v = \partial_t u,$$

then we can rewrite the wave equation as the following system of equations:

$$\begin{aligned}\partial_t u &= v, \\ \partial_t v &= \Delta u.\end{aligned}$$

The original wave equation includes no noise. However, many physical systems are affected by noise. Hence, a modification of the wave equation which includes white noise is also studied:

$$\begin{aligned}\partial_t^2 u &= \Delta u + f(u)\dot{W}, \\ u(0, x) &= u_0(x), \\ \partial_t u(0, x) &= u_1(x).\end{aligned}\tag{1}$$

Note $x \in \mathbb{R}$ and $\dot{W} = \dot{W}(t, x)$ is white noise.

One well-known point is that Lipschitz continuity is sufficient for the uniqueness of SDEs. Thus, many mathematicians have studied whether Hölder continuity can still ensure the uniqueness property of SDEs. Gomez, Lee, Mueller, Neuman, and Salins [Gomez et al. 2017] studied the uniqueness property of the following two-dimensional model of SDEs:

$$\begin{aligned}dX &= Y dt, \\ dY &= |X|^\alpha dB, \\ (X_0, Y_0) &= (x_0, y_0).\end{aligned}\tag{2}$$

The results focused on $f(x) = |x|^\alpha$ since it is a prototype of an equation with Hölder continuous coefficients. Moreover, (2) is a version of (1) when we drop the dependence on x , which allows us to study the modified wave equation with more simplicity. Notice that if we take the differential dY of the first derivative of X , which is Y in the system of equations, it resembles the second derivative in time in the stochastic wave equation.

It was proven in [Gomez et al. 2017] that if $\alpha > \frac{1}{2}$ and $(x_0, y_0) \neq (0, 0)$, then (2) has a unique solution in the strong sense up to the time τ at which (X_t, Y_t) first hits the origin $(0, 0)$.

Since much is still unknown about higher-dimensional SDEs, we wish to continue the study of the uniqueness property of (2) in the three-dimensional case, which is

$$\begin{aligned} dX &= Y dt, \\ dY &= Z dt, \\ dZ &= |X|^\alpha dB, \end{aligned} \tag{3}$$

$$(X_0, Y_0, Z_0) = (x_0, y_0, z_0).$$

Theorem 1. *If $\frac{3}{4} < \alpha < 1$ and $(X_0, Y_0, Z_0) \neq 0$, then (3) has a unique strong solution, up to the time τ at which the solution (X_t, Y_t, Z_t) first hits the value $(0, 0, 0)$ or blows up.*

Moreover, we say the solution (X_t, Y_t, Z_t) blows up in finite time, with positive probability, if there is a random time $\tau < \infty$ such that

$$P\left(\lim_{t \uparrow \tau} |(X_t, Y_t, Z_t)|_{l^\infty} = \infty\right) > 0.$$

2. Proof of Theorem 1

Let (X_t^i, Y_t^i, Z_t^i) , $i = 1, 2$, be two solutions to (3) with $(x_0, y_0, z_0) \neq (0, 0, 0)$; in other words, (X_t^i, Y_t^i, Z_t^i) , $i = 1, 2$, have the same initial condition $(X_0, Y_0, Z_0) \neq (0, 0, 0)$. Note that since (X_t^i, Y_t^i, Z_t^i) , $i = 1, 2$, have the same initial conditions, from now on, we use X_0, Y_0, Z_0 instead of $X_0^{i,n}, Y_0^{i,n}, Z_0^{i,n}$.

Let τ be the first time t that either (X_t^1, Y_t^1, Z_t^1) or (X_t^2, Y_t^2, Z_t^2) hits the origin $(0, 0, 0)$ or blows up. We let τ be infinity if there is no such time.

First, if $X_0 \neq 0$, then (3) will have Lipschitz continuity up to the time that $X_t = 0$, and thus enables uniqueness to hold. Suppose after a certain amount of time X_t hits zero, where Lipschitz continuity no longer holds; then due to the strong Markov property, we begin the process again with $X_0 = 0$. So our goal is to prove pathwise uniqueness between excursions of X up to the time τ starting from $X_0 = 0$.

For any fixed n , let τ_n be the first time that either

$$|(X_t^1, Y_t^1, Z_t^1)|_{l^\infty} \wedge |(X_t^2, Y_t^2, Z_t^2)|_{l^\infty} \leq 2^{-n}$$

or

$$|(X_t^1, Y_t^1, Z_t^1)|_{l^\infty} \vee |(X_t^2, Y_t^2, Z_t^2)|_{l^\infty} \geq 2^n.$$

Here, the infinity norm (also known as the L_∞ -norm, l_∞ -norm, max norm, or uniform norm) of a vector \vec{v} is denoted by $|\vec{v}|_{l_\infty}$ and is defined as the maximum of

the absolute values of its components,

$$|\vec{v}|_{l_\infty} = \max\{|v_i| : i = 1, 2, \dots, n\},$$

and

$$\begin{aligned} a \wedge b &= \min(a, b), \\ a \vee b &= \max(a, b). \end{aligned}$$

If there is no such time, we let τ_n be infinity. Note that $\lim_{n \uparrow \infty} (\tau_n) = \tau$.

Now, for each fixed n , we will show uniqueness up to time τ_n in the system of equations

$$\begin{aligned} dX_t^{i,n} &= Y_t^{i,n} dt, \\ dY_t^{i,n} &= Z_t^{i,n} dt, \\ dZ_t^{i,n} &= |X_t^{i,n}|^\alpha \mathbf{1}_{[0, \tau_n]}(t) dB_t, \end{aligned} \tag{4}$$

$$(X_0, Y_0, Z_0) = (x_0, y_0, z_0).$$

In other words, after the time τ_n , we have $dZ_t^{i,n} = 0$, which makes $Z_t^{i,n}$ become constant. Specifically, given $m, n \in N$, we need

$$(X_t^n, Y_t^n, Z_t^n) = (X_t^m, Y_t^m, Z_t^m)$$

for all $t \leq \tau_n \wedge \tau_m$.

Now, before continuing the proof of uniqueness, by the method of contradiction, we show that the times that X_t hits zero do not accumulate before the time τ_n , almost surely.

For each n , let A_n be the event on which the times that $X_t^{i,n} = 0$, $i = 1$ or $i = 2$, accumulate before τ_n , and assume $P(A_n) > 0$. Then, on A_n , suppose σ_n is an accumulation point of the times t at which $X_t^{i,n} = 0$; i.e., there exists a sequence of times $\rho_{1,n} < \rho_{2,n} < \dots$ that converges to σ_n , and $X_{\rho_{k,n}}^{i,n} = 0$. Hence, on A_n , $\lim_{k \rightarrow \infty} \rho_{k,n} = \sigma_n$.

We have $X_t^{i,n}$ is almost surely continuous, and that $X_{\rho_{k,n}}^{i,n} = 0$ on A_n , so

$$\lim_{\rho_{k,n} \rightarrow \sigma_n} X_{\rho_{k,n}}^{i,n} = X_{\sigma_n}^{i,n} = 0$$

on A_n .

Note that $dX_t^{i,n} = Y_t^{i,n} dt$, and $Y_t^{i,n}$ is almost surely continuous. So if $Y_{\sigma_n}^{i,n} \neq 0$ on A_n , then there exists a random interval $[\sigma_n(\omega) - \epsilon(\omega), \sigma_n(\omega)]$ of positive length for which $X_t^{i,n} \neq 0$ on $[\sigma_n(\omega) - \epsilon(\omega), \sigma_n(\omega)]$. This contradicts the hypothesis of $\rho_{k,n}$ converging to σ_n .

If $Y_{\sigma_n}^{i,n} = 0$, there are two cases,

$$\sigma_n \geq \tau_n \quad \text{and} \quad \sigma_n < \tau_n.$$

If $\sigma_n \geq \tau_n$, then it means that the times at which $X_t^{i,n}$ hit zero do not accumulate before τ_n .

If $\sigma_n < \tau_n$, then with $X_{\sigma_n}^{i,n} = 0$ and $Y_{\sigma_n}^{i,n} = 0$, we have $|Z_{\sigma_n}^{i,n}| > 2^{-n}$. Now suppose $Z_{\sigma_n}^{i,n} > 2^{-n}$, as the case $Z_{\sigma_n}^{i,n} < 2^{-n}$ is approached similarly due to symmetry.

If $Z_{\sigma_n}^{i,n} > 2^{-n}$, since Z_t is almost surely continuous, there exists a time interval $[\sigma_n(\omega) - \epsilon'(\omega), \sigma_n(\omega)]$ on which $Z_t^{i,n} > 2^{-n}/2$. Thus for all $t \in [\sigma_n(\omega) - \epsilon'(\omega), \sigma_n(\omega)]$ we almost surely have

$$Y_t^{i,n} = Y_{\sigma_n}^{i,n} - \int_t^{\sigma_n} Z_s^{i,n} ds < -\frac{2^{-n}}{2}(\sigma_n - t).$$

Hence, integrating over $X_t^{i,n}$ for $t \in [\sigma_n - \epsilon', \sigma_n]$, we have

$$\begin{aligned} X_t^{i,n} &= X_{\sigma_n}^{i,n} - \int_t^{\sigma_n} Y_s^{i,n} ds = - \int_t^{\sigma_n} Y_s^{i,n} ds > \int_t^{\sigma_n} \frac{2^{-n}}{2}(\sigma_n - s) ds \\ &= \frac{2^{-n}}{2} \left(\sigma_n s - \frac{s^2}{2} \right) \Big|_t^{\sigma_n} = \frac{2^{-n}}{4}(\sigma_n - t)^2 > 0 \end{aligned}$$

as $t < \sigma_n$. Hence, this contradicts the hypothesis of $\rho_{k,n}$ converging to σ_n on A_n . So in conclusion, $P(A_n) = 0$, which means the times at which X_t hits zero do not accumulate before the time τ_n .

One more point we need to address before continuing with the proof of uniqueness is the existence of solutions. This problem is resolved in Theorems 21.7 and 21.8 and Lemma 21.17 of [Kallenberg 2002], which prove that for all times $t \geq 0$, solutions of multidimensional SDEs exist with probability 1 provided the coefficients are continuous and bounded.

Specifically, in our problem, since $\alpha \in (0, 1)$, the coefficients of system (3) are continuous and bounded by $(2^n)^\alpha \vee 2^n = 2^n$ up to the τ_n for each n , which satisfies the condition stated in the existence theory in [Kallenberg 2002]. Hence, existence of solution holds for all $t \leq \tau_n$ for all n . Therefore, up to the time $\tau = \sup \tau_n$, existence of solutions is ensured.

From now on, $X_t^{i,n}$ means $X_t^{1,n}$ and $X_t^{2,n}$. We define $Y_t^{i,n}$ and $Z_t^{i,n}$ similarly. Back to the proof of uniqueness, we have, for all $t \in [0, \tau_n]$,

$$|Z_t^{i,n}| \vee |Y_t^{i,n}| \leq 2^n.$$

So

$$|Y_t^{i,n}| = \left| Y_0 + \int_0^t Z_s^{i,n} ds \right| \leq |Y_0| + \int_0^t |Z_s^{i,n}| ds \leq 2^n + 2^n t.$$

Therefore,

$$|X_t^{i,n}| \leq |X_0 + \int_0^t Y_s^{i,n} ds| \leq \int_0^t |Y_s^{i,n}| ds \leq \int_0^t (2^n + 2^n s) ds = 2^n \left(t + \frac{t^2}{2} \right).$$

Now let

$$t_{0,n} = \frac{2^{-2n}}{2}. \tag{5}$$

Then

$$t_{0,n} + \frac{t_{0,n}^2}{2} = \frac{2^{-2n}}{2} + \frac{2^{-2n}}{8} \leq \frac{2^{-2n}}{2} + \frac{2^{-2n}}{2} = 2^{-2n}.$$

So

$$2^n \left(t_{0,n} + \frac{t_{0,n}^2}{2} \right) \leq 2^n \cdot 2^{-2n} = 2^{-n}.$$

Since the quadratic function $2^n(t + t^2/2)$ is increasing when $t \geq 0$, we have

$$|X_t^{i,n}| \leq 2^n \left(t + \frac{t^2}{2} \right) \leq 2^{-n}$$

for all $t \in [0, t_{0,n}]$.

Since $|X_t^{1,n}|$ and $|X_t^{2,n}|$ belong in $[0, 2^{-n}]$ for $t \in [0, t_{0,n}]$, based on the definition of τ_n above, either

$$|Y_0| \geq 2^{-n} \tag{6}$$

or

$$|Z_0| \geq 2^{-n} \tag{7}$$

for each fixed n . This is due to the fact that the solutions have the same initial condition (X_0, Y_0, Z_0) and for each time $t \in [0, t_{n,0}]$, either

$$|Y_t^{i,n}| \geq 2^{-n} \quad \text{or} \quad |Z_t^{i,n}| \geq 2^{-n}.$$

First, we deal with $Y_0 > 0$. Due to symmetry, we can deal with the case $Y_0 < 0$ with similar methods and thus omit the proof.

Now, with $Y_0 > 0$, we look at other subcases based on $Z_0^{i,n}$.

Case I: $Y_0 > 0$, $|Z_0| \leq 2^{-n}$. If $|Z_0| \leq 2^{-n}$, then (6) takes place. We are looking at the case $Y_0 > 0$, and thus $Y_0 > 2^{-n}$. Also, note that $dZ_t^{i,n} = 0$ for all $t > \tau_n$ and $|Z_t^{i,n}| \leq 2^n$ for all $t \in [0, \tau_n]$. Hence, $|Z_t^{i,n}| \leq 2^n$ for all t , which means $Z_t^{i,n} \geq -2^n$ for all t . Next, we have

$$Y_t^{i,n} = Y_0 + \int_0^t Z_s ds \geq 2^{-n} - \int_0^t 2^n ds = 2^{-n} - 2^n t.$$

If

$$0 < t < t_{0,n} = \frac{2^{-2n}}{2},$$

where $t_{0,n}$ is defined as in (5), then

$$2^n t < \frac{2^{-n}}{2}.$$

Thus

$$2^{-n} - 2^n t > \frac{2^{-n}}{2}$$

for all $t \in [0, t_{0,n}]$. In other words, $Y_t^{i,n} > 2^{-n}/2$ for $t \in [0, t_{0,n}]$. So, for all $t \in [0, t_{0,n}]$, we have

$$Y_t^{i,n} \geq \frac{2^{-n}}{2}.$$

Hence

$$X_t^{i,n} \geq X_0^{i,n} + \int_0^t \frac{2^{-n}}{2} ds \geq \frac{2^{-n}}{2} t. \quad (8)$$

Furthermore, based on (8) and $|X_t^{i,n}| \leq 2^{-n}$ for all $t \in [0, t_{0,n}]$, it leads to $t_{0,n} \leq 2$, otherwise $X_t^{i,n} > 2^{-n}$, which means that $t > t_{0,n}$, a contradiction.

Note that

$$\begin{aligned} X_t^{i,n} &= X_0 + \int_0^t Y_s^{i,n} ds, \\ Y_s^{i,n} &= Y_0 + \int_0^s Z_k^{i,n} dk, \\ Z_k^{i,n} &= Z_0 + \int_0^k |X_r^{i,n}|^\alpha \mathbf{1}_{[0, \tau_n]}(t) dB_r. \end{aligned}$$

Thus

$$\begin{aligned} X_t^{i,n} &= X_0 + Y_0 t + \int_0^t \int_0^s \left(Z_0 + \int_0^k |X_r^{i,n}|^\alpha \mathbf{1}_{[0, \tau_n]}(t) dB_r \right) dk ds \\ &= X_0 + Y_0 t + \int_0^t \int_0^s Z_0 dk ds + \int_0^t \int_0^s \int_0^k |X_r^{i,n}|^\alpha \mathbf{1}_{[0, \tau_n]}(t) dB_r dk ds \\ &= X_0 + Y_0 t + Z_0 \frac{t^2}{2} + \int_0^t \int_0^s \int_0^k |X_r^{i,n}|^\alpha \mathbf{1}_{[0, \tau_n]}(t) dB_r dk ds. \end{aligned}$$

Hence

$$(X_t^{1,n} - X_t^{2,n})^2 = \left(\int_0^t \int_0^s \int_0^k (|X_r^{1,n}|^\alpha - |X_r^{2,n}|^\alpha) \mathbf{1}_{[0, \tau_n]}(r) dB_r dk ds \right)^2.$$

Apply the Cauchy-Schwarz inequality twice, we get

$$\begin{aligned} (X_t^{1,n} - X_t^{2,n})^2 &\leq t \int_0^t \left(\int_0^s \int_0^k (|X_r^{1,n}|^\alpha - |X_r^{2,n}|^\alpha) \mathbf{1}_{[0, \tau_n]}(r) dB_r dk \right)^2 ds \\ &\leq t \int_0^t s \int_0^s \left(\int_0^k (|X_r^{1,n}|^\alpha - |X_r^{2,n}|^\alpha) \mathbf{1}_{[0, \tau_n]}(r) dB_r \right)^2 dk ds. \end{aligned}$$

Thus

$$E[(X_t^{1,n} - X_t^{2,n})^2] \leq t E \int_0^t s \int_0^s \left(\int_0^k (|X_r^{1,n}|^\alpha - |X_r^{2,n}|^\alpha) \mathbf{1}_{[0, \tau_n]}(r) dB_r \right)^2 dk ds.$$

By Itô's isometry,

$$\begin{aligned} E[(X_t^{1,n} - X_t^{2,n})^2] &\leq t E \int_0^t s \int_0^s \int_0^k (|X_r^{1,n}|^\alpha - |X_r^{2,n}|^\alpha) \mathbf{1}_{[0, \tau_n]}(r) dr dk ds \\ &\leq t E \int_0^t t \int_0^s \int_0^k (|X_r^{1,n}|^\alpha - |X_r^{2,n}|^\alpha)^2 dr dk ds \end{aligned}$$

$$\begin{aligned} &\leq t^2 E \int_0^t \int_0^s \int_0^k (|X_r^{1,n}|^\alpha - |X_r^{2,n}|^\alpha)^2 dr dk ds \\ &\leq t^2 E \int_0^t \int_0^t \int_0^t (|X_r^{1,n}|^\alpha - |X_r^{2,n}|^\alpha)^2 dr dk ds \\ &= t^4 E \int_0^t (|X_r^{1,n}|^\alpha - |X_r^{2,n}|^\alpha)^2 dr. \end{aligned}$$

Now we apply the mean value theorem for the function $f(x) = x^\alpha$, $0 < \alpha < 1$, and $a < b$:

$$b^\alpha - a^\alpha = \alpha c^{\alpha-1} (b - a) \leq \alpha a^{\alpha-1} (b - a)$$

for $c \in (a, b)$. Then for $r \in [0, t_{0,n}]$, where t_0 is determined in (5), we apply (8):

$$||X_r^{1,n}|^\alpha - |X_r^{2,n}|^\alpha| \leq \alpha \left(\frac{2^{-n}}{2} r\right)^{\alpha-1} ||X_r^{1,n}| - |X_r^{2,n}||. \tag{9}$$

Now let

$$D_t = E[(|X_t^{1,n}| - |X_t^{2,n}|)^2].$$

Since $t_{0,n} \leq 2$, we have for all $t \in [0, t_{0,n}]$

$$D_t \leq E[(X_t^{1,n} - X_t^{2,n})^2] \leq C_n \int_0^t r^{2\alpha-2} D_r dr.$$

for some C_n depending on n . Since $\alpha > \frac{3}{4}$, we have $r^{2\alpha-2}$ is integrable on $[0, t_{0,n}]$. At this stage we apply Gronwall’s lemma:

Lemma. *Let I denote an interval of the real line of the form $[a, \infty)$ or $[a, b]$ or (a, b) with $a < b$. Let β and u be real-valued continuous functions defined on I . If u is differentiable on the interior I° of I (the interval I without the endpoint a and possibly b) and satisfies the differential inequality*

$$u'(t) \leq \beta(t)u(t), \quad t \in I^\circ,$$

then u is bounded by the solution of the corresponding differential equation $v'(t) = \beta(t)v(t)$:

$$u(t) \leq u(a) \exp\left(\int_a^t \beta(s) ds\right)$$

for all $t \in I$.

Hence, with $D_0 = 0$, we have $D_t = 0$ for all $t \in [0, t_{0,n}]$. Therefore, (3) has unique strong solution in $[0, t_{0,n}]$.

Since $\alpha \geq \frac{3}{4}$, we have $2\alpha - 2 \geq -1$. Hence, $r^{2\alpha-2}$ is integrable on $[0, t_{0,n}]$. Note that in this case since $X_t \geq 2^{-n}$, $\eta \leq 1$. Applying Gronwall’s lemma, with $D_0 = 0$, we have $D_t = 0$ for all $t \in [0, t_{0,n}]$. Therefore, (3) has a unique solution in strong sense up to time $t_{0,n}$.

Note that for all $t \in [0, t_{0,n}]$, we have $Y_t^{i,n} > 2^{-n}/2 > 0$. Hence $X_t^{i,n}$ is strictly increasing, which, leads to $X_{t_{0,n}}$ strictly positive. Therefore, by the strong Markov property, we have uniqueness until the time X next hits zero.

Case II: $Y_0 > 0, Z_0 < -2^{-n}$. Since Z_0 starts negative, $Y_t^{i,n}$ decreases for an amount of time. Since Y_0 is positive, let say $Y_0 = \beta > 0$. Note that for all t , we have $|Z_t^{i,n}| < 2^n$, which means $Z_t^{i,n} > -2^n$. First, we have

$$Y_t^{i,n} = \beta + \int_0^t Z_s ds \geq \beta - \int_0^t 2^n ds = \beta - 2^n t.$$

Let

$$t'_{0,n} = \frac{\beta}{2^{n+1}}. \tag{10}$$

If

$$0 < t < t'_{0,n} = \frac{\beta}{2^{n+1}},$$

then

$$2^n t < \frac{\beta}{2};$$

thus

$$\beta - 2^n t > \frac{\beta}{2}$$

for all $t \in [0, t_{0,n}']$. In other words, $Y_t^{i,n} > \beta/2$ for all $t \in [0, t'_{0,n}]$.

So, for all $t \in [0, t_{0,n} \wedge t'_{0,n}]$, where $t_{0,n}$ and $t'_{0,n}$ are determined in (5) and (10) respectively, we have

$$Y_t^{i,n} \geq \frac{\beta}{2}$$

Hence

$$X_t^{i,n} \geq X_0^{i,n} + \int_0^t \frac{\beta}{2} ds \geq \frac{\beta}{2} t.$$

Applying the same method (9) above, we use the mean value theorem for the new lower bound of $X_t^{i,n}$:

$$||X_r^{1,n}|^\alpha - |X_r^{2,n}|^\alpha| \leq \alpha \left(\frac{\beta}{2} r\right)^{\alpha-1} ||X_r^{1,n}| - |X_r^{2,n}||.$$

Hence

$$D_t \leq E[(X_t^{1,n} - X_t^{2,n})^2] \leq C_n \int_0^t r^{2\alpha-2} D_r dr.$$

Again, applying Gronwall's lemma, with $D_0 = 0$, we have $D_t = 0$ for all $t \in [0, t_{0,n} \wedge t'_{0,n}]$. Therefore, (3) has a unique strong solution in $[0, t_{0,n} \wedge t'_{0,n}]$. As in the previous cases, we have $Y_t^{i,n} > \beta/2 > 0$ for all $t \in [t_{0,n} \wedge t'_{0,n}]$, which makes $X_t^{i,n}$ strictly increasing. So $X_{t_{0,n} \wedge t'_{0,n}}$ is strictly positive. Thus, by the strong Markov property, we have uniqueness until the next time X hits zero.

Case III: $Y_0 > 0$, $Z_0 > 2^{-n}$. Now we let T_n be the first time that either $Z_t^{1,n}$ or $Z_t^{2,n}$ hits the value $2^{-n}/2$. Since both $Z_t^{1,n}$ and $Z_t^{2,n}$ are continuous, we have $T_n > 0$ with probability 1. So we now prove uniqueness up to the time $t_{0,n} \wedge T_n$, where $t_{0,n}$ is defined in (5).

Then for all t in $[0, t_{0,n} \wedge T_n]$, we have

$$Z_t^{i,n} \geq \frac{2^{-n}}{2};$$

therefore

$$Y_t^{i,n} \geq Y_0 + \int_0^t \frac{2^{-n}}{2} ds \geq \frac{2^{-n}}{2} t \quad (11)$$

since $Y_0^{i,n} \geq 0$.

Based on (11), for $t \in [0, t_{0,n} \wedge T_n]$

$$X_t^{i,n} \geq X_0 + \int_0^t \frac{2^{-n}}{2} s ds \geq \frac{2^{-n}}{4} t^2. \quad (12)$$

Now we define

$$X_t^{i,n} = \tilde{X}_t^{i,n}, \quad Y_t^{i,n} = \tilde{Y}_t^{i,n}, \quad Z_t^{i,n} = \tilde{Z}_t^{i,n}$$

for $i = 1, 2$ and for $t \leq \tau_n \wedge T_n \wedge t_{0,n}$, where $t_{0,n}$ is defined as in (5) above.

Thus the following system of equations holds up to the stopping time $\tau_n \wedge T_n \wedge t_{0,n}$:

$$\begin{aligned} d\tilde{X}_t^{i,n} &= \tilde{Y}_t^{i,n} dt \\ d\tilde{Y}_t^{i,n} &= \tilde{Z}_t^{i,n} dt \\ d\tilde{Z}_t^{i,n} &= |\tilde{X}_t^{i,n}|^\alpha \mathbf{1}_{[0, \tau_n \wedge T_n \wedge t_{0,n}]}(t) dB_t, \end{aligned} \quad (13)$$

with $(\tilde{X}_0^{i,n}, \tilde{Y}_0^{i,n}, \tilde{Z}_0^{i,n}) = (X_0, Y_0, Z_0)$ for $i = 1, 2$. Furthermore, using (13), $\tilde{X}_t^{i,n}$, $\tilde{Y}_t^{i,n}$, and $\tilde{Z}_t^{i,n}$ can be defined for all times.

Using Itô's isometry as above with $\tilde{X}_t^{i,n}$, $\tilde{Y}_t^{i,n}$, and $\tilde{Z}_t^{i,n}$,

$$\begin{aligned} E[(\tilde{X}_t^{1,n} - \tilde{X}_t^{2,n})^2] &\leq t E \int_0^t s \int_0^s \int_0^k ((|\tilde{X}_r^{1,n}|^\alpha - |\tilde{X}_r^{2,n}|^\alpha) \mathbf{1}_{[0, \tau_n \wedge T_n \wedge t_{0,n}]}(r))^2 dr dk ds \\ &\leq t E \int_0^t t \int_0^s \int_0^k (|\tilde{X}_r^{1,n}|^\alpha - |\tilde{X}_r^{2,n}|^\alpha)^2 \mathbf{1}_{[0, \tau_n \wedge T_n \wedge t_{0,n}]}(r) dr dk ds \\ &\leq t^2 E \int_0^t \int_0^s \int_0^k (|\tilde{X}_r^{1,n}|^\alpha - |\tilde{X}_r^{2,n}|^\alpha)^2 \mathbf{1}_{[0, \tau_n \wedge T_n \wedge t_{0,n}]}(r) dr dk ds \\ &\leq t^2 E \int_0^t \int_0^t \int_0^t (|\tilde{X}_r^{1,n}|^\alpha - |\tilde{X}_r^{2,n}|^\alpha)^2 \mathbf{1}_{[0, \tau_n \wedge T_n \wedge t_{0,n}]}(r) dr dk ds \\ &= t^4 E \int_0^t (|\tilde{X}_r^{1,n}|^\alpha - |\tilde{X}_r^{2,n}|^\alpha)^2 \mathbf{1}_{[0, \tau_n \wedge T_n \wedge t_{0,n}]}(r) dr. \end{aligned}$$

Using (12) and the mean value theorem, for $r \in [0, \tau_n \wedge T_n \wedge t_{0,n}]$, we have

$$||\tilde{X}_r^{1,n}|^\alpha - |\tilde{X}_r^{2,n}|^\alpha| \leq \alpha \left(\frac{2^{-n}}{4} r^2 \right)^{\alpha-1} ||\tilde{X}_r^{1,n}| - |\tilde{X}_r^{2,n}||.$$

Hence

$$E[(\tilde{X}_t^{1,n} - \tilde{X}_t^{2,n})^2] \leq t^4 \alpha^2 \left(\frac{2^{-n}}{4} \right)^{2(\alpha-1)} E \int_0^t r^{4(\alpha-1)} (|\tilde{X}_r^{1,n}| - |\tilde{X}_r^{2,n}|)^2 dr,$$

so if we let

$$D_t = E[(|\tilde{X}_t^{1,n}| - |\tilde{X}_t^{2,n}|)^2],$$

then

$$D_t \leq E[(\tilde{X}_t^{1,n} - \tilde{X}_t^{2,n})^2] \leq C_n \int_0^t r^{4\alpha-4} D_r dr.$$

Again, applying Gronwall's lemma, with $D_0 = 0$, we have $D_t = 0$. Note that at the time $t_{0,n} \wedge T_n$, since we have $Z_t^{i,n} > 0$ for all $t \in [0, t_{0,n} \wedge T_n]$, and also $Y_0 > 0$, it leads to $Y_t^{i,n} > 0$ for all $t \in [0, t_{0,n} \wedge T_n]$. Thus $X_t^{i,n}$ is strictly increasing, which means $X_{t_{0,n} \wedge T_n}$ must be strictly greater than zero. Therefore, by the strong Markov property, we obtain uniqueness of the process until X next hits zero.

Case IV: $Y_0 = 0$. If $Y_0 = 0$, then based on the definition of τ_n , we have $|Z_0| > 2^{-n}$. We will first deal with the case $Z_0 > 2^{-n}$, and the case $Z_0 < 2^{-n}$ is approached the same way due to symmetry. As in Case III, let T_n be the first time that either $Z_t^{1,n}$ or $Z_t^{2,n}$ hits the value $2^{-n}/2$. Due to the continuity of $Z_t^{1,n}$ and $Z_t^{2,n}$, we have $T_n > 0$ with probability 1. So with for all $t \in [0, t_0 \wedge T_n]$, where t_0 is determined in (5), we have

$$Y_t^{i,n} \geq Y_0 + \int_0^t \frac{2^{-n}}{2} ds = \frac{2^{-n}}{2} t.$$

Then

$$X_t^{i,n} \geq X_0 + \int_0^t \frac{2^{-n}}{2} s ds \geq \frac{2^{-n}}{4} t^2.$$

We now apply the same method as in Case III by looking at $\tilde{X}_t^{i,n}$, $\tilde{Y}_t^{i,n}$, and $\tilde{Z}_t^{i,n}$, which are defined as

$$X_t^{i,n} = \tilde{X}_t^{i,n}, \quad Y_t^{i,n} = \tilde{Y}_t^{i,n}, \quad Z_t^{i,n} = \tilde{Z}_t^{i,n}$$

for $i = 1, 2$ and for $t \leq \tau_n \wedge T_n \wedge t_{0,n}$, as $t_{0,n}$ defined as in (5) above.

Thus the following system of equations holds up to the stopping time $\tau_n \wedge T_n \wedge t_{0,n}$:

$$\begin{aligned} d\tilde{X}_t^{i,n} &= \tilde{Y}_t^{i,n} dt \\ d\tilde{Y}_t^{i,n} &= \tilde{Z}_t^{i,n} dt \\ d\tilde{Z}_t^{i,n} &= |\tilde{X}_t^{i,n}|^\alpha \mathbf{1}_{[0, \tau_n \wedge T_n \wedge t_{0,n}]}(t) dB_t, \end{aligned}$$

with $(\tilde{X}_0^{i,n}, \tilde{Y}_0^{i,n}, \tilde{Z}_0^{i,n}) = (X_0, Y_0, Z_0)$ for $i = 1, 2$. Furthermore, using (13), $\tilde{X}_t^{i,n}$, $\tilde{Y}_t^{i,n}$, $\tilde{Z}_t^{i,n}$ can be defined for all time.

Again, using the same strategy in Case III and the mean value theorem, we have

$$||\tilde{X}_r^{1,n}|^\alpha - \tilde{X}_r^{2,n}|^\alpha| \leq \alpha \left(\frac{2^{-n}}{4} r^2\right)^{\alpha-1} ||\tilde{X}_r^{1,n}| - |\tilde{X}_r^{2,n}||.$$

Hence, if we let

$$D_t = E[(|\tilde{X}_t^{1,n}| - |\tilde{X}_t^{2,n}|)^2],$$

then

$$D_t \leq E[(\tilde{X}_t^{1,n} - \tilde{X}_t^{2,n})^2] \leq C_n \int_0^t r^{4\alpha-4} D_r dr.$$

Gronwall’s lemma with $D_0 = 0$ yields $D_t = 0$, completing the proof of **Theorem 1**.

In this case, we also have $Y_t^{i,n} > 2^{-n}/2t > 0$ for all $t \in [0, t_{0,n} \wedge T_n]$. Hence $X_t^{i,n}$ is strictly increasing, which yields $X_{t_{0,n} \wedge T_n}$ strictly positive. So, by the strong Markov property, we have uniqueness up to the time X next hits zero.

Now with uniqueness proved, we actually can even strengthen the proof by showing that $\tau_n^1 = \tau_n^2$ for all n , where τ_n^1 and τ_n^2 respectively stand for the stopping times at the critical values for X_t^1 and X_t^2 . Without loss of generality, suppose $\tau_n^2 > \tau_n^1$: So at the time τ_n^1 , X_t^2 has not yet reached the critical values, which are 2^{-n} or 2^n , as stated above. But since we have uniqueness up to $\tau_n^1 \wedge \tau_n^2$, this implies X_t^1 has also not reached the critical value at the time τ_n^1 , which is a contradiction to the definition of τ_n^1 . Hence, τ_n^1 and τ_n^2 must be equal.

References

[Gomez et al. 2017] A. Gomez, J. J. Lee, C. Mueller, E. Neuman, and M. Salins, “On uniqueness and blowup properties for a class of second order SDEs”, *Electron. J. Probab.* **22** (2017), art. id. 72. MR Zbl

[Hartman 1964] P. Hartman, *Ordinary differential equations*, Wiley, New York, 1964. MR Zbl

[Kallenberg 2002] O. Kallenberg, *Foundations of modern probability*, 2nd ed., Springer, 2002. MR Zbl

[Protter 1990] P. Protter, *Stochastic integration and differential equations: a new approach*, Appl. Math. (NY) **21**, Springer, 1990. MR Zbl

[Watanabe and Yamada 1971] S. Watanabe and T. Yamada, “On the uniqueness of solutions of stochastic differential equations, II”, *J. Math. Kyoto Univ.* **11** (1971), 553–563. MR Zbl

Received: 2019-10-03 Revised: 2020-02-18 Accepted: 2020-05-23

carl.e.mueller@rochester.edu Department of Mathematics, University of Rochester, Rochester, NY, United States

gtruong@u.rochester.edu Department of Mathematics, University of Rochester, Rochester, NY, United States

INVOLVE YOUR STUDENTS IN RESEARCH

Involve showcases and encourages high-quality mathematical research involving students from all academic levels. The editorial board consists of mathematical scientists committed to nurturing student participation in research. Bridging the gap between the extremes of purely undergraduate research journals and mainstream research journals, *Involve* provides a venue to mathematicians wishing to encourage the creative involvement of students.

MANAGING EDITOR

Kenneth S. Berenhaut Wake Forest University, USA

BOARD OF EDITORS

Colin Adams	Williams College, USA	Robert B. Lund	Clemson University, USA
Arthur T. Benjamin	Harvey Mudd College, USA	Gaven J. Martin	Massey University, New Zealand
Martin Bohner	Missouri U of Science and Technology, USA	Mary Meyer	Colorado State University, USA
Amarjit S. Budhiraja	U of N Carolina, Chapel Hill, USA	Frank Morgan	Williams College, USA
Pietro Cerone	La Trobe University, Australia	Mohammad Sal Moslehian	Ferdowsi University of Mashhad, Iran
Scott Chapman	Sam Houston State University, USA	Zuhair Nashed	University of Central Florida, USA
Joshua N. Cooper	University of South Carolina, USA	Ken Ono	Univ. of Virginia, Charlottesville
Jem N. Corcoran	University of Colorado, USA	Yuval Peres	Microsoft Research, USA
Toka Diagana	University of Alabama in Huntsville, USA	Y.-F. S. Pétermann	Université de Genève, Switzerland
Michael Dorff	Brigham Young University, USA	Jonathon Peterson	Purdue University, USA
Sever S. Dragomir	Victoria University, Australia	Robert J. Plemmons	Wake Forest University, USA
Joel Foisy	SUNY Potsdam, USA	Carl B. Pomerance	Dartmouth College, USA
Erin W. Fulp	Wake Forest University, USA	Vadim Ponomarenko	San Diego State University, USA
Joseph Gallian	University of Minnesota Duluth, USA	Bjorn Poonen	UC Berkeley, USA
Stephan R. Garcia	Pomona College, USA	József H. Przytycki	George Washington University, USA
Anant Godbole	East Tennessee State University, USA	Richard Rebarber	University of Nebraska, USA
Ron Gould	Emory University, USA	Robert W. Robinson	University of Georgia, USA
Sat Gupta	U of North Carolina, Greensboro, USA	Javier Rojo	Oregon State University, USA
Jim Haglund	University of Pennsylvania, USA	Filip Saidak	U of North Carolina, Greensboro, USA
Johnny Henderson	Baylor University, USA	Hari Mohan Srivastava	University of Victoria, Canada
Glenn H. Hurlbert	Virginia Commonwealth University, USA	Andrew J. Sterge	Honorary Editor
Charles R. Johnson	College of William and Mary, USA	Ann Trenk	Wellesley College, USA
K. B. Kulasekera	Clemson University, USA	Ravi Vakil	Stanford University, USA
Gerry Ladas	University of Rhode Island, USA	Antonia Vecchio	Consiglio Nazionale delle Ricerche, Italy
David Larson	Texas A&M University, USA	John C. Wierman	Johns Hopkins University, USA
Suzanne Lenhart	University of Tennessee, USA	Michael E. Zieve	University of Michigan, USA
Chi-Kwong Li	College of William and Mary, USA		

PRODUCTION

Silvio Levy, Scientific Editor

Cover: Alex Scorpan

See inside back cover or msp.org/involve for submission instructions. The subscription price for 2020 is US \$205/year for the electronic version, and \$275/year (+\$35, if shipping outside the US) for print and electronic. Subscriptions, requests for back issues and changes of subscriber address should be sent to MSP.

Involve (ISSN 1944-4184 electronic, 1944-4176 printed) at Mathematical Sciences Publishers, 798 Evans Hall #3840, c/o University of California, Berkeley, CA 94720-3840, is published continuously online. Periodical rate postage paid at Berkeley, CA 94704, and additional mailing offices.

Involve peer review and production are managed by EditFlow[®] from Mathematical Sciences Publishers.

PUBLISHED BY

 **mathematical sciences publishers**

nonprofit scientific publishing

<http://msp.org/>

© 2020 Mathematical Sciences Publishers

Hyperbolic triangular prisms with one ideal vertex	361
GRANT S. LAKELAND AND CORINNE G. ROTH	
On the sandpile group of Eulerian series-parallel graphs	381
KYLE WEISHAAR AND JAMES SEIBERT	
Linkages of calcium-induced calcium release in a cardiomyocyte simulated by a system of seven coupled partial differential equations	399
GERSON C. KROIZ, CARLOS BARAJAS, MATTHIAS K. GOBBERT AND BRADFORD E. PEERCY	
Covering numbers and schlicht functions	425
PHILIPPE DROUIN AND THOMAS RANSFORD	
Uniqueness of a three-dimensional stochastic differential equation	433
CARL MUELLER AND GIANG TRUONG	
Sharp sectional curvature bounds and a new proof of the spectral theorem	445
MAXINE CALLE AND COREY DUNN	
Qualitative investigation of cytolytic and noncytolytic immune response in an HBV model	455
JOHN G. ALFORD AND STEPHEN A. MCCOY	
A Cheeger inequality for graphs based on a reflection principle	475
EDWARD GELERNT, DIANA HALIKIAS, CHARLES KENNEY AND NICHOLAS F. MARSHALL	
Sets in \mathbb{R}^d determining k taxicab distances	487
VAJRESH BALAJI, OLIVIA EDWARDS, ANNE MARIE LOFTIN, SOLOMON MCHARO, LO PHILLIPS, ALEX RICE AND BINEYAM TSEGAYE	
Total difference chromatic numbers of graphs	511
RANJAN ROHATGI AND YUFEI ZHANG	
Decline of pollinators and attractiveness of the plants	529
LEILA ALICKOVIC, CHANG-HEE BAE, WILLIAM MAI, JAN RYCHTÁŘ AND DEWEY TAYLOR	