

involve

a journal of mathematics

Conjecture $\mathcal{O}$ holds for some horospherical varieties of Picard
rank 1

Lela Bones, Garrett Fowler, Lisa Schneider and Ryan M. Shifler



Conjecture $\mathcal{O}$ holds for some horospherical varieties of Picard rank 1

Lela Bones, Garrett Fowler, Lisa Schneider and Ryan M. Shifler

(Communicated by Kenneth S. Berenhaut)

Property $\mathcal{O}$ for an arbitrary complex, Fano manifold X is a statement about the eigenvalues of the linear operator obtained from the quantum multiplication of the anticanonical class of X . Conjecture $\mathcal{O}$ is a conjecture that property $\mathcal{O}$ holds for any Fano variety. Pasquier classified the smooth nonhomogeneous horospherical varieties of Picard rank 1 into five classes. Conjecture $\mathcal{O}$ has already been shown to hold for the odd symplectic Grassmannians, which is one of these classes. We will show that conjecture $\mathcal{O}$ holds for two more classes and an example in a third class of Pasquier's list. Perron–Frobenius theory reduces our proofs to be graph-theoretic in nature.

1. Introduction

The purpose of this paper is to prove that conjecture $\mathcal{O}$ holds for some horospherical varieties of Picard rank 1. We recall the precise statement of conjecture $\mathcal{O}$ for varieties of Picard rank 1, following [Galkin et al. 2016, Section 3]. Let F be a Fano variety, let $K := K_F$ be the canonical line bundle of F , let F_D be a fundamental divisor of F , and let

$$c_1(F) := c_1(-K) \in H^2(F)$$

be the anticanonical class. The Fano index of F is r , where r is the greatest integer such that $K_F \cong -rF_D$. The small quantum cohomology ring $(QH^*(F), \star)$ is a graded algebra over $\mathbb{Z}[q]$, where q is the quantum parameter. We define the small quantum cohomology in Section 2.1. Consider the specialization

$$H^\bullet(F) := QH^*(F)|_{q=1}$$

at $q = 1$. The quantum multiplication by the first Chern class $c_1(F)$ induces an endomorphism $\hat{c}_1$ of the finite-dimensional vector space $H^\bullet(F)$:

$$y \in H^\bullet(F) \mapsto \hat{c}_1(y) := (c_1(F) \star y)|_{q=1}.$$

MSC2010: primary 14N35; secondary 14N15, 15B48.

Keywords: quantum cohomology, horospherical, conjecture $\mathcal{O}$.

Set $\delta_0 := \max\{|\delta| : \delta \text{ is an eigenvalue of } \hat{c}_1\}$. Then property $\mathcal{O}$ states the following:

- (1) The real number δ_0 is an eigenvalue of $\hat{c}_1$ of multiplicity 1.
- (2) If δ is any eigenvalue of $\hat{c}_1$ with $|\delta| = \delta_0$, then $\delta = \delta_0 \gamma$ for some r -th root of unity $\gamma \in \mathbb{C}$, where r is the Fano index of F .

Property $\mathcal{O}$ was conjectured to hold for any Fano, complex manifold F in [Galkin et al. 2016]. If a Fano, complex, manifold has property $\mathcal{O}$ then we say that the space satisfies conjecture $\mathcal{O}$. Conjecture $\mathcal{O}$ underlies gamma conjectures I and II of Galkin, Golyshev, and Iritani. The gamma conjectures refine earlier conjectures by Dubrovin on Frobenius manifolds and mirror symmetry. Conjecture $\mathcal{O}$ has already been proved for the homogeneous G/P case in [Cheong and Li 2017], the odd symplectic Grassmannians in [Li et al. 2019], del Pezzo surfaces in [Hu et al. 2019], and projective complete intersections in [Ke 2018]. The Perron–Frobenius theory of nonnegative matrices reduces the proofs that conjecture $\mathcal{O}$ holds for the homogeneous and the odd symplectic Grassmannian cases to be a graph-theoretic check. This is because conjecture $\mathcal{O}$ is largely reminiscent of Perron–Frobenius theory. In this manuscript we will use the same graph-theoretic approach to prove that conjecture $\mathcal{O}$ holds for some smooth horospherical varieties of Picard rank 1.

Next we recall the definition of a horospherical variety following [Gonzales et al. 2018]. Let G be a complex reductive group. A G -variety is a reduced scheme of finite type over the field of complex numbers $\mathbb{C}$, equipped with an algebraic action of G . Let B be a Borel subgroup of G . A G -variety X is called spherical if X has a dense B -orbit. Let X be a G -spherical variety and let H be the stabilizer of a point in the dense G -orbit in X . The variety X is called *horospherical* if H contains a conjugate of the maximal unipotent subgroup of G contained in the Borel subgroup B .

Smooth horospherical varieties of Picard rank 1 were classified in [Pasquier 2009]. These varieties are either homogeneous or can be constructed in a uniform way via a triple $(\text{Type}(G), \omega_Y, \omega_Z)$ of representation-theoretic data, where $\text{Type}(G)$ is the semisimple Lie type of the reductive group G and ω_Y, ω_Z are fundamental weights. See [Pasquier 2009, Section 1.3] for details. Pasquier classified the possible triples into five classes:

- (1) $(B_n, \omega_{n-1}, \omega_n)$ with $n \geq 3$.
- (2) $(B_3, \omega_1, \omega_3)$.
- (3) $(C_n, \omega_m, \omega_{m-1})$ with $n \geq 2$ and $m \in [2, n]$ (the odd symplectic Grassmannians).
- (4) $(F_4, \omega_2, \omega_3)$.
- (5) $(G_2, \omega_1, \omega_2)$.

Proposition 3.6 [Pasquier 2006] showed the triples in the above list are Fano varieties. We are now able to state the main theorem:

Theorem 1. *If F belongs to the classes (1) for $n = 3$, (2), (3), and (5) of Pasquier’s list, then conjecture $\mathcal{O}$ holds for F .*

2. Preliminaries

2.1. Quantum cohomology. The small quantum cohomology is defined as follows. Let $(\alpha_i)_i$ be a basis of $H^*(F)$, the classical cohomology ring, and let $(\alpha_i^\vee)_i$ be the dual basis for the Poincaré pairing. The multiplication is given by

$$\alpha_i \star \alpha_j = \sum_{d \geq 0, k} c_{i,j}^{k,d} q^d \alpha_k,$$

where $c_{i,j}^{k,d}$ are the 3-point, genus-0, Gromov–Witten invariants corresponding to the classes α_i , α_j , and $\alpha_k^\vee$. We will make use of the quantum Chevalley formula, which is the multiplication of a hyperplane class h with another class α_j . Theorem 0.0.3 of [Gonzales et al. 2018] implies that if F belongs to the classes (1) for $n = 3$, (2), or (5) of Pasquier’s list, then there is an explicit quantum Chevalley formula. The explicit quantum Chevalley formula is the key ingredient used to prove property $\mathcal{O}$ holds.

2.2. Sufficient criterion for property $\mathcal{O}$ to hold. We recall the notion of the (oriented) quantum Bruhat graph of a Fano variety F . The vertices of this graph are the basis elements

$$\alpha_i \in H^\bullet(F) := QH^*(F)|_{q=1}.$$

There is an oriented edge $\alpha_i \rightarrow \alpha_j$ if the class α_j appears with positive coefficient (where we consider $q > 0$) in the quantum Chevalley multiplication $h \star \alpha_i$ for some hyperplane class h . Using the Perron–Frobenius theory of nonnegative matrices, conjecture $\mathcal{O}$ reduces to a graph-theoretic check of the quantum Bruhat graph. The techniques involving Perron–Frobenius theory used by Li, Mihalcea, and Shifler [Li et al. 2019] and Cheong and Li [2017] imply the following lemma:

Lemma 2. *Suppose the following conditions hold for a Fano variety F :*

- (1) *The matrix representation of $\hat{c}_1$ is nonnegative.*
- (2) *The quantum Bruhat graph of F is strongly connected.*
- (3) *There exists a cycle of length r , the Fano index, in the quantum Bruhat graph of F .*

Then property $\mathcal{O}$ holds for F . We say the matrix representation of $\hat{c}_1$ is nonnegative if all of the entries are nonnegative.

We refer the reader to [Minc 1988, Section 4.3] for further details on Perron–Frobenius theory.

3. Checking property $\mathcal{O}$ holds

Let X be a horospherical variety. We will simplify our notation where the basis of $H^\bullet(X)$ is $\{1, h, \alpha_i\}_{i \in I}$ for some finite index set I . Observe by [Gonzales et al. 2018] that the anticanonical classes are

$$c_1(X) = \begin{cases} 5h & \text{when } X \text{ is case (1) for } n = 3, \\ 7h & \text{when } X \text{ is case (2),} \\ 4h & \text{when } X \text{ is case (5),} \end{cases}$$

and the Fano indices are

$$r = \begin{cases} 5 & \text{when } X \text{ is case (1) for } n = 3, \\ 7 & \text{when } X \text{ is case (2),} \\ 4 & \text{when } X \text{ is case (5).} \end{cases}$$

The endomorphism $\hat{c}_1$ acting on the basis elements of $H^\bullet(X)$ is determined by the Chevalley formula in the following way:

$$\begin{aligned} \hat{c}_1(\alpha_i) &= 5(h \star \alpha_i)|_{q=1} && \text{when } X \text{ is case (1) for } n = 3, \\ \hat{c}_1(\alpha_i) &= 7(h \star \alpha_i)|_{q=1} && \text{when } X \text{ is case (2), and} \\ \hat{c}_1(\alpha_i) &= 4(h \star \alpha_i)|_{q=1} && \text{when } X \text{ is case (5).} \end{aligned}$$

Each of the following three subsections will show that conjecture $\mathcal{O}$ holds for case (1) for $n = 3$, case (2), and case (5) of Pasquier’s list, respectively. In each subsection we will reformulate the quantum Chevalley formulas stated in [Gonzales et al. 2018], present the quantum Bruhat graph, and argue that each condition of Lemma 2 is satisfied. For each case, we have kept the same format of the equations presented by [Gonzales et al. 2018] with our prescribed basis for ease of identification for the reader. For example, line 3 in Proposition 4.3 of that paper is

$$h * \sigma'_{u_2} = \sigma'_{u_3} + \sigma'_{u'_3} \quad \text{and} \quad h * \sigma'_{u'_2} = 2\sigma'_{u'_3} + \tau_{v_0}.$$

In Proposition 3 below we identify this line with

$$\hat{c}_1(\alpha_1) = 5\alpha_3 + 5\alpha_4 \quad \text{and} \quad \hat{c}_1(\alpha_2) = 10\alpha_3 + 5\alpha_5.$$

3.1. Case (1) for $n = 3$. We will reformulate the quantum Chevalley formula stated in [Gonzales et al. 2018] using the basis $\{1, h, \alpha_1, \alpha_2, \dots, \alpha_{18}\}$.

Proposition 3. *The following equalities hold by [Gonzales et al. 2018, Proposition 4.3]:*

- (1) $\hat{c}_1(1) = 5h$.
- (2) $\hat{c}_1(h) = 10\alpha_1 + 5\alpha_2$.
- (3) $\hat{c}_1(\alpha_1) = 5\alpha_3 + 5\alpha_4$ and $\hat{c}_1(\alpha_2) = 10\alpha_3 + 5\alpha_5$.




Figure 1. The quantum Bruhat graph of the Fano variety X in case (1) for $n = 3$.

- (4) $\hat{c}_1(\alpha_3) = 10\alpha_6 + 5\alpha_7 + 5\alpha_8$, $\hat{c}_1(\alpha_4) = 5\alpha_6 + 10\alpha_7$, and $\hat{c}_1(\alpha_5) = 5\alpha_8$.
- (5) $\hat{c}_1(\alpha_6) = 10\alpha_9 + 5\alpha_{10} + 5\alpha_{11}$, $\hat{c}_1(\alpha_7) = 5\alpha_{10}$, and $\hat{c}_1(\alpha_8) = 5\alpha_{11} + 5 \cdot 1$.
- (6) $\hat{c}_1(\alpha_9) = 5\alpha_{12} + 5\alpha_{13}$, $\hat{c}_1(\alpha_{10}) = 10\alpha_{13} + 5\alpha_{14}$, $\hat{c}_1(\alpha_{11}) = 5\alpha_{12} + 5\alpha_{14} + 5h$.
- (7) $\hat{c}_1(\alpha_{12}) = 5\alpha_{15} + 5\alpha_1$, $\hat{c}_1(\alpha_{13}) = 5\alpha_{15} + 5\alpha_{16}$, and $\hat{c}_1(\alpha_{14}) = 5\alpha_{15} + 5\alpha_2$.
- (8) $\hat{c}_1(\alpha_{15}) = 5\alpha_{17} + 5\alpha_3$ and $\hat{c}_1(\alpha_{16}) = 5\alpha_{17} + 5\alpha_5$.
- (9) $\hat{c}_1(\alpha_{17}) = 5\alpha_{18} + 5\alpha_6 + 5\alpha_8$.
- (10) $\hat{c}_1(\alpha_{18}) = 5\alpha_9 + 5\alpha_{11} + 10 \cdot 1$.

The quantum Bruhat graph is shown in [Figure 1](#). The bold edges indicate a cycle of length $r = 5$, the Fano index.

Lemma 4. *Property $\mathcal{O}$ holds when X is case (1) with $n = 3$ of Pasquier's list.*

Proof. The coefficients that appear in the equations in [Proposition 3](#) are the entries of the matrix representation of $\hat{c}_1$. Therefore, the matrix representation of $\hat{c}_1$ is nonnegative. The quantum Bruhat graph is strongly connected by [Figure 1](#), and the cycle $\alpha_{18}\alpha_{11}\alpha_{14}\alpha_{15}\alpha_{17}\alpha_{18}$ has length $r = 5$. $\square$

3.2. Case (2). Again, we reformulate the quantum Chevalley formula from [\[Gonzales et al. 2018\]](#) using the basis $\{1, h, \alpha_1, \alpha_2, \dots, \alpha_{12}\}$.




Figure 2. The quantum Bruhat graph for case (2).

Proposition 5. *The following equalities hold by [Gonzales et al. 2018, Proposition 4.4]:*

- (1) $\hat{c}_1(1) = 7h$.
- (2) $\hat{c}_1(h) = 7\alpha_1$.
- (3) $\hat{c}_1(\alpha_1) = 14\alpha_2 + 7\alpha_3$.
- (4) $\hat{c}_1(\alpha_2) = 7\alpha_4 + 7\alpha_5$ and $\hat{c}_1(\alpha_3) = 7\alpha_5$.
- (5) $\hat{c}_1(\alpha_4) = 7\alpha_6 + 7\alpha_7$ and $\hat{c}_1(\alpha_5) = 7\alpha_7$.
- (6) $\hat{c}_1(\alpha_6) = 7\alpha_8$ and $\hat{c}_1(\alpha_7) = 7\alpha_8 + 7\alpha_9$.
- (7) $\hat{c}_1(\alpha_8) = 7\alpha_{10}$ and $\hat{c}_1(\alpha_9) = 7\alpha_{10} + 7 \cdot 1$.
- (8) $\hat{c}_1(\alpha_{10}) = 7\alpha_{11} + 7h$.
- (9) $\hat{c}_1(\alpha_{11}) = 7\alpha_{12} + 7\alpha_1$.
- (10) $\hat{c}_1(\alpha_{12}) = 7\alpha_2$.

The quantum Bruhat graph is shown in [Figure 2](#).

Lemma 6. *Property $\mathcal{O}$ holds when X is case (2) of Pasquier’s list.*

Proof. The coefficients that appear in the equations in [Proposition 5](#) are the entries of the matrix representation of $\hat{c}_1$. Therefore, the matrix representation of $\hat{c}_1$ is




Figure 3. The quantum Bruhat graph for case (5).

nonnegative. The quantum Bruhat graph is strongly connected by Figure 2, and the cycle $\alpha_{12}\alpha_2\alpha_4\alpha_6\alpha_8\alpha_{10}\alpha_{11}\alpha_{12}$ has length $r = 7$. $\square$

3.3. Case (5). Again, we reformulate the quantum Chevalley formula from [Gonzales et al. 2018] using the basis $\{1, h, \alpha_1, \alpha_2, \dots, \alpha_{10}\}$.

Proposition 7. *The following equalities hold by [Gonzales et al. 2018, Proposition 4.6]:*

- (1) $\hat{c}_1(1) = 4h$.
- (2) $\hat{c}_1(h) = 12\alpha_1 + 4\alpha_2$.
- (3) $\hat{c}_1(\alpha_1) = 8\alpha_3 + 4\alpha_4$ and $\hat{c}_1(\alpha_2) = 4\alpha_4$.
- (4) $\hat{c}_1(\alpha_3) = 12\alpha_5 + 4\alpha_6$ and $\hat{c}_1(\alpha_4) = 4\alpha_6 + 4 \cdot 1$.
- (5) $\hat{c}_1(\alpha_5) = 4\alpha_7 + 4\alpha_8$ and $\hat{c}_1(\alpha_6) = 8\alpha_7 + 4h$.
- (6) $\hat{c}_1(\alpha_7) = 4\alpha_9 + 4\alpha_1$ and $\hat{c}_1(\alpha_8) = 4\alpha_9 + 4\alpha_2$.
- (7) $\hat{c}_1(\alpha_9) = 4\alpha_{10} + 4\alpha_3 + 4\alpha_4$.
- (8) $\hat{c}_1(\alpha_{10}) = 4\alpha_5 + 4\alpha_6 + 8 \cdot 1$.

The associated quantum Bruhat graph is shown in Figure 3.

Lemma 8. *Property $\mathcal{O}$ holds when X is case (5) of Pasquier's list.*

Proof. The coefficients that appear in the equations in [Proposition 7](#) are the entries of the matrix representation of $\hat{c}_1$. Therefore, the matrix representation of $\hat{c}_1$ is nonnegative. The quantum Bruhat graph is strongly connected by [Figure 3](#), and the cycle $\alpha_{10}\alpha_6\alpha_7\alpha_9\alpha_{10}$ has length $r = 4$. $\square$

[Theorem 1](#) follows from [Lemmas 4, 6, 8](#), and the previously mentioned work done by Li, Mihalcea, Shifler [[Li et al. 2019](#)] for the odd symplectic Grassmannian case.

Acknowledgement

We would like to thank the anonymous referees for their useful comments and suggestions to improve the presentation of this paper.

References

- [Cheong and Li 2017] D. Cheong and C. Li, “On the conjecture $\mathcal{O}$ of GGI for G/P ”, *Adv. Math.* **306** (2017), 704–721. [MR](#) [Zbl](#)
- [Galkin et al. 2016] S. Galkin, V. Golyshev, and H. Iritani, “Gamma classes and quantum cohomology of Fano manifolds: gamma conjectures”, *Duke Math. J.* **165**:11 (2016), 2005–2077. [MR](#) [Zbl](#)
- [Gonzales et al. 2018] R. Gonzales, C. Pech, N. Perrin, and A. Samokhin, “Geometry of horospherical varieties of Picard rank one”, preprint, 2018. [arXiv](#)
- [Hu et al. 2019] J. Hu, H.-Z. Ke, C. Li, and T. Yang, “Gamma conjecture I for del Pezzo surfaces”, preprint, 2019. [arXiv](#)
- [Ke 2018] H.-Z. Ke, “On Conjecture $\mathcal{O}$ for projective complete intersections”, preprint, 2018. [arXiv](#)
- [Li et al. 2019] C. Li, L. C. Mihalcea, and R. M. Shifler, “Conjecture $\mathcal{O}$ holds for the odd symplectic Grassmannian”, *Bull. Lond. Math. Soc.* **51**:4 (2019), 705–714. [MR](#) [Zbl](#)
- [Minc 1988] H. Minc, *Nonnegative matrices*, John Wiley & Sons, New York, 1988. [MR](#) [Zbl](#)
- [Pasquier 2006] B. Pasquier, *Variétés horosphériques de Fano*, Ph.D. thesis, Université Joseph Fourier, 2006, available at <https://tel.archives-ouvertes.fr/tel-00111912/document>.
- [Pasquier 2009] B. Pasquier, “On some smooth projective two-orbit varieties with Picard number 1”, *Math. Ann.* **344**:4 (2009), 963–987. [MR](#) [Zbl](#)

Received: 2018-07-10

Revised: 2019-07-12

Accepted: 2020-07-25

lbones1@gulls.salisbury.edu

*Department of Mathematics, Salisbury University,
Salisbury, MD, United States*

gfwolder2@gulls.salisbury.edu

*Department of Mathematics, Salisbury University,
Salisbury, MD, United States*

lmshneider@salisbury.edu

*Department of Mathematics, Salisbury University,
Salisbury, MD, United States*

rmshifler@salisbury.edu

*Department of Mathematics, Salisbury University,
Salisbury, MD, United States*

INVOLVE YOUR STUDENTS IN RESEARCH

Involve showcases and encourages high-quality mathematical research involving students from all academic levels. The editorial board consists of mathematical scientists committed to nurturing student participation in research. Bridging the gap between the extremes of purely undergraduate research journals and mainstream research journals, *Involve* provides a venue to mathematicians wishing to encourage the creative involvement of students.

MANAGING EDITOR

Kenneth S. Berenhaut Wake Forest University, USA

BOARD OF EDITORS

Colin Adams	Williams College, USA	Robert B. Lund	Clemson University, USA
Arthur T. Benjamin	Harvey Mudd College, USA	Gaven J. Martin	Massey University, New Zealand
Martin Bohner	Missouri U of Science and Technology, USA	Mary Meyer	Colorado State University, USA
Amarjit S. Budhiraja	U of N Carolina, Chapel Hill, USA	Frank Morgan	Williams College, USA
Pietro Cerone	La Trobe University, Australia	Mohammad Sal Moslehian	Ferdowsi University of Mashhad, Iran
Scott Chapman	Sam Houston State University, USA	Zuhair Nashed	University of Central Florida, USA
Joshua N. Cooper	University of South Carolina, USA	Ken Ono	Univ. of Virginia, Charlottesville
Jem N. Corcoran	University of Colorado, USA	Yuval Peres	Microsoft Research, USA
Toka Diagana	University of Alabama in Huntsville, USA	Y.-F. S. Pétermann	Université de Genève, Switzerland
Michael Dorff	Brigham Young University, USA	Jonathon Peterson	Purdue University, USA
Sever S. Dragomir	Victoria University, Australia	Robert J. Plemmons	Wake Forest University, USA
Joel Foisy	SUNY Potsdam, USA	Carl B. Pomerance	Dartmouth College, USA
Erin W. Fulp	Wake Forest University, USA	Vadim Ponomarenko	San Diego State University, USA
Joseph Gallian	University of Minnesota Duluth, USA	Bjorn Poonen	UC Berkeley, USA
Stephan R. Garcia	Pomona College, USA	József H. Przytycki	George Washington University, USA
Anant Godbole	East Tennessee State University, USA	Richard Rebarber	University of Nebraska, USA
Ron Gould	Emory University, USA	Robert W. Robinson	University of Georgia, USA
Sat Gupta	U of North Carolina, Greensboro, USA	Javier Rojo	Oregon State University, USA
Jim Haglund	University of Pennsylvania, USA	Filip Saidak	U of North Carolina, Greensboro, USA
Johnny Henderson	Baylor University, USA	Hari Mohan Srivastava	University of Victoria, Canada
Glenn H. Hurlbert	Virginia Commonwealth University, USA	Andrew J. Sterge	Honorary Editor
Charles R. Johnson	College of William and Mary, USA	Ann Trenk	Wellesley College, USA
K. B. Kulasekera	Clemson University, USA	Ravi Vakil	Stanford University, USA
Gerry Ladas	University of Rhode Island, USA	Antonia Vecchio	Consiglio Nazionale delle Ricerche, Italy
David Larson	Texas A&M University, USA	John C. Wierman	Johns Hopkins University, USA
Suzanne Lenhart	University of Tennessee, USA	Michael E. Zieve	University of Michigan, USA
Chi-Kwong Li	College of William and Mary, USA		

PRODUCTION

Silvio Levy, Scientific Editor

Cover: Alex Scorpan

See inside back cover or msp.org/involve for submission instructions. The subscription price for 2020 is US \$205/year for the electronic version, and \$275/year (+\$35, if shipping outside the US) for print and electronic. Subscriptions, requests for back issues and changes of subscriber address should be sent to MSP.

Involve (ISSN 1944-4184 electronic, 1944-4176 printed) at Mathematical Sciences Publishers, 798 Evans Hall #3840, c/o University of California, Berkeley, CA 94720-3840, is published continuously online. Periodical rate postage paid at Berkeley, CA 94704, and additional mailing offices.

Involve peer review and production are managed by EditFlow® from Mathematical Sciences Publishers.

PUBLISHED BY

 **mathematical sciences publishers**
nonprofit scientific publishing

<http://msp.org/>

© 2020 Mathematical Sciences Publishers

involve

2020

vol. 13

no. 4

Structure constants of $\mathcal{U}(\mathfrak{sl}_2)$	541
ALEXIA GOURLEY AND CHRISTOPHER KENNEDY	
Conjecture $\mathcal{C}$ holds for some horospherical varieties of Picard rank 1	551
LELA BONES, GARRETT FOWLER, LISA SCHNEIDER AND RYAN M. SHIFLER	
Condensed Ricci curvature of complete and strongly regular graphs	559
VINCENT BONINI, CONOR CARROLL, UYEN DINH, SYDNEY DYE, JOSHUA FREDERICK AND ERIN PEARSE	
On equidistant polytopes in the Euclidean space	577
CSABA VINCZE, MÁRK OLÁH AND LETÍCIA LENGYEL	
Polynomial values in Fibonacci sequences	597
ADI OSTROV, DANNY NEFTIN, AVI BERMAN AND REYAD A. ELRAZIK	
Stability and asymptotic analysis of the Föllmer–Schweizer decomposition on a finite probability space	607
SARAH BOESE, TRACY CUI, SAMUEL JOHNSTON, SYLVIE VEGA-MOLINO AND OLEKSII MOSTOVYI	
Eigenvalues of the sum and product of anticommuting matrices	625
VADIM PONOMARENKO AND LOUIS SELSTAD	
Combinatorial random knots	633
ANDREW DUCHARME AND EMILY PETERS	
Conjugation diameter of the symmetric groups	655
ASSAF LIBMAN AND CHARLOTTE TARRY	
Existence of multiple solutions to a discrete boundary value problem with mixed periodic boundary conditions	673
KIMBERLY HOWARD, LONG WANG AND MIN WANG	
Minimal flag triangulations of lower-dimensional manifolds	683
CHRISTIN BIBBY, ANDREW ODESKY, MENG MENG WANG, SHUYANG WANG, ZIYI ZHANG AND HAILUN ZHENG	
Some new Gompertz fractional difference equations	705
TOM CUCHTA AND BROOKE FINCHAM	