

involve

a journal of mathematics

Condensed Ricci curvature of complete
and strongly regular graphs

Vincent Bonini, Conor Carroll, Uyen Dinh,
Sydney Dye, Joshua Frederick and Erin Pearce



Condensed Ricci curvature of complete and strongly regular graphs

Vincent Bonini, Conor Carroll, Uyen Dinh,
 Sydney Dye, Joshua Frederick and Erin Pearce

(Communicated by Kenneth S. Berenhaut)

We study a modified notion of Ollivier's coarse Ricci curvature on graphs introduced by Lin, Lu, and Yau. We establish a rigidity theorem for complete graphs that shows a connected finite simple graph is complete if and only if the Ricci curvature is strictly greater than 1. We then derive explicit Ricci curvature formulas for strongly regular graphs in terms of the graph parameters and the size of a maximal matching in the core neighborhood. As a consequence we are able to derive exact Ricci curvature formulas for strongly regular graphs of girths 4 and 5 using elementary means. An example is provided that shows there is no exact formula for the Ricci curvature for strongly regular graphs of girth 3 that is purely in terms of graph parameters.

1. Introduction

Lott and Villani [2009] and Sturm [2006] discovered a relationship between optimal transport and Ricci curvature on smooth Riemannian manifolds and they pursued a construction of a synthetic notion of Ricci curvature that could be defined independently of differentiable structures. Ollivier [2009; 2010] later introduced a notion of coarse Ricci curvature for Markov chains on metric spaces which has a particularly accessible formulation on graphs. Much work has been done on coarse Ricci curvature on graphs; see [Bauer et al. 2012; Lin et al. 2011; 2014; Jost and Liu 2014; Smith 2014; Bhattacharya and Mukherjee 2015; Bourne et al. 2018; Münch and Wojciechowski 2019].

Ollivier's coarse Ricci curvature on graphs is defined in terms of transport distance of probability measures (see Section 2). We study a modified notion of Ollivier's coarse Ricci curvature on graphs introduced by Lin, Lu, and Yau [Lin et al. 2011]. In this paper we refer to the modified Ricci curvature of [Lin et al. 2011] as the condensed Ricci curvature. Our first result concerning the condensed Ricci

MSC2010: primary 52C99, 53B99; secondary 05C10, 05C81, 05C99.

Keywords: coarse Ricci curvature, strongly regular graphs.

curvature exploits the relationship between Ricci curvature and the eigenvalues of the spectral graph Laplacian to establish a rigidity theorem for complete graphs. This result is stated without proof in [Lin et al. 2011, Example 1].

Theorem 1.1. *A connected finite simple graph $G = (V, E)$ is complete if and only if the condensed Ricci curvature satisfies $\mathbb{k}(x, y) > 1$ for all edges $xy \in E$. In this case, $\mathbb{k}(x, y) = n/(n - 1)$ for all vertices $x, y \in V$.*

We then turn our attention to the derivation of exact formulas for the condensed Ricci curvature on strongly regular graphs. Explicit formulas and curvature bounds for various forms of Ricci curvature have been established for large classes of graphs; see [Lin et al. 2011; 2014; Jost and Liu 2014; Smith 2014; Bhattacharya and Mukherjee 2015; Cushing et al. 2020]. In particular, Bakry–Émery Ricci curvature functions are studied in [Cushing et al. 2020] and explicit formulas for the Bakry–Émery Ricci curvature are derived for Cayley graphs and strongly regular graphs of girths 4 and 5. This work served as our inspiration to establish explicit formulas for the condensed Ricci curvature on strongly regular graphs using elementary means.

Let $G = (V, E)$ be a strongly regular graph with parameters (n, d, α, β) . Here $n = |V|$ is the number of vertices, d is the uniform degree of the vertices, $\alpha \geq 0$ is the number of common neighbors for adjacent vertices, and $\beta \geq 1$ is the number of common neighbors for nonadjacent vertices. Given an edge $xy \in E$, let N_x denote the set of vertices that are adjacent to x , not including y or any vertices that are adjacent to y . Similarly, let N_y denote the set of vertices that are adjacent to y , not including x or any vertices that are adjacent to x (see Section 2 for more details). We compute the following curvature formulas solely from properties of maximum matchings and the regularity properties of strongly regular graphs.

Theorem 1.2. *Suppose $G = (V, E)$ is a strongly regular graph with parameters (n, d, α, β) , and $xy \in E$ with maximum matching \mathcal{M} of size m between N_x and N_y . Then the condensed Ricci curvature satisfies*

$$\mathbb{k}(x, y) = \frac{\alpha + 2}{d} - \frac{|N_x| - m}{d}. \quad (1-1)$$

Remark 1.3. We would like to point out that Theorem 1.2 holds more generally for regular graphs of diameter 2. In this case one takes α to be the number of neighbors common to both x and y . For simplicity in our presentation, we focus on strongly regular graphs.

The explicit formula $\mathbb{k}(x, y) = (\alpha + 2)/d$ was previously established in [Smith 2014, Theorem 6.3] for general graphs with perfect matchings between N_x and N_y . In the setting of strongly regular graphs Theorem 1.2 is thus an extension of the results of [Smith 2014] to the case of maximum matchings. These results give

insight into the importance of understanding matchings between the neighbor sets N_x and N_y in the derivation of exact Ricci curvature formulas. In particular, we are able to exploit this idea to derive the explicit formulas for the Ricci curvature of strongly regular graphs of girths 4 and 5 using elementary means.

For strongly regular graphs $G = (V, E)$ of girth 5, it follows that adjacent vertices share $\alpha = 0$ common neighbors. Moreover, for a given edge $xy \in E$, there are no edges between the neighbor sets N_x and N_y . Then, as a simple consequence of Theorem 1.2, we have the following Ricci curvature formula.

Theorem 1.4. *Suppose $G = (V, E)$ is a strongly regular graph with parameters (n, d, α, β) . If the girth of G is 5, then the condensed Ricci curvature satisfies*

$$\mathbb{k}(x, y) = \frac{3}{d} - 1$$

for all $xy \in E$.

From Theorem 1.4 it is easy to see that the condensed Ricci curvature along any edge of the 5-cycle C_5 is $\mathbb{k} = \frac{1}{2}$, while the condensed Ricci curvature along any edge of the Peterson graph is $\mathbb{k} = 0$. The fact that these are the only nonnegatively curved strongly regular graphs of girth 5 then follows from Theorem 1.4 and the classification of Moore graphs of diameter 2 due to [Hoffman and Singleton 1960].

Corollary 1.5. *The 5-cycle and the Petersen graph are the only strongly regular graphs of girth 5 with nonnegative condensed Ricci curvature along edges.*

For strongly regular graphs $G = (V, E)$ of girth 4, it follows that adjacent vertices share $\alpha = 0$ common neighbors. For a given edge $xy \in E$, we appeal to Hall's theorem and the pigeonhole principle to establish a perfect matching between the neighbor sets N_x and N_y . Then, as a consequence of Theorem 1.2, we have the following Ricci curvature formula.

Theorem 1.6. *Suppose $G = (V, E)$ is a strongly regular graph with parameters (n, d, α, β) . If the girth of G is 4, then the condensed Ricci curvature satisfies*

$$\mathbb{k}(x, y) = \frac{2}{d}$$

for all $xy \in E$. In particular, all strongly regular graphs of girth 4 have positive condensed Ricci curvature.

For strongly regular graphs with girth 3, it turns out that there does not exist an exact formula for the condensed Ricci curvature solely in terms of the graph parameters. This is due to the fact that there are nonisomorphic girth-3 strongly regular graphs with the same graph parameters but different condensed Ricci curvatures. For example, the 4×4 rook's graph and the Shrikhande graph both have parameters $(16, 6, 2, 2)$ but their condensed Ricci curvatures are $\mathbb{k} = \frac{2}{3}$ and $\mathbb{k} = \frac{1}{3}$, respectively. This is due to the fact that for a given edge xy there is a

perfect matching between neighbor sets N_x and N_y for the rook's graph but the maximum matching between N_x and N_y is size $m = 1$ for the Shrikhande graph. For comparison, we would like to point out that the 4×4 rook's graph and the Shrikhande graph were given as an example in [Cushing et al. 2020] to show that Bakry–Émery Ricci curvature functions are not uniquely determined by strongly regular graph parameters.

On the other hand, strongly regular conference graphs exhibit many symmetry properties that we believe always lead to perfect matchings between neighbor sets N_x and N_y . We plan to address conference graphs in a forthcoming paper. For now we make the following conjecture for conference graphs, which can be realized as a direct consequence of Theorem 1.2 provided there are always perfect matchings between N_x and N_y .

Conjecture 1.7. *Let $G = (V, E)$ be a strongly regular conference graph with parameters $(4\beta + 1, 2\beta, \beta - 1, \beta)$, with $\beta \geq 2$. Then the condensed Ricci curvature satisfies*

$$\mathbb{k}(x, y) = \frac{1}{2} + \frac{1}{2\beta}$$

for all $xy \in E$. In particular, all strongly regular conference graphs have positive condensed Ricci curvature strictly greater than $\frac{1}{2}$.

This paper is organized as follows: In Section 2 we formally define transport distance between probability measures on graphs and the condensed Ricci curvature. We briefly discuss our choice of notation and terminology and we state some key results from [Bourne et al. 2018; Bhattacharya and Mukherjee 2015] that simplify the computation of condensed Ricci curvature. In Section 3 we introduce the spectral graph Laplacian and prove Theorem 1.1. In Section 4 we focus on strongly regular graphs and establish our exact curvature formulas.

2. Transport distance and condensed Ricci curvature on graphs

To motivate Ollivier's formulation of coarse Ricci curvature we turn to the setting of smooth Riemannian manifolds. Suppose that x_1 and x_2 are points on an N -dimensional Riemannian manifold M and that m_i is the uniform measure on the respective geodesic ball $B(x_i, \varepsilon)$ of radius ε centered at x_i . Let $\delta = d(x_1, x_2)$ denote the geodesic distance between x_1 and x_2 . If $B(x_1, \varepsilon)$ is parallel transported to $B(x_2, \varepsilon)$ along a geodesic from x_1 to x_2 in the direction of tangent vector v at x_1 , then the transport distance $W(m_1, m_2)$ between the measures and the Ricci curvature on M satisfy the relation [Ollivier 2009]

$$W(m_1, m_2) = \delta \left(1 - \frac{\varepsilon^2}{2(N+2)} \text{Ric}(v, v) + O(\varepsilon^3 + \varepsilon^2\delta) \right) \quad \text{as } \varepsilon, \delta \rightarrow 0.$$

As noted in [Ollivier 2010], this relation allows one to interpret Ricci curvature on Riemannian manifolds as a measure of the average distance traveled between points of geodesic balls versus their centers under parallel transport. Using this relationship to guide intuition, Ollivier defined the Ricci curvature of Markov chains on metric spaces as

$$\text{Ric}(x_1, x_2) = 1 - \frac{W(m_1, m_2)}{d(x_1, x_2)},$$

where $W(m_1, m_2)$ denotes the general transport or Wasserstein distance between measures m_i based at x_i and $d(x_1, x_2)$ is the metric distance between x_1 and x_2 . This definition provides a synthetic notion of Ricci curvature of Markov chains on metric spaces and its geometric validation comes in the form of analogues of the classical theorems of Bonnet–Myers, Lichnerowicz, and Gromov–Lévy [Ollivier 2009; 2010].

Ollivier’s coarse Ricci curvature on graphs is defined in terms of transport distance of probability measures and standard graph distance [Ollivier 2009; 2010]. Let $G = (V, E)$ be a nontrivial, locally finite, undirected, connected, simple graph with standard graph shortest-path distance function

$$\rho : V \times V \rightarrow \mathbb{N} \cup \{0\}.$$

A *probability measure* or *probability mass distribution* on $G = (V, E)$ is a real-valued function $\mu : V \rightarrow [0, 1]$ such that

$$\sum_{v \in V} \mu(v) = 1.$$

Given probability measures μ and ν on G , a *coupling* or *transport plan* between μ and ν is a probability measure $\xi : V \times V \rightarrow [0, 1]$ such that

$$\sum_{w \in V} \xi(v, w) = \mu(v) \quad \text{and} \quad \sum_{v \in V} \xi(v, w) = \nu(w).$$

The terminology transport plan has economic origins and comes from the optimal transport problem, where one interprets a coupling as a plan for transporting mass or goods from one distribution center to another.

The L^1 -Wasserstein or transport distance is a metric on the space of probability measures on G that quantifies the optimal *transport cost* between probability measures on G .

Definition 2.1. The L^1 -Wasserstein or transport distance between probability measures μ and ν on a graph $G = (V, E)$ is given by

$$W(\mu, \nu) = \inf_{\xi \in \chi(\mu, \nu)} \sum_{x \in V} \sum_{y \in V} \rho(x, y) \xi(x, y),$$

where $\chi(\mu, \nu)$ is the set of all transport plans between μ and ν .

We consider a modified notion of Ollivier's coarse Ricci curvature on graphs introduced by Lin, Lu, and Yau [Lin et al. 2011] that is based on probability measures of the form

$$m_x^\varepsilon(y) = \begin{cases} 1 - \varepsilon & y = x, \\ \varepsilon/\deg(x) & y \in \Gamma(x), \\ 0 & \text{otherwise.} \end{cases} \quad (2-1)$$

For $x, y \in V$ the ε -Ollivier Ricci curvature is then defined for $0 \leq \varepsilon \leq 1$ by

$$\kappa_\varepsilon(x, y) = 1 - \frac{W(m_x^\varepsilon, m_y^\varepsilon)}{\rho(x, y)}$$

and the modified Ollivier Ricci curvature is then defined for $x, y \in V$ by

$$\mathbb{k}(x, y) = \lim_{\varepsilon \rightarrow 0} \frac{\kappa_\varepsilon(x, y)}{\varepsilon}.$$

We refer to the modified Ricci curvature as the condensed Ricci curvature. Our definitions differ slightly from those of [Lin et al. 2011] in that we use $1 - \varepsilon$ in place of α as in [Smith 2014; Münch and Wojciechowski 2019]. The reason for this choice in notation and terminology is that we interpret the probability measure (2-1) as encoding an ε -active random walk that becomes stationary as $\varepsilon \rightarrow 0$. One can view this phenomena as a concentration or condensation of measure. This formulation also more closely resembles the notion of coarse Ricci curvature of continuous-time Markov chains given by the derivative

$$\kappa(x, y) = -\frac{d}{dt} \frac{W(m_x^t, m_y^t)}{\rho(x, y)}.$$

Computing Ricci curvature on graphs requires solving the optimal transport problem. By definition a given transport plan between measures m_x^ε and m_y^ε gives an upper bound for the transport distance $W(m_x^\varepsilon, m_y^\varepsilon)$ and therefore a lower bound for condensed Ricci curvature. While an optimal transport plan may be intuitive, realizing that it achieves the transport distance directly from the definition is technically challenging as it requires solving a linear programming problem. Fortunately, there is a dual formulation of this optimization problem in terms of 1-Lipschitz functions that one can use to provide lower bounds to transport distance and consequently upper bounds for condensed Ricci curvature.

Theorem 2.2 (Kantorovich duality theorem). *The L^1 -Wasserstein or transport distance between probability measures μ and ν on a graph $G = (V, E)$ is given by*

$$W(\mu, \nu) = \sup_{f \in \text{Lip}(1)} \sum_{z \in V} f(z)(\mu(z) - \nu(z)),$$

where

$$\text{Lip}(1) = \{f : V \rightarrow \mathbb{R} \mid |f(x) - f(y)| \leq \rho(x, y) \text{ for all } x, y \in V\}$$

is the space of 1-Lipschitz functions on G .

Due to the Kantorovich duality theorem we refer to 1-Lipschitz functions on a graph $G = (V, E)$ as Kantorovich potential functions. With the Kantorovich duality theorem in hand one can compute exact formulas for Ricci curvature by specifying a transport plan and a Kantorovich potential that provide bounds that pinch the condensed Ricci curvature and force equality. Another useful fact for computing condensed Ricci curvature due to [Bourne et al. 2018] is that the Ricci activeness function $\varepsilon \mapsto \mathbb{k}_\varepsilon$ is piecewise linear with two linear parts on regular graphs of degree d and the ε -Ollivier Ricci curvature differs from the condensed Ricci curvature only by scale for all values $0 \leq \varepsilon \leq d/(d+1)$. More precisely, on a regular graph $G = (V, E)$ of degree d , for any $x, y \in V$,

$$\kappa_\varepsilon(x, y) = \varepsilon \mathbb{k}(x, y) \quad \text{for } 0 \leq \varepsilon \leq \frac{d}{d+1}.$$

For simplicity, in our computations we take $\varepsilon = \frac{1}{2}$ and compute the condensed Ricci curvature as

$$\mathbb{k}(x, y) = 2\kappa_{1/2}(x, y).$$

In Riemannian geometry, Ricci curvature is a local quantity so in the context of graphs it seems natural to restrict one's attention to computing Ricci curvature along edges. With our conventions, this restriction allows for further simplification in computing Ricci curvature due to [Bhattacharya and Mukherjee 2015]. For vertices $x \in V$ let

$$\Gamma(x) = \{y \in V \mid \rho(x, y) = 1\} = \{y \in V \mid xy \in E\}$$

denote the neighbor set of x . Given an edge $xy \in E$ we denote the set of common neighbors or *triangle set* of x and y by

$$\nabla_{xy} = \Gamma(x) \cap \Gamma(y).$$

Then we further decompose the 1-step neighborhoods of x and y into disjoint unions

$$\Gamma(x) = N_x \cup \nabla_{xy} \cup \{y\} \quad \text{and} \quad \Gamma(y) = N_y \cup \nabla_{xy} \cup \{x\}$$

where

$$N_x = \Gamma(x) \setminus (\nabla_{xy} \cup \{y\}) \quad \text{and} \quad N_y = \Gamma(y) \setminus (\nabla_{xy} \cup \{x\}).$$

We denote the set of common 2-step neighbors or *pentagon set* of x and y by

$$P_{xy} = \{z \in V \mid \rho(x, z) = 2 \text{ and } \rho(y, z) = 2\}.$$

As in [Bhattacharya and Mukherjee 2015], we refer to the disjoint union of vertices

$$\mathcal{N}_{xy} = \{x\} \cup \{y\} \cup \nabla_{xy} \cup N_x \cup N_y \cup P_{xy} \quad (2-2)$$

as the *core neighborhood* of an edge $xy \in E$. It is immediate that the core neighborhood represents a partition of all vertices with distance less than or equal to 2 from x and y . Moreover, the reduction lemma [Bhattacharya and Mukherjee 2015, Lemma 2.3] shows that when computing the condensed Ricci curvature of an edge $xy \in E$, it is sufficient to consider only the induced subgraph of G lying within the core neighborhood \mathcal{N}_{xy} .

3. A rigidity theorem for complete graphs

In this section we establish a rigidity theorem for complete graphs that shows that a complete graph is the only graph with condensed Ricci curvature strictly greater than 1. Given a connected finite simple graph $G = (V, E)$ with n vertices v_1, \dots, v_n , the adjacency matrix A of G is a symmetric $n \times n$ matrix with entries

$$\begin{cases} a_{ij} = 1 & \text{if } v_i v_j \in E, \\ a_{ij} = 0 & \text{otherwise.} \end{cases}$$

Let d_i denote the degree of vertex v_i and let $D = \text{diag}(d_1, \dots, d_n)$ be the $n \times n$ diagonal matrix of degrees. The spectral graph Laplacian is then defined as

$$L = I - D^{-1/2} A D^{-1/2}.$$

With these conventions it follows that the eigenvalues of the spectral graph Laplacian are nondecreasing with

$$0 = \lambda_0 \leq \lambda_1 \leq \dots \leq \lambda_{n-1},$$

where the multiplicity of the 0 eigenvalue is the number of connected components of G . Hence, a finite graph is connected if and only if $\lambda_1 > \lambda_0 = 0$. Similar to the spectrum of the metric Laplacian on Riemannian manifolds, the spectrum of the spectral Laplacian on graphs encodes geometric and structural properties of a graph.

In the context of Riemannian geometry the Lichnerowicz theorem [1958] is a widely celebrated theorem that provides a lower bound for the first nonzero eigenvalue of the metric Laplacian when the Ricci curvature has a strict positive lower bound. More precisely, if (M^n, g) is an n -dimensional compact Riemannian manifold with

$$\text{Ric}_g \geq (n-1)kg$$

for some positive constant $k > 0$, then the first positive eigenvalue of the metric Laplacian $\lambda_1 \geq nk$. The graph analogue of the Lichnerowicz theorem similarly shows that a positive lower bound on condensed Ricci curvature serves as a lower bound on the first nonzero eigenvalue of the spectral graph Laplacian.

Theorem 3.1 [Lin et al. 2011, Theorem 4.2]. *Suppose $G = (V, E)$ is a connected finite simple graph with condensed Ricci curvature*

$$\mathbb{k}(x, y) \geq \kappa_0 > 0$$

for some positive constant κ_0 and all edges $xy \in E$. Then the first nonzero eigenvalue of the spectral graph Laplacian satisfies $\lambda_1 \geq \kappa_0$.

In order to establish a rigidity theorem for complete graphs we first record some well-known results on the first nonzero eigenvalue of the spectral graph Laplacian.

Theorem 3.2 [Chung 1997, Lemma 1.7]. *Let $G = (V, E)$ be a simple graph on n vertices. Then*

$$\lambda_1 \leq \frac{n}{n-1}$$

with equality if and only if G is complete. Moreover, complete graphs are the only graphs with $\lambda_1 > 1$.

We now prove the following rigidity theorem through direct computation and the aid of Theorems 3.1 and 3.2.

Theorem 3.3. *A connected finite simple graph $G = (V, E)$ is complete if and only if the condensed Ricci curvature satisfies $\mathbb{k}(x, y) > 1$ for all edges $xy \in E$.*

Proof. Let $G = (V, E)$ be the complete graph on n vertices and let $xy \in E$. Then the triangle set ∇_{xy} is equal to $V \setminus \{x, y\}$ since G is complete. Let m_x and m_y denote $\varepsilon = \frac{1}{2}$ -probability measures based at x and y , respectively, defined as in (2-1). Consider the transport plan $\xi : V \times V \rightarrow [0, 1]$ between m_x and m_y given by

$$\xi(v, w) = \begin{cases} \frac{1}{2} - \frac{1}{2(n-1)}, & v = x, w = y, \\ \frac{1}{2(n-1)}, & v = w, \\ 0, & \text{otherwise,} \end{cases}$$

which transports excess mass directly from x to y and leaves the remainder of the distribution fixed. Then by Definition 2.1 the transport distance satisfies

$$W(m_x, m_y) \leq \sum_{v \in V} \sum_{w \in V} \rho(v, w) \xi(v, w) = \xi(x, y) = \frac{1}{2} - \frac{1}{2(n-1)}$$

and therefore the condensed Ricci curvature satisfies

$$\mathbb{k}(x, y) = 2\kappa_{1/2}(x, y) \geq 2\left(1 - \left(\frac{1}{2} - \frac{1}{2(n-1)}\right)\right) = 1 + \frac{1}{n-1} > 1. \quad (3-1)$$

Conversely, let $G = (V, E)$ be a connected finite simple graph with condensed Ricci curvature $\mathbb{k}(x, y) > 1$ for all edges $xy \in E$. Then by the graph Lichnerowicz theorem (Theorem 3.1) of [Lin et al. 2011], it follows $\lambda_1 > 1$ and therefore G is complete by Theorem 3.2. \square

Remark 3.4. Using the 1-Lipschitz function $f : V \rightarrow \mathbb{R}$ defined by

$$f(v) = \begin{cases} 1, & v = x, \\ 0, & \text{otherwise,} \end{cases}$$

it follows from the Kantorovich duality theorem (Theorem 2.2) that the transport distance satisfies

$$W(m_x, m_y) \geq \sum_{v \in V} f(v)(m_x(v) - m_y(v)) = m_x(x) - m_y(x) = \frac{1}{2} - \frac{1}{2(n-1)}$$

and therefore the condensed Ricci curvature satisfies

$$\mathbb{k}(x, y) = 2\kappa_{1/2}(x, y) \leq 2\left(1 - \left(\frac{1}{2} - \frac{1}{2(n-1)}\right)\right) = 1 + \frac{1}{n-1}. \quad (3-2)$$

Hence, from (3-1) and (3-2) we see that the condensed Ricci curvature of the complete graph on n vertices is

$$\mathbb{k}(x, y) = 1 + \frac{1}{n-1} = \frac{n}{n-1}$$

for all edges $xy \in E$ as stated in [Lin et al. 2011, Example 1].

4. Strongly regular graphs

In this section we derive an explicit formula for the condensed Ricci curvature of strongly regular graphs $G = (V, E)$ in terms of the graph parameters and the size of a maximum matching in the core neighborhood. Recall that a simple, undirected, finite graph $G = (V, E)$ is said to be a strongly regular graph with parameters (n, d, α, β) if G has n vertices of constant degree d where any two adjacent vertices share $\alpha \geq 0$ common neighbors and any two nonadjacent vertices share $\beta \geq 1$ common neighbors. Given an edge $xy \in E$, we consider the core neighborhood decomposition (2-2) given by

$$\mathcal{N}_{xy} = \{x\} \cup \{y\} \cup \nabla_{xy} \cup N_x \cup N_y \cup P_{xy}$$

that was introduced in [Bhattacharya and Mukherjee 2015] and where the neighbor sets satisfy

$$N_x = \Gamma(x) \setminus (\nabla_{xy} \cup \{y\}) \quad \text{and} \quad N_y = \Gamma(y) \setminus (\nabla_{xy} \cup \{x\}).$$

Since strongly regular graphs have diameter 2, it follows that $V = \mathcal{N}_{xy}$. Hence, for strongly regular graphs the induced subgraph lying within the core neighborhood of any edge accounts for the entire graph G . Moreover, from strong regularity we see $|\nabla_{xy}| = \alpha$ so that

$$|N_x| = |N_y| = d - \alpha - 1 \quad \text{and} \quad |P_{xy}| = n - 2d - \alpha.$$

Our strategy in the derivation of an explicit formula for the condensed Ricci curvature of strongly regular graphs reformulates the problem of finding candidates for an optimal transport plan and an optimal Kantorovich potential into a maximum matching problem.

Definition 4.1. A *matching* on a finite undirected simple graph $G = (V, E)$ is a subset of edges $\mathcal{M} \subseteq E$ such that no vertex in V is incident to more than one edge in \mathcal{M} . The size of a matching \mathcal{M} on G is the total number of edges in \mathcal{M} . A matching \mathcal{M} on G is said to be a *maximum matching* if it contains the largest possible number of edges. A maximum matching between disjoint subsets of vertices $S, T \subset V$ is said to be a *perfect matching* if every vertex in S and T is incident to exactly one edge of the matching.

Let $G = (V, E)$ be a strongly regular graph with edge $xy \in E$. Given a maximum matching \mathcal{M} between the neighbor sets N_x and N_y , we denote the vertices that are matched in N_x and N_y by M_x and M_y , respectively. If \mathcal{M} is not a perfect matching, we denote the remaining unmatched vertices in N_x and N_y by $U_x = N_x \setminus M_x$ and $U_y = N_y \setminus M_y$, respectively. With this decomposition of N_x and N_y into matched and unmatched vertices, it is fairly straightforward to develop a candidate for an optimal transport plan. However, developing a candidate for an optimal Kantorovich potential requires a deeper understanding of the incidence relations between edges in a given maximum matching and the remaining edges between N_x and N_y .

Definition 4.2. Let \mathcal{M} be a matching on an undirected simple graph $G = (V, E)$. An *alternating path* P in G is a path $P = v_0 v_1 \cdots v_k$ such that the initial vertex v_0 is unmatched and $v_i v_{i+1} \in \mathcal{M}$ if and only if $v_{i-1} v_i \notin \mathcal{M}$ for all $0 < i < k$. An *augmenting path* P in G is an alternating path $P = v_0 v_1 \cdots v_k$ such that the terminal vertex v_k is also unmatched.

We refer to a single vertex as a trivial path. Hence, with this convention any unmatched vertex is by itself a trivial augmenting path. One of the main tools in our construction of a candidate for an optimal Kantorovich potential is Berge's lemma [1957].

Lemma 4.3 (Berge's lemma). *A matching \mathcal{M} on an undirected simple graph $G = (V, E)$ is a maximum matching if and only if G contains no nontrivial augmenting paths.*

Given a matching \mathcal{M} on a graph $G = (V, E)$ and a subset of vertices $U \subseteq V$, we denote the subset of matched vertices in U by $\mathcal{M}(U)$. For a subset of vertices $S \subseteq V$ we denote the subset of alternating paths with initial vertex in S by A_S . For subsets of vertices $S, T \subseteq V$ we denote the set of vertices in T that lie along an alternating path initiated in S by $A_S(T)$. We use the contrapositive of the following lemma to show that certain vertices in the neighbor sets N_x and N_y have no edges

between them. Ultimately, this allows us to ensure our candidate for an optimal Kantorovich potential is 1-Lipschitz.

Lemma 4.4. *Let $H = (V, E)$ be a finite, undirected, bipartite graph with parts S, T . Let \mathcal{M} be a maximum matching on H and suppose $v \in A_S(S)$ and $w \in T$. If $vw \in E$, then $w \in A_S(T)$.*

Proof. First note that since H is bipartite, it follows that an alternating path initiated in S can only terminate in an unmatched vertex in T . Therefore, if $v \in A_S(S)$ is an unmatched vertex, then v must be the initial vertex of an alternating path. But then w must be a matched vertex so that $w \in A_S(T)$ since otherwise $vw \in E$ is an augmenting path, contradicting Berge's lemma as \mathcal{M} is a maximum matching. Now assume that v is a matched vertex and that $P = v_0v_1 \cdots v_iv$ is an alternating path initiated at $v_0 \neq v \in S$ that terminates at $v \in S$. Then since $v_0 \in S$ is an unmatched vertex and H is bipartite, it follows that $v_iv \in \mathcal{M}$ since P is an alternating path. But then $vw \notin \mathcal{M}$ since otherwise v is incident to more than one edge in \mathcal{M} . But then the concatenation $P' = v_0v_1 \cdots v_ivw$ is an alternating path so that $w \in A_S(T)$. \square

We use this final lemma to compute a lower bound for condensed Ricci curvature from our Kantorovich potential in terms of the number of vertices in N_x and N_y along alternating paths initiated in N_y .

Lemma 4.5. *Let $H = (V, E)$ be a finite, undirected, bipartite graph with parts S, T . Suppose \mathcal{M} is a maximum matching of size m on H . Then*

$$|A_S(S)| = |A_S(T)| + |S| - m.$$

Proof. Let $v \in A_S(T)$ and suppose $P = v_0v_1 \cdots v_iv$ is an alternating path initiated at $v_0 \in S$ that terminates at $v \in T$. Then since $v_0 \in S$ is an unmatched vertex and H is bipartite, it follows that $v_iv \notin \mathcal{M}$ since P is an alternating path. If v is an unmatched vertex, then P is an augmenting path contrary to Berge's lemma as \mathcal{M} is a maximum matching. Hence, v must be a matched vertex and therefore since matched vertices are incident to a single edge in the matching, it follows that there exists a unique vertex $w \in S$ such that $vw \in \mathcal{M}$. But then the concatenation $P' = v_0v_1 \cdots vw$ is an alternating path initiated in S that terminates at the matched vertex $w \in \mathcal{M}(S)$. Hence, for each $v \in A_S(T)$ there is a unique $w \in A_S(S) \cap \mathcal{M}(S)$ with $vw \in \mathcal{M}$.

On the other hand, given $w \in A_S(S) \cap \mathcal{M}(S)$ suppose $P = w_0w_1 \cdots w_iw$ is an alternating path initiated in S that terminates at $w \in S$. Then since w is a matched vertex by assumption, it is clear that P is a nontrivial path and that $w_i \in T$ is the unique matched vertex with $w_iw \in \mathcal{M}$. But then $P' = w_0w_1 \cdots w_i$ is an alternating path initiated in S that terminates at $w_i \in T$ so that $w_i \in A_S(T)$. Hence, for each $w \in A_S(S) \cap \mathcal{M}(S)$ there is a unique $v \in A_S(T)$ with $vw \in \mathcal{M}$.

Thus, since $A_S(T) \subseteq \mathcal{M}(T)$, we have a bijection between the subsets $A_S(T)$ and $A_S(S) \cap \mathcal{M}(S)$ and therefore

$$|A_S(T)| = |A_S(S) \cap \mathcal{M}(S)|.$$

Noting $S \setminus \mathcal{M}(S) \subseteq A_S(S)$ since under our conventions all unmatched vertices in S are trivial alternating paths initiated in S , it follows

$$\begin{aligned} |A_S(T)| &= |A_S(S) \cap \mathcal{M}(S)| = |A_S(S) \setminus (S \setminus \mathcal{M}(S))| \\ &= |A_S(S)| - |S \setminus \mathcal{M}(S)| = |A_S(S)| - (|S| - m). \end{aligned}$$

Hence,

$$|A_S(S)| = |A_S(T)| + |S| - m$$

as desired. \square

We are now in a position to derive an explicit formula for condensed Ricci curvature of an edge $xy \in E$ in terms of the graph parameters and the size of a maximum matching between the neighbor sets N_x and N_y .

Theorem 4.6. *Suppose $G = (V, E)$ is a strongly regular graph with parameters (n, d, α, β) , and $xy \in E$ with maximum matching \mathcal{M} of size m between N_x and N_y . Then the condensed Ricci curvature satisfies*

$$\mathbb{k}(x, y) = \frac{\alpha + 2}{d} - \frac{|N_x| - m}{d}.$$

Proof. Let $xy \in E$ and consider the core neighborhood decomposition of G as in (2-2). Let H denote the induced bipartite subgraph consisting of all edges in E between vertices in N_x and N_y . Suppose that \mathcal{M} is a maximum matching on H of size $|\mathcal{M}| = m$. Assuming \mathcal{M} is not a perfect matching, as above we denote the matched and unmatched vertices in N_x and N_y by M_x, M_y and U_x, U_y , respectively. Note that $|U_x| = |U_y|$ since $|N_x| = |N_y|$ and $|M_x| = |M_y|$. For $i = 1, \dots, |N_x| - m$, let $x_i'' \in N_x$ and $y_i'' \in N_y$ denote the unmatched vertices in H and for $j = 1, \dots, m$, let $x_j^m \in N_x$ and $y_j^m \in N_y$ denote the matched vertices in H . Up to reordering of indices we may assume $x_i^m y_i^m \in \mathcal{M}$ for $i = 1, \dots, m$.

Clearly, there are no edges in E shared by two unmatched vertices since \mathcal{M} is a maximum matching so $x_i'' y_j'' \notin E$ for all $1 \leq i, j \leq |N_x| - m$. But then $\rho(x_i'', y_i'') = 2$ in G since x_i'' and y_i'' are not adjacent and G has diameter 2. Hence, for $1 \leq i \leq |N_x| - m$, we may pair unmatched vertices x_i'' with y_i'' along 2-step paths in G . For simplicity in notation write $x_i'' y_i'' \in \mathcal{M}^2$ to denote unmatched vertices $x_i'' \in U_x$ and $y_i'' \in U_y$ that have been paired along a 2-step path. Together these matchings/pairings induce a transport plan that moves excess mass from x to y and that moves mass from $x_i^m \in M_x, x_i'' \in U_x$ directly to their matched/paired neighbors $y_i^m \in M_y, y_i'' \in U_y$ along 1-step and 2-step paths, respectively. More precisely,

consider the transport plan defined by

$$\xi(v, w) = \begin{cases} \frac{1}{2} - \frac{1}{2d}, & v = x, w = y, \\ \frac{1}{2d}, & v = w \in \nabla_{xy} \cup \{x\} \cup \{y\}, \\ \frac{1}{2d}, & vw \in \mathcal{M}, \\ \frac{1}{2d}, & vw \in \mathcal{M}^2, \\ 0, & \text{otherwise.} \end{cases} \quad (4-1)$$

Noting that $|N_x| = |N_y| = d - \alpha - 1$ it follows that the transport distance satisfies

$$\begin{aligned} W(m_x, m_y) &\leq \sum_{v \in V} \sum_{w \in V} \rho(v, w) \xi(v, w) \\ &= \left(\frac{1}{2} - \frac{1}{2d} \right) + m \cdot \frac{1}{2d} + (|N_x| - m) \cdot 2 \cdot \frac{1}{2d} \\ &= \frac{1}{2} + \frac{1}{2d} ((d - \alpha - 2) + (|N_x| - m)). \end{aligned}$$

Thus, the condensed Ricci curvature satisfies

$$\begin{aligned} \mathbb{k}(x, y) &= 2\kappa_{1/2}(x, y) \\ &\geq 2 \left(1 - \left(\frac{1}{2} + \frac{1}{2d} ((d - \alpha - 2) + (|N_x| - m)) \right) \right) \\ &= \frac{\alpha + 2}{d} - \frac{|N_x| - m}{d}. \end{aligned}$$

Next we define a Kantorovich potential function $f : V \rightarrow \mathbb{R}$ with respect to the matching \mathcal{M} by

$$f(z) = \begin{cases} 1, & z = x, \\ 1, & z \in N_x \setminus A_{N_y}(N_x), \\ -1, & z \in A_{N_y}(N_y), \\ 0, & \text{otherwise,} \end{cases} \quad (4-2)$$

where $A_{N_y}(N_x)$ and $A_{N_y}(N_y)$ denote the vertices in N_x and N_y , respectively, that lie along alternating paths initiated in N_y . Taking $T = N_x$ and $S = N_y$ in Lemma 4.4, it follows that there are no edges between $N_x \setminus A_{N_y}(N_x)$ and $A_{N_y}(N_y)$. Hence, f is a 1-Lipschitz function on V .

Let $a_x = |A_{N_y}(N_x)|$ and $a_y = |A_{N_y}(N_y)|$. Then

$$|N_x \setminus A_{N_y}(N_x)| = |N_x| - a_x = d - \alpha - 1 - a_x$$

and by Lemma 4.5

$$a_y = |A_{N_y}(N_y)| = |A_{N_y}(N_x)| + |N_y| - m = a_x + |N_x| - m.$$

Therefore, by the Kantorovich duality theorem, we find that the transport distance satisfies

$$\begin{aligned} W(m_x, m_y) &\geq \sum_{z \in V} f(z)(m_x(z) - m_y(z)) = \left(\frac{1}{2} - \frac{1}{2d}\right) + (d - \alpha - 1 - a_x) \cdot \frac{1}{2d} + a_y \cdot \frac{1}{2d} \\ &= \left(\frac{1}{2} - \frac{1}{2d}\right) + (d - \alpha - 1 - a_x) \frac{1}{2d} + (a_x + |N_x| - m) \frac{1}{2d} \\ &= \frac{1}{2} + \frac{1}{2d} ((d - \alpha - 2) + (|N_x| - m)). \end{aligned}$$

Similar to the calculation (4-2), we have

$$\mathbb{k}(x, y) = 2\kappa_{1/2}(x, y) \leq \frac{\alpha + 2}{d} - \frac{|N_x| - m}{d}. \quad (4-3)$$

Hence, from the inequalities (4-2) and (4-3), it follows

$$\mathbb{k}(x, y) = \frac{\alpha + 2}{d} - \frac{|N_x| - m}{d}.$$

Now if \mathcal{M} is a perfect matching between N_x and N_y , then the size of \mathcal{M} is by definition $m = |N_x| = |N_y| = d - \alpha - 1$. Moreover, since all vertices in N_x and N_y are matched by some edge in \mathcal{M} , it follows that there are no alternating paths initiated in N_x or N_y in the induced bipartite subgraph between vertices in N_x and N_y . But then $A_{N_y}(N_x) = \emptyset$ and $A_{N_x}(N_y) = \emptyset$. Taking all of this into account, we see that the same transport plan (4-1) coupled with the Kantorovich potential (4-2) yields the equality

$$\mathbb{k}(x, y) = \frac{\alpha + 2}{d}$$

in the case of a perfect matching between N_x and N_y . \square

Strongly regular graphs of girths 4 and 5. In this subsection we exploit the adjacency properties of strongly regular graphs $G = (V, E)$ to derive explicit formulas for the condensed Ricci curvature of strongly regular graphs of girths 4 and 5. Let $G = (V, E)$ be a strongly regular graph with parameters (n, d, α, β) . It is well known [Cameron and van Lint 1975] that if G has girth 5, then $\alpha = 0$ and $\beta = 1$, while a strongly regular graph G with girth 4 has $\alpha = 0$ and $\beta \geq 2$. The following theorem is an immediate consequence of Theorem 4.6 and the fact that the girth of a graph is its minimal cycle length.

Theorem 4.7. *Suppose $G = (V, E)$ is a strongly regular graph with parameters (n, d, α, β) . If the girth of G is 5, then the condensed Ricci curvature satisfies*

$$\mathbb{k}(x, y) = \frac{3}{d} - 1$$

for all edges $xy \in E$.

Proof. Let $xy \in E$ and consider the core neighborhood decomposition of G as in (2-2). Let $x_0 \in N_x$ and suppose that $x_0 y_0 \in E$ for some $y_0 \in N_y$. Then $x x_0 y_0 y x$ is a cycle of length 4, contradicting the fact that G has girth 5. Thus, there are no edges between N_x and N_y so the maximum matching between N_x and N_y has size $m = 0$. Therefore, from Theorem 4.6 with $\alpha = 0$ and $m = 0$, we find that the condensed Ricci curvature satisfies

$$\mathbb{k}(x, y) = \frac{3}{d} - 1. \quad \square$$

Remark 4.8. The transportation plan (4-1) and Kantorovich potential (4-2) can be used to compute this curvature formula directly. Since G has girth 5, there are no edges between vertices in N_x and N_y so $\mathcal{M} = \emptyset$ and each vertex in N_x can be paired to a vertex in N_y through a 2-step path. Moreover, if we consider the induced bipartite subgraph H between N_x and N_y , the collection of alternating paths in H that start in N_y consists solely of trivial paths so $A_{N_y}(N_y) = N_y$ and $A_{N_y}(N_x) = \emptyset$.

In the girth 4 case we appeal to Hall's theorem, which can be used to give necessary and sufficient conditions for perfect matchings on bipartite graphs. Let $H = (V, E)$ be a finite, bipartite graph with parts S, T . For any subset $W \subseteq S$, we define

$$\Gamma_T(W) = \{v \in T \mid vw \in E \text{ for some } w \in W\}$$

to be the collection of all vertices $v \in T$ sharing an edge with some vertex in W . More simply put, $\Gamma_T(W)$ is the neighbor set of W in T .

Theorem 4.9 (Hall's theorem). *Let $H = (V, E)$ be a finite, undirected, bipartite graph with parts S, T . Then there is a matching that covers S if and only if*

$$|\Gamma_T(W)| \geq |W|$$

for every subset of vertices $W \subseteq S$.

With Hall's theorem in hand we are able to establish a perfect matching between the neighbor sets N_x and N_y for strongly regular graphs with girth 4 as a consequence of the pigeonhole principle.

Theorem 4.10. *Suppose $G = (V, E)$ is a strongly regular graph with parameters (n, d, α, β) . If the girth of G is 4, then the condensed Ricci curvature satisfies*

$$\mathbb{k}(x, y) = \frac{2}{d}$$

for all edges $xy \in E$.

Proof. Let $xy \in E$ and consider the core neighborhood decomposition of G as in (2-2). Then since $\alpha = 0$, it follows that G is triangle free. Suppose that $x_i \in N_x$. Then since x_i is nonadjacent to y , it follows x_i and y have $\beta \geq 2$ common neighbors.

Clearly, one such neighbor is x so x_i has $\beta - 1 \geq 1$ neighbors in N_y since $\nabla_{xy} = \emptyset$. Similarly, each vertex $y_j \in N_y$ has $\beta - 1 \geq 1$ neighbors in N_x .

Now consider the induced bipartite subgraph $H = G[N_x \cup N_y]$ consisting of all edges in E between vertices in N_x and N_y . For sake of contradiction suppose that $X \subseteq N_x$ with $|X| = q > |\Gamma_{N_y}(X)| = k$. Since each vertex $x_i \in X \subseteq N_x$ has $\beta - 1 \geq 1$ neighbors in N_y , it follows that there are $(\beta - 1)q$ edges between X and $\Gamma_{N_y}(X)$. Therefore, by the generalized pigeonhole principle there exists some $w \in \Gamma_{N_y}(X)$ with

$$|\Gamma(w)| \geq \left\lceil \frac{(\beta - 1)q}{k} \right\rceil > \left\lceil \frac{(\beta - 1)q}{q} \right\rceil = \beta - 1$$

since $q > k$. This contradicts the fact that every vertex in N_y has exactly $\beta - 1$ neighbors in N_x . Thus, $|X| \leq |\Gamma_{N_y}(X)|$ for all subsets $X \subset N_x$ and therefore since $|N_x| = |N_y|$ it follows that there is a perfect matching in H by Hall's theorem. Noting that $\alpha = 0$, it follows from Theorem 4.6 with $\alpha = 0$ and $m = |N_x|$ that the condensed Ricci curvature satisfies

$$\mathbb{k}(x, y) = \frac{2}{d}. \quad \square$$

Acknowledgements

The authors were partially supported by NSF FURST grant DMS-1620552, the Cal Poly Frost Fund, CSU Fresno, and Cal Poly, San Luis Obispo. The authors would like to thank CSU Fresno for their hospitality during their summer participation in the FURST program.

References

- [Bauer et al. 2012] F. Bauer, J. Jost, and S. Liu, “Ollivier–Ricci curvature and the spectrum of the normalized graph Laplace operator”, *Math. Res. Lett.* **19**:6 (2012), 1185–1205. MR Zbl
- [Berge 1957] C. Berge, “Two theorems in graph theory”, *Proc. Nat. Acad. Sci. U.S.A.* **43** (1957), 842–844. MR Zbl
- [Bhattacharya and Mukherjee 2015] B. B. Bhattacharya and S. Mukherjee, “Exact and asymptotic results on coarse Ricci curvature of graphs”, *Discrete Math.* **338**:1 (2015), 23–42. MR Zbl
- [Bourne et al. 2018] D. P. Bourne, D. Cushing, S. Liu, F. Münch, and N. Peyerimhoff, “Ollivier–Ricci idleness functions of graphs”, *SIAM J. Discrete Math.* **32**:2 (2018), 1408–1424. MR Zbl
- [Cameron and van Lint 1975] P. J. Cameron and J. H. van Lint, *Graph theory, coding theory and block designs*, London Math. Soc. Lecture Note Series **19**, Cambridge University Press, 1975. MR Zbl
- [Chung 1997] F. R. K. Chung, *Spectral graph theory*, CBMS Regional Conference Series in Mathematics **92**, American Math. Soc., Providence, RI, 1997. MR Zbl
- [Cushing et al. 2020] D. Cushing, S. Liu, and N. Peyerimhoff, “Bakry–Émery curvature functions on graphs”, *Canad. J. Math.* **72**:1 (2020), 89–143. MR Zbl

- [Hoffman and Singleton 1960] A. J. Hoffman and R. R. Singleton, “On Moore graphs with diameters 2 and 3”, *IBM J. Res. Develop.* **4** (1960), 497–504. MR Zbl
- [Jost and Liu 2014] J. Jost and S. Liu, “Ollivier’s Ricci curvature, local clustering and curvature-dimension inequalities on graphs”, *Discrete Comput. Geom.* **51**:2 (2014), 300–322. MR Zbl
- [Lichnerowicz 1958] A. Lichnerowicz, *Géométrie des groupes de transformations*, Travaux et Recherches Mathématiques, III, Dunod, Paris, 1958. MR Zbl
- [Lin et al. 2011] Y. Lin, L. Lu, and S.-T. Yau, “Ricci curvature of graphs”, *Tohoku Math. J. (2)* **63**:4 (2011), 605–627. MR Zbl
- [Lin et al. 2014] Y. Lin, L. Lu, and S.-T. Yau, “Ricci-flat graphs with girth at least five”, *Comm. Anal. Geom.* **22**:4 (2014), 671–687. MR Zbl
- [Lott and Villani 2009] J. Lott and C. Villani, “Ricci curvature for metric-measure spaces via optimal transport”, *Ann. of Math. (2)* **169**:3 (2009), 903–991. MR Zbl
- [Münch and Wojciechowski 2019] F. Münch and R. K. Wojciechowski, “Ollivier Ricci curvature for general graph Laplacians: heat equation, Laplacian comparison, non-explosion and diameter bounds”, *Adv. Math.* **356** (2019), 106759, 45. MR Zbl
- [Ollivier 2009] Y. Ollivier, “Ricci curvature of Markov chains on metric spaces”, *J. Funct. Anal.* **256**:3 (2009), 810–864. MR Zbl
- [Ollivier 2010] Y. Ollivier, “A survey of Ricci curvature for metric spaces and Markov chains”, pp. 343–381 in *Probabilistic approach to geometry*, edited by M. Kotani et al., Adv. Stud. Pure Math. **57**, Math. Soc. Japan, Tokyo, 2010. MR Zbl
- [Smith 2014] J. D. H. Smith, “Ricci curvature, circulants, and a matching condition”, *Discrete Math.* **329** (2014), 88–98. MR Zbl
- [Sturm 2006] K.-T. Sturm, “On the geometry of metric measure spaces, I”, *Acta Math.* **196**:1 (2006), 65–131. MR Zbl

Received: 2019-08-26

Revised: 2020-03-11

Accepted: 2020-08-06

vbonini@calpoly.edu

Mathematics Department, California Polytechnic State University, San Luis Obispo, CA, United States

ccarro07@calpoly.edu

Mathematics Department, California Polytechnic State University, San Luis Obispo, CA, United States

udinh@calpoly.edu

Mathematics Department, California Polytechnic State University, San Luis Obispo, CA, United States

sydney.dye24@gmail.com

Mathematics Department, California Polytechnic State University, San Luis Obispo, CA, United States

jmfreder@calpoly.edu

Mathematics Department, California Polytechnic State University, San Luis Obispo, CA, United States

epearse@calpoly.edu

Mathematics Department, California Polytechnic State University, San Luis Obispo, CA, United States

involve

msp.org/involve

INVOLVE YOUR STUDENTS IN RESEARCH

Involve showcases and encourages high-quality mathematical research involving students from all academic levels. The editorial board consists of mathematical scientists committed to nurturing student participation in research. Bridging the gap between the extremes of purely undergraduate research journals and mainstream research journals, *Involve* provides a venue to mathematicians wishing to encourage the creative involvement of students.

MANAGING EDITOR

Kenneth S. Berenhaut Wake Forest University, USA

BOARD OF EDITORS

Colin Adams	Williams College, USA	Robert B. Lund	Clemson University, USA
Arthur T. Benjamin	Harvey Mudd College, USA	Gaven J. Martin	Massey University, New Zealand
Martin Bohner	Missouri U of Science and Technology, USA	Mary Meyer	Colorado State University, USA
Amarjit S. Budhiraja	U of N Carolina, Chapel Hill, USA	Frank Morgan	Williams College, USA
Pietro Cerone	La Trobe University, Australia	Mohammad Sal Moslehian	Ferdowsi University of Mashhad, Iran
Scott Chapman	Sam Houston State University, USA	Zuhair Nashed	University of Central Florida, USA
Joshua N. Cooper	University of South Carolina, USA	Ken Ono	Univ. of Virginia, Charlottesville
Jem N. Corcoran	University of Colorado, USA	Yuval Peres	Microsoft Research, USA
Toka Diagana	University of Alabama in Huntsville, USA	Y.-F. S. Pétermann	Université de Genève, Switzerland
Michael Dorff	Brigham Young University, USA	Jonathon Peterson	Purdue University, USA
Sever S. Dragomir	Victoria University, Australia	Robert J. Plemmons	Wake Forest University, USA
Joel Foisy	SUNY Potsdam, USA	Carl B. Pomerance	Dartmouth College, USA
Errin W. Fulp	Wake Forest University, USA	Vadim Ponomarenko	San Diego State University, USA
Joseph Gallian	University of Minnesota Duluth, USA	Bjorn Poonen	UC Berkeley, USA
Stephan R. Garcia	Pomona College, USA	József H. Przytycki	George Washington University, USA
Anant Godbole	East Tennessee State University, USA	Richard Rebarber	University of Nebraska, USA
Ron Gould	Emory University, USA	Robert W. Robinson	University of Georgia, USA
Sat Gupta	U of North Carolina, Greensboro, USA	Javier Rojo	Oregon State University, USA
Jim Haglund	University of Pennsylvania, USA	Filip Saidak	U of North Carolina, Greensboro, USA
Johnny Henderson	Baylor University, USA	Hari Mohan Srivastava	University of Victoria, Canada
Glenn H. Hurlbert	Virginia Commonwealth University, USA	Andrew J. Sterge	Honorary Editor
Charles R. Johnson	College of William and Mary, USA	Ann Trenk	Wellesley College, USA
K. B. Kulasekera	Clemson University, USA	Ravi Vakil	Stanford University, USA
Gerry Ladas	University of Rhode Island, USA	Antonia Vecchio	Consiglio Nazionale delle Ricerche, Italy
David Larson	Texas A&M University, USA	John C. Wierman	Johns Hopkins University, USA
Suzanne Lenhart	University of Tennessee, USA	Michael E. Zieve	University of Michigan, USA
Chi-Kwong Li	College of William and Mary, USA		

PRODUCTION

Silvio Levy, Scientific Editor


Cover: Alex Scorpan

See inside back cover or msp.org/involve for submission instructions. The subscription price for 2020 is US \$205/year for the electronic version, and \$275/year (+\$35, if shipping outside the US) for print and electronic. Subscriptions, requests for back issues and changes of subscriber address should be sent to MSP.

Involve (ISSN 1444-4184 electronic, 1444-4176 printed) at Mathematical Sciences Publishers, 798 Evans Hall #3840, c/o University of California, Berkeley, CA 94720-3840, is published continuously online. Periodical rate postage paid at Berkeley, CA 94704, and additional mailing offices.

Involve peer review and production are managed by EditFlow® from Mathematical Sciences Publishers.

PUBLISHED BY

 **mathematical sciences publishers**
nonprofit scientific publishing

<http://msp.org/>

© 2020 Mathematical Sciences Publishers

involve

2020

vol. 13

no. 4

Structure constants of $\mathcal{U}(\mathfrak{sl}_2)$	541
ALEXIA GOURLEY AND CHRISTOPHER KENNEDY	
Conjecture \mathcal{C} holds for some horospherical varieties of Picard rank 1	551
LELA BONES, GARRETT FOWLER, LISA SCHNEIDER AND RYAN M. SHIFLER	
Condensed Ricci curvature of complete and strongly regular graphs	559
VINCENT BONINI, CONOR CARROLL, UYEN DINH, SYDNEY DYE, JOSHUA FREDERICK AND ERIN PEARSE	
On equidistant polytopes in the Euclidean space	577
CSABA VINCZE, MÁRK OLÁH AND LETÍCIA LENGUEL	
Polynomial values in Fibonacci sequences	597
ADI OSTROV, DANNY NEFTIN, AVI BERMAN AND REYAD A. ELRAZIK	
Stability and asymptotic analysis of the Föllmer–Schweizer decomposition on a finite probability space	607
SARAH BOESE, TRACY CUI, SAMUEL JOHNSTON, SYLVIE VEGA-MOLINO AND OLEKSII MOSTOVYI	
Eigenvalues of the sum and product of anticommuting matrices	625
VADIM PONOMARENKO AND LOUIS SELSTAD	
Combinatorial random knots	633
ANDREW DUCHARME AND EMILY PETERS	
Conjugation diameter of the symmetric groups	655
ASSAF LIBMAN AND CHARLOTTE TARRY	
Existence of multiple solutions to a discrete boundary value problem with mixed periodic boundary conditions	673
KIMBERLY HOWARD, LONG WANG AND MIN WANG	
Minimal flag triangulations of lower-dimensional manifolds	683
CHRISTIN BIBBY, ANDREW ODESKY, MENG MENG WANG, SHUYANG WANG, ZIYI ZHANG AND HAILUN ZHENG	
Some new Gompertz fractional difference equations	705
TOM CUCHTA AND BROOKE FINCHAM	