

involve

a journal of mathematics

Existence of multiple solutions
to a discrete boundary value problem
with mixed periodic boundary conditions

Kimberly Howard, Long Wang and Min Wang



Existence of multiple solutions to a discrete boundary value problem with mixed periodic boundary conditions

Kimberly Howard, Long Wang and Min Wang

(Communicated by Johnny Henderson)

A second-order discrete boundary value problem with mixed periodic boundary conditions is studied. Sufficient conditions on the multiplicity of solutions in a weak sense are obtained by using the critical point theory. An example is given to demonstrate the applications of our results as well.

1. Introduction

We consider a nonlinear boundary value problem (BVP) consisting of a second-order difference equation

$$-\Delta^2 u(t-1) = f(t, u(t)), \quad t \in [2, N]_{\mathbb{Z}}, \quad (1-1)$$

and a pair of mixed periodic boundary conditions (BCs)

$$u(0) = -u(N), \quad \Delta u(0) = \Delta u(N), \quad (1-2)$$

where

- $N \geq 3$ is a positive integer and $[a, b]_{\mathbb{Z}}$ denotes the discrete interval $\{a, a+1, \dots, b\}$ for any integers a and b with $a \leq b$;
- Δ is the forward difference operator defined by $\Delta u(t) = u(t+1) - u(t)$ and $\Delta^2 u(t) = \Delta(\Delta u(t))$; and
- $f : [2, N]_{\mathbb{Z}} \times \mathbb{R} \rightarrow \mathbb{R}$ is continuous in its second variable.

By a solution of BVP (1-1), (1-2), we mean a function $u : [0, N+1]_{\mathbb{Z}} \rightarrow \mathbb{R}$ that satisfies (1-1) and (1-2).

Remark 1.1. It is notable that in this paper, the solution is only required to satisfy (1-1) on $[2, N]_{\mathbb{Z}}$ instead of $[1, N]_{\mathbb{Z}}$. We call it *a solution in a weak sense*. Assume (1-1) is defined on $[1, N]_{\mathbb{Z}}$. By applying the standard variational technique in the

MSC2010: 34B15, 39A23.

Keywords: discrete boundary value problem, mixed periodic boundary conditions, variational methods.

traditional way, let

$$\begin{aligned}\tilde{H} &= \{u : [0, N+1]_{\mathbb{Z}} \rightarrow \mathbb{R} \mid u \text{ satisfies the BC (1-2)}\}, \\ \tilde{J}(u) &= \frac{1}{2} \sum_{t=1}^N (\Delta u(t-1))^2 - \sum_{t=1}^N \int_0^{u(t)} f(t, s) ds.\end{aligned}$$

If $u \in \tilde{H}$ is a critical point of $\tilde{J}(u)$, then u must satisfy $\Delta u(0) = \Delta u(N) = 0$ and $u(0) = -u(N)$, which become a special case of the antiperiodic BCs studied in the literature; see for example [Lyons and Neugebauer 2015a; Kuang and Yang 2018]. We are not interested in such simple BCs. This is the motivation to consider the solution in a weak sense. The reader is referred to [Kong and Wang $\geq$ 2020, Remark 2.2] for more discussions on the solution in a weak sense.

BVPs have been an active research area for decades. Both continuous and discrete BVPs subject to various BCs have been investigated by numerous scholars. The reader is referred to [Davis et al. 2016; Feng et al. 2018; Garcia and Neugebauer 2019; Graef et al. 2008; 2013; 2017; 2018; Henderson and Luca 2018; 2019; Kuang and Yang 2018; Lyons and Neugebauer 2015a; 2015b; Liang and Weng 2007; Liu et al. 2018; Kong et al. 2017] for recent advances in BVPs. Although BVPs with either periodic BCs or antiperiodic BCs have been extensively studied, to the best of our knowledge, there are relatively few works on the problems with mixed periodic BCs, i.e., a combination of periodic and antiperiodic BCs. For continuous BVPs, Garcia and Neugebauer [2019] studied the existence of solutions for the BVPs consisting of the differential equation

$$u'' = f(t, u, u'), \quad 0 < t < 1,$$

and two sets of mixed periodic BCs

$$u(0) + u(1) = 0, \quad u'(0) - u'(1) = 0,$$

or

$$u(0) - u(1) = 0, \quad u'(0) + u'(1) = 0.$$

For discrete BVPs, Kong and Wang [2020] considered the existence of solutions (in a weak sense) for the BVP

$$\begin{aligned}-\Delta^2 u(t-1) &= f(u(t)), \quad t \in [2, N]_{\mathbb{Z}}, \\ u(0) &= -u(N), \quad \Delta u(0) = \Delta u(N),\end{aligned}\tag{1-3}$$

by the mountain pass lemma. In [Kong and Wang $\geq$ 2020], the existence of solutions (in a weak sense) for the BVP

$$\begin{aligned}-\Delta(r(t-1)\Delta u(t-1)) &= f(t, u(t)), \quad t \in [2, N]_{\mathbb{Z}}, \\ u(0) &= u(N), \quad r(0)\Delta u(0) = -r(N)\Delta u(N),\end{aligned}$$

was studied by Clark's theorem. Note that as commented in [Kong and Wang ≥ 2020], there was a typo in (1.1) in [Kong and Wang 2020]. The domain was mistakenly written as $t \in [1, N]_{\mathbb{Z}}$ and should be replaced by $t \in [2, N]_{\mathbb{Z}}$ as seen in (1-3) above.

We will extend [Kong and Wang 2020] by considering a more general problem. It is easy to see that BVP (1-1), (1-2) covers BVP (1-3) as a special case. By using the linking theorem, a new set of sufficient conditions for the existence of solutions will be derived. Our work will supplement the existing results and contribute to the application of the variational method to the BVPs with mixed periodic BCs.

This paper is organized as follows: after this introduction, the preliminary results are given in Section 2. Section 3 contains the main results. An example is given in Section 3 as well.

2. Preliminaries

We present some definitions and lemmas needed for the proof of our main result.

Definition 2.1. Assume U is a Banach space. We say that a functional $J \in C^1(U, \mathbb{R})$ satisfies the Palais–Smale (PS) condition if every sequence $\{u_n\} \subset U$, such that $J(u_n)$ is bounded and $J'(u_n) \rightarrow 0$ as $n \rightarrow \infty$, has a convergent subsequence. The sequence $\{u_n\}$ is called a PS sequence.

The following lemma, which is the well-known linking theorem, plays a crucial role in the proof of our main results; see for example [Rabinowitz 1986, Theorem 5.3].

Lemma 2.2. Let U be a real Banach space, $U = U_1 \oplus U_2$, where U_1 is finite-dimensional. Suppose that $J \in C^1(U, \mathbb{R})$ satisfies the PS condition and the following:

- (J1) There are positive constants c and ρ such that $J|_{\partial B_\rho(0) \cap U_2} \geq c$.
- (J2) There are $\mu \in \partial B_1(0) \cap U_2$ and a positive constant $\hat{c} \geq \rho$ such that $J|_{\partial \Omega} \leq 0$, where

$$\Omega = (\bar{B}_{\hat{c}}(0) \cap U_1) \oplus \{s\mu \mid 0 < s < \hat{c}\}.$$

Then J possesses a critical value $c_0 \geq c$, where

$$c_0 = \inf_{d \in \Gamma} \sup_{u \in \Omega} J(d(u)) \quad (2-1)$$

and $\Gamma = \{d \in C(\bar{\Omega}, U) \mid d|_{\partial \Omega} = I\}$, where I denotes the identity operator.

In the rest of the paper, we will use the Banach space U defined by

$$U = \{u : [0, N+1]_{\mathbb{Z}} \rightarrow \mathbb{R} \mid u(0) = -u(N), \Delta u(0) = \Delta u(N), u(1) = 0\}.$$

By [Kong and Wang 2020, Remark 2.1], U is an $(N-1)$ -dimensional Banach space with the norm defined by

$$\|u\| = \left(\sum_{t=1}^N u^2(t) \right)^{\frac{1}{2}}.$$

Let $\tilde{f} : [1, N]_{\mathbb{Z}} \times \mathbb{R} \rightarrow \mathbb{R}$ and $\tilde{F} : [1, N]_{\mathbb{Z}} \times \mathbb{R} \rightarrow \mathbb{R}$ be defined by

$$\tilde{f}(t, x) = \begin{cases} 0, & t = 1, \\ f(t, x), & t \in [2, N-1]_{\mathbb{Z}}, \\ f(t, x) + 2x, & t = N, \end{cases} \quad (2-2)$$

$$\tilde{F}(t, x) = \int_0^x \tilde{f}(t, s) ds. \quad (2-3)$$

Clearly, both $\tilde{f}(t, x)$ and $\tilde{F}(t, x)$ are continuous in x . Define $J : U \rightarrow \mathbb{R}$ by

$$J(u) = \frac{1}{2} \sum_{t=1}^N (\Delta u(t-1))^2 - \sum_{t=1}^N \tilde{F}(t, u(t)). \quad (2-4)$$

A critical point of J is a point u at which J' , the Fréchet derivative of J , vanishes, and the value of J at u is then called a critical value of J ; see for example [Rabinowitz 1986]. The following lemma links the critical point of J and the solution of BVP (1-1), (1-2).

Lemma 2.3. *The function $u \in U$ is a critical point of J if and only if u is a solution of BVP (1-1), (1-2) in a weak sense.*

Proof. For any $u \in U$, from (2-2)–(2-4), we obtain

$$J(u) = \frac{1}{2} \sum_{t=1}^N (\Delta u(t-1))^2 - \sum_{t=2}^N \int_0^{u(t)} f(t, s) ds - 2 \int_0^{u(N)} s ds.$$

It is easy to see that J is continuously differentiable. For any $v \in U$, we have that $J'(u)$ at $u \in U$ is given by

$$\langle J'(u), v \rangle = \sum_{t=1}^N \Delta u(t-1) \Delta v(t-1) - \sum_{t=2}^N f(t, u(t)) v(t) - 2u(N)v(N).$$

Using the summation by parts formula, we have

$$\begin{aligned} \sum_{t=1}^N \Delta u(t-1) \Delta v(t-1) &= \Delta u(N)v(N) - \Delta u(0)v(0) - \sum_{t=1}^N \Delta^2 u(t-1)v(t) \\ &= 2\Delta u(N)v(N) - \sum_{t=1}^N \Delta^2 u(t-1)v(t). \end{aligned}$$

Then it follows that

$$\begin{aligned}
 \langle J'(u), v \rangle &= 2\Delta u(N)v(N) - \sum_{t=1}^N \Delta^2 u(t-1)v(t) - \sum_{t=2}^N f(t, u(t))v(t) - 2u(N)v(N) \\
 &= -2\Delta u(0)v(0) - \sum_{t=1}^N \Delta^2 u(t-1)v(t) - \sum_{t=2}^N f(t, u(t))v(t) - 2u(0)v(0) \\
 &= -\sum_{t=1}^N \Delta^2 u(t-1)v(t) - \sum_{t=2}^N f(t, u(t))v(t) \\
 &= -\sum_{t=2}^N [\Delta^2 u(t-1) + f(t, u(t))]v(t).
 \end{aligned}$$

Therefore $\langle J'(u), v \rangle = 0$ if and only if (1-1) holds. $\square$

Now we consider an equivalent form of J . Let

$$A = \begin{bmatrix} 2 & -1 & 0 & \cdots & \cdots & 0 & 1 \\ -1 & 2 & -1 & 0 & \cdots & \cdots & 0 \\ 0 & -1 & 2 & -1 & 0 & \cdots & 0 \\ & \ddots & \ddots & \ddots & \ddots & \ddots & \\ 0 & \cdots & 0 & -1 & 2 & -1 & 0 \\ 0 & \cdots & \cdots & 0 & -1 & 2 & -1 \\ 1 & 0 & \cdots & 0 & 0 & -1 & 2 \end{bmatrix}_{N \times N}, \quad N \geq 3. \quad (2-5)$$

Then it is easy to verify that for any $u \in U$,

$$J(u) = \frac{1}{2}u^T Au - \sum_{t=1}^N \tilde{F}(t, u(t)), \quad (2-6)$$

where $(\cdot)^T$ denotes the transpose and $u = (0, u(2), \dots, u(N))^T$. Note that by [Kong and Wang 2020, Remark 2.2], A has N positive eigenvalues $0 < \lambda_1 \leq \lambda_2 \leq \dots \leq \lambda_N$ such that for any $u \in \mathbb{R}^N$, we have $\lambda_1 \|u\|^2 \leq u^T Au \leq \lambda_N \|u\|^2$.

3. Main results

The following theorem is our first main result.

Theorem 3.1. *Let $\lambda_1, \dots, \lambda_N$ be the eigenvalues of A defined by (2-5). Assume the function f from (1-1) satisfies $f(t, 0) = 0$, $t \in [2, N]_{\mathbb{Z}}$. Additionally, the function $\tilde{F}(t, x)$ defined by (2-3) satisfies the following assumptions:*

(F1) *There exist two constants $\delta_1 > 0$ and $a_1 \in (0, \frac{1}{2}\lambda_1)$ such that $\tilde{F}(t, x) \leq a_1 x^2$ for $|x| \leq \delta_1$ and $t \in [2, N]_{\mathbb{Z}}$.*

(F2) *There exist two constants $a_2 \in (\frac{1}{2}\lambda_N, \infty)$ and $a_3 > 0$ such that $\tilde{F}(t, x) \geq a_2x^2 - a_3$ for all $(t, x) \in [2, N]_{\mathbb{Z}} \times \mathbb{R}$.*

Then BVP (1-1), (1-2) admits at least three solutions, one being trivial, and two being nontrivial.

The following lemma will be needed to prove Theorem 3.1.

Lemma 3.2. *Assume (F2) of Theorem 3.1 is satisfied. Then the functional J defined by (2-4) (or (2-6)) satisfies the PS condition.*

Proof. For any sequence $\{u_k\} \subseteq U$ with $J'(u_k) \rightarrow 0$ as $k \rightarrow \infty$ and $-B \leq J(u_k) \leq B$, where B is a positive constant, we claim $\{u_k\}$ is bounded. We will prove it by contradiction.

Assume $\{u_k\} \subseteq U$ is an unbounded sequence. Then there exists a subsequence $\{u_m\}$ of $\{u_k\}$ such that $\|u_m\| \rightarrow \infty$ and $J'(u_m) \rightarrow 0$ as $m \rightarrow \infty$, and $-B \leq J(u_m) \leq B$ for $m = 1, 2, \dots$. From (F2) of Theorem 3.1 and (2-6), when m is large enough, we have

$$\begin{aligned} J(u_m) &= \frac{1}{2}u_m^T A u_m - \sum_{t=1}^N \tilde{F}(t, u_m(t)) \\ &\leq \frac{1}{2}\lambda_N \|u_m\|^2 - \sum_{t=2}^N \tilde{F}(t, u_m(t)) \leq \frac{1}{2}\lambda_N \|u_m\|^2 - \sum_{t=2}^N (a_2(u_m(t))^2 - a_3) \\ &\leq \frac{1}{2}\lambda_N \|u_m\|^2 - a_2 \|u_m\|^2 + a_3 N = (\frac{1}{2}\lambda_N - a_2) \|u_m\|^2 + a_3 N. \end{aligned} \quad (3-1)$$

Hence $J(u_m) \rightarrow -\infty$ as $m \rightarrow \infty$, and this contradicts the fact that $\{J(u_m)\}$ is bounded. Thus, $\{u_k\}$ is bounded. Since U is a finite-dimensional space, we know $\{u_k\}$ has a convergent subsequence. Therefore, J satisfies the PS condition. $\square$

Now we are ready to prove Theorem 3.1.

Proof of Theorem 3.1. It is obvious that $u \equiv 0$ is a solution of BVP (1-1), (1-2).

By (3-1), we can show $J(u) \rightarrow -\infty$ as $\|u\| \rightarrow \infty$ for any $u \in U$. Using the fact that $J(u)$ is bounded above in U , define $\bar{J} = \sup_{u \in U} J(u)$. Therefore, we have a sequence $\{u_k\}$ on U such that $\bar{J} = \lim_{k \rightarrow \infty} J(u_k)$. From the proof of Lemma 3.2, it follows that $\{u_k\}$ is bounded. Then $\{u_k\}$ has a convergent subsequence defined by $\{u_{k_n}\}$. Let $\bar{u} = \lim_{n \rightarrow \infty} u_{k_n}$, and it is easy to see that $\bar{u} \in U$ with $J(\bar{u}) = \bar{J}$. Therefore $\bar{u}$ is a critical point of J .

To apply Lemma 2.2, we need to separate U into two subspaces P and Q with $U = P \oplus Q$. Let $\{\eta_1, \dots, \eta_N\} \subset \mathbb{R}^N$ be the orthonormal eigenvectors associated with $\{\lambda_1, \dots, \lambda_N\}$. Define $P = \text{span}\{\eta_1, \dots, \eta_M\}$ and $Q = \text{span}\{\eta_{M+1}, \dots, \eta_N\}$, for some $M \in [2, N-1]_{\mathbb{Z}}$, such that $U = P \oplus Q$, $P \cap U \neq \{0\}$, and $Q \cap U \neq \{0\}$. Clearly, both P and Q are finite-dimensional spaces. Moreover, for any $u \in P$, there

exist $b_1, \dots, b_M \in \mathbb{R}$ such that $u = \sum_{i=1}^M b_i \eta_i$ and $\|u\|^2 = \sum_{i=1}^M b_i^2$; for any $u \in Q$, there exist $b_{M+1}, \dots, b_N \in \mathbb{R}$ such that $u = \sum_{i=M+1}^N b_i \eta_i$ and $\|u\|^2 = \sum_{i=M+1}^N b_i^2$.

Since U is finite-dimensional, for any $u \in U$, there exist positive numbers κ_1 and κ_2 such that $\kappa_1 < \kappa_2$ and

$$\kappa_1 \max\{|u(1)|, |u(2)|, \dots, |u(N)|\} \leq \|u\| \leq \kappa_2 \max\{|u(1)|, |u(2)|, \dots, |u(N)|\}.$$

Observe for any $u \in Q$, such that $\|u\| = \kappa_1 \delta_1$ and $\max\{|u(1)|, |u(2)|, \dots, |u(N)|\} \leq \delta_1$,

$$\begin{aligned} J(u) &= \frac{1}{2} u^T A u - \sum_{t=1}^N \tilde{F}(t, u(t)) \geq \frac{1}{2} \lambda_1 \|u\|^2 - \sum_{t=2}^N \tilde{F}(t, u(t)) \\ &\geq \frac{1}{2} \lambda_1 \|u\|^2 - \sum_{t=2}^N a_1 (u(t))^2 = \frac{1}{2} \lambda_1 \|u\|^2 - a_1 \|u\|^2 \\ &= \left(\frac{1}{2} \lambda_1 - a_1\right) (\kappa_1 \delta_1)^2 > 0. \end{aligned} \quad (3-2)$$

Then assumption (J1) of Lemma 2.2 is satisfied. It is easy to see (3-2) also implies $\bar{J} > 0$. So $\bar{u}$ is a nontrivial solution of BVP (1-1), (1-2).

Next we verify (J2) of Lemma 2.2. Choose $\mu \in Q$ and a positive constant $\hat{c}$ such that $\|\mu\| = 1$ and

$$\hat{c} > \max \left\{ \kappa_1 \delta_1, \frac{2a_3 N}{2a_2 - \lambda_N} \right\}.$$

Let $\Omega = (B_{\hat{c}}(0) \cap P) \oplus \{s\mu \mid 0 < s < \hat{c}\}$. Note that for any $v \in P$

$$\begin{aligned} J(s\mu + v) &= \frac{1}{2} (s\mu + v)^T A (s\mu + v) - \sum_{t=1}^N \tilde{F}(t, s\mu(t) + v(t)) \\ &\leq \frac{1}{2} \lambda_N s^2 + \frac{1}{2} \lambda_N \|v\|^2 - \sum_{t=2}^N (a_2 (s\mu(t) + v(t))^2 - a_3) \\ &\leq \frac{1}{2} \lambda_N s^2 + \frac{1}{2} \lambda_N \|v\|^2 - a_2 s^2 - a_2 \|v\|^2 + a_3 N \\ &= \left(\frac{1}{2} \lambda_N - a_2\right) s^2 + \left(\frac{1}{2} \lambda_N - a_2\right) \|v\|^2 + a_3 N \\ &\leq \left(\frac{1}{2} \lambda_N - a_2\right) \|v\|^2 + a_3 N. \end{aligned}$$

Therefore, $J|_{\partial\Omega} \leq 0$. Hence (J2) holds.

Then by Lemma 2.2, we know $J(u)$ has a critical value $c_0 \geq \left(\frac{1}{2} \lambda_1 - a_1\right) (\kappa_1 \delta_1)^2$. Let $\tilde{u} \in U$ be a critical point corresponding to c_0 ; i.e., $J(\tilde{u}) = c_0$. If $\bar{u} \neq \tilde{u}$, then by Lemma 2.3, $\bar{u}$ and $\tilde{u}$ are two different solutions of BVP (1-1), (1-2). If $\bar{u} = \tilde{u}$, by (2-1), $\sup_{u \in U} J(u) = \inf_{d \in \Gamma} \sup_{u \in \Omega} J(d(u))$. Choose distinct $d_1, d_2 \in \Gamma$ such that

$$\{d_1(u) \mid u \in \Omega \setminus \partial\Omega\} \cap \{d_2(u) \mid u \in \Omega \setminus \partial\Omega\} = \emptyset.$$

Then

$$\sup_{u \in \Omega} J(d_1(u)) = \sup_{u \in \Omega} J(d_2(u)) = \sup_{u \in U} J(u).$$

Hence there exist $u_1, u_2 \in \Omega$ such that $d_1(u_1)$ and $d_2(u_2)$ are two different critical points of J .

Therefore, we have shown BVP (1-1), (1-2) admits at least three solutions. $\square$

The following corollary is a direct consequence of Theorem 3.1.

Corollary 3.3. Assume the function f from (1-1) satisfies $f(t, 0) = 0$, $t \in [2, N]_{\mathbb{Z}}$, and $\tilde{F}(t, x)$ defined by (2-3) is of the form

$$\tilde{F}(t, x) = a(t)|x|^{r(t)}, \quad a(t) > 0, \quad r(t) > 2, \quad t \in [2, N]_{\mathbb{Z}}.$$

Then BVP (1-1), (1-2) admits at least three solutions, one being trivial, and two being nontrivial.

We conclude the paper with the following example.

Example 3.4. Consider BVP (1-1), (1-2) with $N = 4$ and f defined by

$$f(t, x) = \begin{cases} \frac{2}{5}x^3, & t = 2, 3, \\ \frac{2}{5}x^3 - 2x, & t = 4. \end{cases}$$

By (2-2) and (2-3), we know

$$\tilde{F}(t, x) = \frac{1}{10}x^4, \quad t = 2, 3, 4.$$

Then by Corollary 3.3, BVP (1-1), (1-2) admits at least three solutions.

Acknowledgement

This research was supported by the Mentor Protégé Research Program from the College of Science and Mathematics at Kennesaw State University.

References

- [Davis et al. 2016] J. M. Davis, P. W. Eloe, J. R. Graef, and J. Henderson, “Positive solutions for a singular fourth order nonlocal boundary value problem”, *Int. J. Pure Appl. Math.* **109**:1 (2016), 67–84.
- [Feng et al. 2018] Y. Feng, J. R. Graef, L. Kong, and M. Wang, “The forward and inverse problems for a fractional boundary value problem”, *Appl. Anal.* **97**:14 (2018), 2474–2484. MR Zbl
- [Garcia and Neugebauer 2019] A. E. Garcia and J. T. Neugebauer, “Solutions of boundary value problems at resonance with periodic and antiperiodic boundary conditions”, *Involve* **12**:1 (2019), 171–180. MR Zbl
- [Graef et al. 2008] J. R. Graef, L. Kong, and H. Wang, “Existence, multiplicity, and dependence on a parameter for a periodic boundary value problem”, *J. Differential Equations* **245**:5 (2008), 1185–1197. MR Zbl

- [Graef et al. 2013] J. R. Graef, L. Kong, and M. Wang, “Multiple solutions to a periodic boundary value problem for a nonlinear discrete fourth order equation”, *Adv. Dyn. Syst. Appl.* **8**:2 (2013), 203–215. MR Zbl
- [Graef et al. 2017] J. R. Graef, L. Kong, Q. Kong, and M. Wang, “On a fractional boundary value problem with a perturbation term”, *J. Appl. Anal. Comput.* **7**:1 (2017), 57–66. MR
- [Graef et al. 2018] J. R. Graef, S. Heidarkhani, L. Kong, and M. Wang, “Existence of solutions to a discrete fourth order boundary value problem”, *J. Difference Equ. Appl.* **24**:6 (2018), 849–858. MR Zbl
- [Henderson and Luca 2018] J. Henderson and R. Luca, “Positive solutions for a system of coupled fractional boundary value problems”, *Lith. Math. J.* **58**:1 (2018), 15–32. MR Zbl
- [Henderson and Luca 2019] J. Henderson and R. Luca, “Existence of positive solutions for a system of semipositone coupled discrete boundary value problems”, *J. Difference Equ. Appl.* **25**:4 (2019), 516–541. MR Zbl
- [Kong and Wang 2020] L. Kong and M. Wang, “Existence of solutions for a second order discrete boundary value problem with mixed periodic boundary conditions”, *Appl. Math. Lett.* **102** (2020), art. id. 106138. MR Zbl
- [Kong and Wang $\geq$ 2020] L. Kong and M. Wang, “Multiple and particular solutions of a second order discrete boundary value problem with mixed periodic boundary conditions”, submitted.
- [Kong et al. 2017] L. Kong, J. Parsley, K. Rizzo, and N. Russell, “Anti-periodic solutions for a higher order difference equation with p -Laplacian”, *J. Appl. Anal.* **23**:2 (2017), 111–125. MR Zbl
- [Kuang and Yang 2018] J. Kuang and Y. Yang, “Variational approach to anti-periodic boundary value problems involving the discrete p -Laplacian”, *Bound. Value Probl.* **2018** (2018), art. id. 86. MR
- [Liang and Weng 2007] H. Liang and P. Weng, “Existence and multiple solutions for a second-order difference boundary value problem via critical point theory”, *J. Math. Anal. Appl.* **326**:1 (2007), 511–520. MR Zbl
- [Liu et al. 2018] X. Liu, T. Zhou, and H. Shi, “Existence and multiple solutions to a discrete fourth order boundary value problem”, *Adv. Difference Equ.* **2018** (2018), art. id. 427. MR Zbl
- [Lyons and Neugebauer 2015a] J. W. Lyons and J. T. Neugebauer, “A difference equation with anti-periodic boundary conditions”, *Dyn. Contin. Discrete Impuls. Syst. Ser. A Math. Anal.* **22**:1 (2015), 47–60. MR Zbl
- [Lyons and Neugebauer 2015b] J. W. Lyons and J. T. Neugebauer, “Existence of an antisymmetric solution of a boundary value problem with antiperiodic boundary conditions”, *Electron. J. Qual. Theory Differ. Equ.* **2015** (2015), art. id. 72. MR Zbl
- [Rabinowitz 1986] P. H. Rabinowitz, *Minimax methods in critical point theory with applications to differential equations*, CBMS Region. Conf. Ser. Math. **65**, Amer. Math. Sci., Providence, RI, 1986. MR Zbl

Received: 2020-03-09 Revised: 2020-03-27 Accepted: 2020-04-28

khowar43@students.kennesaw.edu *Department of Mathematics, Kennesaw State University, Marietta, GA, United States*

lwang17@kennesaw.edu *Department of Mathematics, Kennesaw State University, Marietta, GA, United States*

min.wang@kennesaw.edu *Department of Mathematics, Kennesaw State University, Marietta, GA, United States*

involve

msp.org/involve

INVOLVE YOUR STUDENTS IN RESEARCH

Involve showcases and encourages high-quality mathematical research involving students from all academic levels. The editorial board consists of mathematical scientists committed to nurturing student participation in research. Bridging the gap between the extremes of purely undergraduate research journals and mainstream research journals, *Involve* provides a venue to mathematicians wishing to encourage the creative involvement of students.

MANAGING EDITOR

Kenneth S. Berenhaut Wake Forest University, USA

BOARD OF EDITORS

Colin Adams	Williams College, USA	Robert B. Lund	Clemson University, USA
Arthur T. Benjamin	Harvey Mudd College, USA	Gaven J. Martin	Massey University, New Zealand
Martin Bohner	Missouri U of Science and Technology, USA	Mary Meyer	Colorado State University, USA
Amarjit S. Budhiraja	U of N Carolina, Chapel Hill, USA	Frank Morgan	Williams College, USA
Pietro Cerone	La Trobe University, Australia	Mohammad Sal Moslehian	Ferdowsi University of Mashhad, Iran
Scott Chapman	Sam Houston State University, USA	Zuhair Nashed	University of Central Florida, USA
Joshua N. Cooper	University of South Carolina, USA	Ken Ono	Univ. of Virginia, Charlottesville
Jem N. Corcoran	University of Colorado, USA	Yuval Peres	Microsoft Research, USA
Toka Diagana	University of Alabama in Huntsville, USA	Y.-F. S. Pétermann	Université de Genève, Switzerland
Michael Dorff	Brigham Young University, USA	Jonathon Peterson	Purdue University, USA
Sever S. Dragomir	Victoria University, Australia	Robert J. Plemmons	Wake Forest University, USA
Joel Foisy	SUNY Potsdam, USA	Carl B. Pomerance	Dartmouth College, USA
Errin W. Fulp	Wake Forest University, USA	Vadim Ponomarenko	San Diego State University, USA
Joseph Gallian	University of Minnesota Duluth, USA	Bjorn Poonen	UC Berkeley, USA
Stephan R. Garcia	Pomona College, USA	József H. Przytycki	George Washington University, USA
Anant Godbole	East Tennessee State University, USA	Richard Rebarber	University of Nebraska, USA
Ron Gould	Emory University, USA	Robert W. Robinson	University of Georgia, USA
Sat Gupta	U of North Carolina, Greensboro, USA	Javier Rojo	Oregon State University, USA
Jim Haglund	University of Pennsylvania, USA	Filip Saidak	U of North Carolina, Greensboro, USA
Johnny Henderson	Baylor University, USA	Hari Mohan Srivastava	University of Victoria, Canada
Glenn H. Hurlbert	Virginia Commonwealth University, USA	Andrew J. Sterge	Honorary Editor
Charles R. Johnson	College of William and Mary, USA	Ann Trenk	Wellesley College, USA
K. B. Kulasekera	Clemson University, USA	Ravi Vakil	Stanford University, USA
Gerry Ladas	University of Rhode Island, USA	Antonia Vecchio	Consiglio Nazionale delle Ricerche, Italy
David Larson	Texas A&M University, USA	John C. Wierman	Johns Hopkins University, USA
Suzanne Lenhart	University of Tennessee, USA	Michael E. Zieve	University of Michigan, USA
Chi-Kwong Li	College of William and Mary, USA		

PRODUCTION

Silvio Levy, Scientific Editor

Cover: Alex Scorpan

See inside back cover or msp.org/involve for submission instructions. The subscription price for 2020 is US \$205/year for the electronic version, and \$275/year (+\$35, if shipping outside the US) for print and electronic. Subscriptions, requests for back issues and changes of subscriber address should be sent to MSP.

Involve (ISSN 1444-4184 electronic, 1444-4176 printed) at Mathematical Sciences Publishers, 798 Evans Hall #3840, c/o University of California, Berkeley, CA 94720-3840, is published continuously online. Periodical rate postage paid at Berkeley, CA 94704, and additional mailing offices.

Involve peer review and production are managed by EditFlow® from Mathematical Sciences Publishers.

PUBLISHED BY

 **mathematical sciences publishers**
nonprofit scientific publishing

<http://msp.org/>

© 2020 Mathematical Sciences Publishers

involve

2020

vol. 13

no. 4

Structure constants of $\mathcal{U}(\mathfrak{sl}_2)$	541
ALEXIA GOURLEY AND CHRISTOPHER KENNEDY	
Conjecture $\mathcal{C}$ holds for some horospherical varieties of Picard rank 1	551
LELA BONES, GARRETT FOWLER, LISA SCHNEIDER AND RYAN M. SHIFLER	
Condensed Ricci curvature of complete and strongly regular graphs	559
VINCENT BONINI, CONOR CARROLL, UYEN DINH, SYDNEY DYE, JOSHUA FREDERICK AND ERIN PEARSE	
On equidistant polytopes in the Euclidean space	577
CSABA VINCZE, MÁRK OLÁH AND LETÍCIA LENGUEL	
Polynomial values in Fibonacci sequences	597
ADI OSTROV, DANNY NEFTIN, AVI BERMAN AND REYAD A. ELRAZIK	
Stability and asymptotic analysis of the Föllmer–Schweizer decomposition on a finite probability space	607
SARAH BOESE, TRACY CUI, SAMUEL JOHNSTON, SYLVIE VEGA-MOLINO AND OLEKSII MOSTOVYI	
Eigenvalues of the sum and product of anticommuting matrices	625
VADIM PONOMARENKO AND LOUIS SELSTAD	
Combinatorial random knots	633
ANDREW DUCHARME AND EMILY PETERS	
Conjugation diameter of the symmetric groups	655
ASSAF LIBMAN AND CHARLOTTE TARRY	
Existence of multiple solutions to a discrete boundary value problem with mixed periodic boundary conditions	673
KIMBERLY HOWARD, LONG WANG AND MIN WANG	
Minimal flag triangulations of lower-dimensional manifolds	683
CHRISTIN BIBBY, ANDREW ODESKY, MENG MENG WANG, SHUYANG WANG, ZIYI ZHANG AND HAILUN ZHENG	
Some new Gompertz fractional difference equations	705
TOM CUCHTA AND BROOKE FINCHAM	