

# involve

a journal of mathematics

Some new Gompertz fractional difference equations

Tom Cuchta and Brooke Fincham





# Some new Gompertz fractional difference equations

Tom Cuchta and Brooke Fincham

(Communicated by Martin J. Bohner)

We introduce three new fractional Gompertz difference equations using the Riemann–Liouville discrete fractional calculus. These three models are based a nonfractional Gompertz difference equation, and they differ depending on whether a fractional operator replaces the difference operator, the integral operator defining the logarithm, or both simultaneously. An explicit solution to one of them is achieved with restricted parameters and recurrence relation solutions are derived for all three. Finally, we fit these models to data to compare them with a previously published discrete fractional Gompertz model and the continuous model.

## 1. Introduction

The Gompertz function dates back to the study of human mortality conducted by Benjamin Gompertz [1825]. Since then, it has been used in numerous ways, particularly as a population growth model; e.g., see [Alves et al. 2019; Easton 1999; Jane et al. 2020; Laird 1964; Pezzini et al. 2019; Winsor 1932]. The Gompertz differential equation is

$$y' = -ry \ln\left(\frac{y}{K}\right),$$

which has closed-form general solution as the iterated exponential  $y(t) = Ke^{Ce^{-rt}}$ .

In contrast to the continuous Gompertz model, we are concerned with a discrete theory. The fundamental operator of the “forward” difference calculus is the  $\Delta$  operator that obeys  $\Delta f(t) = f(t+1) - f(t)$ . An enormous amount of research in discrete analogues to the theory of differential equations has been conducted for these operators, and there are many well-known popular introductions to it such as [Bohner and Peterson 2001; Elaydi 2005; Kelley and Peterson 2010]. There are numerous difference equations that have been called a “Gompertz difference equation” in the

---

*MSC2020:* primary 39A60; secondary 26A33, 92D25, 33E12.

*Keywords:* discrete fractional calculus, Gompertz, parameter estimation, data fitting.

literature, such as in [Akin et al. 2020; Nobile et al. 1982; Satoh 2000]. We now consider the  $\Delta$ -Gompertz model from [Bohner 2005, (12)], defined by

$$\Delta y = (\ominus r)(a + \tilde{L}_y)y, \quad y(t_0) = y_0, \quad (1)$$

where  $r$  and  $a$  are constants and

$$\tilde{L}_y(t, t_0) = \sum_{k=t_0}^{t-1} \frac{y(k+1) - y(k)}{y(k)} \quad (2)$$

is the  $\Delta$ -logarithm, originally defined in [Bohner 2005, (3)]. The model (1) originally used time-dependent  $r$  and  $a$ , but we will not consider that generality here.

We will generalize (1) to a model using fractional sum and difference operators. Fractional differential calculus dates back to the early days of the theory of calculus [Ross 1975]. The fundamental idea of fractional calculus is the extension of derivatives and integrals to noninteger order. The fractional discrete calculus dates back to [Kuttner 1957], with an explosion of interest in more recent times [Abdeljawad 2011; Atici and Eloe 2009; Dzieliński et al. 2010; Ferreira 2013; Miller and Ross 1989; Wu et al. 2015]; see in particular the recent monograph [Goodrich and Peterson 2015]. An analogue of the Gompertz differential equation using fractional differential operators has also been developed and studied [Solís-Pérez et al. 2019; Frunzo et al. 2019]. Fractional Gompertz difference equations are not a new idea; for example, see [Wang et al. 2014; Bolton et al. 2015; Atıcı and Şengül 2010], which use various fractional difference operators. We will take one model from the literature for comparison to the models we will propose: Atıcı et al. defined a fractional  $\Delta$ -difference equation in [Atıcı et al. 2017, (3)] by

$$(c - b \log y(t))y(t) = \begin{cases} \Delta^\nu y^\sigma(t - \nu), & \nu \in (0, 1.5) \setminus \{1\}, \\ \Delta y(t), & \nu = 1, \end{cases} \quad (3)$$

where  $\Delta^\nu$  denotes the discrete Riemann–Liouville  $\Delta$ -fractional difference operator. We emphasize that (3) uses the classical logarithm and not a logarithm resembling (2). This is the fundamental change we have made by choosing (1) as the basis for our models. The equation (3) has the following form, convenient for numerical computation [Atıcı et al. 2017, p. 323]:

$$\begin{aligned} & y(t+1) \\ &= \begin{cases} (c - b \log(y(t)))y(t) - \sum_{s=0}^{t-1} \frac{\Gamma(t-\nu-s)}{\Gamma(t-s+1)\Gamma(-\nu)} y(s+1), & \nu \in (0, 1.5) \setminus \{1\}, \\ (c+1 - b \log(y(t)))y(t), & \nu = 1. \end{cases} \end{aligned} \quad (4)$$

The recurrence (4) was used to fit the model (3) to data collected on mice tumors.

### 2. Preliminaries and definitions

For  $a, b \in \mathbb{R}$  with  $a - b$  a positive integer, we define as in [Goodrich and Peterson 2015, p. 1], henceforth abbreviated [GP15], the sets

$$\mathbb{N}_a := \{a, a + 1, a + 2, a + 3, \dots\}, \quad \mathbb{N}_a^b := \{a, a + 1, a + 2, a + 3, \dots, b\}.$$

Throughout, we use the usual convention for empty products that  $\prod_{j=a}^b f(j) = 1$  and for empty sums  $\sum_{j=a}^b f(j) = 0$  whenever  $b < a$ . The backwards jump operator  $\rho : \mathbb{N}_a \rightarrow \mathbb{N}_{a-1}$  is given by [GP15, p. 150], where  $\rho(t) = t - 1$  for  $t > 0$  and  $\rho(0) = 0$ . The backwards difference operator is [GP15, p. 150]  $\nabla y(t) = y(t) - y(\rho(t))$ . If  $f : \mathbb{N}_{a+1} \rightarrow \mathbb{R}$ ,  $a \in \mathbb{N}_a$ , and  $b \in \mathbb{N}_{a+1}$ , then the discrete  $\nabla$ -integral of  $f$  is the summation defined by

$$\int_a^b f(t) \nabla t = \sum_{t=a+1}^b f(t)$$

[GP15, Definition 3.31].

The gamma function  $\Gamma : (0, \infty) \rightarrow [1, \infty)$  is defined by

$$\Gamma(t) = \int_0^\infty x^{t-1} e^{-x} dx$$

[Artin 1964, (2.1)]. Integration by parts shows that the functional equation

$$\Gamma(t + 1) = t\Gamma(t)$$

holds [Artin 1964, (2.2)], which extends the domain of  $\Gamma$  to  $\mathbb{R} \setminus \{0, -1, -2, \dots\}$ . For  $r \in \mathbb{Z}^+$  and  $t \in \mathbb{N}_0$ , the rising factorial function is

$$t^{\bar{r}} = \frac{\Gamma(t + r)}{\Gamma(t)}$$

[GP15, Definition 3.3]. The gamma function is primarily used here to define the  $\nabla$ -Taylor monomials,  $H_\mu$ , which are defined for  $\mu \neq -1, -2, -3, \dots$  and  $t \in \mathbb{N}_a$  by [GP15, Definition 3.56],

$$H_\mu(t, a) = \frac{(t - a)^{\bar{\mu}}}{\Gamma(\mu + 1)} = \frac{\Gamma(t - a + \mu)}{\Gamma(\mu + 1)\Gamma(t - a)},$$

so if  $\mu = 0$ , then we observe  $H_0(t, a) = 1$ . In particular, we observe the following useful identity that we will occasionally use:

$$H_\mu(t, t - 1) = \frac{\Gamma(t - (t - 1) + \mu)}{\Gamma(\mu + 1)\Gamma(t - (t - 1))} = \frac{\Gamma(\mu + 1)}{\Gamma(\mu + 1)\Gamma(1)} = 1.$$

Given  $p : \mathbb{N}_a \rightarrow \mathbb{R}$  such that  $1 - p(t) \neq 0$  for all  $t$ , we define the  $\boxminus$  operator by

$$(\boxminus p)(t) = \frac{-p(t)}{1 - p(t)}$$

[GP15, (3.10)]. In particular, if  $p(t) = r$  is a constant function, then

$$(\boxminus r)(t) = \frac{-r}{1-r}$$

is also a constant function. For  $p : \mathbb{N}_{a+1} \rightarrow \mathbb{R}$  obeying  $1 - p(t) \neq 0$ , the  $\nabla$ -exponential function  $E_p$  is given by [GP15, (3.7)]

$$E_p(t, s) = \begin{cases} \prod_{\tau=s+1}^t 1/(1-p(\tau)), & t \in \mathbb{N}_s, \\ \prod_{\tau=t+1}^s [1-p(\tau)], & t \in \mathbb{N}_a^{s-1}. \end{cases} \quad (5)$$

The following result confirms that the  $\nabla$ -exponential solves a first-order difference equation.

**Theorem 1** [GP15, Theorem 3.6]. *Assume  $p : \mathbb{N}_{a+1} \rightarrow \mathbb{R}$  such that  $1 - p(t) \neq 0$ , and let  $t \in \mathbb{N}_{a+1}$ . The  $\nabla$ -exponential  $y(t) = E_p(t, s)$  is the unique solution of the initial value problem*

$$\nabla y(t) = p(t)y(t), \quad y(s) = 1.$$

If  $f : \mathbb{N}_{a+1} \rightarrow \mathbb{R}$ ,  $\nu > 0$ , and  $t \in \mathbb{N}_a$ , then the  $\nu$ -th order  $\nabla$ -fractional sum of  $f$  at  $t$  is defined by [GP15, Definition 3.58]

$$\nabla_a^{-\nu} f(t) = \int_a^t H_{\nu-1}(t, \rho(s)) f(s) \nabla s,$$

where we let  $\nabla_a^{-0} f(t) = f(t)$ . Let  $f : \mathbb{N}_a \rightarrow \mathbb{R}$ ,  $\nu > 0$ ,  $t \in \mathbb{N}_{a+1}$ , and  $N \in \mathbb{N}_1$ , where  $N - 1 < \nu \leq N$ . The  $\nu$ -th order  $\nabla$ -fractional difference of  $f$  at  $t$  is similarly defined by [GP15, (3.32)]

$$\nabla_a^{\nu} f(t) = \int_a^t H_{-\nu-1}(t, \rho(s)) f(s) \nabla s. \quad (6)$$

From the definition, it is easy to see that  $\nabla_a^{\nu} f(a+1) = f(a+1)$ . The following result shows how fractional differences and summations compose.

**Theorem 2** [GP15, Theorem 3.109]. *If  $f : \mathbb{N}_a \rightarrow \mathbb{R}$  and  $\nu, \mu > 0$ , then*

$$\nabla_a^{\nu} \nabla_a^{-\mu} f(t) = \nabla_a^{\nu-\mu} f(t).$$

The fractional difference operator subtracts from the order of the  $\nabla$ -Taylor monomials as  $\nabla_{t_0}^{\nu} H_{\mu}(t, t_0) = H_{\mu-\nu}(t, t_0)$  [GP15, Theorem 3.93(ii)], and so for any constant  $C$

$$\nabla_a^{\nu} C = C \nabla_a^{\nu} H_0(t, a) = C H_{-\nu}(t, a).$$

In order to solve fractional initial value problems in closed form, the  $\nabla$ -convolution is needed. Let  $f, g : \mathbb{N}_{a+1} \rightarrow \mathbb{R}$ , and define the  $\nabla$ -convolution by [GP15, Definition 3.77]

$$(f * g)(t) = \int_a^t f(t - \rho(\tau) + a)g(\tau) \nabla\tau.$$

The  $\nabla$ -Mittag-Leffler function,  $E_{p,\alpha,\beta}$ , where  $|p| < 1$ ,  $\alpha > 0$ ,  $\beta \in \mathbb{R}$ , and  $t \in \mathbb{N}_a$ , is defined by [GP15, Definition 3.98]

$$E_{p,\alpha,\beta}(t, a) = \sum_{k=0}^{\infty} p^k H_{\alpha k + \beta}(t, a).$$

The following result is a discrete fractional analogue of the variation of constants formula for first-order initial value problems.

**Theorem 3** [GP15, Theorem 3.104]. *If  $f : \mathbb{N}_a \rightarrow \mathbb{R}$ ,  $|c| < 1$ , and  $0 < \nu < 1$ , then the unique solution of the fractional initial value problem*

$$\begin{cases} \nabla_{\rho(a)}^\nu x(t) + cx(t) = f(t), & t \in \mathbb{N}_{a+1}, \\ x(a) = A, & A \in \mathbb{R}, \end{cases}$$

is given by

$$x(t) = [E_{-c,\nu,\nu-1}(\cdot, \rho(a)) * f(\cdot)](t) + [A(c + 1) - f(a)]E_{-c,\nu,\nu-1}(t, \rho(a)). \tag{7}$$

The recurrence (4) came from applying a  $\Delta$ -analogue [Atıcı and Şengül 2010, p. 322] of the following lemma to (3).

**Lemma 4.** *If  $f : \mathbb{N}_{s+1} \rightarrow \mathbb{R}$ , then for  $t \in \mathbb{N}_{s+1}$*

$$\nabla_s^\nu f(t) = f(t) + \sum_{k=s+1}^{t-1} \frac{\Gamma(t - k - \nu)f(k)}{\Gamma(-\nu)\Gamma(t - k + 1)}.$$

*Proof.* Calculating directly from the definition,

$$\begin{aligned} \nabla_s^\nu f(t) &= \int_s^t H_{-\nu-1}(t, \rho(\tau))f(\tau) \nabla\tau = \sum_{k=s+1}^t H_{-\nu-1}(t, \rho(k))f(k) \\ &= H_{-\nu-1}(t, \rho(t))f(t) + \sum_{k=s+1}^{t-1} H_{-\nu-1}(t, \rho(k))f(k) \\ &= f(t) + \sum_{k=s+1}^{t-1} \frac{\Gamma(t - k - \nu)f(k)}{\Gamma(-\nu)\Gamma(t - k + 1)}, \end{aligned}$$

completing the proof. □

We now define the various  $\nabla$ -logarithms we will use. The natural  $\nabla$ -analogue of (2) is

$$L_y(t, t_0) = \int_{t_0}^t \frac{y^\nabla(\tau)}{y(\tau)} \nabla\tau, \quad t \in \mathbb{N}_{t_0+1}.$$

We define the first fractional  $\nabla_\nu$ -logarithm  $L_{y,\nu}$  by

$$L_{y,\nu}(t, t_0) = \left( \nabla_{t_0}^{-\nu} \frac{y^\nabla}{y} \right)(t), \quad t \in \mathbb{N}_{t_0+1}. \quad (8)$$

we define the second fractional  $\nabla_\nu$ -logarithm  $\ell_{y,\nu}$  by

$$\ell_{y,\nu}(t, t_0) = \int_{t_0}^t \frac{\nabla_{t_0}^\nu y(\tau)}{y(\tau)} \nabla\tau, \quad t \in \mathbb{N}_{t_0+1}, \quad (9)$$

and the third  $\nabla_\nu$ -logarithm  $\Lambda_{y,\nu}$  by

$$\Lambda_{y,\nu}(t, t_0) = \left( \nabla_{t_0}^{-\nu} \frac{\nabla_{t_0}^\nu y}{y} \right)(t), \quad t \in \mathbb{N}_{t_0+1}. \quad (10)$$

### 3. Three Gompertz fractional $\nabla$ -difference equations

The natural  $\nabla$ -analogue of (1) is

$$\nabla y = (\ominus r)(a + L_y)y, \quad y(t_0) = y_0, \quad (11)$$

whose unique solution is easily found using the same method used in [Cuchta and Streipert 2020]. There are many ways in which a  $\nabla$ -fractional analogue of (11) could be defined, depending on which operators in the equation are made fractional. We will consider three such analogues in this article. In light of (3), we will allow the order of our fractional derivative to lie in the interval  $(0, 1.5)$ , where  $\nu = 1$  corresponds to the nonfractional  $\nabla$  difference. First, we preserve the nonfractional difference on  $y$  from (11), but we use the fractional  $\nabla$ -logarithm (8) to obtain

$$\nabla y = (\ominus r)y(a + L_{y,\nu}), \quad y(t_0) = y_0. \quad (12)$$

If instead the difference on  $y$  in (11) is a fractional difference, but the  $\nabla$ -logarithm (9) is used, then we obtain

$$\nabla_{t_0}^\nu y(t) = (\ominus r)y(a + \ell_{y,\nu}), \quad y(t_0 + 1) = y_0. \quad (13)$$

Finally, if we make the difference on  $y$  in (11) a fractional difference and use the fractional  $\nabla$ -logarithm (10), then we obtain

$$\nabla_{t_0}^\nu y(t) = (\ominus r)y(a + \Lambda_{y,\nu}), \quad y(t_0 + 1) = y_0. \quad (14)$$

Note that all these logarithms reduce to  $L_y$  as  $\nu \rightarrow 1$ . We now find a closed-form solution of (12) in terms of the  $\nabla$ -Mittag-Leffler function.

**Theorem 5.** *If  $0 < \nu < 1$ ,  $r < \frac{1}{2}$ , and, for  $t \in \mathbb{N}_{t_0}$ ,*

$$\sum_{k=t_0+1}^t E_{\ominus r, \nu, \nu-1}(t-k+1+t_0, t_0) H_{-\nu}(k, t_0) \neq -1 - \frac{1-r}{ar}, \tag{15}$$

*then the initial value problem (12) has the unique solution  $y(t) = y_0 E_p(t, t_0)$  for  $t \in \mathbb{N}_{t_0}$ , where*

$$p(t) = (\ominus r)a + (\ominus r)(E_{\ominus r, \nu, \nu-1}(\cdot, t_0) * a H_{-\nu}(\cdot, t_0))(t).$$

*Proof.* If  $z(t) = a + L_{y, \nu}(t, t_0)$ , then

$$(\nabla_{t_0}^\nu z)(t) = a H_{-\nu}(t, t_0) + \frac{\nabla y(t)}{y(t)}. \tag{16}$$

Now compute

$$\begin{aligned} a + L_{y, \nu}(t_0 + 1, t_0) &= a + \left( \nabla_{t_0}^{-\nu} \frac{\nabla y}{y} \right)(t_0 + 1) \\ &= a + \int_{t_0}^{t_0+1} H_{\nu-1}(t_0 + 1, \rho(s)) \frac{\nabla y(s)}{y(s)} \nabla s \\ &= a + H_{\nu-1}(t_0 + 1, t_0) \frac{\nabla y(t_0 + 1)}{y(t_0 + 1)} = a + 1 - \frac{y_0}{y(t_0 + 1)}. \end{aligned}$$

We expand the difference equation (12) at  $t = t_0 + 1$  to get

$$y(t_0 + 1) - y_0 = -\frac{r}{1-r} y(t_0 + 1) \left( a + 1 - \frac{y_0}{y(t_0 + 1)} \right),$$

and hence

$$y(t_0 + 1) = \frac{y_0}{1 + ra}.$$

Immediately, we obtain  $z(t_0 + 1) = a + L_{y, \nu}(t_0 + 1, t_0) = a(1 - r)$ . Using the definition of  $z$ , (16), and dividing the difference equation (12) by  $y$ , the following initial value problem for  $z$  is derived:

$$\nabla_{t_0}^\nu z + (-\ominus r)z = a H_{-\nu}(t, t_0), \quad z(t_0 + 1) = a(1 - r). \tag{17}$$

Theorem 3 implies that (17) has the unique solution

$$z(t) = a + (E_{\ominus r, \nu, \nu-1}(\cdot, t_0) * a H_{-\nu}(\cdot, t_0))(t).$$

Therefore, we have the initial value problem  $\nabla y = (\ominus r)zy$ ,  $y(t_0) = y_0$ . We need to ensure that the function  $h : \mathbb{N}_{t_0} \rightarrow \mathbb{R}$  given by  $h(t) = (\ominus r)z(t)$  obeys  $h(t) \neq 1$ . Expanding out the definition of  $h$  reveals

$$1 \neq \left( \frac{-r}{1-r} \right) \left( a + ak \sum_{k=t_0}^t E_{\ominus r, \nu, \nu-1}(t-k+1+t_0, t_0) H_{-\nu}(k, t_0) \right),$$

and by algebraic rearrangement, this becomes the condition (15). Thus, we may appeal to Theorem 1 to complete the proof.  $\square$

Theorem 5 provides an impractical solution for numerical computation for a variety of reasons. In particular, the  $\nabla$ -Mittag-Leffler function requires the evaluation of an infinite series, the order is restricted to  $0 < \nu < 1$ , and condition (15) is not trivial to verify. So, we now derive a recurrence relation for numerical computation of the solution of (12).

**Theorem 6.** *The initial value problem (12) is algebraically equivalent to the recurrence*

$$y(t) = y_0 \prod_{k=t_0+1}^t \frac{1}{1 - (\ominus r)(a + z(k))}, \quad t \in \mathbb{N}_{t_0},$$

where

$$z(t) = -ra - (1 - r) \sum_{k=t_0+1}^{t-1} \frac{\Gamma(t - k - \nu)z(k)}{\Gamma(-\nu)\Gamma(t - k + 1)}, \quad t \in \mathbb{N}_{t_0+1}. \quad (18)$$

*Proof.* If  $z(t) = L_{y,\nu}(t, t_0)$ , then

$$\nabla_{t_0}^\nu z(t) = \frac{y^\nabla(t)}{y(t)} = (\ominus r)(a + z(t)).$$

Apply Lemma 4 to the left-hand side to get

$$z(t) + \sum_{k=t_0+1}^{t-1} \frac{\Gamma(t - k - \nu)z(k)}{\Gamma(-\nu)\Gamma(t - k + 1)} = (\ominus r)(a + z(t)). \quad (19)$$

We obtain (18) by rearranging (19) and solving for  $z(t)$ . Substituting  $z(t)$  back into (12), we observe

$$\nabla y(t) = (\ominus r)(a + z(t))y(t), \quad y(t_0) = y_0.$$

By Theorem 1, we have  $y(t) = y_0 E_{(\ominus r)(a+z(t))}(t, t_0)$ , and applying (5) completes the proof.  $\square$

**Theorem 7.** *The initial value problem (13) is algebraically equivalent to the following recurrence for  $t > t_0 + 1$ :*

$$y(t) = \frac{-\sum_{k=t_0+1}^{t-1} H_{-\nu-1}(t, k-1)y(k)}{1-r+r(a+1+\sum_{k=t_0+1}^{t-1} \frac{1}{y(k)} \sum_{j=t_0+1}^k H_{-\nu-1}(k, j-1)y(j))}, \quad t \in \mathbb{N}_{t_0+2}.$$

*Proof.* Expand the fractional derivative and the logarithm in (13) to obtain

$$\int_{t_0}^t H_{-\nu-1}(t, \rho(s))y(s) \nabla s = (\ominus r)y(t) \left( a + \int_{t_0}^t \frac{\nabla_{t_0}^\nu y(\tau)}{y(\tau)} \nabla \tau \right).$$

Separating the final term of the  $\nabla$ -integrals and rearranging algebraically yields

$$\begin{aligned}
 y(t) &= \frac{((\ominus r) - 1) \int_{t_0}^{t-1} H_{-\nu-1}(t, \rho(s))y(s) \nabla s}{1 - (\ominus r)(a + 1 + \int_{t_0}^{t-1} \frac{\nabla_{t_0}^\nu y(\tau)}{y(\tau)} \nabla \tau)} \\
 &= \frac{((\ominus r) - 1) \int_{t_0}^{t-1} H_{-\nu-1}(t, \rho(s))y(s) \nabla s}{1 - (\ominus r)(a + 1 + \int_{t_0}^{t-1} \frac{1}{y(\tau)} \int_{t_0}^\tau H_{-\nu-1}(\tau, \rho(s))y(s) \nabla s \nabla \tau)},
 \end{aligned}$$

and writing the  $\nabla$ -integrals as summations, replacing  $(\ominus r)$  with  $-r/(1 - r)$ , and routine algebraic simplification completes the proof.  $\square$

**Theorem 8.** *The initial value problem (14) is algebraically equivalent to the recurrence*

$$y(t) = \frac{-\sum_{k=t_0+1}^{t-1} H_{-\nu-1}(t, k-1)y(k)}{1-r+r(a+1+\sum_{k=t_0+1}^{t-1} \frac{H_{-\nu-1}(t,k-1)}{y(k)} \sum_{j=t_0+1}^k H_{-\nu-1}(k, j-1)y(j))}, \quad t \in \mathbb{N}_{t_0+2}. \tag{20}$$

*Proof.* Expand (14) using definitions to get

$$\int_{t_0}^t H_{-\nu-1}(t, \rho(\tau))y(\tau) \nabla \tau = (\ominus r)y(t) \left( a + \int_{t_0}^t H_{\nu-1}(t, \rho(\tau)) \frac{\nabla_{t_0}^\nu y(\tau)}{y(\tau)} \nabla \tau \right).$$

Separate the final terms from the  $\nabla$ -integrals and rearrange algebraically to obtain

$$\begin{aligned}
 y(t) &= \frac{((\ominus r) - 1) \int_{t_0}^{t-1} H_{-\nu-1}(t, \rho(\tau))y(\tau) \nabla \tau}{1 - (\ominus r)a - (\ominus r) - (\ominus r) \int_{t_0}^{t-1} H_{\nu-1}(t, \rho(\tau)) \frac{\nabla_{t_0}^\nu y(\tau)}{y(\tau)} \nabla \tau} \\
 &= \frac{((\ominus r) - 1) \int_{t_0}^{t-1} H_{-\nu-1}(t, \rho(\tau))y(\tau) \nabla \tau}{1 - (\ominus r)(a + 1 + \int_{t_0}^{t-1} \frac{H_{\nu-1}(t,\rho(\tau))}{y(\tau)} \int_{t_0}^\tau H_{-\nu-1}(\tau, \rho(s))y(s) \nabla s \nabla \tau)},
 \end{aligned}$$

and writing the  $\nabla$ -integrals as summations and applying the definition of  $(\ominus r)$  completes the proof.  $\square$

#### 4. Curve fitting and comparison of models

We will now present curve fittings of our three models (12), (13), and (14), the model (3), and the continuous model to data that had previously fit with (continuous) Gompertz curves. In order to test the fit of our models, we have programmed<sup>1</sup> each model into the Python programming language and used the `curve_fit` function from `scipy.optimize`. According to its documentation, the `curve_fit` function ultimately implements the Levenberg–Marquardt algorithm for solving nonlinear least squares problems. To implement the solutions of our models, we used the

<sup>1</sup><https://github.com/tomcuchta/cuchtafinchamdiscretefractionalcompertz>

	$\nu$	$a$	$r$	$y_0$	RSS	SE
(12)	$5.58 \cdot 10^{-1}$	$-2.38 \cdot 10^{-1}$	$2.33 \cdot 10^0$	$1.78 \cdot 10^2$	$8.49 \cdot 10^6$	$2.43 \cdot 10^2$
(13)	$1.13 \cdot 10^0$	$-1.52 \cdot 10^0$	$4.89 \cdot 10^{-1}$	$2.07 \cdot 10^2$	$8.61 \cdot 10^6$	$2.44 \cdot 10^2$
(14)	$8.77 \cdot 10^{-1}$	$-1.73 \cdot 10^0$	$4.87 \cdot 10^{-1}$	$2.08 \cdot 10^2$	$8.60 \cdot 10^6$	$2.44 \cdot 10^2$
	$\nu$	$c$	$b$	$y_0$	RSS	SE
(3)	$5.88 \cdot 10^{-1}$	$4.46 \cdot 10^0$	$6.79 \cdot 10^{-1}$	$7.45 \cdot 10^1$	$8.61 \cdot 10^6$	$2.44 \cdot 10^2$
	$L_\infty$	$G$	$t_0$		RSS	SE
ctn	$5.85 \cdot 10^2$	$3.81 \cdot 10^{-1}$	$1.79 \cdot 10^0$		$8.56 \cdot 10^6$	$2.43 \cdot 10^2$

**Table 1.** Results of curve fittings to data from [Hilling et al. 2016b] and their goodness of fit. The continuous curve was given there by the three-parameter formula  $L_\infty e^{-e^{-G(t-t_0)}}$ .

recurrences in Theorems 6, 7, and 8 directly. The initial condition argument  $t_0$  was always chosen so that the data point with smallest independent variable agreed with the minimum of the domain of the solution. The effect this has is to horizontally shift the solutions so that they all “begin” at the same value of the independent variable in a given data set. Because those recurrence relations are self-referential at every iteration, they are computationally inefficient without optimizing the code in some way. So, we used the technique of memoization, meaning that we stored all newly generated values for a solution in a dictionary data structure to reference when computing the next value of the solution.

We present the parameters found, and as done in [Atıcı et al. 2017], we computed the residual sum of squared errors (RSS) and the standard error (SE). The RSS is defined by


$$\text{RSS} = \sum_{i=1}^n (y_i - Y(t_i))^2,$$

where  $n$  denotes the number of data points,  $(t_i, y_i)$  denotes the data points themselves, and  $Y(t_i)$  denotes the prediction of the fitted model at  $t_i$ , and SE is given by

$$\text{SE} = \sqrt{\frac{\text{RSS}}{n - k}},$$

where  $n$  denotes the total number of data points and  $k$  is the number of parameters of the model.

The article [Hilling et al. 2016b] concerns the relationship between the age (in years) and length (in millimeters) of 155 *ictalurus punctatus* (“channel catfish”) caught in Cheat Lake in West Virginia. The lengths of the fish were measured, and a technique to determine their age was performed. Their data set is publicly accessible



**Figure 1.** Left: Optimal models for the data from [Hilling et al. 2016b]. Right: Optimal models for the data from [Yang et al. 2019b]. Male (solid curve) and female (dashed curve) tail-length data. The curve for (12) is difficult to see because it is below the other curves.

male tail-length data						
	$\nu$	$a$	$r$	$y_0$	RSS	SE
(12)	$8.92 \cdot 10^{-1}$	$-8.34 \cdot 10^{-1}$	$9.55 \cdot 10^{-2}$	$2.94 \cdot 10^0$	$2.72 \cdot 10^4$	$5.33 \cdot 10^0$
(13)	$1.13 \cdot 10^0$	$-1.53 \cdot 10^0$	$5.79 \cdot 10^{-2}$	$2.85 \cdot 10^0$	$2.72 \cdot 10^4$	$5.33 \cdot 10^0$
(14)	$8.25 \cdot 10^{-1}$	$-1.87 \cdot 10^0$	$1.39 \cdot 10^{-1}$	$3.56 \cdot 10^0$	$2.72 \cdot 10^4$	$5.33 \cdot 10^0$
	$\nu$	$c$	$b$	$y_0$	RSS	SE
(3)	$2.30 \cdot 10^{-2}$	$1.15 \cdot 10^0$	$1.02 \cdot 10^{-1}$	$3.07 \cdot 10^0$	$2.72 \cdot 10^4$	$5.33 \cdot 10^0$
	$A$	$B$	$K$		RSS	SE
ctn	$1.17 \cdot 10^1$	$1.27 \cdot 10^0$	$4.52 \cdot 10^{-2}$		$2.72 \cdot 10^4$	$5.33 \cdot 10^0$

female tail-length data						
	$\nu$	$a$	$r$	$y_0$	RSS	SE
(12)	$8.60 \cdot 10^{-1}$	$-6.76 \cdot 10^{-1}$	$1.03 \cdot 10^{-1}$	$2.89 \cdot 10^0$	$1.54 \cdot 10^4$	$4.35 \cdot 10^0$
(13)	$1.12 \cdot 10^0$	$-1.51 \cdot 10^0$	$4.62 \cdot 10^{-2}$	$2.84 \cdot 10^0$	$1.54 \cdot 10^4$	$4.35 \cdot 10^0$
(14)	$7.56 \cdot 10^{-1}$	$-1.72 \cdot 10^0$	$1.82 \cdot 10^{-1}$	$3.56 \cdot 10^0$	$1.54 \cdot 10^4$	$4.36 \cdot 10^0$
	$\nu$	$c$	$b$	$y_0$	RSS	SE
(3)	$3.58 \cdot 10^{-2}$	$1.14 \cdot 10^0$	$1.23 \cdot 10^{-1}$	$3.35 \cdot 10^0$	$1.54 \cdot 10^4$	$4.35 \cdot 10^0$
	$A$	$B$	$K$		RSS	SE
ctn	$1.05 \cdot 10^1$	$1.18 \cdot 10^0$	$3.95 \cdot 10^{-2}$		$1.53 \cdot 10^4$	$4.35 \cdot 10^0$

**Table 2.** Results of curve fittings to data from [Yang et al. 2019b] and their goodness of fit. The continuous curve was given there by the three-parameter formula  $Ae^{-Be^{-Kt}}$ .

[Hilling et al. 2016a] and contains columns for the age and length (both positive integers), which we used without modification. Their analysis of the various growth curves concluded that the von Bertalanffy growth (proportional to  $1 - e^{-k(t-t_0)}$ , commonly used in length-based methods for fish population estimates [Pauly and Morgan 1987]) was the model of best fit among those considered. The results of our work are summarized in Table 1 and visualized in Figure 1, left. We see that the five models have essentially identical RSS, indicating that they all fit equally well. The models (12), (14), and (3) have optimal  $\nu$  parameter between 0.55 and 0.87, while the optimal  $\nu$  for (13) is larger than 1. The fit of (12) is not monotone, while the other four models are visually quite similar.

In [Yang et al. 2019b], the logistic, Gompertz, and von Bertalanffy growth curves were fit to data collected about *plestiodon elegans* (a type of skink) pertaining to some of their physical traits — specifically, the tail length of thirteen male and eleven female individuals were measured once a week for 85 weeks. Their data set is publicly accessible [Yang et al. 2019a] and contains columns for the animal identification number, week number (both positive integers), and tail length in centimeters (accurate to two decimal places), which we used without modification. The study concluded that the continuous Gompertz model was the best fit for both sexes for tail length data. We have fit the models to male and female tail-length data, and our results are summarized in Table 2, and visualized in Figure 1, right. We observe that the RSS and SE are nearly identical for all curve fittings, indicating that the models fit similarly well. The models (12) and (14) have  $\nu$ -value between 0.74 and 0.89, (13) always has a  $\nu$  value larger than 1, and the model (3) always has small  $\nu$ .

## 5. Conclusion

We have investigated three new natural fractional analogues of the Gompertz  $\nabla$ -difference equation. We have found a closed form solution of one of them in terms of the  $\nabla$ -Mittag-Leffler function, but with necessarily restricted parameters. We derived recurrence relations for all three. We have fit the models to data sets and compared the fit to the existing fractional Gompertz model by Atıcı et al. It appears that both of the data sets were fit nearly equally well by all four models according to RSS and SE, and so using biological considerations or more advanced statistical techniques would be necessary to decide which model is best. We observed that the four discrete fractional models under consideration did not perform better than their continuous counterparts in terms of RSS or SE.

There are numerous directions of future research involving these models. Finding simple sufficient conditions for the regressivity of the solution of the first new fractional model is of great interest, not only for the fractional model but also for the original nonfractional model. Many qualitative properties remain unexplored,

including analysis of stability and oscillation of their solutions. One interesting fractional generalization that we did not consider is taking the orders of the fractional difference and fractional sum in the third new fractional logarithm to be different. Such a model would gain an extra parameter, and so it may fit better in terms of RSS and SE. When comparing models with different numbers of parameters, a more robust statistical analysis should be conducted, e.g., the Akaike information criterion method.

Our choice of fractional derivative had some consequences that complicated some aspects of the work. Most importantly, in the proof of Theorem 5, the fractional derivative of  $z(t) = a + L_{y,v}(t, t_0)$  was computed yielding

$$(\nabla_{t_0}^{\nu} z)(t) = aH_{-\nu}(t, t_0) + \frac{\nabla y(t)}{y(t)}.$$

Since we used the Riemann–Liouville fractional difference, the derivative of the constant  $a$  was not zero, and we consequently had to use a convolution via Theorem 3 to express the closed-form solution of the first new model. This raises the general question of defining fractional Gompertz equations using the Caputo fractional differences (and others), where the fractional derivative of a constant is zero. Such models may be easier to analyze analytically, but may have other unforeseen downsides.

### Acknowledgments

This research was made possible by NASA West Virginia Space Grant Consortium, Training Grant #NNX15AI01H. We also thank the referees for their valuable recommendations that improved the quality of this manuscript.

### References

- [Abdeljawad 2011] T. Abdeljawad, “On Riemann and Caputo fractional differences”, *Comput. Math. Appl.* **62**:3 (2011), 1602–1611. MR Zbl
- [Akin et al. 2020] E. Akin, N. Nesliye Pelen, I. Uğur Tiryaki, and F. Yalcin, “Parameter identification for Gompertz and logistic dynamic equations”, *PLoS ONE* **15**:4 (2020), art. id. e0230582.
- [Alves et al. 2019] W. J. Alves, E. B. Malheiros, N. K. Sakomura, E. P. da Silva, G. da Silva Viana, C. A. G. M. de Paula Reis, and R. M. Suzuki, “*In vivo* description of body growth and chemical components of egg-laying pullets”, *Livestock Sci.* **220** (2019), 221–229.
- [Artin 1964] E. Artin, *The gamma function*, Holt, Rinehart and Winston, New York, 1964. MR Zbl
- [Atıcı and Şengül 2010] F. M. Atıcı and S. Şengül, “Modeling with fractional difference equations”, *J. Math. Anal. Appl.* **369**:1 (2010), 1–9. MR Zbl
- [Atıcı and Eloe 2009] F. M. Atıcı and P. W. Eloe, “Initial value problems in discrete fractional calculus”, *Proc. Amer. Math. Soc.* **137**:3 (2009), 981–989. MR Zbl
- [Atıcı et al. 2017] F. M. Atıcı, M. Atıcı, M. Belcher, and D. Marshall, “A new approach for modeling with discrete fractional equations”, *Fund. Inform.* **151**:1-4 (2017), 313–324. MR

- [Bohner 2005] M. Bohner, “The logarithm on time scales”, *J. Difference Equ. Appl.* **11**:15 (2005), 1305–1306. MR Zbl
- [Bohner and Peterson 2001] M. Bohner and A. Peterson, *Dynamic equations on time scales: an introduction with applications*, Birkhäuser, Boston, MA, 2001. MR Zbl
- [Bolton et al. 2015] L. Bolton, A. H. J. J. Cloot, S. W. Schoombie, and J. P. Slabbert, “A proposed fractional-order Gompertz model and its application to tumour growth data”, *Math. Med. Biol.* **32**:2 (2015), 187–207. MR Zbl
- [Cuchta and Streipert 2020] T. Cuchta and S. Streipert, “Dynamic Gompertz model”, *Appl. Math. Inf. Sci.* **14**:1 (2020), 9–17. MR
- [Dzieliński et al. 2010] A. Dzieliński, D. Sierociuk, and G. Sarwas, “Some applications of fractional order calculus”, *Bull. Polish Acad. Sci. Tech. Sci.* **58**:4 (2010), 583–592. Zbl
- [Easton 1999] D. M. Easton, “X-ray survival as Gompertz growth in number killed”, *J. Theoret. Biol.* **196**:1 (1999), 1–8.
- [Elaydi 2005] S. Elaydi, *An introduction to difference equations*, 3rd ed., Springer, 2005. MR Zbl
- [Ferreira 2013] R. A. C. Ferreira, “Existence and uniqueness of solution to some discrete fractional boundary value problems of order less than one”, *J. Difference Equ. Appl.* **19**:5 (2013), 712–718. MR Zbl
- [Frunzo et al. 2019] L. Frunzo, R. Garra, A. Giusti, and V. Luongo, “Modeling biological systems with an improved fractional Gompertz law”, *Commun. Nonlinear Sci. Numer. Simul.* **74** (2019), 260–267. MR Zbl
- [Gompertz 1825] B. Gompertz, “On the nature of the function expressive of the law of human mortality, and on a new mode of determining the value of life contingencies”, *Phil. Trans. R. Soc. Lond.* **115** (1825), 513–583.
- [Goodrich and Peterson 2015] C. Goodrich and A. C. Peterson, *Discrete fractional calculus*, Springer, 2015. MR Zbl
- [Hilling et al. 2016a] C. Hilling, S. Welsh, and D. Smith, “Data from: age, growth and fall diet of channel catfish in Cheat Lake, West Virginia”, Dryad, dataset, 2016, available at <https://doi.org/10.5061/dryad.8583t>.
- [Hilling et al. 2016b] C. D. Hilling, S. A. Welsh, and D. M. Smith, “Age, growth and fall diet of channel catfish in Cheat Lake, West Virginia”, *J. Fish. Wildl. Manag.* **7**:2 (2016), 304–314.
- [Jane et al. 2020] S. A. Jane, F. A. Fernandes, E. M. Silva, J. A. Muniz, T. J. Fernandes, and G. V. Pimentel, “Adjusting the growth curve of sugarcane varieties using nonlinear models”, *Ciência Rural* **50**:3 (2020), art. id. e20190408.
- [Kelley and Peterson 2010] W. G. Kelley and A. C. Peterson, *The theory of differential equations: classical and qualitative*, 2nd ed., Springer, 2010. MR Zbl
- [Kuttner 1957] B. Kuttner, “On differences of fractional order”, *Proc. London Math. Soc.* (3) **7** (1957), 453–466. MR Zbl
- [Laird 1964] A. K. Laird, “Dynamics of tumour growth”, *British J. Cancer* **18**:3 (1964), 490–502.
- [Miller and Ross 1989] K. S. Miller and B. Ross, “Fractional difference calculus”, pp. 139–152 in *Univalent functions, fractional calculus, and their applications* (Kōriyama, 1988), edited by H. M. Srivastava and S. Owa, Horwood, Chichester, 1989. MR Zbl
- [Nobile et al. 1982] A. G. Nobile, L. M. Ricciardi, and L. Sacerdote, “On Gompertz growth model and related difference equations”, *Biol. Cybernetics* **42** (1982), 221–229. Zbl
- [Pauly and Morgan 1987] D. Pauly and G. R. Morgan (editors), *Length-based methods in fisheries research* (Sicily, Italy, 1985), ICLARM Conference Proceedings **13**, International Center for Living Aquatic Resources Management, Safat, Kuwait, 1987.

- [Pezzini et al. 2019] R. V. Pezzini, A. C. Filho, F. Carini, C. T. Bandeira, J. A. Kleinpaul, and D. L. Silveira, “Modeling the growth of sudangrass cultivars at sowing times”, *J. Agric. Sci.* **11**:14 (2019), 84–95.
- [Ross 1975] B. Ross, “A brief history and exposition of the fundamental theory of fractional calculus”, pp. 1–36 in *Fractional calculus and its applications*, edited by B. Ross, Lecture Notes in Mathematics **457**, Springer, 1975. Zbl
- [Satoh 2000] D. Satoh, “A discrete Gompertz equation and a software reliability growth model”, *IEICE Trans. Info. Sys.* **E83-D**:7 (2000), 1508–1513.
- [Solís-Pérez et al. 2019] J. E. Solís-Pérez, J. F. Gómez-Aguilar, R. F. Escobar-Jiménez, L. Torres, and V. H. Olivares-Peregrino, “Parameter estimation of fractional Gompertz model using cuckoo search algorithm”, pp. 81–95 in *Fractional derivatives with Mittag-Leffler kernel*, edited by J. F. Gómez et al., Stud. Syst. Decis. Control **194**, Springer, 2019. MR Zbl
- [Wang et al. 2014] J. Wang, H. Xiang, and F. Chen, “Existence of positive solutions for a discrete fractional boundary value problem”, *Adv. Difference Equ.* (2014), art. id. 253. MR Zbl
- [Winsor 1932] C. P. Winsor, “The Gompertz curve as a growth curve”, *Proc. Natl. Acad. Sci. USA* **18**:1 (1932), 1–8. Zbl
- [Wu et al. 2015] G.-C. Wu, D. Baleanu, S.-D. Zeng, and Z.-G. Deng, “Discrete fractional diffusion equation”, *Nonlinear Dynam.* **80**:1-2 (2015), 281–286. MR Zbl
- [Yang et al. 2019a] C. Yang, J. Zhao, R. E. Diaz, and N. Lyu, “Data from: development of sexual dimorphism in two sympatric skinks with different growth rates”, Dryad, dataset, 2019, available at <https://doi.org/10.5061/dryad.nb25502>.
- [Yang et al. 2019b] C. Yang, J. Zhao, R. E. Diaz, and N. Lyu, “Development of sexual dimorphism in two sympatric skinks with different growth rates”, *Ecol. Evol.* **9**:13 (2019), 7752–7760.

Received: 2020-04-29    Revised: 2020-08-06    Accepted: 2020-08-10

tcuchta@fairmontstate.edu    *Department of Computer Science and Mathematics,  
Fairmont State University, Fairmont, WV, United States*

brookefincham@gmail.com    *Department of Computer Science and Mathematics,  
Fairmont State University, Fairmont, WV, United States*



# involve

msp.org/involve

## INVOLVE YOUR STUDENTS IN RESEARCH

*Involve* showcases and encourages high-quality mathematical research involving students from all academic levels. The editorial board consists of mathematical scientists committed to nurturing student participation in research. Bridging the gap between the extremes of purely undergraduate research journals and mainstream research journals, *Involve* provides a venue to mathematicians wishing to encourage the creative involvement of students.

### MANAGING EDITOR

Kenneth S. Berenhaut Wake Forest University, USA

### BOARD OF EDITORS

Colin Adams	Williams College, USA	Robert B. Lund	Clemson University, USA
Arthur T. Benjamin	Harvey Mudd College, USA	Gaven J. Martin	Massey University, New Zealand
Martin Bohner	Missouri U of Science and Technology, USA	Mary Meyer	Colorado State University, USA
Amarjit S. Budhiraja	U of N Carolina, Chapel Hill, USA	Frank Morgan	Williams College, USA
Pietro Cerone	La Trobe University, Australia	Mohammad Sal Moslehian	Ferdowsi University of Mashhad, Iran
Scott Chapman	Sam Houston State University, USA	Zuhair Nashed	University of Central Florida, USA
Joshua N. Cooper	University of South Carolina, USA	Ken Ono	Univ. of Virginia, Charlottesville
Jem N. Corcoran	University of Colorado, USA	Yuval Peres	Microsoft Research, USA
Toka Diagana	University of Alabama in Huntsville, USA	Y.-F. S. Pétermann	Université de Genève, Switzerland
Michael Dorff	Brigham Young University, USA	Jonathon Peterson	Purdue University, USA
Sever S. Dragomir	Victoria University, Australia	Robert J. Plemmons	Wake Forest University, USA
Joel Foisy	SUNY Potsdam, USA	Carl B. Pomerance	Dartmouth College, USA
Errin W. Fulp	Wake Forest University, USA	Vadim Ponomarenko	San Diego State University, USA
Joseph Gallian	University of Minnesota Duluth, USA	Bjorn Poonen	UC Berkeley, USA
Stephan R. Garcia	Pomona College, USA	József H. Przytycki	George Washington University, USA
Anant Godbole	East Tennessee State University, USA	Richard Rebarber	University of Nebraska, USA
Ron Gould	Emory University, USA	Robert W. Robinson	University of Georgia, USA
Sat Gupta	U of North Carolina, Greensboro, USA	Javier Rojo	Oregon State University, USA
Jim Haglund	University of Pennsylvania, USA	Filip Saidak	U of North Carolina, Greensboro, USA
Johnny Henderson	Baylor University, USA	Hari Mohan Srivastava	University of Victoria, Canada
Glenn H. Hurlbert	Virginia Commonwealth University, USA	Andrew J. Sterge	Honorary Editor
Charles R. Johnson	College of William and Mary, USA	Ann Trenk	Wellesley College, USA
K. B. Kulasekera	Clemson University, USA	Ravi Vakil	Stanford University, USA
Gerry Ladas	University of Rhode Island, USA	Antonia Vecchio	Consiglio Nazionale delle Ricerche, Italy
David Larson	Texas A&M University, USA	John C. Wierman	Johns Hopkins University, USA
Suzanne Lenhart	University of Tennessee, USA	Michael E. Zieve	University of Michigan, USA
Chi-Kwong Li	College of William and Mary, USA		

### PRODUCTION

Silvio Levy, Scientific Editor

Cover: Alex Scorpan

See inside back cover or [msp.org/involve](http://msp.org/involve) for submission instructions. The subscription price for 2020 is US \$205/year for the electronic version, and \$275/year (+\$35, if shipping outside the US) for print and electronic. Subscriptions, requests for back issues and changes of subscriber address should be sent to MSP.

Involve (ISSN 1944-4184 electronic, 1944-4176 printed) at Mathematical Sciences Publishers, 798 Evans Hall #3840, c/o University of California, Berkeley, CA 94720-3840, is published continuously online. Periodical rate postage paid at Berkeley, CA 94704, and additional mailing offices.

Involve peer review and production are managed by EditFLOW<sup>®</sup> from Mathematical Sciences Publishers.

PUBLISHED BY

 **mathematical sciences publishers**  
nonprofit scientific publishing

<http://msp.org/>

© 2020 Mathematical Sciences Publishers

# involve

2020 vol. 13 no. 4

Structure constants of $\mathcal{U}(\mathfrak{sl}_2)$	541
ALEXIA GOURLEY AND CHRISTOPHER KENNEDY	
Conjecture $\mathcal{C}$ holds for some horospherical varieties of Picard rank 1	551
LELA BONES, GARRETT FOWLER, LISA SCHNEIDER AND RYAN M. SHIFLER	
Condensed Ricci curvature of complete and strongly regular graphs	559
VINCENT BONINI, CONOR CARROLL, UYEN DINH, SYDNEY DYE, JOSHUA FREDERICK AND ERIN PEARSE	
On equidistant polytopes in the Euclidean space	577
CSABA VINCZE, MÁRK OLÁH AND LETÍCIA LENGYEL	
Polynomial values in Fibonacci sequences	597
ADI OSTROV, DANNY NEFTIN, AVI BERMAN AND REYAD A. ELRAZIK	
Stability and asymptotic analysis of the Föllmer–Schweizer decomposition on a finite probability space	607
SARAH BOESE, TRACY CUI, SAMUEL JOHNSTON, SYLVIE VEGA-MOLINO AND OLEKSII MOSTOVYI	
Eigenvalues of the sum and product of anticommuting matrices	625
VADIM PONOMARENKO AND LOUIS SELSTAD	
Combinatorial random knots	633
ANDREW DUCHARME AND EMILY PETERS	
Conjugation diameter of the symmetric groups	655
ASSAF LIBMAN AND CHARLOTTE TARRY	
Existence of multiple solutions to a discrete boundary value problem with mixed periodic boundary conditions	673
KIMBERLY HOWARD, LONG WANG AND MIN WANG	
Minimal flag triangulations of lower-dimensional manifolds	683
CHRISTIN BIBBY, ANDREW ODESKY, MENGMENG WANG, SHUYANG WANG, ZIYI ZHANG AND HAILUN ZHENG	
Some new Gompertz fractional difference equations	705
TOM CUCHTA AND BROOKE FINCHAM	