

involve

a journal of mathematics

Counting profile strings from rectangular tilings

Anthony Petrosino, Alissa Schembor and Kathryn Haymaker



Counting profile strings from rectangular tilings

Anthony Petrosino, Alissa Schembor and Kathryn Haymaker

(Communicated by Arthur T. Benjamin)

We study the enumeration of profiles, which are outlines that occur when tiling a rectangular board with squares, dominoes, and trominoes. Profiles of length m correspond to a special subset of the set $\{0, 1, 2, 3\}^m$, called profile strings. Profiles and their corresponding strings first appeared in the enumeration of the tilings of rectangular $2 \times n$ and $3 \times n$ boards with squares, dominoes, and trominoes. Profiles also play a role in enumerating the tilings of an $m \times n$ board for any fixed $m \geq 2$. We describe how profiles arise when enumerating tilings, and we prove that the number of profile strings of length m equals $m \cdot 3^{m-1}$.

1. Background

A *tiling* of an $m \times n$ board using square tiles (1×1), domino tiles (1×2 or 2×1), and I -tromino tiles (1×3 or 3×1) is a configuration of tiles that covers every position in the board. In this paper we will use the word “tromino” to mean exclusively an I -tromino. Figure 1 shows a tiling of a 3×4 board using two squares, two dominoes, and two trominoes. Throughout this paper, the variable x denotes the placement of a square tile, y denotes the placement of a domino tile, and z denotes the placement of a tromino tile.

Read [1982] and Haymaker and Robertson [2017] determined recursions for the number of tilings of a $2 \times n$ board with squares and dominoes, and squares, dominoes, and trominoes, respectively, using the technique of building a tiling from the leftmost, topmost untiled position. The outline of the right-hand side of a partially tiled board is called a *profile*, and counting the number of profiles of length m is the main goal of this paper.

We first define a building process that begins on the leftmost, topmost open square.

Definition 1. Consider a blank or partially tiled $m \times n$ board. A *building move* is when a tile is placed in the leftmost column that has an uncovered square, in the topmost square that is uncovered.

MSC2020: 05A19, 05B45.

Keywords: rectangular tilings, enumeration, induction.




Figure 1. An example tiling of a 3×4 board with squares, dominoes, and trominoes.

There is a one-to-one correspondence between tilings of an $m \times n$ rectangular board and sequences of building moves that result in a complete $m \times n$ tiling [Read 1982]. That is, every tiling can be obtained from a unique sequence of building moves.

Definition 2. The *target square* of a partial tiling being built using building moves is the topmost square in the leftmost column with an uncovered square.

For an example of a tiling obtained using building moves, see Figure 2. In each partial tiling in Figure 2, the “ \times ” denotes the position of the target square.

Definition 3. The *m*-row *profile* of a partial tiling of an $m \times n$ board is the outline of the rightmost edge of the partial tiling created by building moves. A *profile string* of an *m*-row profile is a string of length *m* where positions are assigned values of 0, 1, 2, or 3 depending on how many squares to the right of the rightmost fully tiled column are covered with tiles. We define the *length* of the profile to be *m*, the number of rows in the profile.

For example, the profile of the second partial tiling shown in Figure 2 has corresponding profile string 110.

Definition 4. The *length-*m* A profile* is the profile whose outline is a vertical line. The *A profile string* is $\underbrace{0 \cdots 0}_m$, the all-zero string of length *m*.




Figure 2. Sequence of profile strings from the building process for this 3×4 tiling: $000 \rightarrow 110 \rightarrow 002 \rightarrow 302 \rightarrow 201 \rightarrow 110 \rightarrow 000$.

Next, we give an example of how building moves result in a sequence of profiles and profile strings.

Example 5 (illustrating profile strings with the 3×4 tiling). [Figure 2](#) begins with an untiled 3×4 board, placing the target square in the leftmost-topmost position, indicated by the “ \times ”. This profile is represented by the profile string 000 (the A profile). This is followed by the placement of a vertical domino, causing the target square to move down to the bottom row, and resulting in a profile string of 110. Next, a horizontal tromino is placed on top of the target square position, causing the target square to now move to the top row. Although this profile string appears to be 113, it should be noted that strings are denoted by the number of squares beyond the rightmost fully tiled column. Because there is one fully tiled column, the profile string reduces to 002. The process of constructing a tiling arrangement proceeds with the placement of squares, dominoes, or trominoes on top of the target square. Once the construction is complete, it will result in a 000 profile string, which is exactly how the example began.

Remark 6. The profile of a partially tiled board is the *shape* of the right-hand side of the board; the same shape/profile can result from different tile configurations. In fact, for any profile, it is possible to construct the profile shape using only “horizontal” tiles. To see this, consider a partial tiling that contains a vertical domino, for example. A specific example is the partial tiling on the far left of the middle row in [Figure 2](#). This partial tiling can be changed into a different partial tiling with only horizontal tiles by replacing the vertical domino with two squares stacked vertically. The outline of the right side of the board remains the same; i.e., the profile remains the same. Similarly, if a partial tiling contains a vertical tromino, that piece can be replaced by three vertically stacked squares, yielding the same profile. This property of profiles becomes important in [Section 3](#) when we count profiles using a construction method with only horizontal tiles.

2. Profile-type strings

To describe the shape of profiles in a numerical sense, we define profile-type strings. We use profile-type strings in order to further analyze the behavior of profiles resulting from the tiling of an $m \times n$ board. In [Section 3](#), we will show that profile-type strings give a combinatorial description of all possible profiles that could result from the tiling of an $m \times n$ board using square tiles, domino tiles, and I -tromino tiles.

Definition 7. A string x_1, x_2, \dots, x_m in $\{0, 1, 2, 3\}^m$ is called a *profile-type* string if it satisfies the following conditions:

- (1) There exists an $i \in \{1, 2, \dots, m\}$ such that $x_i = 0$.
- (2) If $x_i = 0$, then $x_j \neq 3$ for all $j > i$.

In words, a profile-type string is a string with at least one 0, where no 3 occurs to the right of a 0 in the string.

For example, the following are profile-type strings with $m = 5$: 31202, 11302, 00122. On the other hand, the following strings of length 5 are not profile-type strings: 12321, 00130, 00123.

The $m = 2$ case contains six profile-type strings

$$\begin{aligned} &00, 20, 10, \\ &30, 01, 02, \end{aligned} \tag{2-1}$$

while the $m = 3$ case contains 27 profile-type strings

$$\begin{aligned} &000, 300, 200, 110, 100, 330, 320, 310, 230, \\ &220, 210, 002, 001, 120, 130, 101, 202, 201, \\ &011, 010, 102, 302, 301, 012, 022, 021, 020. \end{aligned} \tag{2-2}$$

Proposition 8. *There are 108 profile-type strings for the $m = 4$ case.*

Proof. We enumerate the profile-type strings in $\{0, 1, 2, 3\}^4$ as follows. Let a, b, c, d represent a profile-type string. Then we count the following cases:

- (1) If $a = 0$, then $b, c, d \in \{0, 1, 2\}$ by [Definition 7](#), so there are $3^3 = 27$ such strings.
- (2) If $a \in \{1, 2, 3\}$ and $b = 0$, then $c, d \in \{0, 1, 2\}$, so there are 27 such strings.
- (3) If $a, b \in \{1, 2, 3\}$ and $c = 0$, then $d \in \{0, 1, 2\}$, so there are 27 such strings.
- (4) If $a, b, c \in \{1, 2, 3\}$ and $d = 0$, there are 27 strings of this type, given the options for a, b , and c .

The total number of such strings is $4(27) = 108$. □

Theorem 9. *The number of profile-type strings of length m is equal to $m \cdot 3^{m-1}$.*

Proof. We will go through each position of a length- m profile-type string, anchor a 0 onto each position, and see how many different values are possible for all of the positions other than the fixed position. If there is a 0 anchored on a fixed position, we will say that there cannot be any 0 on any position before the fixed position, in order to avoid repeats in the enumeration.

Let x_i represent the i -th entry in the profile-type string.

The cases begin as follows:

- (1) Use x_1 as the anchor, with $x_1 = 0$; then $x_2, \dots, x_m \in \{0, 1, 2\}$ as there cannot be a 3 following a 0 in a profile-type string, by definition.
- (2) Next use x_2 as the anchor position, so $x_2 = 0$; then $x_1 \in \{1, 2, 3\}$ as there are no restrictions on the first position, besides not having 0 to avoid double

counting. Further, $x_3, \dots, x_m \in \{0, 1, 2\}$ as there cannot be a 3 following a 0 in a profile-type string, by definition.

In general, for $k = 1, \dots, m$, the k -th case is to anchor $x_k = 0$, and assume $x_i \in \{1, 2, 3\}$ for each i satisfying $1 \leq i < k$. Therefore by the definition of profile-type strings, $x_j \in \{0, 1, 2\}$ for each j satisfying $k < j \leq m$.

Each of these m cases has 3^{m-1} strings: there are three options for every position, other than in the fixed position in which we anchored a 0; as there are $m - 1$ positions with three options (every position but the fixed position in a length- m profile-type string), there are 3^{m-1} options. There are m cases, so the total number of profile-type strings is $m \cdot 3^{m-1}$. □

3. Counting profiles

In this section, we prove that the profile-type strings of length m are in one-to-one correspondence with the profile strings of length m . That is, we can drop the word “type” in the name profile-type strings!

As a motivating example, we first look at the case of $m = 2$.

Example 10 (profiles and profile-type strings for $m = 2$). Notice that the length-2 strings that satisfy the conditions in Definition 7 are given in (2-1).

On the other hand, the profiles with two rows are in Figure 3, and their corresponding profile strings are

$$A : 00, \quad B : 20, \quad C : 10, \quad D : 30, \quad E : 01, \quad F : 02.$$

By inspection, we see that the profile strings for profiles A–F in Figure 3 are precisely the profile-type strings listed in (2-1).

Definition 11. A *profile construction* is the process of sequentially adding squares, dominoes, and trominoes to the A-profile until the desired profile results. The order in which squares, dominoes, and trominoes are added must correspond with proper




Figure 3. This figure shows all six profiles for $m = 2$. We see that there is a one-to-one correspondence between these profiles and the profile-type strings for $m = 2$.




Figure 4. Deconstruction of the length-3 profile 202 to the A -profile in the -1 column position.

building moves; that is, at each step a square, domino, or tromino must be added onto the target square.

Definition 12. A *profile deconstruction* is the process of sequentially removing squares, dominoes, and trominoes from a profile until all that is left is the A -profile. The order in which squares, dominoes, and trominoes are removed must correspond with the construction of a profile; that is, squares, dominoes, and trominoes must be removed based on where the previous target square was located.

Remark 13. Profile deconstructions are not unique; there are different ways to deconstruct the same profile.

Example 14 (deconstruction and reconstruction with an additional row). We will shortly see an induction proof that involves adding a row to a profile. In this example we show a deconstruction of the profile with string 202 to the A -profile in column -1 . Then we add a row and construct the profile with string 2020, as a motivating example for the inductive step.

In [Figure 4](#), the initial profile will be deconstructed to the -1 column position. In order to do so, the deconstruction first removes a tromino from the bottom row, followed by the removal of a square from the middle row, and finally the removal of a tromino from the top row. This allows for the figure to be fully deconstructed to the A -profile in column -1 .

In [Figure 5](#), an additional row is then added to the profile. This begins the process of reconstructing the profile 2020 by adding the following (in this particular order, from top to bottom): tromino, square, tromino, square.




Figure 5. Construction of the $m = 4$ profile 2020 by undoing the deconstruction in [Figure 4](#), with an additional row.

Lemma 15. *Every profile of length m can be deconstructed to the A -profile of length m in column -1 .*

Proof. The following algorithm deconstructs a profile of length m to the A -profile in the -1 column. Let $\mathcal{P}_m \subseteq \{0, 1, 2, 3\}^m$ be the set of profile strings of length m .

Input: $x_1, x_2, \dots, x_m \in \mathcal{P}_m$. The algorithm runs through $k = m, m - 1, \dots, 1$. If $x_k \in \{0, 1, 2\}$, remove a length- $(x_k + 1)$ -tile from the k -th row and proceed to row $k - 1$. If $x_k = 3$, remove a combined square and tromino from the k -th row and proceed to row $k - 1$. The algorithm terminates when $k = 1$. The resulting profile is an A -profile at the -1 column. \square

To reconstruct the profile from A at the -1 column, place the removed pieces from top to bottom; in the rows consisting of a square followed by a tromino, place the squares first. Then in the 0 column, place the trominoes from top to bottom. These steps comprise two phases of reconstruction.

Two-phase reconstruction, starting with the A -profile in the -1 column:

- **Phase 1:** If $x_k \in \{0, 1, 2\}$, then place a length- $(x_k + 1)$ tile in row k . If $x_k = 3$, phase 1 is to place a square tile in row k .
- **Phase 2:** Place trominoes in all rows where $x_k = 3$.

Notice that this two-phase process is designed to follow the rules of building moves and target squares, while resulting in the desired profile.

Example 16 (two-phase reconstruction of 3120).

- **Phase 1:** In the first portion of [Figure 6](#), we see the result of phase 1 of construction to obtain the profile with string 3120. Note that for the desired profile, $x_1 = 3$, $x_2 = 1$, $x_3 = 2$, and $x_4 = 0$. Begin phase 1 by placing a square (length-1 tile) in row 1, as $x_1 = 3$. Next, place a domino (length-2 tile) in row 2, as $x_2 = 1$. Next, place a tromino (length-3 tile) in row 3, as $x_3 = 2$. Next, place a square (length-1 tile) in row 4, as $x_4 = 0$.
- **Phase 2:** Place a tromino (length-3 tile) in row 1, as $x_1 = 3$. Phase 2 is also shown in [Figure 6](#).

Next we give a lemma that contributes to establishing the one-to-one correspondence between profiles and profile-type strings.

Lemma 17. *If x_1, \dots, x_m, x_{m+1} is a profile-type string of length $m + 1$ where at least one of x_1, \dots, x_m equals 0, then the string $\sigma = x_1, x_2, \dots, x_m$ is a profile-type string of length m .*

Proof. Assume that x_1, \dots, x_m, x_{m+1} is a profile-type string, and at least one of x_1, \dots, x_m equals 0. Let σ denote the string x_1, x_2, \dots, x_m . Condition (1) in [Definition 7](#) holds for the string σ , since it contains at least one zero.




Figure 6. Two phase construction of the profile with string 3120, starting at the -1 column position.

To show that condition (2) in Definition 7 holds, we proceed by contradiction. Suppose that the string σ is not a profile-type string of length m because it violates condition (2). If σ violates condition (2), then there exist i, j such that $1 \leq i < j \leq m$, and $x_i = 0$ and $x_j = 3$ in σ . The entries x_i and x_j are identical in the original string $x_1, x_2, \dots, x_m, x_{m+1}$, which implies that it also violates condition (2). This is a contradiction as we assumed that the original string was a profile-type string. \square

The main result of this paper is next.

Theorem 18. *The m -row profiles are in one-to-one correspondence with the profile-type strings of length m .*

Proof. The theorem contains two claims:

- (1) An m -row profile string is a profile-type string of length m (i.e., a string that satisfies Definition 7).
- (2) A profile-type string of length m corresponds to an m -row profile.

To prove the first claim, let x_1, \dots, x_m be a profile string of length m corresponding to a profile P . By the definition of profile strings, at least one entry x_j must be 0, so the string satisfies condition (1) in Definition 7. Suppose that $x_j = 0$ and $x_k = 3$ for some $k > j$. Then in the construction of the profile P , a square, domino, or tromino must have been placed in the k -th row, while the j -th row contained an empty square in a position farther up and to the left. This contradicts the building move process, so no such j and k exist.

To prove the second claim, that every profile-type string corresponds to a profile string, we consider two cases and use induction.

We use induction on m . The base case is $m = 2$. See Example 10 for details of the one-to-one correspondence in the base case.

The inductive hypothesis is that there is a one-to-one correspondence between m -row profiles and profile-type strings of length m .

We will show the correspondence holds for $m + 1$.

Case 1: Suppose that the profile-type string $S = x_1, x_2, \dots, x_{m+1}$ has $x_{m+1} = 0$, and $x_j \neq 0$ for all $j = 1, \dots, m$. Then we construct a profile with the profile string S as follows:

Algorithm for constructing a length- $(m+1)$ profile, Case 1:

- (1) Start with the A -profile of length $m + 1$.
- (2) From top to bottom, place tiles of length x_i horizontally in row i for $i = 1, 2, \dots, m$.
- (3) Leave row $m + 1$ unchanged.

The resulting profile has corresponding string $x_1, x_2, \dots, x_m, x_{m+1}$, where x_{m+1} is the only entry that equals 0. (Notice that this case does not involve the inductive hypothesis.)

Case 2: For this case, consider the profile-type string $S = x_1, x_2, \dots, x_m, x_{m+1}$, where at least one of x_1, \dots, x_m equals 0.

Thus x_1, \dots, x_m is a profile-type string of length m by [Lemma 17](#), so by the inductive hypothesis there is a profile P of length m that corresponds to it.

Algorithm for constructing a length- $(m+1)$ profile, Case 2:

- (1) Deconstruct the profile P to the A -profile.
- (2) Add a row to the A -profile. We are now working with an $(m+1) \times n$ array.
- (3) Declare the column that the A -profile is on as column -1 . The outline of the resulting profile will be at most four columns to the right of this column.
- (4) Implement phase 1 of the profile reconstruction process for x_1, \dots, x_m in the first m rows. As $x_{m+1} \in \{0, 1, 2\}$, place a length- $(x_{m+1}+1)$ tile in row $m + 1$.
- (5) Implement phase 2 of the profile reconstruction process for x_1, \dots, x_m in the first m rows. (Since $x_{m+1} \neq 3$, there is no phase 2 for row $m + 1$).

This algorithm shows that there is a profile P with profile string S , proving our claim that every profile-type string S corresponds to a profile. \square

Corollary 19. *The number of m -row profiles is $m \cdot 3^{m-1}$.*

Proof. By [Theorem 18](#), the profiles are in one-to-one correspondence with profile-like strings, and by [Theorem 9](#) there are $m \cdot 3^{m-1}$ profile-type strings of length m . \square

Remark 20. Notice that the case where the only zero in the string S is $x_{m+1} = 0$ does not rely on the deconstruction and reconstruction of x_1, \dots, x_m . This is because constructing a profile where only the final entry is 0 requires only one phase, working from top to bottom and placing exactly the length- x_i tile in row i .

4. Conclusions and open questions

In this paper we proved the one-to-one correspondence of profile-type strings of length m with profile strings of length m , partly using induction. This understanding allowed for the creation of a basic formula for the number of profiles in terms of m for any given $m \times n$ board.

An open question that stems from this work is: how can we generalize the profile-type string definition for tilings with more types of tiles, for example, L -trominoes or quadrominoes?

Acknowledgments

The authors would like to thank Richard Filingeri for help with programming for counting profiles for small m .

References

- [Haymaker and Robertson 2017] K. Haymaker and S. Robertson, “Counting colorful tilings of rectangular arrays”, *J. Integer Seq.* **20**:5 (2017), art. id. 17.5.8. [MR](#) [Zbl](#)
- [Read 1982] R. C. Read, “The dimer problem for narrow rectangular arrays: a unified method of solution, and some extensions”, *Aequationes Math.* **24**:1 (1982), 47–65. [MR](#) [Zbl](#)

Received: 2020-06-04

Revised: 2020-08-16

Accepted: 2020-08-17

apetros1@villanova.edu

Villanova University, Villanova, PA, United States

aschembo@villanova.edu

Villanova University, Villanova, PA, United States

kathryn.haymaker@villanova.edu

Villanova University, Villanova, PA, United States

INVOLVE YOUR STUDENTS IN RESEARCH

Involve showcases and encourages high-quality mathematical research involving students from all academic levels. The editorial board consists of mathematical scientists committed to nurturing student participation in research. Bridging the gap between the extremes of purely undergraduate research journals and mainstream research journals, *Involve* provides a venue to mathematicians wishing to encourage the creative involvement of students.

MANAGING EDITOR

Kenneth S. Berenhaut Wake Forest University, USA

BOARD OF EDITORS

Colin Adams	Williams College, USA	Robert B. Lund	Clemson University, USA
Arthur T. Benjamin	Harvey Mudd College, USA	Gaven J. Martin	Massey University, New Zealand
Martin Bohner	Missouri U of Science and Technology, USA	Mary Meyer	Colorado State University, USA
Amarjit S. Budhiraja	U of N Carolina, Chapel Hill, USA	Frank Morgan	Williams College, USA
Pietro Cerone	La Trobe University, Australia	Mohammad Sal Moslehian	Ferdowsi University of Mashhad, Iran
Scott Chapman	Sam Houston State University, USA	Zuhair Nashed	University of Central Florida, USA
Joshua N. Cooper	University of South Carolina, USA	Ken Ono	Univ. of Virginia, Charlottesville
Jem N. Corcoran	University of Colorado, USA	Yuval Peres	Microsoft Research, USA
Toka Diagana	University of Alabama in Huntsville, USA	Y.-F. S. Pétermann	Université de Genève, Switzerland
Michael Dorff	Brigham Young University, USA	Jonathon Peterson	Purdue University, USA
Sever S. Dragomir	Victoria University, Australia	Robert J. Plemmons	Wake Forest University, USA
Joel Foisy	SUNY Potsdam, USA	Carl B. Pomerance	Dartmouth College, USA
Erin W. Fulp	Wake Forest University, USA	Vadim Ponomarenko	San Diego State University, USA
Joseph Gallian	University of Minnesota Duluth, USA	Bjorn Poonen	UC Berkeley, USA
Stephan R. Garcia	Pomona College, USA	József H. Przytycki	George Washington University, USA
Anant Godbole	East Tennessee State University, USA	Richard Rebarber	University of Nebraska, USA
Ron Gould	Emory University, USA	Robert W. Robinson	University of Georgia, USA
Sat Gupta	U of North Carolina, Greensboro, USA	Javier Rojo	Oregon State University, USA
Jim Haglund	University of Pennsylvania, USA	Filip Saidak	U of North Carolina, Greensboro, USA
Johnny Henderson	Baylor University, USA	Hari Mohan Srivastava	University of Victoria, Canada
Glenn H. Hurlbert	Virginia Commonwealth University, USA	Andrew J. Sterge	Honorary Editor
Charles R. Johnson	College of William and Mary, USA	Ann Trenk	Wellesley College, USA
K. B. Kulasekera	Clemson University, USA	Ravi Vakil	Stanford University, USA
Gerry Ladas	University of Rhode Island, USA	Antonia Vecchio	Consiglio Nazionale delle Ricerche, Italy
David Larson	Texas A&M University, USA	John C. Wierman	Johns Hopkins University, USA
Suzanne Lenhart	University of Tennessee, USA	Michael E. Zieve	University of Michigan, USA
Chi-Kwong Li	College of William and Mary, USA		

PRODUCTION

Silvio Levy, Scientific Editor


Cover: Alex Scorpan

See inside back cover or msp.org/involve for submission instructions. The subscription price for 2020 is US \$205/year for the electronic version, and \$275/year (+\$35, if shipping outside the US) for print and electronic. Subscriptions, requests for back issues and changes of subscriber address should be sent to MSP.

Involve (ISSN 1944-4184 electronic, 1944-4176 printed) at Mathematical Sciences Publishers, 798 Evans Hall #3840, c/o University of California, Berkeley, CA 94720-3840, is published continuously online. Periodical rate postage paid at Berkeley, CA 94704, and additional mailing offices.

Involve peer review and production are managed by EditFlow[®] from Mathematical Sciences Publishers.

PUBLISHED BY

 **mathematical sciences publishers**

nonprofit scientific publishing

<http://msp.org/>

© 2020 Mathematical Sciences Publishers

involve

2020

vol. 13

no. 5

Set-valued domino tableaux and shifted set-valued domino tableaux	721
FLORENCE MAAS-GARIÉPY AND REBECCA PATRIAS	
The first digit of the discriminant of Eisenstein polynomials as an invariant of totally ramified extensions of p-adic fields	747
CHAD AWTREY, ALEXANDER GAURA, SEBASTIAN PAULI, SANDI RUDZINSKI, ARIEL UY AND SCOTT ZINZER	
Counting pseudo progressions	759
JAY CUMMINGS, QUIN DARCY, NATALIE HOBSON, DREW HORTON, KEITH RHODEWALT, MORGAN THROCKMORTON AND RY ULMER-STRACK	
Growth series for graphs	781
WALTER LIU AND RICHARD SCOTT	
Peg solitaire in three colors on graphs	791
TARA C. DAVIS, ALEXSIS DE LAMERE, GUSTAVO SOPENA, ROBERTO C. SOTO, SONALI VYAS AND MELISSA WONG	
Disagreement networks	803
FLORIN CATRINA AND BRIAN ZILLI	
Rings whose subrings have an identity	823
GREG OMAN AND JOHN STROUD	
Simple graphs of order 12 and minimum degree 6 contain K_6 minors	829
RYAN ODENEAL AND ANDREI PAVELESCU	
Mixed volume of small reaction networks	845
NIDA OBATAKE, ANNE SHIU AND DILRUBA SOFIA	
Counting profile strings from rectangular tilings	861
ANTHONY PETROSINO, ALISSA SCHEMBOR AND KATHRYN HAYMAKER	
Isomorphisms of graded skew Clifford algebras	871
RICHARD G. CHANDLER AND NICHOLAS ENGEL	
Eta-quotients of prime or semiprime level and elliptic curves	879
MICHAEL ALLEN, NICHOLAS ANDERSON, ASIMINA HAMAKIOTES, BEN OLTSIK AND HOLLY SWISHER	