

involve

a journal of mathematics

Isomorphisms of graded skew Clifford algebras

Richard G. Chandler and Nicholas Engel



Isomorphisms of graded skew Clifford algebras

Richard G. Chandler and Nicholas Engel

(Communicated by Vadim Ponomarenko)

A regular algebra of global dimension $n + 1$ is often called a quantum \mathbb{P}^n . In 2011, Nafari, Vancliff and Zhang showed that graded skew Clifford algebras (GSCAs) could be used to classify most quadratic quantum \mathbb{P}^2 s. Some time later, Chandler, Tomlin and Vancliff used their work with certain families of GSCAs to develop a conjecture on the quantum space of a generic quadratic quantum \mathbb{P}^3 . These results suggest that (the isomorphism classes of) GSCAs are likely to play a fundamental role in the classification of quadratic quantum \mathbb{P}^3 s. In this article, we will discuss some of these results on GSCAs and discuss new results on isomorphisms between GSCAs.

Introduction

The notion of a graded skew Clifford algebra was defined in [Cassidy and Vancliff 2010] as a generalization of graded Clifford algebras (see [Le Bruyn 1995]). Therein, it was shown that GSCAs had many properties analogous to those of graded Clifford algebras. In addition, several families of quadratic regular graded skew Clifford algebras of global dimension 4 that were not twists of graded Clifford algebras were produced. It was shown that many of the algebras in these families had a one-dimensional line scheme [Shelton and Vancliff 2002a; 2002b] and a point scheme [Artin et al. 1990] consisting of twenty distinct points; hence, these algebras became candidates for generic quantum \mathbb{P}^3 s (in the language of [Chandler and Vancliff 2015]).

The line scheme and point scheme of some of these families were explicitly computed in [Chandler and Vancliff 2015; Tomlin and Vancliff 2018]. The line scheme of the generic member of the family studied in [Chandler and Vancliff 2015] was found to consist of a spatial elliptic curve, four planar elliptic curves and two nonsingular conics; the line scheme of the generic member of the family studied in [Tomlin and Vancliff 2018] was found to consist of four planar elliptic curves and four nonsingular conics. In both cases, the computations suggested that

MSC2020: 16S37, 16S38.

Keywords: graded skew Clifford algebras, μ -symmetry, isomorphisms.

the nonsingular conics were coming from a spatial elliptic curve in which one of the defining polynomials factored. This suggests the following conjecture.

Conjecture 0.1 [Chandler and Vancliff 2015]. There exists a class of generic quadratic quantum \mathbb{P}^3 s in which any representative member has a line scheme that is isomorphic to the union of two spatial elliptic curves and four planar elliptic curves.

At present, no regular algebra with such a line scheme is known. However, given the results in [Chandler and Vancliff 2015; Tomlin and Vancliff 2018], it is possible that some GSCA may satisfy this property and thus be candidates for generic quantum \mathbb{P}^3 s.

In [Nafari et al. 2011] it was shown that most quadratic quantum \mathbb{P}^2 s may be constructed as a twist by an automorphism (in the sense of [Artin et al. 1991]) of either a graded skew Clifford algebra or of an Ore extension (see [Goodearl and Warfield 2004]) of a regular graded skew Clifford algebra of global dimension 2. The only regular algebras of global dimension 3 that were not classified in this way are some that have a point scheme of an elliptic curve (classified as type E and an open subset of type A in [Artin and Schelter 1987]). It is currently unclear as to whether an analogous result holds for quantum \mathbb{P}^3 s.

These results suggest that GSCAs are likely to play some role in the classification of quantum \mathbb{P}^3 s (and hence the isomorphism classes of such algebras will be of interest).

The article is outlined as follows. In Section 1, we will introduce preliminary notions that will be relevant to the discussions in the article. In Section 2, we will give our main results. In particular, we will discuss two types of isomorphisms: one in which the degree-1 generators of the algebras are identified and the degree-2 generators are related via matrix multiplication and another in which the degree-2 generators of the algebras are identified and the degree-1 generators are related via a diagonal mapping.

1. Preliminaries

Throughout this article, we adopt the following notation:

- \mathbb{k} denotes an algebraically closed field with $\text{char}(\mathbb{k}) \neq 2$.
- \mathbb{k}^\times denotes the nonzero elements of \mathbb{k} .
- $\mathbb{M}_n(\mathbb{k})$ denotes the ring of $n \times n$ matrices with entries in \mathbb{k} .
- M_{ij} denotes the ij -th entry of the matrix M and M^T denotes the transpose of M .
- If S is a positively graded, connected \mathbb{k} -algebra, then S_n denotes the homogeneous degree- n elements of S .
- If V is a \mathbb{k} -vector space, then V^n denotes the vector space $\underbrace{V \times V \times \cdots \times V}_{n \text{ times}}$.

Definition 1.1 [Cassidy and Vancliff 2010]. Let $\mu_{ij} \in \mathbb{k}^\times$, where $1 \leq i, j \leq n$, such that $\mu_{ij}\mu_{ji} = 1$ for all $i \neq j$ and $\mu_{ii} = 1$ for all i . We write $\mu = (\mu_{ij}) \in \mathbb{M}_n(\mathbb{k})$. A matrix $M \in \mathbb{M}_n(\mathbb{k})$ is called μ -symmetric if $M_{ij} = \mu_{ij}M_{ji}$ for all i, j .

The set of all $n \times n$ μ -symmetric matrices forms a \mathbb{k} -vector space, denoted by $\text{Sym}_n^\mu(\mathbb{k})$. The standard basis of $\text{Sym}_n^\mu(\mathbb{k})$ is

$$\mathfrak{B} = \{E_{ii} : 1 \leq i \leq n\} \cup \{E_{ij} + \mu_{ji}E_{ji} : 1 \leq i < j \leq n\},$$

where $E_{ij} \in \mathbb{M}_n(\mathbb{k})$ denotes the matrix with a 1 in the ij -th position and zeroes elsewhere.

In [Cassidy and Vancliff 2010], it was shown that the notion of identifying a quadratic form and a symmetric matrix could be generalized to μ -symmetric matrices as follows. Let $\mu \in \mathbb{M}_n(\mathbb{k})$ be as in Definition 1.1 and $M \in \text{Sym}_n^\mu(\mathbb{k})$. Let S denote the \mathbb{k} -algebra on z_1, z_2, \dots, z_n with defining relations $z_j z_i = \mu_{ij} z_i z_j$ for all $i, j = 1, \dots, n$ so that S is a skew polynomial ring. Then the image of $q = z^T M z$, where $z = [z_1 z_2 \cdots z_n]^T$, in S is a quadratic form. Furthermore, if $q = \sum_{i \leq j} a_{ij} z_i z_j \in S_2$, one may associate to q an $M \in \text{Sym}_n^\mu(\mathbb{k})$ defined by $M_{ii} = a_{ii}$ for all i and, for $i < j$, $M_{ij} = a_{ij}/2$.

Definition 1.2 [Cassidy and Vancliff 2010]. Let $\mu \in \mathbb{M}_n(\mathbb{k})$ be as in Definition 1.1 and $M_1, \dots, M_n \in \text{Sym}_n^\mu(\mathbb{k})$. A graded skew Clifford algebra $A = A(\mu, M_1, \dots, M_n)$ associated to μ and M_1, \dots, M_n is a graded \mathbb{k} -algebra on degree-1 generators x_1, \dots, x_n and on degree-2 generators y_1, \dots, y_n with defining relations given by

- $x_i x_j + \mu_{ij} x_j x_i = \sum_{k=1}^n (M_k)_{ij} y_k$ for all $i, j = 1, \dots, n$; and,
- any additional (degree-3 or degree-4) relations necessary in order to guarantee the existence of a normalizing sequence that spans $\bigoplus_{k=1}^n \mathbb{k} y_k$.

Note that if a graded skew Clifford algebra is regular, then it will only have relations of the form $x_i x_j + \mu_{ij} x_j x_i = \sum_{k=1}^n (M_k)_{ij} y_k$ for all $i, j = 1, \dots, n$.

2. Main results

We first examine the case where the degree-1 generators of each algebra are identified and the degree-2 generators are related via a mapping by matrix multiplication. Our algebras will be \mathbb{k} -algebras whose defining relations are of the form $x_i x_j + \mu_{ij} x_j x_i = \sum_{k=1}^n (M_k)_{ij} y_k$. Each such algebra will map onto a graded skew Clifford algebra since the defining relations of a graded skew Clifford algebra will include these relations and (possibly) additional relations necessary to guarantee the existence of a normalizing sequence, as required by Definition 1.2.

Theorem 2.1. *Let $\mu \in \mathbb{M}_n(\mathbb{k})$ be as in Definition 1.1 and let*

$$M_1, M_2, \dots, M_n, N_1, N_2, \dots, N_n \in \text{Sym}_n^\mu(\mathbb{k}).$$

Let A be a \mathbb{k} -algebra on degree-1 generators x_1, x_2, \dots, x_n and degree-2 generators y_1, y_2, \dots, y_n with defining relations

$$x_i x_j + \mu_{ij} x_j x_i = \sum_{k=1}^n (M_k)_{ij} y_k \quad \text{for all } i, j.$$

Similarly, let B be a \mathbb{k} -algebra on degree-1 generators X_1, X_2, \dots, X_n and degree-2 generators Y_1, Y_2, \dots, Y_n with defining relations

$$X_i X_j + \mu_{ij} X_j X_i = \sum_{k=1}^n (N_k)_{ij} Y_k \quad \text{for all } i, j.$$

Given a nonsingular matrix $P \in M_n(\mathbb{k})$, A is isomorphic to B under a map $\varphi: A \rightarrow B$ defined by

$$\begin{bmatrix} x_1 \\ x_2 \\ \vdots \\ x_n \end{bmatrix} \xrightarrow{\varphi} \begin{bmatrix} X_1 \\ X_2 \\ \vdots \\ X_n \end{bmatrix} \quad \text{and} \quad \begin{bmatrix} y_1 \\ y_2 \\ \vdots \\ y_n \end{bmatrix} \xrightarrow{\varphi} P \begin{bmatrix} Y_1 \\ Y_2 \\ \vdots \\ Y_n \end{bmatrix}$$

if and only if $N = P^T M$, where $N = [N_1 \ N_2 \ \dots \ N_n]^T$, $M = [M_1 \ M_2 \ \dots \ M_n]^T \in (\text{Sym}_n^\mu(\mathbb{k}))^n$.

Proof. Let \mathfrak{B} denote the standard \mathbb{k} -basis of $\text{Sym}_n^\mu(\mathbb{k})$. We may write the defining relations of A as the matrix equation

$$\begin{bmatrix} 2x_1^2 \\ x_1 x_2 + \mu_{12} x_2 x_1 \\ \vdots \\ 2x_n^2 \end{bmatrix} = \begin{bmatrix} (M_1)_{11} & (M_2)_{11} & \dots & (M_n)_{11} \\ (M_1)_{12} & (M_2)_{12} & \dots & (M_n)_{12} \\ \vdots & \vdots & \ddots & \vdots \\ (M_1)_{nn} & (M_2)_{nn} & \dots & (M_n)_{nn} \end{bmatrix} \begin{bmatrix} y_1 \\ y_2 \\ \vdots \\ y_n \end{bmatrix}.$$

Applying the map φ to the above equation we obtain the defining relations of B :

$$\begin{bmatrix} 2X_1^2 \\ X_1 X_2 + \mu_{12} X_2 X_1 \\ \vdots \\ 2X_n^2 \end{bmatrix} = \begin{bmatrix} (M_1)_{11} & (M_2)_{11} & \dots & (M_n)_{11} \\ (M_1)_{12} & (M_2)_{12} & \dots & (M_n)_{12} \\ \vdots & \vdots & \ddots & \vdots \\ (M_1)_{nn} & (M_2)_{nn} & \dots & (M_n)_{nn} \end{bmatrix} P \begin{bmatrix} Y_1 \\ Y_2 \\ \vdots \\ Y_n \end{bmatrix}.$$

Hence

$$\begin{bmatrix} (M_1)_{11} & (M_2)_{11} & \dots & (M_n)_{11} \\ (M_1)_{12} & (M_2)_{12} & \dots & (M_n)_{12} \\ \vdots & \vdots & \ddots & \vdots \\ (M_1)_{nn} & (M_2)_{nn} & \dots & (M_n)_{nn} \end{bmatrix} P = \begin{bmatrix} (N_1)_{11} & (N_2)_{11} & \dots & (N_n)_{11} \\ (N_1)_{12} & (N_2)_{12} & \dots & (N_n)_{12} \\ \vdots & \vdots & \ddots & \vdots \\ (N_1)_{nn} & (N_2)_{nn} & \dots & (N_n)_{nn} \end{bmatrix}.$$

The k -th columns of the matrices in the previous equation are the \mathfrak{B} -coordinate vector of M_k and N_k , respectively. Thus,

$$\begin{aligned}
 [N_1 \ \cdots \ N_n] &= [M_1 \ \cdots \ M_n] P \implies \begin{bmatrix} N_1 \\ \vdots \\ N_n \end{bmatrix} = P^T \begin{bmatrix} M_1 \\ \vdots \\ M_n \end{bmatrix} \\
 &\implies N = P^T M. \quad \square
 \end{aligned}$$

Example 2.2. Let

$$\mu = \begin{bmatrix} 1 & i \\ -i & 1 \end{bmatrix}, \quad M_1 = \begin{bmatrix} 1 & 0 \\ 0 & 1 \end{bmatrix}, \quad M_2 = \begin{bmatrix} 0 & i \\ 1 & 0 \end{bmatrix}, \quad N_1 = \begin{bmatrix} 2 & -i \\ -1 & 2 \end{bmatrix}, \quad N_2 = \begin{bmatrix} i & i \\ 1 & i \end{bmatrix}.$$

Then $A = A(\mu, M_1, M_2)$ is given by

$$A = \frac{\mathbb{k}\langle x_1, x_2, y_1, y_2 \rangle}{\langle 2x_1^2 - y_1, 2x_2^2 - y_1, x_1x_2 + ix_2x_1 - iy_2 \rangle}$$

and $B = B(\mu, N_1, N_2)$ is given by

$$B = \frac{\mathbb{k}\langle X_1, X_2, Y_1, Y_2 \rangle}{\langle 2X_1^2 - 2Y_1 - iY_2, 2X_2^2 - 2Y_1 - iY_2, X_1X_2 + iX_2X_1 + iY_1 - iY_2 \rangle}.$$

One can compute that

$$\begin{cases} N_1 = 2M_1 - M_2, \\ N_2 = iM_1 + M_2 \end{cases} \implies \begin{bmatrix} N_1 \\ N_2 \end{bmatrix} = \begin{bmatrix} 2 & -1 \\ i & 1 \end{bmatrix} \begin{bmatrix} M_1 \\ M_2 \end{bmatrix}.$$

Hence,

$$P = \begin{bmatrix} 2 & i \\ -1 & 1 \end{bmatrix}.$$

The above theorem implies that $A \cong B$ under the map $x_i \mapsto X_i$ for all i , $y_1 \mapsto 2Y_1 + iY_2$ and $y_2 \mapsto -Y_1 + Y_2$, which is easily verified.

Corollary 2.3. Let μ, A, B, M, N, φ and P be as in [Theorem 2.1](#). If $q_k, p_k \in S$ denote the quadratic forms associated to M_k and N_k , respectively, then A is isomorphic to B under φ if and only if

$$\begin{bmatrix} p_1 \\ p_2 \\ \vdots \\ p_n \end{bmatrix} = P^T \begin{bmatrix} q_1 \\ q_2 \\ \vdots \\ q_n \end{bmatrix}.$$

Proof. The result follows immediately from multiplying each entry of the vectors N and M in the equation $N = P^T M$ by z^T and z on the left and right, respectively. \square

Example 2.4. Continuing [Example 2.2](#), the quadratic forms associated to M_1, M_2, N_1 and N_2 are $q_1 = z_1^2 + z_2^2, q_2 = 2iz_1z_2, p_1 = 2z_1^2 - 2iz_1z_2 + 2z_2^2$ and $p_2 = iz_1^2 + 2iz_1z_2 + iz_2^2$, respectively. It is easily verified that

$$\begin{cases} p_1 = 2q_1 - q_2, \\ p_2 = iq_1 + q_2 \end{cases} \implies \begin{bmatrix} p_1 \\ p_2 \end{bmatrix} = \begin{bmatrix} 2 & -i \\ i & 1 \end{bmatrix} \begin{bmatrix} q_1 \\ q_2 \end{bmatrix}.$$

We now examine the case where the degree-2 generators of each algebra are identified and the degree-1 generators of the algebra are related via a diagonal mapping.

Theorem 2.5. Let $\mu \in \mathbb{M}_n(\mathbb{k})$ be as in [Definition 1.1](#) and let

$$M_1, M_2, \dots, M_n, N_1, N_2, \dots, N_n \in \text{Sym}_n^\mu(\mathbb{k}).$$

Let A be a \mathbb{k} -algebra on degree-1 generators x_1, x_2, \dots, x_n and degree-2 generators y_1, y_2, \dots, y_n with defining relations

$$x_i x_j + \mu_{ij} x_j x_i = \sum_{k=1}^n (M_k)_{ij} y_k \quad \text{for all } i, j.$$

Similarly, let B be a \mathbb{k} -algebra on degree-1 generators X_1, X_2, \dots, X_n and degree-2 generators Y_1, Y_2, \dots, Y_n with defining relations

$$X_i X_j + \mu_{ij} X_j X_i = \sum_{k=1}^n (N_k)_{ij} Y_k \quad \text{for all } i, j.$$

Then $\varphi : A \rightarrow B$ is an isomorphism defined $x_i \mapsto \alpha_i X_i$ and $y_i \mapsto Y_i$ for $\alpha_i \in \mathbb{k}^\times$ and for all i if and only if $(N_k)_{ij} = (M_k)_{ij} / (\alpha_i \alpha_j)$ for all i, j, k .

Proof. Applying the map φ to the defining relations of A yields the defining relations of B :

$$\alpha_i \alpha_j X_i X_j + \mu_{ij} \alpha_i \alpha_j X_j X_i = \sum_{k=1}^n (M_k)_{ij} Y_k \implies X_i X_j + \mu_{ij} X_j X_i = \sum_{k=1}^n \frac{(M_k)_{ij}}{\alpha_i \alpha_j} Y_k.$$

Hence, $(N_k)_{ij} = (M_k)_{ij} / (\alpha_i \alpha_j)$. □

Example 2.6. Let

$$\mu = \begin{bmatrix} 1 & i \\ -i & 1 \end{bmatrix}, \quad M_1 = \begin{bmatrix} 1 & 0 \\ 0 & 1 \end{bmatrix}, \quad M_2 = \begin{bmatrix} 0 & i \\ 1 & 0 \end{bmatrix}, \quad N_1 = \begin{bmatrix} 1 & 0 \\ 0 & -1 \end{bmatrix}, \quad N_2 = \begin{bmatrix} 0 & -1 \\ i & 0 \end{bmatrix}.$$

Then $A = A(\mu, M_1, M_2)$ is given by

$$A = \frac{\mathbb{k}\langle x_1, x_2, y_1, y_2 \rangle}{\langle 2x_1^2 - y_1, 2x_2^2 - y_1, x_1x_2 + ix_2x_1 - iy_2 \rangle}$$

and $B = B(\mu, N_1, N_2)$ is given by

$$B = \frac{\mathbb{k}\langle X_1, X_2, Y_1, Y_2 \rangle}{\langle 2X_1^2 - Y_1, 2X_2^2 + Y_1, X_1X_2 + iX_2X_1 + Y_2 \rangle}.$$

In checking to see if $(N_k)_{ij} = (M_k)_{ij}/(\alpha_i\alpha_j)$, we arrive at the equations

$$\begin{cases} 1 = 1/\alpha_1^2, \\ -1 = 1/\alpha_2^2, \\ -1 = i/(\alpha_1\alpha_2), \\ i = 1/(\alpha_1\alpha_2) \end{cases} \implies \begin{cases} \alpha_1 = \pm 1, \\ \alpha_2 = \mp i. \end{cases}$$

The above theorem implies that $A \cong B$ under both the maps φ_+ and φ_- defined by $\varphi_{\pm} : x_1 \mapsto \pm X_1, x_2 \mapsto \mp i X_2$ and $y_k \mapsto Y_k$, for all k .

Corollary 2.7. *Let $\mu, A, B, M_1, M_2, \dots, M_n, N_1, N_2, \dots, N_n$, and φ be as in Theorem 2.5. If $q_k = \sum_{i \leq j} a_{kij} z_i z_j \in S$ and $p_k = \sum_{i \leq j} b_{kij} z_i z_j \in S$ denote the quadratic forms associated to M_k and N_k , respectively, then A is isomorphic to B under φ if and only if $b_{kij} = a_{kij}/(\alpha_i\alpha_j)$.*

Proof. The result follows immediately from $(N_k)_{ij} = (M_k)_{ij}/(\alpha_i\alpha_j)$ and the definition of the quadratic form in [Cassidy and Vancliff 2010]. □

Example 2.8. Continuing 2.6, the quadratic forms associated to M_1, M_2, N_1 and N_2 are $q_1 = z_1^2 + z_2^2, q_2 = 2iz_1z_2, p_1 = z_1^2 - z_2^2$ and $p_2 = -2z_1z_2$, respectively. It is easily verified that $b_{kij} = a_{kij}/(\alpha_i\alpha_j)$ holds for these forms.

The reader should note that the more general case in which the degree-1 generators of an algebra are related via a mapping by matrix multiplication is much more complicated. Such maps can transform graded skew Clifford algebras into isomorphic algebras whose defining relations cannot be directly written in the form $x_i x_j + \mu_{ij} x_j x_i = \sum_{k=1}^n (M_k)_{ij} y_k$ without changing the presentation of the algebra, as illustrated in the following example.

Example 2.9. Let

$$\mu = \begin{bmatrix} 1 & i \\ -i & 1 \end{bmatrix}, \quad M_1 = \begin{bmatrix} 1 & 0 \\ 0 & 1 \end{bmatrix}, \quad M_2 = \begin{bmatrix} 0 & i \\ 1 & 0 \end{bmatrix}.$$

Then $A = A(\mu, M_1, M_2)$ is given by

$$A = \frac{\mathbb{k}\langle x_1, x_2, y_1, y_2 \rangle}{\langle 2x_1^2 - y_1, 2x_2^2 - y_1, x_1x_2 + ix_2x_1 - iy_2 \rangle}.$$

Consider the isomorphism $\varphi : A \rightarrow \varphi(A)$ defined by $x_1 \mapsto X_1 - X_2$, $x_2 \mapsto X_2$, and $y_k \mapsto Y_k$ for all k . Applying φ to the defining relations of A yields the relations

$$\begin{aligned} 2X_1^2 + 2X_2^2 - 2X_1X_2 - 2X_2X_1 - Y_1 &= 0, \\ 2X_2^2 - Y_1 &= 0, \\ X_1X_2 + iX_2X_1 - (1+i)X_2^2 - iY_2 &= 0. \end{aligned}$$

These relations cannot all be written in the form $X_iX_j + \mu_{ij}X_jX_i = \sum_{k=1}^n (N_k)_{ij}Y_k$ without a change of presentation; hence, the image of the algebra does not present as a graded skew Clifford algebra.

References

- [Artin and Schelter 1987] M. Artin and W. F. Schelter, “Graded algebras of global dimension 3”, *Adv. in Math.* **66**:2 (1987), 171–216. [MR](#) [Zbl](#)
- [Artin et al. 1990] M. Artin, J. Tate, and M. Van den Bergh, “Some algebras associated to automorphisms of elliptic curves”, pp. 33–85 in *The Grothendieck Festschrift, Vol. I*, edited by P. Cartier et al., Progr. Math. **86**, Birkhäuser, Boston, MA, 1990. [MR](#) [Zbl](#)
- [Artin et al. 1991] M. Artin, J. Tate, and M. Van den Bergh, “Modules over regular algebras of dimension 3”, *Invent. Math.* **106**:2 (1991), 335–388. [MR](#) [Zbl](#)
- [Cassidy and Vancliff 2010] T. Cassidy and M. Vancliff, “Generalizations of graded Clifford algebras and of complete intersections”, *J. Lond. Math. Soc.* (2) **81**:1 (2010), 91–112. Correction in **90**:2 (2014), 631–636. [MR](#) [Zbl](#)
- [Chandler and Vancliff 2015] R. G. Chandler and M. Vancliff, “The one-dimensional line scheme of a certain family of quantum \mathbb{P}^3 s”, *J. Algebra* **439** (2015), 316–333. [MR](#) [Zbl](#)
- [Goodearl and Warfield 2004] K. R. Goodearl and R. B. Warfield, Jr., *An introduction to noncommutative Noetherian rings*, 2nd ed., London Mathematical Society Student Texts **61**, Cambridge University Press, 2004. [MR](#) [Zbl](#)
- [Le Bruyn 1995] L. Le Bruyn, “Central singularities of quantum spaces”, *J. Algebra* **177**:1 (1995), 142–153. [MR](#) [Zbl](#)
- [Nafari et al. 2011] M. Nafari, M. Vancliff, and J. Zhang, “Classifying quadratic quantum \mathbb{P}^2 s by using graded skew Clifford algebras”, *J. Algebra* **346** (2011), 152–164. [MR](#) [Zbl](#)
- [Shelton and Vancliff 2002a] B. Shelton and M. Vancliff, “Schemes of line modules, I”, *J. London Math. Soc.* (2) **65**:3 (2002), 575–590. [MR](#) [Zbl](#)
- [Shelton and Vancliff 2002b] B. Shelton and M. Vancliff, “Schemes of line modules, II”, *Comm. Algebra* **30**:5 (2002), 2535–2552. [MR](#) [Zbl](#)
- [Tomlin and Vancliff 2018] D. Tomlin and M. Vancliff, “The one-dimensional line scheme of a family of quadratic quantum \mathbb{P}^3 s”, *J. Algebra* **502** (2018), 588–609. [MR](#) [Zbl](#)

Received: 2020-06-22

Revised: 2020-08-07

Accepted: 2020-08-10

richard.chandler@untDallas.edu

*Department of Mathematics and Information Sciences,
University of North Texas at Dallas, Dallas, TX, United States*

nicholas.engel@mavs.uta.edu

*Department of Mathematics, University of Texas at Arlington,
Arlington, TX, United States*

INVOLVE YOUR STUDENTS IN RESEARCH

Involve showcases and encourages high-quality mathematical research involving students from all academic levels. The editorial board consists of mathematical scientists committed to nurturing student participation in research. Bridging the gap between the extremes of purely undergraduate research journals and mainstream research journals, *Involve* provides a venue to mathematicians wishing to encourage the creative involvement of students.

MANAGING EDITOR

Kenneth S. Berenhaut Wake Forest University, USA

BOARD OF EDITORS

Colin Adams	Williams College, USA	Robert B. Lund	Clemson University, USA
Arthur T. Benjamin	Harvey Mudd College, USA	Gaven J. Martin	Massey University, New Zealand
Martin Bohner	Missouri U of Science and Technology, USA	Mary Meyer	Colorado State University, USA
Amarjit S. Budhiraja	U of N Carolina, Chapel Hill, USA	Frank Morgan	Williams College, USA
Pietro Cerone	La Trobe University, Australia	Mohammad Sal Moslehian	Ferdowsi University of Mashhad, Iran
Scott Chapman	Sam Houston State University, USA	Zuhair Nashed	University of Central Florida, USA
Joshua N. Cooper	University of South Carolina, USA	Ken Ono	Univ. of Virginia, Charlottesville
Jem N. Corcoran	University of Colorado, USA	Yuval Peres	Microsoft Research, USA
Toka Diagana	University of Alabama in Huntsville, USA	Y.-F. S. Pétermann	Université de Genève, Switzerland
Michael Dorff	Brigham Young University, USA	Jonathon Peterson	Purdue University, USA
Sever S. Dragomir	Victoria University, Australia	Robert J. Plemmons	Wake Forest University, USA
Joel Foisy	SUNY Potsdam, USA	Carl B. Pomerance	Dartmouth College, USA
Erin W. Fulp	Wake Forest University, USA	Vadim Ponomarenko	San Diego State University, USA
Joseph Gallian	University of Minnesota Duluth, USA	Bjorn Poonen	UC Berkeley, USA
Stephan R. Garcia	Pomona College, USA	József H. Przytycki	George Washington University, USA
Anant Godbole	East Tennessee State University, USA	Richard Rebarber	University of Nebraska, USA
Ron Gould	Emory University, USA	Robert W. Robinson	University of Georgia, USA
Sat Gupta	U of North Carolina, Greensboro, USA	Javier Rojo	Oregon State University, USA
Jim Haglund	University of Pennsylvania, USA	Filip Saidak	U of North Carolina, Greensboro, USA
Johnny Henderson	Baylor University, USA	Hari Mohan Srivastava	University of Victoria, Canada
Glenn H. Hurlbert	Virginia Commonwealth University, USA	Andrew J. Sterge	Honorary Editor
Charles R. Johnson	College of William and Mary, USA	Ann Trenk	Wellesley College, USA
K. B. Kulasekera	Clemson University, USA	Ravi Vakil	Stanford University, USA
Gerry Ladas	University of Rhode Island, USA	Antonia Vecchio	Consiglio Nazionale delle Ricerche, Italy
David Larson	Texas A&M University, USA	John C. Wierman	Johns Hopkins University, USA
Suzanne Lenhart	University of Tennessee, USA	Michael E. Zieve	University of Michigan, USA
Chi-Kwong Li	College of William and Mary, USA		

PRODUCTION

Silvio Levy, Scientific Editor

Cover: Alex Scorpan

See inside back cover or msp.org/involve for submission instructions. The subscription price for 2020 is US \$205/year for the electronic version, and \$275/year (+\$35, if shipping outside the US) for print and electronic. Subscriptions, requests for back issues and changes of subscriber address should be sent to MSP.

Involve (ISSN 1944-4184 electronic, 1944-4176 printed) at Mathematical Sciences Publishers, 798 Evans Hall #3840, c/o University of California, Berkeley, CA 94720-3840, is published continuously online. Periodical rate postage paid at Berkeley, CA 94704, and additional mailing offices.

Involve peer review and production are managed by EditFlow[®] from Mathematical Sciences Publishers.

PUBLISHED BY

 **mathematical sciences publishers**

nonprofit scientific publishing

<http://msp.org/>

© 2020 Mathematical Sciences Publishers

involve

2020

vol. 13

no. 5

Set-valued domino tableaux and shifted set-valued domino tableaux	721
FLORENCE MAAS-GARIÉPY AND REBECCA PATRIAS	
The first digit of the discriminant of Eisenstein polynomials as an invariant of totally ramified extensions of p-adic fields	747
CHAD AWTREY, ALEXANDER GAURA, SEBASTIAN PAULI, SANDI RUDZINSKI, ARIEL UY AND SCOTT ZINZER	
Counting pseudo progressions	759
JAY CUMMINGS, QUIN DARCY, NATALIE HOBSON, DREW HORTON, KEITH RHODEWALT, MORGAN THROCKMORTON AND RY ULMER-STRACK	
Growth series for graphs	781
WALTER LIU AND RICHARD SCOTT	
Peg solitaire in three colors on graphs	791
TARA C. DAVIS, ALEXSIS DE LAMERE, GUSTAVO SOPENA, ROBERTO C. SOTO, SONALI VYAS AND MELISSA WONG	
Disagreement networks	803
FLORIN CATRINA AND BRIAN ZILLI	
Rings whose subrings have an identity	823
GREG OMAN AND JOHN STROUD	
Simple graphs of order 12 and minimum degree 6 contain K_6 minors	829
RYAN ODENEAL AND ANDREI PAVELESCU	
Mixed volume of small reaction networks	845
NIDA OBATAKE, ANNE SHIU AND DILRUBA SOFIA	
Counting profile strings from rectangular tilings	861
ANTHONY PETROSINO, ALISSA SCHEMBOR AND KATHRYN HAYMAKER	
Isomorphisms of graded skew Clifford algebras	871
RICHARD G. CHANDLER AND NICHOLAS ENGEL	
Eta-quotients of prime or semiprime level and elliptic curves	879
MICHAEL ALLEN, NICHOLAS ANDERSON, ASIMINA HAMAKIOTES, BEN OLTSIK AND HOLLY SWISHER	