

```

gap> g:= SymmetricGroup( 4 );
Sym( [ 1 .. 4 ] )
gap> tbl:= CharacterTable( g ); HasIrr( tbl );
i5 : betti(t,Weights=>{1,0})
false
      0 1 2 3 4
o5 = total: 1 4 13 14 4
      0: 1 . . .
      1: . 2 2 4 2
      2: . 2 5 6 .
      3: . . 4 . 2
      4: . . . 4 .
      5: . . 2 . .
gap> tblmod2:= CharacterTable( tbl, 2 );
      BrauerTable( Sym( [ 1 .. 4 ] ), 2 )
gap> tblmod2 = CharacterTable( tbl, 2 );
true
gap> tblmod2 = BrauerTable( tbl, 2 );
true
o5 : BrauerTable
i6 : betti(t,Weights=>{0,1})
      0 1 2 3 4
o6 = total: 1 4 13 14 4
      0: 1 . . .
      1: . 2 . .
      2: . 2 . .
      3: . . 4 . 2
      4: . . . 4 .
      5: . . 2 . .
gap> libtbl:= CharacterTable( "M" );
      CharacterTable( "M" )
gap> CharacterTableRegular( libtbl, 2 );
      BrauerTable( "M", 2 )
gap> BrauerTable( libtbl, 2 );
fail
gap> CharacterTable( "Symmetric", 4 );
      CharacterTable( "Sym(4)" )
i7 : t1 = betti(t,Weights=>{1,1})
gap> ComputedBrauerTables( tbl );
      [ , BrauerTable( Sym( [ 1 .. 4 ] ), 2 ), ]
      ring r1 = 32003,(x,y,z),ds;
      int a,b,c,t=11,5,3,0;
      poly f = x^a*y^b+z^(3*c)+x^(c+2)*y^(c-1)+x^(c-2)*y^c*(y^2+t*x)^2;
      option(noprot);
      timer=1;
      ring r2 = 32003,(x,y,z),dp;
      poly f=imap(r1,f);
      ideal j=jacob(f);
      vdim(std(j));
==> 536
      vdim(std(j+f));
==> 195
      timer=0; // reset timer
o7 : BettiTally
i8 : peek t1
o8 = BettiTally{ (0, {0, 0}, 0) => 1 }
      (1, {2, 2}, 4) => 2
      (1, {3, 3}, 6) => 2
      (2, {3, 7}, 10) => 2
      (2, {4, 4}, 8) => 1
      (2, {4, 5}, 9) => 4
      (2, {5, 4}, 9) => 4
      (2, {7, 3}, 10) => 2
      (3, {4, 7}, 11) => 4
      (3, {7, 4}, 11) => 4
      (4, {5, 7}, 12) => 2
      (4, {7, 5}, 12) => 2

```

Journal of Software for Algebra and Geometry

Computing quasidegrees of A-graded modules

ROBERTO BARRERA

Computing quasidegrees of A -graded modules

ROBERTO BARRERA

ABSTRACT: We describe the main functions of the Macaulay2 package `Quasidegrees.m2`. The purpose of this package is to compute the quasidegree set of a finitely generated $\mathbb{Z}^d$ -graded module presented as the cokernel of a monomial matrix. We provide examples with motivation coming from A -hypergeometric systems.

1. INTRODUCTION. Throughout, $R = \mathbb{k}[x_1, \dots, x_n]$ is a $\mathbb{Z}^d$ -graded polynomial ring over a field $\mathbb{k}$ and $\mathfrak{m} = \langle x_1, \dots, x_n \rangle$ denotes the homogeneous maximal ideal in R . Let $M = \bigoplus_{\beta \in \mathbb{Z}^d} M_\beta$ be a $\mathbb{Z}^d$ -graded R -module. The *true degree set* of M is

$$\text{tdeg}(M) = \{\beta \in \mathbb{Z}^d \mid M_\beta \neq 0\}.$$

The *quasidegree set* of M , denoted $\text{qdeg}(M)$, is the Zariski closure in $\mathbb{C}^d$ of $\text{tdeg}(M)$.

The purpose of the Macaulay2 package `Quasidegrees.m2` (provided as an online supplement) is to compute the quasidegree set of a finitely generated $\mathbb{Z}^d$ -graded module presented as the cokernel of a monomial matrix. By a monomial matrix, we mean a matrix where each entry is either zero or a monomial in R . The initial motivation for `Quasidegrees.m2` was to compute the quasidegree sets of certain local cohomology modules supported at $\mathfrak{m}$ of $\mathbb{Z}^d$ -graded R -modules, so there are some methods in the package specific to local cohomology. Recall that the *i -th local cohomology module* of M with support at the ideal $I \subset R$ is the i -th right derived functor of the left exact I -torsion functor

$$\Gamma_I(M) = \{m \in M \mid I^t m = 0 \text{ for some } t \in \mathbb{N}\}$$

on the category of R -modules.

By the vanishing theorems of local cohomology [Eisenbud 1995], the quasidegree sets of the local cohomology modules supported at $\mathfrak{m}$ of M can be seen as measuring how far the module is from being Cohen–Macaulay. From the A -hypergeometric systems point of view, the quasidegree set of the non-top local cohomology modules supported at $\mathfrak{m}$ of R/I_A , where I_A is the toric ideal associated

MSC2010: 13D45, 13P20, 33C70.

Keywords: quasidegree, hypergeometric system, local cohomology.

`Quasidegrees.m2` version 1.0

to A in R , determine the parameters β where the A -hypergeometric system $H_A(\beta)$ has rank higher than expected (see Section 3).

2. QUASIDEGREES. The main function of `Quasidegrees.m2` is `quasidegrees`, which computes the quasidegree set of a module that is presented by a monomial matrix.

We use the idea of standard pairs of monomial ideals to compute the quasidegree set of a $\mathbb{Z}^d$ -graded R -module. Given a monomial x^u and a subset $Z \subset \{x_1, \dots, x_n\}$, the pair (x^u, Z) indexes the monomials $x^u \cdot x^v$ where $\text{supp}(x^v) \subset Z$. A *standard pair* of a monomial ideal $I \subset R$ is a pair (x^u, Z) satisfying:

- (1) $\text{supp}(x^u) \cap Z = \emptyset$.
- (2) All of the monomials indexed by (x^u, Z) are outside of I .
- (3) (x^u, Z) is maximal in the sense that $(x^u, Z) \not\subseteq (x^v, Y)$ for any other pair (x^v, Y) satisfying the first two conditions.

To compute the quasidegree set of M we first find a monomial presentation of M so that M is the cokernel of a monomial matrix ϕ . We then compute the standard pairs of the ideals generated by the rows of ϕ and to each standard pair we associate the degrees of the corresponding variables. Algorithm 1 below is implemented in `Quasidegrees.m2`. The input is an R -module presented by a monomial matrix

$$\phi : R^s \rightarrow R^t.$$

As in Macaulay2, we write the degree of the k -th factor of R^t next to the k -th row of the matrix ϕ .

In the Macaulay2 implementation of the algorithm, we represent the output as a list of pairs (u, Z) with $u \in \mathbb{Q}^d$ and $Z \subset \mathbb{Q}^d$, where the pair (u, Z) represents the plane

$$u + \sum_{v \in Z} \mathbb{C} \cdot v.$$

Input: R -module M presented by monomial matrix $\phi = \alpha_i [c_{j,k} x^{u_{j,k}}] : R^s \rightarrow R^t$

Output: `qdeg(M)`

$Q = \emptyset$

for $1 \leq k \leq t$ **do**

$SP = \{\text{standard pairs of } \langle c_{k,1} x^{u_{k,1}}, c_{k,2} x^{u_{k,2}}, \dots, c_{k,s} x^{u_{k,s}} \rangle\}$

$Q = Q \cup \{\deg(x^u) + \alpha_k + \sum_{x_i \in F} \mathbb{C} \cdot \deg(x_i) \mid (x^u, Z) \in SP\}$

end for

return Q

Algorithm 1. Compute `qdeg(M)`.

The union of these planes over all such pairs in the output is the quasidegree set of M .

The following is an example of `Quasidegrees.m2` computing the quasidegree set of an R -module:

```
i1 : R=QQ[x,y,z]
o1 = R
o1 : PolynomialRing
i2 : I=ideal(x*y,y*z)
o2 = ideal (x*y, y*z)
o2 : Ideal of R
i3 : M=R^1/I
o3 = cokernel | xy yz |
                                     1
o3 : R-module, quotient of R
i4 : Q = quasidegrees M
o4 = {{0, {| 1 |}}, {0, {| 1 |, | 1 |}}}}
o4 : List
```

The above example displays a caveat of `quasidegrees` in that there may be some redundancies in the output. By a redundancy, we mean when one plane in the output is contained in another. The redundancy above is clear:

$$\text{qdeg}(\mathbb{k}[x, y, z]/\langle xy, yz \rangle) = \mathbb{C} = \{z_1 + z_2 \in \mathbb{C} \mid z_1, z_2 \in \mathbb{C}\}.$$

The function `removeRedundancy` gets rid of redundancies in the list of planes:

```
i5 : removeRedundancy Q
o5 = {{0, {| 1 |, | 1 |}}}}
o5 : List
```

3. QUASIDEGREES AND HYPERGEOMETRIC SYSTEMS. In this section, we discuss the motivation for `Quasidegrees.m2` and the methods therein which aid us in our studies. Let $A = [a_1 \ a_2 \ \cdots \ a_n]$ be an integer $(d \times n)$ -matrix with $\mathbb{Z}A = \mathbb{Z}^d$ and such that the cone over its columns is pointed. There is a natural $\mathbb{Z}^d$ -grading of R by the columns of A given by $\deg(x_j) = a_j$, the j -th column of A . A module that is homogeneous with respect to this grading is said to be A -graded. By the assumptions on A , R is positively graded by A , that is, the only polynomials of degree 0 are the constants. Given such a matrix A and a polynomial ring R in n variables, the method `toGradedRing` gives R an A -grading. For example, let

$$A = \begin{pmatrix} 1 & 1 & 1 & 1 & 1 \\ 0 & 0 & 1 & 1 & 0 \\ 0 & 1 & 1 & 0 & -2 \end{pmatrix}.$$

We make the A -graded polynomial ring $\mathbb{Q}[x_1, x_2, x_3, x_4, x_5]$:

```

i6 : A=matrix{{1,1,1,1,1},{0,0,1,1,0},{0,1,1,0,-2}}
o6 = | 1 1 1 1 1 |
      | 0 0 1 1 0 |
      | 0 1 1 0 -2 |
      3         5
o6 : Matrix ZZ  <--- ZZ
i7 : R=QQ[x_1..x_5]
o7 = R
o7 : PolynomialRing
i8 : R=toGradedRing(A,R)
o8 = R
o8 : PolynomialRing
i9 : describe R
o9 = QQ[x , x , x , x , x , Degrees => {{1}, {1}, {1}, {1}, {1 }},
      1   2   3   4   5           {0} {0} {1} {1} {0 }
                                {0} {1} {1} {0} {-2}
      Heft=>{1, 2:0},MonomialOrder=>{MonomialSize=>32},DegreeRank=>3]
                                {GRevLex=>{5:1}}
                                {Position=>Up}

```

The *toric ideal associated to A in R* is the binomial ideal

$$I_A = \langle \mathbf{x}^u - \mathbf{x}^v : A\mathbf{u} = A\mathbf{v} \rangle.$$

The method `toricIdeal` computes the toric ideal associated to A in the ring R . We continue with the A and R from the above example and compute the toric ideal I_A associated to A in R :

```

i10 : I=toricIdeal(A,R)
o10 = ideal (x x  - x x , x x  - x x , x x  - x x x , x  - x x )
           1 3    2 4    1 4    3 5    1 4    2 3 5    1    2 5
o10 : Ideal of R

```

We now introduce A -hypergeometric systems. Given a matrix $A \in \mathbb{Z}^{d \times n}$ as above and a $\beta \in \mathbb{C}^d$, the A -hypergeometric system with parameter $\beta \in \mathbb{C}^d$ [Saito et al. 2000], denoted $H_A(\beta)$, is the system of partial differential equations:

$$\frac{\partial^{|v|}}{\partial \mathbf{x}^v} \phi(\mathbf{x}) = \frac{\partial^{|u|}}{\partial \mathbf{x}^u} \phi(\mathbf{x}) \quad \text{for all } \mathbf{u}, \mathbf{v}, A\mathbf{u} = A\mathbf{v},$$

$$\sum_{j=1}^n a_{ij} x_j \frac{\partial}{\partial x_j} \phi(\mathbf{x}) = \beta_i \phi(\mathbf{x}), \quad \text{for } i = 1, \dots, d.$$

Such systems are sometimes called *GKZ-hypergeometric systems*. The function `gkz` in the Macaulay2 package `Dmodules` computes this system as an ideal in the Weyl algebra. The *rank* of $H_A(\beta)$ is

$$\text{rank}(H_A(\beta)) = \dim_{\mathbb{C}} \left\{ \begin{array}{l} \text{germs of holomorphic solutions of } H_A(\beta) \\ \text{near a generic nonsingular point} \end{array} \right\}.$$

The function `holonomicRank` in `Dmodules` computes the rank of an A -hypergeometric system. In general, rank is not a constant function of β . Denote $\text{vol}(A)$ to be $d!$ times the Euclidean volume of $\text{conv}(A \cup \{0\})$, the convex hull of the columns of A and the origin in $\mathbb{R}^d$. The following theorem gives the parameters β for which $\text{rank}(H_A(\beta))$ is higher than expected:

Theorem 3.1 [Matusevich et al. 2005]. *Let $H_A(\beta)$ be an A -hypergeometric system with parameter β . If $\beta \in \text{qdeg}(\bigoplus_{i=0}^{d-1} H_{\mathfrak{m}}^i(R/I_A))$ then $\text{rank}(H_A(\beta)) > \text{vol}(A)$. Otherwise, $\text{rank}(H_A(\beta)) = \text{vol}(A)$.*

Since Theorem 3.1 was the initial motivation for `Quasidegrees.m2`, the package has a method `quasidegreesLocalCohomology` (abbreviated `qlc`) to compute the quasidegree set of the local cohomology modules $H_{\mathfrak{m}}^i(R/I_A)$. If the input is an integer i and the R -module R/I_A , then the method computes $\text{qdeg}(H_{\mathfrak{m}}^i(R/I_A))$. If the input is only the module R/I_A , the method computes the quasidegree set in Theorem 3.1.

We use graded local duality to compute the local cohomology modules of a finitely generated A -graded R -module supported at the maximal ideal $\mathfrak{m}$:

Theorem 3.2 (graded local duality [Bruns and Herzog 1993; Miller 2002]). *Given an A -graded R -module M , there is an A -graded vector space isomorphism*

$$\text{Ext}_R^{n-i}(M, R)_{\alpha} \cong \text{Hom}_{\mathbb{k}}(H_{\mathfrak{m}}^i(M)_{-\alpha-\varepsilon_A}, \mathbb{k}),$$

where $\mathfrak{m} = \langle x_1, \dots, x_n \rangle$ and $\varepsilon_A = \sum_{j=1}^n a_j$.

The algorithm implemented for `quasidegreesLocalCohomology` is essentially Algorithm 1 applied to the `Ext`-modules of M with the additional twist of ε_A coming from local duality. For our purposes, we exploit the fact that the higher syzygies of R/I_A are generated by monomials in R^m (see [Miller and Sturmfels 2005], Chapter 9).

Continuing our running example, we use `quasidegreesLocalCohomology` to compute the quasidegree set of $\bigoplus_{i=0}^{d-1} H_{\mathfrak{m}}^i(R/I_A)$:

```
i11 : M=R^1/I
o11 = cokernel | x_1x_3-x_2x_4 x_1x_4^2-x_3^2x_5
x_1^2x_4-x_2x_3x_5 x_1^3-x_2^2x_5 |
1
o11 : R-module, quotient of R
```

```

i12 : quasidegreesLocalCohomology M
o12 = {{| 0 |, {| 1 |}}}
      | 0 |   | 0 |
      | 1 |   | -2 |
o12 : List

```

Thus

$$\text{qdeg}\left(\bigoplus_{i=0}^{d-1} H_m^i(R/I_A)\right) = \begin{bmatrix} 0 \\ 0 \\ 1 \end{bmatrix} + \mathbb{C} \cdot \begin{bmatrix} 1 \\ 0 \\ -2 \end{bmatrix}. \quad (1)$$

As a check, we use the methods `gkz` and `holonomicRank` from the package `Dmodules` to compute $\text{rank}(H_A(0))$ and $\text{rank}(H_A(\beta))$ for two different β in (1) and demonstrate a rank jump:

```

i13 : holonomicRank gkz(A,{0,0,0}) -- vol A in this case
o13 = 4
i14 : holonomicRank gkz(A,{0,0,1})
o14 = 5
i15 : holonomicRank gkz(A,{3/2,0,-2})
o15 = 5

```

SUPPLEMENT. The online supplement contains version 1.0 of `Quasidegrees.m2`.

REFERENCES.

- [Bruns and Herzog 1993] W. Bruns and J. Herzog, *Cohen–Macaulay rings*, Cambridge Studies in Advanced Mathematics **39**, Cambridge University Press, 1993. MR Zbl
- [Eisenbud 1995] D. Eisenbud, *Commutative algebra: with a view toward algebraic geometry*, Graduate Texts in Mathematics **150**, Springer, 1995. MR Zbl
- [Matusevich et al. 2005] L. F. Matusevich, E. Miller, and U. Walther, “Homological methods for hypergeometric families”, *J. Amer. Math. Soc.* **18**:4 (2005), 919–941. MR Zbl
- [Miller 2002] E. Miller, “Graded Greenlees–May duality and the Čech hull”, pp. 233–253 in *Local cohomology and its applications* ((Guanajuato, 1999)), edited by G. Lyubeznik, Lecture Notes in Pure and Appl. Math. **226**, Dekker, New York, 2002. MR Zbl
- [Miller and Sturmfels 2005] E. Miller and B. Sturmfels, *Combinatorial commutative algebra*, Graduate Texts in Mathematics **227**, Springer, 2005. MR Zbl
- [Saito et al. 2000] M. Saito, B. Sturmfels, and N. Takayama, *Gröbner deformations of hypergeometric differential equations*, Algorithms and Computation in Mathematics **6**, Springer, 2000. MR Zbl

RECEIVED: 27 Jun 2015

REVISED: 13 Oct 2018

ACCEPTED: 26 Feb 2019

ROBERTO BARRERA:

rbarrera@txstate.edu

Department of Mathematics, Texas State University, San Marcos, TX, United States



<i>Strongly stable ideals and Hilbert polynomials</i>	1
Davide Alberelli and Paolo Lella	
<i>DiffAlg: a Differential algebra package</i>	11
Manuel Dubinsky, César Massri, Ariel Molinuevo and Federico Quallbrunn	
<i>Matroids: a Macaulay2 package</i>	19
Justin Chen	
<i>Computing quasidegrees of A-graded modules</i>	29
Roberto Barrera	
<i>An algorithm for enumerating difference sets</i>	35
Dylan Peifer	
<i>Hyperplane arrangements in CoCoA</i>	43
Elisa Palezzato and Michele Torielli	
<i>Numerical implicitization</i>	55
Justin Chen and Joe Kileel	
<i>Random Monomial Ideals: a Macaulay2 package</i>	65
Sonja Petrović, Despina Stasi and Dane Wilburne	
<i>Calculations involving symbolic powers</i>	71
Ben Drabkin, Eloísa Grifo, Alexandra Seceleanu and Branden Stone	
<i>The gfanlib interface in Singular and its applications</i>	81
Anders Jensen, Yue Ren and Hans Schönemann	

