

```

gap> g:= SymmetricGroup( 4 );
Sym( [ 1 .. 4 ] )
gap> tbl:= CharacterTable( g );; HasIrr( tbl );
i5 : betti(t,Weights=>{1,0})
false
      0 1 2 3 4 gap> tblmod2:= CharacterTable( tbl, 2 );
o5 = total: 1 4 13 14 4 BrauerTable( Sym( [ 1 .. 4 ] ), 2 )
      0: 1 . . . .
      1: . 2 2 4 2 gap> tblmod2 = CharacterTable( tbl, 2 );
      2: . 2 5 6 . true
      3: . . 4 . 2
      4: . . . 4 . gap> tblmod2 = BrauerTable( tbl, 2 );
      5: . . 2 . . true
gap> tblmod2 = BrauerTable( tbl, 2 );
o5 : BrauerTable( Sym( [ 1 .. 4 ] ), 2 )
i6 : betti(t,Weights=>{0,1})
      0 1 2 3 4 gap> libtbl:= CharacterTable( "M" );
o6 = total: 1 4 13 14 4 CharacterTable( "M" )
      0: 1 . . . . gap> CharacterTableRegular( libtbl, 2 );
      1: . 2 2 . 2 BrauerTable( "M", 2 );
      2: . 2 5 6 . BrauerTable( libtbl, 2 );
      3: . . 4 . 2 gap> BrauerTable( libtbl, 2 );
      4: . . . 4 . fail
      5: . . 2 . .
gap> CharacterTable( "Symmetric", 4 );
o6 : BettiTally CharacterTable( "Sym(4)" )
i7 : t1 = betti(t,Weights=>{1,1})
gap> ComputedBrauerTables( tbl );
      0 1 2 3 4 [ , BrauerTable( Sym( [ 1 .. 4 ] ), 2 ) ]
o7 = total: 1 4 13 14 4
      0: 1 . . . .
      1: . . . . .
      2: . . . . .
      3: . 2 . . .
      4: . . . . .
      5: . 2 . . .
      6: . . 1 . .
      7: . . 8 6 .
      8: . . 4 8 4
ring r1 = 32003,(x,y,z),ds;
int a,b,c,t=11,5,3,0;
poly f = x^a+y^b+z^(3*c)+x^(c+2)*y^(c-1)+x^(c-2)*y^c*(y^2+t*x)^2;
option(noprot);
timer=1;
ring r2 = 32003,(x,y,z),dp;
poly f=imap(r1,f);
ideal j=jacob(f);
vdim(std(j));
==> 536
vdim(std(j+f));
==> 195
timer=0; // reset timer
o7 : BettiTally
i8 : peek t1
o8 = BettiTally{(0, {0, 0}, 0) => 1 }
      (1, {2, 2}, 4) => 2
      (1, {3, 3}, 6) => 2
      (2, {3, 7}, 10) => 2
      (2, {4, 4}, 8) => 1
      (2, {4, 5}, 9) => 4
      (2, {5, 4}, 9) => 4
      (2, {7, 3}, 10) => 2
      (3, {4, 7}, 11) => 4
      (3, {5, 5}, 10) => 6
      (3, {7, 3}, 10) => 2
      (4, {5, 7}, 12) => 2
      (4, {7, 5}, 12) => 2

```

Journal of Software for Algebra and Geometry

The Schur-Veronese package in Macaulay2

JULIETTE BRUCE, DANIEL ERMAN, STEVE GOLDSTEIN AND JAY YANG

The Schur–Veronese package in Macaulay2

JULIETTE BRUCE, DANIEL ERMAN, STEVE GOLDSTEIN AND JAY YANG

ABSTRACT: This note introduces the *Macaulay2* package *SchurVeronese*, which gathers together data about Veronese syzygies and makes it readily accessible in *Macaulay2*. In addition to standard Betti tables, the package includes information about the Schur decompositions of the various spaces of syzygies. The package also includes a number of functions useful for manipulating and studying this data.

In [Bruce et al. 2020] the authors used a combination of high-throughput and high-performance computation and numerical techniques to compute the Betti tables of \mathbb{P}^2 under the d -fold Veronese embedding, as well as the Betti tables of the pushforwards of line bundles $\mathcal{O}_{\mathbb{P}^2}(b)$ under that embedding, for a number of values of b and d . These computations resulted in new data, such as Betti tables, multigraded Betti numbers, and Schur Betti numbers. (For $b = 0$, most the cases had been previously computed in [Castricky et al.].) This note introduces the *SchurVeronese* package for *Macaulay2*, which makes this data readily accessible via *Macaulay2* for further experimentation and study.

1. VERONESE SYZYGIES. Throughout this section we fix $n \in \mathbb{N}$ and let $S = \mathbb{C}[x_0, x_1, \dots, x_n]$ be the polynomial ring with the standard grading. The d -th Veronese module of S twisted by b is

$$S(b; d) := \bigoplus_{i \in \mathbb{Z}} S_{di+b}.$$

If $b = 0$, then $S(0; d)$ is the Veronese subring of S , and if $b \neq 0$ then $S(b; d)$ is an $S(0; d)$ -module. Moreover, if we set $R = \text{Sym}(S_d)$ to be the symmetric algebra on S_d , then we may consider $S(b; d)$ as a graded R -module. Geometrically, if $b = 0$ this corresponds to the homogenous coordinate ring of \mathbb{P}^n under the d -fold embedding $\mathbb{P}^n \rightarrow \mathbb{P}^{\binom{n+d}{d}-1}$, and for other b it corresponds to the pushforward of $\mathcal{O}_{\mathbb{P}^n}(b)$ under the d -fold embedding.

Our interest is in studying the syzygies of $S(b; d)$. See the introduction of [Bruce et al. 2020] for background on Veronese syzygies including a summary of known results. Throughout this paper, we set $K_{p,q}(\mathbb{P}^n, b; d) := \text{Tor}_p^R(S(b; d), \mathbb{C})_{p+q}$, which is isomorphic to the vector space of degree $p+q$ syzygies of $S(b; d)$ of homological degree p . Using the standard conventions for graded Betti numbers, the rank

Bruce received support from the NSF GRFP under grant DGE-1256259, NSF grant DMS-1502553, NSF MSPRF DMS-2002239, and from the Graduate School and the Office of the Vice Chancellor for Research and Graduate Education at the University of Wisconsin–Madison with funding from the Wisconsin Alumni Research Foundation. She is grateful for the support of the Mathematical Sciences Research Institute in Berkeley, California, where she was in residence for the Fall 2020 semester. Erman received support from NSF grant DMS-1601619. Yang received support from NSF grant DMS-1502553.

MSC2010: 13D02.

Keywords: syzygies, free resolutions, Veronese syzygies.

SchurVeronese version 1.1

of the vector space $K_{p,q}$ corresponds to the Betti number $\beta_{p,p+q}$, and we write $\beta_{p,p+q}(S(b; d)) := \dim \operatorname{Tor}_p^R(S(b; d), \mathbb{C})_{p+q} = \dim K_{p,q}(\mathbb{P}^n, b; d)$. Following the usual *Macaulay2* notation, the Betti table of $S(b; d)$ will be the table where $\beta_{p,p+q}(S(b; d))$ is placed in the (p, q) -spot.

Outside of the case $n = 1$, the Betti tables of $S(b; d)$ are unknown even for modest values of d . There is not even a conjecture about what the Betti table of $S(b; d)$ should be for $n = 2$ and $d \geq 7$.

This package provides an array of computed data about $S(b; d)$ in the case $n = 2$ and for $0 \leq b < d \leq 8$ (though the data are incomplete for some of the larger values of d). While computing this data, including the Schur functor decompositions, took substantial time, the resulting data are concise and easy to work with in *Macaulay2*. The bulk of this package thus consists of these output data, which are included as auxiliary files. The functions provided in this package make this data accessible in a user-friendly way. Our hope is that this will allow those interested in Veronese syzygies to make headway on formulating conjectures and proving results in this area. Moreover, as new cases of Veronese syzygies are computed, these can easily be incorporated into future versions of the package.

2. AN OVERVIEW OF THE DATA. When computing data for $S(b; d)$ we always work under the hypothesis that $0 \leq b < d$, as the Betti table of $S(b; d)$ and $S(b + d; d)$ differ only by a vertical shift. We have included data for the cases $n = 1$ and $d \leq 10$, although this can also easily be computed using the Eagon–Northcott complex. The main data are for the cases $n = 2$ and $0 \leq b < d \leq 8$. In [Bruce et al. 2020], we obtained full computations for $d \leq 6$; moreover since those algorithms worked in parallel with respect to multidegrees, we obtained incomplete data for some cases where $d = 7, 8$, and we have included these partial data in this package as well.

The algorithms in [Bruce et al. 2020] are a mix of symbolic and numeric algebra. Thus some entries in the data are not provably correct, while others are. One can determine precisely when $K_{p,q} \neq 0$ by combining [Ein and Lazarsfeld 2012, Remark 6.5], [Green 1984b, Theorem 2.2], and [Green 1984a, Theorem 2.c.6]. Our computation of a $K_{p,q}$ -group (and all related data such as the Schur functor decomposition) will be provably correct if and only if $K_{p+1,q-1}$ and $K_{p-1,q+1}$ both vanish; in cases where this does not occur, the data for $K_{p,q}$ may have been computed numerically, and thus may not be provably correct. For a longer discussion of potential numerical error issues, see [Bruce et al. 2020, §5.2].

3. TOTAL BETTI TABLES. The Betti table for $S(b; d)$ can be called up using the `totalBettiTally` command. For example, the Betti table of $S(2; 4)$ when $n = 2$ is produced below.

```
i6 : totalBettiTally(4,2,0)
o6 = total:  0  1  2  3  4  5  6  7  8  9  10 11 12
            1  75 536 1947 4488 7095 7920 6237 3344 1089 175 24 3
            0:  1
            1:  . 75 536 1947 4488 7095 7920 6237 3344 1089 120 . .
            2:  . . . . . . . . . . . 55 24 3
o6 : BettiTally
```

Note that this is purely numeric: the package does not produce a minimal free resolution; the function simply returns the Betti numbers obtained by a previous computation. The command `totalBetti` is similar, but expresses the Betti numbers simply as a hash table.

There is also a distinction between the indexing conventions. When working with hash tables, we follow the more concise $K_{p,q}$ indexing conventions, instead of the $\beta_{p,p+q}$ indexing conventions used for Betti tallies. Thus, for instance, in the above example, the Betti number $\beta_{2,3}$ would correspond to key $(2, \{3\}, 3)$ in the Betti tally, but in the hash table `totalBetti` it corresponds to key $(2, 1)$:

```
i4 :E = totalBetti(4,2,0);
i5 : E#(2,1)
o5 = 536
```

If one tries to call a Betti table outside of the acceptable range of n, b, d , we return an error message.

```
i10 : totalBettiTally(4,3,0)
o10 = Need n = 1 or 2
```

As noted above, there were instances where we were able to partially compute Betti tables, for instance in the case of the 7-uple embedding of \mathbb{P}^2 . In those cases, we have recorded the entries that we know, and we mark the unknown entries with “infinity”. For example:

```
i14 : B = totalBetti(7,2,0);
i15 : B#(4,1)
o15 = 1031184
i16 : B#(20,1)
o16 = infinity
o16 : InfiniteNumber
```

Thus, in this case, we see $\dim K_{4,1}(\mathbb{P}^2, 2; 7) = 1031184$, but were unable to compute $\dim K_{20,1}(\mathbb{P}^2, 2; 7)$.

4. SCHUR DECOMPOSITION. When $n = 2$ and $d \geq 5$, the Betti tables of $S(b; d)$ are often unwieldy to work with, as they and their entries tend to be quite large. For example, the Betti table of $S(0; 6)$ has 26 columns and many of the entries are on the order of 10^7 .

A more concise way of recording the syzygies would be to take into account the symmetries coming from representation theory. The natural linear action of $GL_{n+1}(\mathbb{C})$ on S induces an action on each vector space $K_{p,q}(\mathbb{P}^n, b; d)$. We can thus decompose this as a direct sum of Schur functors of total weight $d(p+q) + b$, i.e.,

$$K_{p,q}(\mathbb{P}^n, b; d) = \bigoplus_{|\lambda|=d(p+q)+b} \mathcal{S}_\lambda(\mathbb{C}^{n+1})^{\oplus m_{p,\lambda}(\mathbb{P}^n, b; d)},$$

with $m_{p,\lambda}(\mathbb{P}^n, b; d)$ being the Schur Betti numbers and \mathcal{S}_λ being the Schur functor corresponding to the partition λ [Fulton and Harris 1991, p. 76]. The Schur Betti numbers can be accessed via the `schurBetti` command, which returns a hash table whose keys correspond to pairs (p, q) for which $K_{p,q}(\mathbb{P}^n, b; d) \neq 0$, and whose values are lists corresponding to the Schur decomposition of this syzygy module.

For example, let us consider $K_{2,1}(\mathbb{P}^2, 0; 4)$, which is a vector space of dimension 536. As a representation of $GL_3(\mathbb{C})$, it turns out to be the sum of 9 distinct Schur functors, each appearing with multiplicity 1:

$$K_{2,1}(\mathbb{P}^2, 0; 4) = \mathcal{S}_{(9,2,1)} \oplus \mathcal{S}_{(8,4,0)} \oplus \mathcal{S}_{(8,3,1)} \oplus \mathcal{S}_{(7,5,0)} \oplus \mathcal{S}_{(7,4,1)} \oplus \mathcal{S}_{(7,3,2)} \oplus \mathcal{S}_{(6,5,1)} \oplus \mathcal{S}_{(6,4,2)} \oplus \mathcal{S}_{(5,4,1)}.$$

```
i26 : (schurBetti(4,2,0))#(2,1)
o26 = {({9, 2, 1}, 1), ({8, 4, 0}, 1), ({8, 3, 1}, 1), ({7, 5, 0}, 1),
-----
({7, 4, 1}, 1), ({7, 3, 2}, 1), ({6, 5, 1}, 1), ({6, 4, 2}, 1), ({5, 4, 3}, 1)}
o8 : List
```

From this, it is easy to compute statistics such as the number of representations and the number of distinct representations appearing in the Schur decomposition of $K_{p,q}(n, b; d)$. The *SchurVeronese* package provides commands for these. For instance, in our example above we see that:

```
i11 : (numDistinctRepsBetti(4,2,0))#(2,1)
o11 = 9
```

We can also display the number of representations appearing in each entry of the Betti table. In the following example, the first table counts distinct Schur functors and the second counts the number of Schur functors with multiplicity:

```
i29 : makeBettiTally numDistinctRepsBetti(4,2,0)
o29 = total: 0 1 2 3 4 5 6 7 8 9 10 11 12
             1 2 9 17 23 23 26 25 21 13 3 1 1
0: 1
1: . 2 9 17 23 23 26 25 21 13 3 1 1
2: . . . . . . . . . 2 1 1

i30 : makeBettiTally numRepsBetti(4,2,0)
o30 = total: 0 1 2 3 4 5 6 7 8 9 10 11 12
             1 2 9 28 55 79 86 69 38 14 3 1 1
0: 1
1: . 2 9 28 55 79 86 69 38 14 3 1 1
2: . . . . . . . . . 2 1 1
```

Thus, $K_{4,1}(\mathbb{P}^2, 0; 4)$ is the sum of 55 irreducible representations, 23 of which are distinct.

5. MULTIGRADED BETTI NUMBERS. One can also specialize the action of $GL_{n+1}(\mathbb{C})$ to the torus action via $(\mathbb{C}^*)^{n+1}$. This gives a decomposition of $K_{p,q}(\mathbb{P}^n, b; d)$ into a sum of \mathbb{Z}^{n+1} -graded vector spaces of total weight $d(p+q)+b$. Specifically, writing $\mathbb{C}(-\mathbf{a})$ for the vector space \mathbb{C} together with the $(\mathbb{C}^*)^{n+1}$ -action given by $(\lambda_0, \lambda_1, \dots, \lambda_n) \cdot \mu = \lambda_0^{a_0} \lambda_1^{a_1} \cdots \lambda_n^{a_n} \mu$, we have

$$K_{p,q}(\mathbb{P}^n, b; d) = \bigoplus_{\substack{\mathbf{a} \in \mathbb{Z}^{n+1} \\ |\mathbf{a}| = d(p+q)+b}} \mathbb{C}(-\mathbf{a})^{\oplus \beta_{p,\mathbf{a}}(\mathbb{P}^n, b; d)}$$

as \mathbb{Z}^{n+1} -graded vector spaces, or equivalently as $(\mathbb{C}^*)^{n+1}$ representations.

The *SchurVeronese* package produces these multigraded Betti numbers for a number of examples via the `multiBetti` command. As `schurBetti` does, this command returns a hash table whose keys correspond to pairs (p, q) for which $K_{p,q}(\mathbb{P}^n, b; d) \neq 0$, and whose values are multigraded Hilbert polynomials encoding the multigraded decomposition of $K_{p,q}(n, b; d)$. More specifically, the value of `(multiBetti(d,n,b))#(p,q)` is the polynomial

$$\sum_{\substack{\mathbf{a} \in \mathbb{Z}^{n+1} \\ |\mathbf{a}| = d(p+q)+b}} \beta_{p,\mathbf{a}}(n, b; d) \mathbf{t}^{\mathbf{a}}$$

where $\mathbf{t}^{\mathbf{a}}$ denotes $t_0^{a_0} t_1^{a_1} \cdots t_n^{a_n}$.

For example, $K_{12,2}(2, 0; 4)$ is the 3-dimensional \mathbb{Z}^3 -graded vector space

$$K_{12,2}(2, 0; 4) \cong \mathbb{C}(-19, 19, 18) \oplus \mathbb{C}(-19, 18, 19) \oplus \mathbb{C}(-18, 19, 19).$$

The following code computes this, illustrating that the multigraded Hilbert function for $K_{12,2}(2, 0; 4)$ is

```

 $t_0^{19} t_1^{19} t_2^{18} + t_0^{19} t_1^{18} t_2^{19} + t_0^{18} t_1^{19} t_2^{19}.$ 
i4 : (multiBetti(4,2,0))#(12,2)
o4 =  $t_0^{19} t_1^{19} t_2^{18} + t_0^{19} t_1^{18} t_2^{19} + t_0^{18} t_1^{19} t_2^{19}$ 
o4 : QQ[t_0, t_1, t_2]

```

SUPPLEMENT. The online supplement contains version 1.1 of *SchurVeronese*.

ACKNOWLEDGMENTS. We thank Claudiu Raicu and Gregory G. Smith for useful conversations.

REFERENCES.

- [Bruce et al. 2020] J. Bruce, D. Erman, S. Goldstein, and J. Yang, “Conjectures and Computations about Veronese Syzygies”, *Experimental Mathematics* **29**:4 (2020), 398–413.
- [Castricky et al.] W. Castryck, F. Cools, J. Demeyer, and A. Lemmens, “Computing graded Betti tables of toric surfaces”. arXiv
- [Ein and Lazarsfeld 2012] L. Ein and R. Lazarsfeld, “Asymptotic syzygies of algebraic varieties”, *Inventiones Mathematicae* **190** (2012), 603–646.
- [Fulton and Harris 1991] W. Fulton and J. Harris, *Representation theory*, vol. 129, Graduate Texts in Mathematics, Springer-Verlag, New York, 1991. MR
- [Green 1984a] M. L. Green, “Koszul cohomology and the geometry of projective varieties, I”, *J. Differential Geom.* **19**:1 (1984), 125–171.
- [Green 1984b] M. L. Green, “Koszul cohomology and the geometry of projective varieties, II”, *J. Differential Geom.* **20**:1 (1984), 279–289.

RECEIVED: 28 May 2019

REVISED: 18 Feb 2021

ACCEPTED: 5 May 2021

JULIETTE BRUCE:

juliette.bruce@berkeley.edu

Department of Mathematics, University of California, Berkeley, Berkeley, CA, United States

DANIEL ERMAN:

derman@math.wisc.edu

Department of Mathematics, University of Wisconsin, Van Vleck Hall, Madison, WI, United States

STEVE GOLDSTEIN:

sgoldstein@wisc.edu

Botany Department and Department of Biostatistics and Medical Informatics, University of Wisconsin, Madison, WI, United States

JAY YANG:

jkYang@umn.edu

School of Mathematics, University of Minnesota, Minneapolis, MN, United States

<i>Phylogenetic trees</i>	1
Hector Baños, Nathaniel Bushek, Ruth Davidson, Elizabeth Gross, Pamela E. Harris, Robert Krone, Colby Long, Allen Stewart and Robert Walker	
<i>Software for doing computations in graded Lie algebras</i>	9
Clas Löfwall and Samuel Lundqvist	
<i>The relative canonical resolution: Macaulay2-package, experiments and conjectures</i>	15
Christian Bopp and Michael Hoff	
<i>The FrobeniusThresholds package for Macaulay2</i>	25
Daniel J. Hernández, Karl Schwede, Pedro Teixeira and Emily E. Witt	
<i>Computing theta functions with Julia</i>	41
Daniele Agostini and Lynn Chua	
<i>Decomposable sparse polynomial systems</i>	53
Taylor Brysiewicz, Jose Israel Rodriguez, Frank Sottile and Thomas Yahl	
<i>A package for computations with sparse resultants</i>	61
Giovanni Staglianò	
<i>ExteriorModules: a package for computing monomial modules over an exterior algebra</i>	71
Luca Amata and Marilena Crupi	
<i>The Schur–Veronese package in Macaulay2</i>	83
Juliette Bruce, Daniel Erman, Steve Goldstein and Jay Yang	
<i>admcycles - a Sage package for calculations in the tautological ring of the moduli space of stable curves</i>	89
Vincent Delecroix, Johannes Schmitt and Jason van Zelm	
<i>Coding theory package for Macaulay2</i>	113
Taylor Ball, Eduardo Camps, Henry Chimal-Dzul, Delio Jaramillo-Velez, Hiram López, Nathan Nichols, Matthew Perkins, Ivan Soprunov, German Vera-Martínez and Gwyn Whieldon	
<i>Threaded Gröbner bases: a Macaulay2 package</i>	123
Sonja Petrović and Shahrzad Zelenberg	
<i>Standard pairs of monomial ideals over nonnormal affine semigroups in SageMath</i>	129
Byeongsu Yu	
<i>Computations with rational maps between multi-projective varieties</i>	143
Giovanni Staglianò	