


NISSUNA UMANA INVESTIGAZIONE SI PUO DIMANDARE VERA SCIENZA
S'ESSA NON PASSA PER LE MATEMATICHE DIMOSTRAZIONI
LEONARDO DA VINCI



vol. 6

no. 3

2018

MATHEMATICS AND MECHANICS
of
Complex Systems

SANTOSH KUMAR AND DINBANDHU MANDAL

**GENERATION OF SH-TYPE WAVES DUE TO SHEARING STRESS
DISCONTINUITY IN AN ANISOTROPIC LAYER OVERLYING AN
INITIALLY STRESSED
ELASTIC HALF-SPACE**





GENERATION OF SH-TYPE WAVES DUE TO SHEARING STRESS DISCONTINUITY IN AN ANISOTROPIC LAYER OVERLYING AN INITIALLY STRESSED ELASTIC HALF-SPACE

SANTOSH KUMAR AND DINBANDHU MANDAL

The paper investigates the generation of SH-type waves due to a sudden application of a stress discontinuity which moves after creation at the anisotropic layer of finite thickness overlying an initially stressed isotropic half-space. The displacements are obtained in exact form by the method due to Cagniard modified by de Hoop. Two cases of shearing stress discontinuities are considered. The numerical results are obtained for a particular model and discussed by plotting graphs for displacement component with the elapsed time of the disturbance for different values of initial stress and also for different values of initial time at which pulses arrive.

1. Introduction

The notion of initial stress is essential to the study of seismic wave propagation. Biot [1940] is largely responsible for the notion's introduction and initial applications to elastic wave propagation; in [Biot 1965] he further developed the notion of initial stress. Many authors have used that book as fundamental to the study of wave propagation in an initially stressed medium. Abd-Alla and Ahmed [1999] analyzed Love waves propagation in a non-homogeneous orthotropic elastic layer under initial stress overlying semi-infinite medium. Khurana [2001] considered the effect of initial stress on the propagation of Love wave. Further significant steps were taken in [Das and Dey 1968; 1970, Dey 1971; Dey and Addy 1978; Chattopadhyay and De 1981; Chattopadhyay and Kar 1978; Majhi et al. 2016; 2017] to cite but a few works.

In addition to initial stress, shearing stress discontinuity also plays a vital role in the study of seismic wave propagation. Pal [1983] considered the problem of generation and propagation of SH-type waves due to non-uniformly moving stress

Communicated by Holm Altenbach.

MSC2010: 35L05, 74J15.

Keywords: SH-type waves, initial stress, shearing stress discontinuity, Cagniard–de Hoop technique.

discontinuity in layered anisotropic elastic half-space using Garvin's [1956] techniques, which are a modification of Cagniard's [1939] technique. Pal and Kumar [2000] considered the generation of SH waves by a moving stress discontinuity in an anisotropic soil layer over an elastic half-space using the Cagniard–de Hoop special reduction technique [de Hoop 1960]. Next de Hoop [2002] considered the reflection and transmission properties of an elastic interfacial bonding of two semi-infinite solids, investigated for the simplest possible case of a line-source excited two-dimensional SH-wave. Pal and Mandal [2014] studied the generation of SH-type waves due to sudden application of a stress discontinuity which moves after creation at the sandy layer of finite thickness overlying an isotropic and inhomogeneous elastic half-space. Mandal et al. [2014] considered the disturbance and propagation of SH-type waves in an anisotropic soil layer overlying an inhomogeneous elastic half-space by a moving stress discontinuity. All authors have considered the effect of initial stress and shearing stress discontinuity separately but haven't considered the initial stress and shearing stress discontinuity together.

In the present problem our intention is to investigate the two dimensional problem of generation of SH-type waves at the free surface of an anisotropic layer due to an impulsive stress discontinuity moving with uniform velocity along the interface of initially stressed isotropic medium. The displacement is calculated numerically for two particular distances on the surface for two different types of the discontinuity in the shearing stress for different value of initial stress. It involves Laplace and Fourier transform and the inversion is based on Garvin's [1956] method. The problem discussed may be of importance in connection with the propagation of cracks in the layer. Two cases of stress discontinuity are considered and the numerical results are shown graphically.

2. Formulation of problem

We consider an anisotropic elastic layer of thickness h with elastic constants L_1, N_1 and density ρ_1 over an initially stressed isotropic half-space with elastic constant μ_2 and density ρ_2 . The interface of these two media is considered at $z = 0$ whereas free surface is at $z = -h$. Here, z axis is directed vertically downward and x axis is assumed in the direction of the propagation of wave with velocity c . For SH-type of waves the displacement does not depend on y and if (u, v, w) be the displacement at any point $P(x, y, z)$ into the medium then $u = w = 0$ and v are function of x, z and t . The two equations of motion are identically satisfied. The geometrical configuration is depicted in Figure 1.

The equation of motion for the anisotropic layer (Medium I) without body forces is given by

$$N_1 \frac{\partial^2 v_1}{\partial x^2} + L_1 \frac{\partial^2 v_1}{\partial z^2} = \rho_1 \frac{\partial^2 v_1}{\partial t^2}. \quad (1)$$




Figure 1. Geometry of problem.

The stress strain relation is given by $(\tau_{xy})_I = N_1 \frac{\partial v_1}{\partial x}$ and $(\tau_{yz})_I = L_1 \frac{\partial v_1}{\partial z}$.

The equation of motion for initially stressed isotropic half-space (Medium II) without body forces is given by

$$\frac{\partial(\tau_{xy})_{II}}{\partial x} + \frac{\partial(\tau_{yz})_{II}}{\partial z} - \frac{P}{2} \frac{\partial v_2}{\partial x^2} = \frac{\partial^2}{\partial t^2}(\rho_2 v_2). \tag{2}$$

The stress strain relation is given by $(\tau_{xy})_{II} = \mu_2 \frac{\partial v_2}{\partial x}$ and $(\tau_{yz})_{II} = \mu_2 \frac{\partial v_2}{\partial z}$.

The boundary conditions are

$$(\tau_{yz})_I = 0 \quad \text{at } z = -h, \tag{3}$$

$$v_1 = v_2 \quad \text{at } z = 0, \tag{4}$$

$$(\tau_{yz})_I - (\tau_{yz})_{II} = S(x, t)H(t) \quad \text{at } z = 0, \tag{5}$$

where $S(x, t)$ is a function of x and t ; $H(t)$ is the Heaviside unit function of time t .

3. Method of solution

The above problem can readily be solved by using the Laplace and Fourier transforms combined with the modified Cagniard–de Hoop [1960] method. The Laplace transform with respect to t and the Fourier transform with respect to x are defined by

$$\bar{v} \cong \int_{-\infty}^{\infty} e^{-i\xi x} dx \int_0^{\infty} e^{-pt} v(x, z; t) dt. \tag{6}$$

We can easily get the upper and lower layer with $v_2 \rightarrow 0$ as $z \rightarrow \infty$ in the form

$$\bar{v}_1(x, z; p) = \int_{-\infty}^{\infty} (A \cosh s_1 z + B \sinh s_1 z) e^{i\xi x} d\xi, \tag{7}$$

$$\bar{v}_2 = \int_{-\infty}^{\infty} C e^{(i\xi x - s_2 z)} d\xi, \tag{8}$$

where the constants A , B and C are to be determined from the boundary conditions

(3)–(5):

$$s_1 = \left(\phi_1^2 \xi^2 + \frac{p^2}{\beta_1^2} \right)^{\frac{1}{2}}, \quad s_2 = \left(\phi_2^2 \xi^2 + \frac{p^2}{\beta_2^2} \right)^{\frac{1}{2}}, \tag{9}$$

where $\beta_1^2 = \frac{L_1}{\rho_1}$, $\beta_2^2 = \frac{\mu_2}{\rho_2}$, $\phi_1^2 = \frac{N_1}{L_1}$ and $\phi_2^2 = 1 - \frac{P}{2\mu_2}$.

It follows from the boundary conditions (3) and (4) that

$$A = C, \quad A \cosh s_1 h = B \sinh s_1 h. \tag{10}$$

Case I. Let

$$S(x, t) = \begin{cases} Q, & a \leq x \leq b + Vt, \\ 0, & \text{elsewhere,} \end{cases} \tag{11}$$

where Q is constant.

This definition of stress discontinuity shows that it is created in the region $x = a$ to $x = b$ and then expands with the uniform velocity V in the x direction. In particular, when $a = b = 0$, the discontinuity is created at the origin and expands with uniform velocity V in the x direction.

From the boundary condition (5) one gets, with the help of (11),

$$BL_1 s_1 + C\mu_2 s_2 = \frac{Q}{2\pi p} \left[\frac{e^{-i\xi a} - e^{-i\xi b}}{i\xi} + \frac{e^{-i\xi b}}{i\xi + \frac{p}{V}} \right]. \tag{12}$$

Solving for A , B and C from (10) and (12), we get the displacement function at the free surface at $z = -h$ in the form

$$\begin{aligned} \bar{v}_1(x, -h; p) &= \frac{Q}{2\pi p} \int_{-\infty}^{\infty} \frac{e^{i\xi x}}{(L_1 s_1 \sinh s_1 h + \mu_2 s_2 \cosh s_1 h)} \left[\frac{e^{-i\xi a} - e^{-i\xi b}}{i\xi} + \frac{e^{-i\xi b}}{i\xi + \frac{p}{V}} \right] d\xi \\ &= \frac{Q}{\pi p} \int_{-\infty}^{\infty} \frac{e^{(i\xi x - h s_1)}}{(L_1 s_1 + \mu_2 s_2)} \left[\frac{e^{-i\xi a} - e^{-i\xi b}}{i\xi} + \frac{e^{-i\xi b}}{i\xi + \frac{p}{V}} \right] (1 - Ke^{-2hs_1})^{-1} d\xi, \end{aligned} \tag{13}$$

where

$$K = \frac{L_1 s_1 - \mu_2 s_2}{L_1 s_1 + \mu_2 s_2} < 1, \tag{14}$$

represents the reflection coefficient of SH-waves incident from the sandy medium at the interface between two half-spaces. The coefficients of different power of K in series of (13) are associated with the pulses undergoing repeated reflection in the upper layer. Using the inverse Laplace transform, we can rewrite (13) in a convenient form:

$$v_1(x, -h; p) = L^{-1}(I_1 + I_2 + I_3) = L^{-1}(I_1) + L^{-1}(I_2) + L^{-1}(I_3), \tag{15}$$

where I_1, I_2 and I_3 are defined in the Appendix. The inverse Laplace transforms of I_1, I_2 and I_3 are obtained by following [Garvin 1956]; details are in the Appendix.

Equation (15) gives the exact value of the surface displacement field $v_1(x, -h, t)$ surface.

Case II. Let

$$S(x, t) = Qh\delta(x - Vt), \tag{16}$$

where Q is a constant and $\delta(x - Vt)$ is Dirac’s delta function of argument $(x - Vt)$. A term h is included on the right-hand side of (16) so as to give S as the dimension of a stress.

The boundary condition (5) gives

$$BL_1s_1 + C\mu_2s_2 = \frac{Qh}{2\pi V(i\xi + \frac{p}{V})}. \tag{17}$$

Solving for A, B and C from (10) and (17) one gets

$$\bar{v}_1(x, -h; p) = \frac{Qh}{\pi V} \int_{-\infty}^{\infty} \frac{e^{(i\xi x - hs_1)}(1 + Ke^{-2hs_1} + K^2e^{-4hs_1} + \dots)}{(i\xi + \frac{p}{V})(L_1s_1 + \mu_2s_2)} d\xi \tag{18}$$

Proceeding similarly as in Case I we obtain

$$v_1(x, -h, t) = \frac{2Q\beta_1h}{\pi L_1} \sum_{n=1,3,5,\dots} \int_0^t G_n[\zeta_n(\lambda)] d\lambda, \tag{19}$$

where

$$G_n[\zeta_n(t)] = \text{Re} \left[\left\{ 1 + \phi_1^2 \zeta_n^2(t) \right\}^{\frac{1}{2}} + \mu (\varepsilon^2 + \phi_2^2 \zeta_n^2(t))^{\frac{1}{2}} \right]^{-1} \times \left[\frac{\beta_1}{V} + i\zeta_n(t) \right]^{-1} \\ \times K^{\frac{n-1}{2}} [\zeta_n(t)] \frac{d\zeta_n(t)}{dt} H \left[t - (x^2 + n^2h^2)^{\frac{1}{2}} \beta_1^{-1} \right] \tag{20}$$

and

$$\zeta_n(t) = \frac{\beta_1}{x^2 + n^2h^2\phi_1^2} \left[itx + nh \left\{ t^2 - (x^2 + n^2h^2\phi_1^2)\beta_1^{-2} \right\}^{\frac{1}{2}} \right], \quad n = 1, 3, 5, \dots \tag{21}$$

If the stress discontinuity is taken as $H(x) - H(x - Vt)$ in place of $\delta(x - Vt)$ the corresponding expression on the right-hand side of (18) will differ only by a constant factor from I_3 (with $a = b = 0$).

4. Numerical results and discussion

For numerical results, we have taken data for anisotropic layer and initially stressed half-space from [Babuska and Cara 1991]:

$$\begin{aligned} N_1 &= 175 \text{ GPa}, & L_1 &= 202 \text{ GPa}, & \rho_1 &= 4408 \text{ kg/m}^3 \\ \mu_2 &= 91.6 \text{ GPa}, & \rho_2 &= 3582 \text{ kg/m}^3. \end{aligned}$$

The values of $K_1 v_1(x, -h, t)$ for $x = 7h$ and $x = 14h$ have been plotted against $\tau_1 = \tau - \tau_0$, where τ_0 denotes the time at which the disturbance arrives at the point of observation with $K_1 = \frac{\pi L_1}{2Q\beta_1 h}$, $\tau = \frac{t\beta_1}{h}$ is the time of the disturbance to arrive from source to initial point. The value of τ at $x = 7h$ is $(7^2 + n^2\phi_1^2)^{1/2}$, $n = 1, 3, 5, \dots$ and $\tau_0 = 7.06$ at $n = 1$ and the value of τ at $x = 14h$ is $(14^2 + n^2\phi_1^2)^{1/2}$, $n = 1, 3, 5, \dots$ and $\tau_0 = 14.03$ at $n = 1$.

When $x = 7h$, for six initial values, we have

$$K_1 v_1(x, -h, t) = \sum_{n=1,3,5,7,9,11} A^0(\theta_n) \cosh^{-1}\left(\tau / \sqrt{7^2 + n^2\phi_1^2}\right) H\left(\tau - \sqrt{7^2 + n^2\phi_1^2}\right),$$

where

$$A^0(\theta_n) = \text{Re} \frac{[(1 - \phi_1^2 \cos^2 \theta_n)^{\frac{1}{2}} - \mu(\varepsilon^2 - \phi_2^2 \cos^2 \theta_n)^{\frac{1}{2}}]^{\frac{n-1}{2}} \sin \theta_n}{[(1 - \phi_1^2 \cos^2 \theta_n)^{\frac{1}{2}} + \mu(\varepsilon^2 - \phi_2^2 \cos^2 \theta_n)^{\frac{1}{2}}]^{\frac{n-3}{2}} (\frac{\beta_1}{V} - \cos \theta_n)};$$

represents the reflection coefficient of SH-type waves incident from the anisotropic medium to initially stressed isotropic half-space. When $x = 14h$, for six initial values, we have the same expression for $K_1 v_1(x, -h, t)$, with 7^2 replaced by 14^2 .

Figures 2 and 3 show graphs of the variation of displacement with elapsed time τ_1 for different values of initial stress and for different values of initial time at which pulses arrive.

In Figure 2, top, the graph is plotted for the disturbance effect for $x = 7h$ and $x = 14h$ and fixed value of $P = 1$ GPa. From the figure it can be observed that all the curves start from the origin with sharp changes in their slope and after sometime the curves get smooth. Also, it reflects that the disturbance is more prominent for early arrival of pulses i.e. for $x = 7h$ has more jumping effect than for $x = 14h$. If the place of observation has more distance from the source the impact of pluses is less. In Figure 2, bottom, the graph is plotted for the disturbance effect for $x = 7h$ and $x = 14h$ and fixed value of $P = 10$ GPa. The nature of the curves remains the same, but the magnitude of disturbance increases to a large extent as initial stress increases.

In Figure 3, top, the graph is plotted for the disturbance for $x = 7h$ and different values of P (1 GPa, 10 GPa and 100 GPa). The nature of the curves is oscillating




Figure 2. Variation of $K_1 v_1(x, -h, t)$ with τ_1 for $x = 7h$ and $x = 14h$, for $P = 1$ GPa (top) and $P = 10$ GPa.

and after some time the curves become smooth and steady. As we increase the values of initial stress, the jumping effect increases. In Figure 3, bottom, the graph is plotted for the disturbance for $x = 14h$ and the same values of P . The nature of the curves remains the same, but the effect of disturbance is reduced to a large extent due to the late arrival of the pulses.

5. Conclusions

The generation of SH-type waves at the free surface of an anisotropic layer due to an impulsive stress discontinuity moving with uniform velocity along the interface of initially stressed isotropic medium has been considered. The displacement is calculated numerically for two particular distances on the surface for two different




Figure 3. Variation of $K_1 v_1(x, -h, t)$ with τ_1 for $x = 7h$ (top) and $x = 14h$ (bottom) for different values of P .

types of the discontinuity in the shearing stress for different value of initial stress. It involves Laplace and Fourier transform and the inversion is based on Garvin's [1956] method. The numerical results are obtained for a particular model. From the figures it can be observed that initial stress and initial time at which pulses arrive has a significant effect. From the graph it is visible that that the displacement factor starts oscillating and after sometime it gets stable. The results are more comprising with the real scenario as we see that the disturbance arrives at the surface, it shakes the surface for a while and slowly gets stable. Also, from the figures it is visible that if the observer is nearer to the source then pulses arrive early and produce more disturbance and if the medium is highly pre-stressed then it produces more disturbance.

Appendix

We have

$$\begin{aligned}
 I_1 &= \frac{Q}{\pi p} \int_{-\infty}^{\infty} \frac{(1 - Ke^{-2hs_1})^{-1}}{i\xi(L_1s_1 + \mu_2s_2)} e^{(i\xi x_1 - hs_1)} d\xi, \\
 I_2 &= \frac{Q}{\pi p} \int_{-\infty}^{\infty} \frac{(1 - Ke^{-2hs_1})^{-1}}{i\xi(L_1s_1 + \mu_2s_2)} e^{(i\xi x_2 - hs_1)} d\xi, \\
 I_3 &= \frac{Q}{\pi p} \int_{-\infty}^{\infty} \frac{(1 - Ke^{-2hs_1})^{-1}}{(i\xi + \frac{p}{V})(L_1s_1 + \mu_2s_2)} e^{(i\xi x_2 - hs_1)} d\xi,
 \end{aligned}$$

with $x_1 = x - a$ and $x_2 = x - b$. In order to evaluate the Laplace inversion integral, we have used Garvin's method; see [Garvin 1956] for discussion of the contour integration and mapping.

Next for non-dimensionalisation, we substitute $\xi = \frac{\zeta p}{\beta_1}$, $\mu = \frac{\mu_2}{L_1}$, $\frac{\beta_1}{\beta_2} = \varepsilon$ in the integral above so that $s_1 = \frac{p}{\beta_1}(1 + \phi_1^2 \zeta^2)^{\frac{1}{2}}$, $s_2 = \frac{p}{\beta_2}(\varepsilon^2 + \phi_2^2 \zeta^2)^{\frac{1}{2}}$ and

$$K = \frac{(1 + \phi_1^2 \zeta^2)^{\frac{1}{2}} - \mu(\varepsilon^2 + \phi_2^2 \zeta^2)^{\frac{1}{2}}}{(1 + \phi_1^2 \zeta^2)^{\frac{1}{2}} + \mu(\varepsilon^2 + \phi_2^2 \zeta^2)^{\frac{1}{2}}}.$$

Thus we obtain

$$I_1 = \frac{2Q}{\pi p} \text{Im} \int_0^{\infty} \frac{e^{(i\xi x_1 - hs_1)}(1 + Ke^{-2hs_1} + K^2 e^{-4hs_1} + \dots)}{\xi(L_1s_1 + \mu_2s_2)} d\xi. \tag{A.1}$$

The first term in I_1 is

$$I_{1,1} = \frac{2Q\beta_1}{\pi p L_1} \text{Im} \int_0^{\infty} \frac{\exp\left[-p\left\{-i\zeta x_1 + h(1 + \phi_1^2 \zeta^2)^{\frac{1}{2}}\right\}/\beta_1\right]}{p\zeta\left[(1 + \phi_1^2 \zeta^2)^{\frac{1}{2}} + \mu(\varepsilon^2 + \phi_2^2 \zeta^2)^{\frac{1}{2}}\right]} d\zeta. \tag{A.2}$$

The integrand (A.2) has singularities at $\zeta = 0, \pm \frac{i}{\phi_1}, \pm \frac{i\varepsilon}{\phi_2}$. Let $t = \left\{-i\zeta x_1 + h(1 + \phi_1^2 \zeta^2)^{\frac{1}{2}}\right\}/\beta_1$. Then by inversion $\zeta(t) = \frac{\beta_1}{x_1^2 + \phi_1^2 h^2} \left[itx_1 + h\{t^2 - (x_1^2 + \phi_1^2 h^2)\beta_1^{-2}\}^{\frac{1}{2}} \right]$.

The mapping of the ζ -plane into the t -plane is shown in Figure 4.

Making the reference to the Figure 4 and the paper of [Pal 1983], we find

$$L^{-1}I_{1,1} = \frac{2Q\beta_1}{\pi L_1} \int_0^t (t - \lambda_1) G_{1,1}[\zeta_{1,1}(\lambda_1)] d\lambda_1, \tag{A.3}$$

where $L[tH(t)] = \frac{1}{p^2}$ and

$$\begin{aligned}
 G_{1,1}[\zeta_{1,1}(t)] &= \text{Im} \left[(1 + \phi_1^2 \zeta_{1,1}^2)^{\frac{1}{2}} + \mu(\varepsilon^2 + \phi_2^2 \zeta_{1,1}^2)^{\frac{1}{2}} \right]^{-1} \zeta_{1,1}^{-1}(t) \\
 &\times \frac{d\zeta_{1,1}(t)}{dt} H \left[t - \left\{ \eta(x_1^2 + h^2 \phi_1^2) \right\}^{\frac{1}{2}} \beta_1^{-1} \right].
 \end{aligned}$$




Figure 4. The t -plane showing the mapping and the contour of integration.

Since for $\frac{\phi_1 h}{\beta_1} < t < \frac{(x_1^2 + h^2 \phi_1^2)^{\frac{1}{2}}}{\beta_1}$, $\left[(1 + \phi_1^2 \zeta_{1,1}^2)^{\frac{1}{2}} + \mu(\varepsilon^2 + \phi_2^2 \zeta_{1,1}^2)^{\frac{1}{2}} \right] \zeta_{1,1}^{-1}(t) \times \frac{d\zeta_{1,1}(t)}{dt}$ is real. In general,

$$L^{-1} I_{1,n} = \frac{2Q\beta_1}{\pi L_1} \int_0^t (t - \lambda_1) G_{1,n}[\zeta_{1,n}(\lambda_1)] d\lambda_1, \tag{A.4}$$

where

$$G_{1,n}[\zeta_{1,n}(t)] = \text{Im} \left[\left\{ 1 + \phi_1^2 \zeta_{1,n}^2(t) \right\}^{\frac{1}{2}} + \mu(\varepsilon^2 + \phi_2^2 \zeta_{1,n}^2(t))^{\frac{1}{2}} \right]^{-1} \zeta_{1,n}^{-1}(t) \\ \times K^{\frac{n-1}{2}}[\zeta_{1,n}(t)] \frac{d\zeta_{1,n}(t)}{dt} H \left[t - \left\{ (x_1^2 + h^2 \phi_1^2) \right\}^{\frac{1}{2}} \beta_1^{-1} \right],$$

and

$$\zeta_{1,n}(t) = \frac{\beta_1}{x_1^2 + n^2 h^2 \phi_1^2} \left[itx_1 + nh \left\{ t^2 - (x_1^2 + n^2 h^2 \phi_1^2) \beta_1^{-2} \right\}^{\frac{1}{2}} \right], \quad n = 1, 3, 5, \dots$$

So that

$$L^{-1} I_1 = \sum_{n=1,3,5,\dots} L^{-1} I_{1,n}. \tag{A.5}$$

Similarly

$$L^{-1} I_2 = \sum_{n=1,3,5,\dots} L^{-1} I_{2,n}, \tag{A.6}$$

where x_1 is replaced by x_2 .

Proceeding in the same way, we get

$$L^{-1}I_3 = \sum_{n=1,3,5,\dots} L^{-1}I_{3,n}, \tag{A.7}$$

where

$$L^{-1}I_{3,n} = \frac{2Q\beta_1}{\pi L_1} \int_0^t (t - \lambda_1) G_{3,n}[\zeta_{2,n}(\lambda_1)] d\lambda_1 \tag{A.8}$$

$$G_{3,n}[\zeta_{2,n}(t)] = \text{Re} \left[\left\{ 1 + \phi_1^2 \zeta_{2,n}^2(t) \right\}^{\frac{1}{2}} + \mu \left(\varepsilon^2 + \phi_2^2 \zeta_{2,n}^2(t) \right)^{\frac{1}{2}} \right]^{-1} \times \left[\frac{\beta_1}{V} + i \zeta_{2,n}(t) \right]^{-1} \\ \times K^{\frac{n-1}{2}} [\zeta_{2,n}(t)] \frac{d\zeta_{2,n}(t)}{dt} H \left[t - (x_2^2 + n^2 h^2 \phi_1^2)^{\frac{1}{2}} \beta_1^{-1} \right],$$

and $\zeta_{2,n}(t)$ is given by Appendix with x_2 in place of x_1 .

Acknowledgement

Kumar, is grateful to the I.S.M., Dhanbad for Financial support in the form of a Research Fellowship during Ph.D. and to my guide Prof. P. C. Pal for their immense support and guidance. The authors declare that they have no conflict of interest.

References

- [Abd-Alla and Ahmed 1999] A. M. Abd-Alla and S. M. Ahmed, "Propagation of Love waves in a non-homogeneous orthotropic elastic layer under initial stress overlying semi-infinite medium", *Appl. Math. Comput.* **106**:2 (1999), 265–275.
- [Babuska and Cara 1991] V. Babuska and M. Cara, *Seismic anisotropy in the earth*, Kluwer, Dordrecht, 1991.
- [Biot 1940] M. A. Biot, "The influence of initial stress on elastic wave", *J. Appl. Phys.* **8** (1940), 522–530.
- [Biot 1965] M. A. Biot, *Mechanics of incremental deformation*, Wiley, New York, 1965.
- [Cagniard 1939] L. Cagniard, *Reflection et refraction des ondes seismiques progressive*, Gauthier-Villars, Paris, 1939.
- [Chattopadhyay and De 1981] A. Chattopadhyay and R. K. De, "Propagation of Love type waves in a visco elastic initially stressed layer overlying a visco-elastic half-space with irregular interface", *Rev. Roum. Sci. Tech. Mech. Appl.* **26**:3 (1981), 449–460.
- [Chattopadhyay and Kar 1978] A. Chattopadhyay and B. K. Kar, "On the dispersion curves of Love types waves in an initially stressed crustal layer having an irregular interface", *Geophys. Res. Bull.* **16**:1 (1978), 13–23.
- [Das and Dey 1968] S. C. Das and S. Dey, "Note on gravity waves", *Indian J. Engin. Math.* **1**:2 (1968), 155–160.
- [Das and Dey 1970] S. C. Das and S. Dey, "Edge wave under initial stress", *Appl. Sci. Res.* **22**:1 (1970), 382–389.
- [de Hoop 1960] A. T. de Hoop, "A modification of Cagniard's method for solving seismic pulse problem", *Appl. Sci. Res. B* **8**:1 (1960), 349–356.

- [de Hoop 2002] A. T. de Hoop, “Reflection and transmission of a transient, elastic, line-source excited SH wave by a planer, elastic bonding surface in solid”, *Int. J. Solids Struct.* **39**:21-22 (2002), 5379–5391.
- [Dey 1971] S. Dey, “Wave propagation in two layered medium under initial stresses”, *J. Pure Appl. Geophys. (Switzerland)* **90**:1 (1971), 38–52.
- [Dey and Addy 1978] S. Dey and S. K. Addy, “Love waves under initial stresses in a visco-elastic medium overlying an elastic half-space”, *Gerlands Beiträge Geophys.* **87**:4 (1978), 306–311.
- [Garvin 1956] W. W. Garvin, “Exact transient solution of the Buried line source problem”, *Proc. R. Soc. Lond. A* **234**:1199 (1956), 528–541.
- [Khurana 2001] P. Khurana, “Love wave propagation in a prestressed medium”, *Indian J. Pure Appl. Math.* **32**:8 (2001), 1201–1207.
- [Majhi et al. 2016] S. Majhi, P. C. Pal, and S. Kumar, “Love waves in a layered functionally graded piezoelectric structure under initial stress”, *Waves Random Complex Media* **26** (2016), 535–552.
- [Majhi et al. 2017] S. Majhi, P. C. Pal, and S. Kumar, “Reflection and transmission of plane SH-waves in an initially stressed inhomogeneous anisotropic magnetoelastic medium”, *J. Seismolog.* **21** (2017), 155–163.
- [Mandal et al. 2014] P. C. Pal, D. Mandal, and S. Kumar, “Disturbance of SH-type waves due to moving stress discontinuity in an anisotropic soil layer overlying an inhomogeneous elastic half-space”, *Sadhana* **39**:2 (2014), 451–465.
- [Pal 1983] P. C. Pal, “Generation of SH-type waves due to non-uniformly moving stress-discontinuity in layered anisotropic elastic half-space”, *Acta Mech.* **49**:3-4 (1983), 209–220.
- [Pal and Kumar 2000] P. C. Pal and L. Kumar, “Generation of SH waves by a moving stress discontinuity in an anisotropic soil layer over an elastic half-space”, *Acta Geophys.* **48**:4 (2000), 465–478.
- [Pal and Mandal 2014] P. C. Pal and D. Mandal, “Generation of SH-type waves due to shearing stress discontinuity in a sandy layer overlying an isotropic and inhomogeneous elastic half-space”, *Acta Geophys.* **62**:1 (2014), 44–58.

Received 21 Oct 2017. Revised 9 Feb 2018. Accepted 26 Apr 2018.

SANTOSH KUMAR: santosh453@gmail.com

Department of Mathematics, VVIT, Vidya Vihar Institute of Technology, Purnea 854303, India

DINBANDHU MANDAL: dinbandhumandal@gmail.com

Department of Mathematics, Chaibasa Engineering College, West Singhbhum 833215, India



MATHEMATICS AND MECHANICS OF COMPLEX SYSTEMS

msp.org/memocs

EDITORIAL BOARD

ANTONIO CARCATERRA	Università di Roma "La Sapienza", Italia
ERIC A. CARLEN	Rutgers University, USA
FRANCESCO DELL'ISOLA	(CO-CHAIR) Università di Roma "La Sapienza", Italia
RAFFAELE ESPOSITO	(TREASURER) Università dell'Aquila, Italia
ALBERT FANNJIANG	University of California at Davis, USA
GILLES A. FRANCFORT	(CO-CHAIR) Université Paris-Nord, France
PIERANGELO MARCATI	Università dell'Aquila, Italy
JEAN-JACQUES MARIGO	École Polytechnique, France
PETER A. MARKOWICH	DAMTP Cambridge, UK, and University of Vienna, Austria
MARTIN OSTOJA-STARZEWSKI	(CHAIR MANAGING EDITOR) Univ. of Illinois at Urbana-Champaign, USA
PIERRE SEPPECHER	Université du Sud Toulon-Var, France
DAVID J. STEIGMANN	University of California at Berkeley, USA
PAUL STEINMANN	Universität Erlangen-Nürnberg, Germany
PIERRE M. SUQUET	LMA CNRS Marseille, France

MANAGING EDITORS

MICOL AMAR	Università di Roma "La Sapienza", Italia
CORRADO LATTANZIO	Università dell'Aquila, Italy
ANGELA MADEO	Université de Lyon-INSA (Institut National des Sciences Appliquées), France
MARTIN OSTOJA-STARZEWSKI	(CHAIR MANAGING EDITOR) Univ. of Illinois at Urbana-Champaign, USA

ADVISORY BOARD

ADNAN AKAY	Carnegie Mellon University, USA, and Bilkent University, Turkey
HOLM ALTENBACH	Otto-von-Guericke-Universität Magdeburg, Germany
MICOL AMAR	Università di Roma "La Sapienza", Italia
HARM ASKES	University of Sheffield, UK
TEODOR ATANACKOVIĆ	University of Novi Sad, Serbia
VICTOR BERDICHEVSKY	Wayne State University, USA
GUY BOUCHITTÉ	Université du Sud Toulon-Var, France
ANDREA BRAIDES	Università di Roma Tor Vergata, Italia
ROBERTO CAMASSA	University of North Carolina at Chapel Hill, USA
MAURO CARFORE	Università di Pavia, Italia
ERIC DARVE	Stanford University, USA
FELIX DARVE	Institut Polytechnique de Grenoble, France
ANNA DE MASI	Università dell'Aquila, Italia
GIANPIETRO DEL PIERO	Università di Ferrara and International Research Center MEMOCS, Italia
EMMANUELE DI BENEDETTO	Vanderbilt University, USA
BERNOLD FIEDLER	Freie Universität Berlin, Germany
IRENE M. GAMBA	University of Texas at Austin, USA
DAVID Y. GAO	Federation University and Australian National University, Australia
SERGEY GAVRILYUK	Université Aix-Marseille, France
TIMOTHY J. HEALEY	Cornell University, USA
DOMINIQUE JEULIN	École des Mines, France
ROGER E. KHAYAT	University of Western Ontario, Canada
CORRADO LATTANZIO	Università dell'Aquila, Italy
ROBERT P. LIPTON	Louisiana State University, USA
ANGELO LUONGO	Università dell'Aquila, Italia
ANGELA MADEO	Université de Lyon-INSA (Institut National des Sciences Appliquées), France
JUAN J. MANFREDI	University of Pittsburgh, USA
CARLO MARCHIORO	Università di Roma "La Sapienza", Italia
GÉRARD A. MAUGIN	Université Paris VI, France
ROBERTO NATALINI	Istituto per le Applicazioni del Calcolo "M. Picone", Italy
PATRIZIO NEFF	Universität Duisburg-Essen, Germany
ANDREY PIATNITSKI	Narvik University College, Norway, Russia
ERRICO PRESUTTI	Università di Roma Tor Vergata, Italy
MARIO PULVIRENTI	Università di Roma "La Sapienza", Italia
LUCIO RUSSO	Università di Roma "Tor Vergata", Italia
MIGUEL A. F. SANJUAN	Universidad Rey Juan Carlos, Madrid, Spain
PATRICK SELVADURAI	McGill University, Canada
ALEXANDER P. SEYRANIAN	Moscow State Lomonosov University, Russia
MIROSLAV ŠILHAVÝ	Academy of Sciences of the Czech Republic
GUIDO SWEERS	Universität zu Köln, Germany
ANTOINETTE TORDSILLAS	University of Melbourne, Australia
LEV TRUSKINOVSKY	École Polytechnique, France
JUAN J. L. VELÁZQUEZ	Bonn University, Germany
VINCENZO VESPRI	Università di Firenze, Italia
ANGELO VULPIANI	Università di Roma La Sapienza, Italia

MEMOCS (ISSN 2325-3444 electronic, 2326-7186 printed) is a journal of the International Research Center for the Mathematics and Mechanics of Complex Systems at the Università dell'Aquila, Italy.

Cover image: "Tangle" by © John Horigan; produced using the *Context Free* program (contextfreeart.org).

PUBLISHED BY



mathematical sciences publishers
nonprofit scientific publishing

<http://msp.org/>

© 2018 Mathematical Sciences Publishers

The variational structure of classical plasticity Gianpietro Del Piero	137
Far-reaching Hellenistic geographical knowledge hidden in Ptolemy's data Lucio Russo	181
Generation of SH-type waves due to shearing stress discontinuity in an anisotropic layer overlying an initially stressed elastic half-space Santosh Kumar and Dinbandhu Mandal	201
Strain gradient and generalized continua obtained by homogenizing frame lattices Houssam Abdoul-Anziz and Pierre Seppecher	213
On the effect of phase transition on the manifold dimensionality: application to the Ising model Elena Lopez, Adrien Scheuer, Emmanuelle Abisset-Chavanne and Francisco Chinesta	251

MEMOCS is a journal of the International Research Center for
the Mathematics and Mechanics of Complex Systems
at the Università dell'Aquila, Italy.

