

# Moscow Journal of Combinatorics and Number Theory

2019

vol. 8 no. 2

A family of four-variable expanders with quadratic growth

Mehdi Makhul



# A family of four-variable expanders with quadratic growth

Mehdi Makhul

We prove that if  $g(x, y)$  is a polynomial of degree  $d$  that is not a polynomial of only  $y$ , then for any finite set  $A \subset \mathbb{R}$

$$|X| \gg_d |A|^2, \quad \text{where } X := \left\{ \frac{g(a_1, b_1) - g(a_2, b_2)}{b_2 - b_1} : a_1, a_2, b_1, b_2 \in A \right\}.$$

We will see this bound is also tight for some polynomial  $g(x, y)$ .

## 1. Introduction

Throughout this paper, when we write  $X \gg Y$ , this means that  $X \geq cY$  for some absolute constant  $c > 0$ .

The sum set of a subset  $A \subset \mathbb{R}$  is defined as  $A + A := \{a + b : a, b \in A\}$ . The product set is defined in a similar way,  $AA := \{ab : a, b \in A\}$ .

The Erdős–Szemerédi conjecture [1983] states that, for all  $\epsilon > 0$  and for any finite set  $A \subset \mathbb{N}$ ,

$$\max\{|A + A|, |AA|\} \geq c(\epsilon)|A|^{2-\epsilon}.$$

It is natural to extend this conjecture to other settings (such as  $\mathbb{R}$ ), and also to change the polynomials  $F(x, y) = x + y$  and  $F(x, y) = xy$  defining the sum and product sets to other polynomials or rational functions. In recent years much research has been done in this direction.

For many such functions, the images of sets are known to always grow. For example, the authors of [Murphy et al. 2015] have studied several multivariable polynomials, including the function

$$G(x_1, x_2, x_3, x_4, x_5) = x_1(x_2 + x_3 + x_4 + x_5).$$

More precisely they showed that, for any finite set  $A \subset \mathbb{R}$ ,

$$|A(A + A + A + A)| \gg \frac{|A|^2}{\log |A|},$$

where  $A(A + A + A + A) := \{x_1(x_2 + x_3 + x_4 + x_5) : x_i \in A\}$ .

In [Murphy et al. 2017], the authors studied a more complicated function, namely

$$H(x_1, x_2, x_3, x_4, x_5) = (x_1 + x_2 + x_3 + x_4)^2 + \log x_5.$$

Makhul was supported by the Austrian Science Fund (FWF): W1214-N15, Project DK9.

MSC2010: primary 11B30; secondary 11B75.

Keywords: Bisector, expander functions.

They showed that, for any finite  $A \subset \mathbb{R}$ ,

$$|\{(a_1 + a_2 + a_3 + a_4)^2 + \log a_5 : a_i \in A\}| \gg \frac{|A|^2}{\log |A|}.$$

In the same circle of ideas, Balog and Roche-Newton [2015] investigated the rational function

$$F(x_1, x_2, x_3, x_4) = \frac{x_1 + x_2}{x_3 + x_4},$$

showing that for any finite set  $A \subset \mathbb{R}$ , we have

$$|F(A, A, A, A)| \geq 2|A|^2 - 1.$$

Our result is a generalization of the method of [Murphy et al. 2015, Corollary 3.1], where they used the Szemerédi–Trotter theorem to prove that for any finite set  $A \subset \mathbb{R}$

$$\left| \frac{A - A}{A - A} \right| \gg |A|^2.$$

A stronger version of this result, with a multiplicative constant 1, follows from an earlier geometric result of Ungar [1982].

In this article we consider a certain class of rational functions of four variables. Suppose that  $g(x, y)$  is a polynomial of two variables of degree  $d$ . Let

$$F(x_1, x_2, y_1, y_2) = \frac{g(x_1, y_1) - g(x_2, y_2)}{y_2 - y_1}$$

be a four-variable rational function in terms of  $x_1, x_2, y_1, y_2$ . The main theorem of this paper is the following result concerning the growth of  $F$ .

**Theorem 1.1.** *Suppose that  $g(x, y)$  is a polynomial of degree  $d$ , that it is not a polynomial of only  $y$ , and that  $A \subset \mathbb{R}$  is a finite set. Then*

$$|X| \gg_d |A|^2, \quad \text{where } X := \left\{ \frac{g(a_1, b_1) - g(a_2, b_2)}{b_2 - b_1} : a_1, a_2, b_1, b_2 \in A \right\}.$$

Notice that the following example shows that the condition that  $g(x, y)$  cannot be a polynomial of only  $y$  is necessary.

**Example 1.2.** Suppose that  $g(x, y) = y^2$  and  $A = \{1, 2, \dots, n\}$ . Then

$$X = \left\{ \frac{b_1^2 - b_2^2}{b_2 - b_1} : b_1, b_2 \in A \right\}$$

equals  $-\{b_2 + b_1 : b_i \in A\}$  and has cardinality  $O(n)$ .

On the other hand, it is known that for some polynomials  $g$ , the result of Theorem 1.1 is tight. For example, if we define  $g(x_1, y_1) = x_1$  then Theorem 1.1 recovers the result of [Murphy et al. 2015; Ungar

1982]. This is known to be tight, since for the set  $A = \{1, \dots, N\}$ ,

$$\left| \frac{A - A}{A - A} \right| = O(N^2).$$

However, we are not aware of any other polynomials  $g$  for which the bound in [Theorem 1.1](#) is tight, and whether or not the bound can be improved for some particular  $g$  is an interesting question.

Our main result has some similarities with a result of Raz, Sharir and Solymosi [[Raz et al. 2015](#)] concerning the growth of two-variable polynomials. Their result states that, if  $F$  is a two-variable polynomial with bounded degree, then for any  $A, B \subset \mathbb{R}$  with  $|A| = |B| = n$ ,

$$|F(A, B)| \gg_d n^{4/3},$$

provided that  $F$  satisfies a nondegeneracy condition. This condition states that  $F$  cannot be of one of the following forms:

- (1)  $F(u, v) = f(g(u) + h(v))$ .
- (2)  $F(u, v) = f(g(u) \cdot h(v))$ .

This result gave an improvement upon an earlier result of Elekes and Ronyai [[2000](#)].

**The Szemerédi–Trotter theorem.** The essential ingredient used to prove our result is a corollary of the Szemerédi–Trotter theorem [[1983](#)], which gives a bound for the number of lines in the plane containing at least a fixed number of points  $k$  from a given finite set, that is, the number of  $k$ -rich lines.

**Theorem 1.3.** *Let  $P$  be a finite set of points and let  $L$  be a finite set of lines. Then the number of incidences  $I(P, L) := \{(p, \ell) \in P \times L : p \in \ell\}$  has the upper bound*

$$I(P, L) \ll |P|^{2/3} |L|^{2/3} + |P| + |L|.$$

More precisely,

$$I(P, L) \leq 4|P|^{2/3} |L|^{2/3} + 4|P| + |L|.$$

If each line in  $L$  appears at most  $d$  times for some constant  $d$ , then a generalization of the Szemerédi–Trotter theorem states that

$$I(P, L) \leq 4d|P|^{2/3} |L|^{2/3} + 4d|P| + d|L|.$$

The main idea of the following corollary is known in literature; we present here a slightly improved version which we could not find in the literature in the form we need.

**Corollary 1.4.** *Let  $k, n \geq 2$  be natural numbers and fix  $d \in \mathbb{N}$  such that  $4d + 1 \leq k \leq d\sqrt{n}$ . Let  $\mathcal{L}$  be a set of  $n$  lines in the plane, and let  $t_{\geq k}$  denote the number of points in the plane contained in at least  $k$  lines of  $\mathcal{L}$ , where each line appears with multiplicity at most  $d$ . Then*

$$t_{\geq k} = O_d\left(\frac{n^2}{k^3}\right).$$

Notice that if  $\mathcal{L}$  is a set of  $n$  lines in the plane such that each line appears at most  $d$  times for some constant  $d$ , then for computing  $t_k$ ,  $k$  should be greater than or equal to  $4d + 1$ . To see this, suppose that

$P_k$  is the set of  $k$ -rich points. Then we have  $k|P_k| \leq 4d|P_k|^{2/3}|L|^{2/3} + 4d|P_k| + d|L|$ . This implies

$$(k - 4d)|P_k| \leq 4d|P_k|^{2/3}|L|^{2/3} + d|L|.$$

Hence we may assume  $k \geq 4d + 1$ , otherwise the inequality gives nothing.

*Proof of Corollary 1.4.* Let  $P_k$  be the set of  $k$ -rich points. Since each line appears at most  $d$  times we have

$$\frac{k|P_k|}{d} \ll |P_k|^{2/3}|L|^{2/3} + |L|,$$

so  $k^3|P_k| \ll d^3|L|^2$  or otherwise  $|P_k| \ll d|L|/k$ . Plugging these bounds back into the Szemerédi–Trotter theorem gives

$$I(P_k, L) \ll |L|^{2/3} \left( \frac{d^3|L|^2}{k^3} \right)^{2/3} + |L|^{2/3} \left( \frac{d|L|}{k} \right)^{2/3} + |L| + \frac{d^3|L|^2}{k^3} + \frac{d|L|}{k}.$$

Since  $k > 4d$  we can ignore last two summands and we obtain

$$I(P_k, L) \ll \frac{d^2|L|^2}{k^2} + \left( \frac{d}{k} \right)^{2/3} |L|^{4/3} + |L|.$$

Note that we have

$$\frac{d^2|L|^2}{k^2} \geq \left( \frac{d}{k} \right)^{2/3} |L|^{4/3}$$

if  $k \leq d\sqrt{n}$ . □

## 2. Main results

Suppose that  $A, B \subset \mathbb{R}$  are finite, and  $g(x_1, y_1)$  is a polynomial of degree  $d$ . We associate an element of  $A \times B$  with a line via

$$A \times B \ni (a, b) \longleftrightarrow l_{a,b} : y = bx - g(a, b).$$

Consider  $\mathcal{L} = \{l_{a,b} : a, b \in A \times B\}$  as a multiset. Then  $\mathcal{L}$  is a set of  $|A||B|$  lines, such that each line appears at most  $d$  times. We also define the quantity

$$n(x, y) = |\{(a, b) \in A \times B : (x, y) \in l_{a,b}\}|,$$

which is interpreted geometrically as the number of lines of  $\mathcal{L}$  that pass through  $(x, y)$ .

**Lemma 2.1.** *Suppose that  $d \in \mathbb{N}$  is fixed. Suppose that  $A, B, X \subset \mathbb{R}$  are finite and satisfy*

$$|X| \leq \frac{|A||B|}{4d^2},$$

with  $0 \notin X$ . Then

$$\sum_{x \in X} \sum_y n^2(x, y) \ll |A|^{3/2} |B|^{3/2} |X|^{1/2}. \quad (1)$$

*Proof.* The set of  $t$ -rich points is given by

$$R_t := \{(x, y) \in \mathbb{R}^2 : n(x, y) \geq t\}.$$

We first show that

$$|R_t| \ll \frac{|A|^2|B|^2}{t^3}.$$

We begin by bounding  $n(x, y)$  for a given point  $(x, y)$ . For fixed  $b_0 \in B$  we obtain a line with slope  $b_0$  passing through  $(x, y)$  and a one-variable polynomial equation  $g(a, b_0)$ . Since each line is determined uniquely, by its slope and one point on it (for fixed  $b_0$  and  $(x, y)$  the equation  $g(a, b_0) = 0$  has at most  $d$  distinct solutions), we have

$$n(x, y) \leq d|B|.$$

With a similar argument for fixed  $a \in A$  we obtain a univariate polynomial equation. Since each line is determined uniquely by its  $y$ -intercept and one point on it we have

$$n(x, y) \leq d|A|.$$

These together imply

$$n(x, y) \leq d(\min\{|A|, |B|\}) \leq (d|A|d|B|)^{1/2} = d|\mathcal{L}|^{1/2}.$$

This implies there are no points incident to more than  $d\sqrt{|\mathcal{L}|}$  lines in  $\mathcal{L}$ , and by applying [Corollary 1.4](#) we get

$$|R_t| \ll \frac{|\mathcal{L}|^2}{t^3} \leq \frac{|A|^2|B|^2}{t^3}.$$

Let  $\Delta > 2d$  be an integer to be specified later. We have

$$\sum_{x \in X} \sum_y n^2(x, y) \leq \sum_{x \in X} \sum_{n(x, y) \leq \Delta} n^2(x, y) + \sum_{\substack{(x, y) \\ n(x, y) > \Delta}} n^2(x, y). \quad (2)$$

The first term is bounded by  $\Delta|A||B||X|$ ; in fact

$$\sum_{x \in X} \sum_{n(x, y) \leq \Delta} n^2(x, y) \leq \Delta \sum_{x \in X} \sum_y n(x, y) = \Delta|A||B| \sum_{x \in X} 1 = \Delta|A||B||X|. \quad (3)$$

For the second term we have

$$\begin{aligned} \sum_{\substack{(x, y) \\ n(x, y) > \Delta}} n^2(x, y) &= \sum_{j \geq 1} \sum_{2^{j-1}\Delta \leq n(x, y) \leq 2^j\Delta} n^2(x, y) \\ &\ll \sum_{j \geq 1} \frac{|A|^2|B|^2}{(2^j\Delta)^3} \cdot (2^j\Delta)^2 = \frac{|A|^2|B|^2}{\Delta} \sum_{j \geq 1} \frac{1}{2^j} = \frac{|A|^2|B|^2}{\Delta}. \end{aligned} \quad (4)$$

For an optimal choice, set

$$\Delta = \left\lceil \frac{(|A||B|)^{1/2}}{|X|^{1/2}} \right\rceil > 2d.$$

Combining the bounds from (2) and (3) and (4), it follows that

$$\sum_x \sum_y n^2(x, y) \ll |A|^{3/2}|B|^{3/2}|X|^{1/2}.$$

□



*Proof of Theorem 1.1.* Consider

$$\begin{aligned} |Y| &= \left| \left\{ (x, a_1, a_2, b_1, b_2) : x = \frac{g(a_1, b_1) - g(a_2, b_2)}{b_1 - b_2}, a_i, b_i \in A \right\} \right| \\ &= \left| \left\{ (x, a_1, a_2, b_1, b_2) : b_1 x - g(a_1, b_1) = b_2 x - g(a_2, b_2) \right\} \right| \\ &= \sum_{x \in X} \sum_y n^2(x, y) \ll |A|^3 |X|^{1/2}. \end{aligned}$$

On the other hand,  $|Y| \geq |A|^4$ . Thus we obtain

$$|A|^4 \ll |A|^3 |X|^{1/2}, \quad \text{and hence} \quad |X| \gg |A|^2. \quad \square$$

Notice that the proof of Theorem 1.1 fails when  $g(x, y)$  is a polynomial of only  $y$ . In fact if  $g(x, y) = h(y)$  for some polynomial  $h$ , then  $\mathcal{L} = \{l_{a,b} : a, b \in A \times B\}$  is a set of  $|A||B|$  lines such that each line appears at least  $|A|$  times (and at most  $d|A|$  times). On the other hand the generalization of the Szemerédi–Trotter theorem and its corollary hold when each line appears at most  $d$  times, where  $d$  is independent of  $|P|$  and  $|L|$  in Theorem 1.3.

**Corollary 2.2.** *Suppose that  $P = A \times A$  is a set of  $|A|^2$  points. Let  $l$  be the  $y$ -axis. Suppose that  $B(P)$  is the set of all bisectors determined by  $P$ . Then  $|B \cap l| \gg |A|^2$ .*

*Proof.* By a simple calculation we can see that the equation of the bisector determined by two points  $(x_1, y_1)$  and  $(x_2, y_2)$  in the  $(s, t)$ -plane is

$$s = \frac{2(x_1 - x_2)t + (x_2^2 - x_1^2) + (y_2^2 - y_1^2)}{2(y_2 - y_1)}.$$

Inserting  $t = 0$ , the hitting point has coordinate

$$\left( 0, \frac{(x_2^2 - x_1^2) + (y_2^2 - y_1^2)}{2(y_2 - y_1)} \right).$$

Setting  $g(x, y) = -\frac{1}{2}(x^2 + y^2)$ , we obtain the result by Theorem 1.1.  $\square$

As we mentioned, this bound is tight for some polynomials, for instance  $g(x, y) = x$ . However, we expect that if  $F(x_1, x_2, y_1, y_2)$  is a generic rational function satisfying the condition of Theorem 1.1 we have  $|X| \gg |A|^3$ .

## Acknowledgements

I would like to thank Oliver Roche-Newton for bringing this problem to my attention and for several helpful conversations.

## References

- [Balog and Roche-Newton 2015] A. Balog and O. Roche-Newton, “New sum-product estimates for real and complex numbers”, *Discrete Comput. Geom.* **53**:4 (2015), 825–846. [MR](#) [Zbl](#)
- [Elekes and Rónyai 2000] G. Elekes and L. Rónyai, “A combinatorial problem on polynomials and rational functions”, *J. Combin. Theory Ser. A* **89**:1 (2000), 1–20. [MR](#) [Zbl](#)

- [Erdős and Szemerédi 1983] P. Erdős and E. Szemerédi, “On sums and products of integers”, pp. 213–218 in *Studies in pure mathematics*, edited by P. Erdős, Birkhäuser, Basel, 1983. [MR](#) [Zbl](#)
- [Murphy et al. 2015] B. Murphy, O. Roche-Newton, and I. Shkredov, “Variations on the sum-product problem”, *SIAM J. Discrete Math.* **29**:1 (2015), 514–540. [MR](#) [Zbl](#)
- [Murphy et al. 2017] B. Murphy, O. Roche-Newton, and I. D. Shkredov, “Variations on the sum-product problem, II”, *SIAM J. Discrete Math.* **31**:3 (2017), 1878–1894. [MR](#) [Zbl](#)
- [Raz et al. 2015] O. E. Raz, M. Sharir, and J. Solymosi, “On triple intersections of three families of unit circles”, *Discrete Comput. Geom.* **54**:4 (2015), 930–953. [MR](#) [Zbl](#)
- [Szemerédi and Trotter 1983] E. Szemerédi and W. T. Trotter, Jr., “Extremal problems in discrete geometry”, *Combinatorica* **3**:3-4 (1983), 381–392. [MR](#) [Zbl](#)
- [Ungar 1982] P. Ungar, “ $2N$  noncollinear points determine at least  $2N$  directions”, *J. Combin. Theory Ser. A* **33**:3 (1982), 343–347. [MR](#) [Zbl](#)

Received 11 May 2018. Revised 17 Jul 2018.

MEHDI MAKHUL:

[mmakhul@risc.jku.at](mailto:mmakhul@risc.jku.at)

Johann Radon Institute for Computational and Applied Mathematics (RICAM), Austrian Academy of Sciences, Linz and  
Research Institute for Symbolic Computation (RISC), Johannes Kepler University, Linz, Austria



# Moscow Journal of Combinatorics and Number Theory

[msp.org/moscow](http://msp.org/moscow)

## EDITORS-IN-CHIEF

- Yann Bugeaud    Université de Strasbourg (France)  
[bugeaud@math.unistra.fr](mailto:bugeaud@math.unistra.fr)
- Nikolay Moshchevitin    Lomonosov Moscow State University (Russia)  
[moshchevitin@gmail.com](mailto:moshchevitin@gmail.com)
- Andrei Raigorodskii    Moscow Institute of Physics and Technology (Russia)  
[mraigor@yandex.ru](mailto:mraigor@yandex.ru)
- Ilya D. Shkredov    Steklov Mathematical Institute (Russia)  
[ilya.shkredov@gmail.com](mailto:ilya.shkredov@gmail.com)

## EDITORIAL BOARD

- Iskander Aliev    Cardiff University (United Kingdom)
- Vladimir Dolnikov    Moscow Institute of Physics and Technology (Russia)
- Nikolay Dolbilin    Steklov Mathematical Institute (Russia)
- Oleg German    Moscow Lomonosov State University (Russia)
- Michael Hoffman    United States Naval Academy
- Grigory Kabatiansky    Russian Academy of Sciences (Russia)
- Roman Karasev    Moscow Institute of Physics and Technology (Russia)
- Gyula O. H. Katona    Hungarian Academy of Sciences (Hungary)
- Alex V. Kontorovich    Rutgers University (United States)
- Maxim Korolev    Steklov Mathematical Institute (Russia)
- Christian Krattenthaler    Universität Wien (Austria)
- Antanas Laurinćikas    Vilnius University (Lithuania)
- Vsevolod Lev    University of Haifa at Oranim (Israel)
- János Pach    EPFL Lausanne (Switzerland) and Rényi Institute (Hungary)
- Rom Pinchasi    Israel Institute of Technology – Technion (Israel)
- Alexander Razborov    Institut de Mathématiques de Luminy (France)
- Joël Rivat    Université d'Aix-Marseille (France)
- Tanguy Rivoal    Institut Fourier, CNRS (France)
- Damien Roy    University of Ottawa (Canada)
- Vladislav Salikhov    Bryansk State Technical University (Russia)
- Tom Sanders    University of Oxford (United Kingdom)
- Alexander A. Sapozhenko    Lomonosov Moscow State University (Russia)
- József Solymosi    University of British Columbia (Canada)
- Andreas Strömbergsson    Uppsala University (Sweden)
- Benjamin Sudakov    University of California, Los Angeles (United States)
- Jörg Thuswaldner    University of Leoben (Austria)
- Kai-Man Tsang    Hong Kong University (China)
- Maryna Viazovska    EPFL Lausanne (Switzerland)
- Barak Weiss    Tel Aviv University (Israel)

## PRODUCTION

- Silvio Levy    (Scientific Editor)  
[production@msp.org](mailto:production@msp.org)

Cover design: Blake Knoll, Alex Scorpan and Silvio Levy

See inside back cover or [msp.org/moscow](http://msp.org/moscow) for submission instructions.

The subscription price for 2019 is US \$310/year for the electronic version, and \$365/year (+\$20, if shipping outside the US) for print and electronic. Subscriptions, requests for back issues and changes of subscriber address should be sent to MSP.

Moscow Journal of Combinatorics and Number Theory (ISSN 2640-7361 electronic, 2220-5438 printed) at Mathematical Sciences Publishers, 798 Evans Hall #3840, c/o University of California, Berkeley, CA 94720-3840 is published continuously online. Periodical rate postage paid at Berkeley, CA 94704, and additional mailing offices.

MJCNT peer review and production are managed by EditFlow® from MSP.

PUBLISHED BY

 **mathematical sciences publishers**  
nonprofit scientific publishing  
<http://msp.org/>

© 2019 Mathematical Sciences Publishers

---

A simple proof of the Hilton–Milner theorem	97
PETER FRANKL	
On the quotient set of the distance set	103
ALEX IOSEVICH, DOOWON KOH and HANS PARSHALL	
Embeddings of weighted graphs in Erdős-type settings	117
DAVID M. SOUKUP	
Identity involving symmetric sums of regularized multiple zeta-star values	125
TOMOYA MACHIDE	
Matiyasevich-type identities for hypergeometric Bernoulli polynomials and poly-Bernoulli polynomials	137
KEN KAMANO	
A family of four-variable expanders with quadratic growth	143
MEHDI MAKHUL	
The Lind–Lehmer Constant for $\mathbb{Z}_2^r \times \mathbb{Z}_4^s$	151
MICHAEL J. MOSSINGHOFF, VINCENT PIGNO and CHRISTOPHER PINNER	
Lattices with exponentially large kissing numbers	163
SERGE VLĂDUȚ	
A note on the set $A(A + A)$	179
PIERRE-YVES BIENVENU, FRANÇOIS HENNECART and ILYA SHKREDOV	
On a theorem of Hildebrand	189
CARSTEN DIETZEL	