

# Moscow Journal of Combinatorics and Number Theory

2020  
vol. 9 no. 3

**Weakly distinguishing graph polynomials on addable properties**

Johann A. Makowsky and Vsevolod Rakita





# Weakly distinguishing graph polynomials on addable properties

Johann A. Makowsky and Vsevolod Rakita

A graph polynomial  $P$  is weakly distinguishing if for almost all finite graphs  $G$  there is a finite graph  $H$  that is not isomorphic to  $G$  with  $P(G) = P(H)$ . It is weakly distinguishing on a graph property  $\mathcal{C}$  if for almost all finite graphs  $G \in \mathcal{C}$  there is  $H \in \mathcal{C}$  that is not isomorphic to  $G$  with  $P(G) = P(H)$ . We give sufficient conditions on a graph property  $\mathcal{C}$  for the characteristic, clique, independence, matching, and domination and  $\xi$  polynomials, as well as the Tutte polynomial and its specializations, to be weakly distinguishing on  $\mathcal{C}$ . One such condition is to be addable and small in the sense of C. McDiarmid, A. Steger and D. Welsh (2005). Another one is to be of genus at most  $k$ .

1. Introduction and preliminaries	333
2. Comparing graph polynomials	335
3. Addable properties and graphs of genus $k$	337
4. The characteristic polynomial	339
5. The domination polynomial	341
6. The $\xi$ polynomial	342
7. $\xi$ -invariants	344
8. Conclusion	347
Acknowledgements	348
References	348

## 1. Introduction and preliminaries

Unless otherwise stated, we only consider simple (finite, loopless, undirected graphs with no parallel edges) graphs with vertices labeled  $1, \dots, n$ . For a graph  $G = (V(G), E(G))$  and  $A \subseteq E(G)$ , we denote by  $G\langle A \rangle$  the graph  $(V(G), A)$ , and by  $k(A)$  the number of connected components of  $G\langle A \rangle$ . A graph property is a family of graphs that is closed under isomorphisms. For a graph property  $\mathcal{C}$ , denote by  $\mathcal{C}(n)$  the graphs of order  $n$  in  $\mathcal{C}$ . We only consider properties such that  $\mathcal{C}(n)$  is nonempty for all sufficiently large  $n$ .

Let  $P$  be a graph polynomial. We say that two non isomorphic graphs  $G$  and  $H$  are  $P$ -mates if  $P(G) = P(H)$ , and that  $G$  is  $P$  unique if it has no  $P$  mates.  $P$  is *trivial* if all graphs  $G, H$  are  $P$ -mates.  $P$  is *complete* if all graphs  $G$  are  $P$ -unique.

In this paper we investigate conditions which imply that almost all graphs in a graph property  $\mathcal{C}$  have a  $P$ -mate. More formally, we give the following definitions:

---

V. Rakita was partially supported by a Jacobs Scholarship for excellent students.

MSC2010: primary 05C31; secondary 05C10, 05C30, 05C69, 05C80.

Keywords: graph polynomials, random graphs, Bollobás–Pebody–Riordan conjecture, addable graph classes.

Let  $P$  be a graph polynomial, and denote by  $\mathcal{G}(n)$  the family of graphs of order  $n$  with vertices labeled  $1, \dots, n$ , and by  $U_P(n)$  the set of  $P$  unique graphs of order  $n$ .

**Definition 1.1.**  $P$  is *weakly distinguishing* if

$$\lim_{n \rightarrow \infty} \frac{|U_P(n)|}{|\mathcal{G}(n)|} = 0$$

and  $P$  is *almost complete* if

$$\lim_{n \rightarrow \infty} \frac{|U_P(n)|}{|\mathcal{G}(n)|} = 1$$

In [Bollobás et al. 2000], Bollobás, Pebody and Riordan conjectured:

**Conjecture 1** (BPR conjecture). The chromatic and Tutte polynomials are almost complete.

In [Makowsky and Zhang 2019], the analogous question for  $r$ -regular hypergraphs was considered, and for  $r \geq 3$  the conjecture was refuted. In [Noy 2003] it was observed, as a remark in the conclusions, that the independence polynomial  $\text{In}(G; x)$ , discussed in Section 7, is weakly distinguishing on all finite graphs. In [Makowsky and Rakita 2019], it was proven that an infinite number of graph polynomials, among them the independence, clique and harmonious polynomials, are weakly distinguishing.

A natural way to approach the question whether a graph polynomial  $P$  is weakly distinguishing, almost complete or otherwise, is to ask, given a graph property  $\mathcal{C}$ , whether almost all graphs in  $\mathcal{C}$  are  $P$  unique in  $\mathcal{C}$ . Let  $\mathcal{C}$  be a graph property and denote by  $U_{P,\mathcal{C}}(n) = \{G \in \mathcal{C}(n) : G \text{ has no } P\text{-mate in } \mathcal{C}\}$ .

**Definition 1.2.**  $P$  is *weakly distinguishing on  $\mathcal{C}$*  if

$$\lim_{n \rightarrow \infty} \frac{|U_{P,\mathcal{C}}(n)|}{|\mathcal{C}(n)|} = 0,$$

and  $P$  is *almost complete on  $\mathcal{C}$*  if

$$\lim_{n \rightarrow \infty} \frac{|U_{P,\mathcal{C}}(n)|}{|\mathcal{C}(n)|} = 1.$$

In this paper we prove that many well studied graph polynomials are weakly distinguishing on infinitely many graph properties  $\mathcal{C}$ , listed in Example 3.2. However, all these properties  $\mathcal{C}$  are small (in the sense that they are sets of measure 0 in the collection of all graphs), so these results do not imply that any of the above polynomials are weakly distinguishing or almost complete for arbitrary properties  $\mathcal{C}$ .

The graph polynomials which we show are weakly distinguishing for  $\mathcal{C}$  include the characteristic polynomial, the domination polynomial, and the  $\xi$ -polynomial, which is a generalization of the both the matching and the Tutte polynomial. These three polynomials are mutually incomparable in distinctive power by Proposition 2.3. Once we know that  $\xi$  is weakly distinguishing for a graph property  $\mathcal{C}$ , this holds also for all the graph polynomials which are substitution instances of  $\xi$  in  $\mathcal{C}$ . This includes the matching polynomials, the independence polynomial, the chromatic polynomial, the Tutte polynomial, the Euler polynomial, and many others; see Section 7.

This paper is organized as follows: In Section 2 we introduce a method to compare distinctive power of graph polynomials and formulate how to use it (Lemma A). In Section 3, we give a graph theoretic background on addable properties, and graphs of genus at most  $k$ , and formulate our main tools, Lemma B and Lemma C. In Sections 4 and 5 we prove our results: Theorems 4.4 and 4.7 for the characteristic

polynomial, Theorems 5.5 and 5.8 for the domination polynomial. In Section 6 we prove the same for the  $\xi$ -polynomial and in Section we apply Lemma A to derive the corresponding results for the matching polynomials, the independence polynomial, the chromatic polynomial, the Tutte polynomial, the Euler polynomial, and many others. In Section 8 we draw conclusions and present open problems.

## 2. Comparing graph polynomials

In this section we provide a tool (Lemma A) which allows us to show that many graph polynomials are weakly distinguishing.

**Definition 2.1.** Let  $\mathcal{C}$  be a graph property,  $P$  and  $Q$  be two graph polynomials and  $G$  and  $H$  two finite graphs.

- (i)  $G$  and  $H$  are *similar* if they have the same number of vertices, edges and connected components.
- (ii)  $P <_{d.p.}^{\mathcal{C}} Q$  or  $Q$  is at least as distinctive as  $P$  in  $\mathcal{C}$  if for all graphs  $G, H \in \mathcal{C}$ ,  $Q(G; \bar{x}) = Q(H; \bar{x})$  implies  $P(G; \bar{y}) = P(H; \bar{y})$ .
- (iii)  $P \sim_{d.p.}^{\mathcal{C}} Q$  or  $P$  and  $Q$  are of the same distinctive power in  $\mathcal{C}$  if  $P <_{d.p.}^{\mathcal{C}} Q$  and  $Q <_{d.p.}^{\mathcal{C}} P$ .
- (iv)  $P <_{s.d.p.}^{\mathcal{C}} Q$  or  $P$  and  $Q$  are of the same distinctive power in  $\mathcal{C}$  on similar graphs if for all similar graphs  $G, H \in \mathcal{C}$ ,  $Q(G; \bar{x}) = Q(H; \bar{x})$  implies  $P(G; \bar{y}) = P(H; \bar{y})$ .
- (v)  $P \sim_{s.d.p.}^{\mathcal{C}} Q$  if  $P <_{s.d.p.}^{\mathcal{C}} Q$  and  $Q <_{s.d.p.}^{\mathcal{C}} P$ .
- (vi) For all graph properties  $\mathcal{C}$   $P <_{d.p.}^{\mathcal{C}} Q$  implies  $P <_{s.d.p.}^{\mathcal{C}} Q$ .

If  $\mathcal{C}$  consists of all finite graphs, we omit it.

The partial preorders  $<_{d.p.}$  and  $<_{s.d.p.}$  between graph polynomials (or general graph invariants) are studied extensively in [Makowsky et al. 2019]. A complete (trivial) graph polynomial is a maximal (minimal) element with respect to  $<_{d.p.}$

**Example 2.2.** (i) The chromatic polynomial  $\chi(G; x)$  and the Tutte polynomial satisfy  $\chi(G; x) <_{s.d.p.} T(G; x, y)$  but not  $\chi(G; x) <_{d.p.} T(G; x, y)$ ; see Section 7, because  $T(G; x, y)$  does not determine the order of  $G$  in the presence of isolated vertices.

- (ii) The characteristic polynomial  $P_A(G; x)$  and the matching polynomial (defect aka acyclic)  $\mu(G; x)$  from Section 7 are  $d.p.$ -equivalent on forests; see [Godsil and Gutman 1981].
- (iii) Let  $\bar{G}$  be the (loopless) complement graph of a simple graph  $G$ , and  $P(G; \bar{x})$  be a (possibly multivariate) graph polynomial. Put  $\bar{P}(G; \bar{x}) = P(\bar{G}, \bar{x})$ . Then  $\bar{P}(G; \bar{x}) \sim_{d.p.} P(G, \bar{x})$ . If we relativize this to a graph property  $\mathcal{C}$ ,  $\bar{P}(G; \bar{x}) \sim_{d.p.}^{\mathcal{C}} P(G, \bar{x})$  holds provided  $\mathcal{C}$  is closed under complement graphs.

**Proposition 2.3.** The following graph polynomials are pairwise incomparable by  $<_{s.d.p.}$ :

- (i) The chromatic polynomial  $\chi(G; x)$  and the independence polynomial  $\text{In}(G; x)$  from Section 7.
- (ii) The characteristic polynomial  $P_A(G, x)$  from Sections 4 and the chromatic polynomial  $\chi(G; x)$ .
- (iii) The characteristic polynomial  $P_A(G; x)$  and the independence polynomial  $\text{In}(G; x)$ .
- (iv) The characteristic polynomial  $P_A(G, x)$  and the domination polynomial  $\text{Dom}(G; x)$ . from Sections 4 and 5.



**Figure 1.**  $P_A$ -mates.



**Figure 2.** A path of length 5.

(v) *The characteristic polynomial and the  $\xi$ -polynomial from Section 6.*

(vi) *The domination polynomial and the  $\xi$ -polynomial.*

*Proof.* The first three examples are from [Makowsky et al. 2019].

For (iv)-(vi) we first note that all graphs of order less than 8 and all trees of order less than 10 are  $\xi$ -unique; see [Trinks 2012].

(A) Consider the graphs  $C_4 \sqcup K_1$  and  $S_5$  in Figure 1. We have

$$P_A(S_5, x) = P_A(C_4 \sqcup K_1, x) = x^2(x+2)(x-2), \text{ Dom}(S_5, 1) = 1, \text{ Dom}(C_4 \sqcup K_1, 1) = 0,$$

and  $\xi(S_5, x) \neq \xi(C_4 \sqcup K_1, x)$ , since they are of order 5.

(B) Let  $P_5$  and  $\hat{P}_5$  be the graphs of order 5 shown in Figure 2. Then  $\text{Dom}(P_5, x) = \text{Dom}(\hat{P}_5, x)$ , since every dominating set of  $\hat{P}_5$  is also a dominating set of  $P_5$ . Further,

$$P_A(P_5, x) = -x(x-1)(x+1)(x^2-3) \neq P_A(\hat{P}_5, x) = -x(x^2-x-3)(x^2+x-1),$$


hence  $P_A(P_5, x) \neq P_A(\hat{P}_5, x)$ . Thus  $\xi(P_5, x) \neq \xi(\hat{P}_5, x)$ , since both  $P_5$  and  $\hat{P}_5$  are  $\xi$ -unique.

(C) The graphs  $G_1$  and  $G_2$  of Figure 3 are of order 10. Therefore  $\xi(G_1, x) = \xi(G_2, x)$ . We also have

$$\begin{aligned} \text{Dom}(G_1; x) &= x^{10} + 10x^9 + 40x^8 + 82x^7 + 92x^6 + 56x^5 + 16x^4, \\ \text{Dom}(G_2; x) &= x^{10} + 10x^9 + 41x^8 + 86x^7 + 94x^6 + 48x^5 + 9x^4, \\ P_A(G_1; x) &= x^2(x^4 - x^3 - 4x^2 + 2x + 3)(x^4 + x^3 - 4x^2 - 2x + 3), \\ P_A(G_2; x) &= x^2(x-1)(x+1)(x^2-2)(x^4 - 5x^2 + 3), \end{aligned}$$

Hence  $\text{Dom}(G_1, x) \neq \text{Dom}(G_2, x)$  and  $P_A(G_1, x) \neq P_A(G_2, x)$ .

Now (A) and (B) proves (iv), (A) and (C) proves (v), and (B) and (C) proves (vi). □



**Figure 3.**  $\xi$ -mates.

In Section 7 we use the following observation:

**Lemma A.** *Let  $P(G; \bar{x})$  and  $Q(G; \bar{y})$  two graph polynomials and  $\mathcal{C}$  and  $\mathcal{D}$  be graph properties with  $\mathcal{D} \subseteq \mathcal{C}$ . Assume that  $P <_{s.d.p}^{\mathcal{C}} Q$  and  $Q$  is weakly distinguishing in  $\mathcal{C}$ . Then also  $P$  is weakly distinguishing in  $\mathcal{C}$  but not necessarily in  $\mathcal{D}$ .*

*Proof.* Clearly,  $P <_{s.d.p}^{\mathcal{C}} Q$  implies that  $U_P(n) \subseteq U_Q(n)$ . Hence

$$\lim_{n \rightarrow \infty} \frac{|U_P(n)|}{|\mathcal{G}(n)|} \leq \lim_{n \rightarrow \infty} \frac{|U_Q(n)|}{|\mathcal{G}(n)|}. \quad \square$$

### 3. Addable properties and graphs of genus $k$

We discuss properties on which the Tutte, domination, matching,  $\xi$ , clique and characteristic polynomials are weakly distinguishing. These properties were studied in [McDiarmid, Steger, Welsh 2005; McDiarmid 2008].

#### *Small addable classes.*

**Definition 3.1.** A graph property  $\mathcal{A}$  is *decomposable* if it is closed under disjoint union, and for all  $G \in \mathcal{A}$  every component of  $G$  is in  $\mathcal{A}$ .

A graph property  $\mathcal{A}$  is *bridge addable* if for each graph  $G \in \mathcal{A}$  and every two vertices  $u, v$  in different components of  $G$  the graph obtained by adding an edge between  $u$  and  $v$  is also in  $\mathcal{A}$ .

A graph property  $\mathcal{A}$  is *addable* if it is decomposable and bridge addable.

**Example 3.2.** The following properties are easily seen to be addable:

- planar graphs,
- outerplanar graphs,
- series-parallel graphs,
- graphs with tree width at most  $k$  for  $k \geq 2$ ,
- $k$ -colorable graphs for  $k \geq 2$ ,
- graphs with no cycles of length greater than  $k$ ,
- graphs with no  $K_k$  minor for  $k \geq 2$ .

Let  $\mathcal{A}$  be a graph property. If  $\mathcal{A}$  is minor closed (that is, closed under deletion of vertices and edges, and under contraction of edges), the Graph Minor theorem says that  $\mathcal{A}$  is characterized by a finite set of forbidden minors (see [Lovász 2006] for more on graph minors and the graph minor theorem). We can characterize addable minor closed graph properties in terms of their forbidden minors:

**Proposition 3.3** [McDiarmid 2009]. *Let  $\mathcal{A}$  be a minor closed graph property. Then  $\mathcal{A}$  is addable if and only if each excluded minor of  $\mathcal{A}$  is 2-connected.*

Note that any nonempty, minor closed addable graph property  $\mathcal{A}$  contains all forests, as it contains the graph with a single vertex, and is closed under taking disjoint unions of graphs in  $\mathcal{A}$  and under adding edges between different components.

In addition to being addable, we will require the properties we consider to be small:

**Definition 3.4.** Let  $\mathcal{A}$  be a graph property, and denote by  $\mathcal{A}_n$  the graphs of order  $n$  in  $\mathcal{A}$ . We say that a graph property  $\mathcal{A}$  is small if there exists a constant  $a > 0$  such that  $|\mathcal{A}_n| \leq a^n n!$  for all sufficiently large  $n$ .

The following result is convenient:

**Theorem 3.5** [Norine et al. 2006]. *Let  $\mathcal{C}$  be a proper minor closed graph property. Then  $\mathcal{C}$  is small.*

Let  $H$  be a graph on vertex set  $\{1, \dots, h\}$  and let  $G$  be a graph on the vertex set  $\{1, \dots, n\}$  where  $n > h$ . Let  $W \subseteq V(G)$  with  $|W| = h$ , and let the root  $r_W$  denote the least element in  $W$ . We say that  $W$  is a pendant appearance of  $H$  in  $G$  if (a) the increasing bijection from  $\{1, \dots, h\}$  to  $W$  gives an isomorphism between  $H$  and the induced subgraph  $G[W]$  of  $G$ , and (b) there exists exactly one edge between  $W$  and  $V(G) - W$ , and this edge is incident with the root  $r_W$ . Our method for proving graph polynomials are weakly distinguishing will rely on the following theorem:

**Theorem 3.6** [McDiarmid, Steger, Welsh 2005]. *Let  $\mathcal{C}$  be a non-empty, addable, and small graph property, let  $H \in \mathcal{C}$  be a connected graph, and let  $R_n$  be a random graph selected uniformly at random from the graphs of order  $n$  in  $\mathcal{C}$ . Denote by  $f_H(R_n)$  the number of pendant appearances of  $H$  in  $R_n$ . Then there are constants  $\alpha > 0, n_0 \in \mathbb{N}$  such that for all  $n > n_0$ ,*

$$\mathbb{P}[f_H(R_n) \leq \alpha n] < e^{-\alpha n}$$

For our purposes, we shall only need a weaker corollary of Theorem 3.6:

**Corollary 3.7.** *Let  $\mathcal{A}$  be an addable proper minor closed graph property,  $H$  a fixed connected graph in  $\mathcal{A}$ . Denote by  $\mathcal{A}'_n$  the set of all  $n$  vertex graphs  $G \in \mathcal{A}$  that have at least one pendant appearance of  $H$ . Then  $\lim_{n \rightarrow \infty} |\mathcal{A}'_n|/|\mathcal{A}_n| = 1$ .*

All the properties listed in Example 3.2 are addable and minor closed, hence the corollary applies to them. Using this corollary, we get:

**Lemma B.** *Let  $P$  be a graph polynomial, and  $\mathcal{A}$  a small addable graph property. If there is a fixed  $H \in \mathcal{A}$  such that every graph  $G \in \mathcal{A}$  with a pendant appearance of  $H$  has a  $P$  mate  $G' \in \mathcal{A}$ , then  $P$  is weakly distinguishing on  $\mathcal{A}$ .*

*Proof.* By Corollary 3.7, almost all graphs  $G \in \mathcal{A}$  have a pendant appearance of  $H$ , and hence almost all graphs  $G \in \mathcal{A}$  have a  $P$ -mate, so  $P$  is weakly distinguishing on  $\mathcal{A}$ .  $\square$

**Graphs with genus at most  $k$ .** Let  $G$  be a graph. The genus of  $G$ , denoted  $g(G)$ , is the minimal genus of a surface in which  $G$  can be embedded. (For more on graphs embedded in surfaces see [Lando and Zvonkin 2004; Mohar and Thomassen 2001], for example.) It is easy to see that  $g(G)$  is well defined, as for all  $G$ ,  $g(G) \leq |E(G)|$ . For  $k \in \mathbb{N}$ , denote by  $\mathcal{C}_k$  the property of graphs with genus at most  $k$ . Note that in general  $\mathcal{C}_k$  is not addable - for instance, the genus of  $K_5$ , the complete graph with 5 vertices, is 1, but the genus of a disjoint union of two copies of  $K_5$  has genus 2- so we can not apply Theorem 3.6 to  $\mathcal{C}_k$ . However, the same result does hold for  $\mathcal{C}_k$  with a slight modification:

**Theorem 3.8** [McDiarmid 2008]. *Let  $H$  be a connected graph,  $k \in \mathbb{N}$ , and let  $R_n$  be a random graph selected uniformly at random from the graphs of order  $n$  in  $\mathcal{C}_k$ . Denote by  $f_H(R_n)$  the number of pendant appearances of  $H$  in  $R_n$ . Then there are constants  $\alpha > 0, n_0 \in \mathbb{N}$  such that for all  $n > n_0$ ,*

$$\mathbb{P}[f_H(R_n) \leq \alpha n] < e^{-\alpha n}$$

Again, we only need a weaker corollary of this theorem:

**Corollary 3.9.** *Let  $H$  be a fixed connected planar graph,  $k \in \mathbb{N}$  and let  $\mathcal{A} = \mathcal{C}_k$  and let  $\mathcal{A}'_n$  be the set of all  $n$  vertex graphs  $G \in \mathcal{A}$  that have at least one pendant appearance of  $H$ . Then  $\lim_{n \rightarrow \infty} |\mathcal{A}'_n|/|\mathcal{A}_n| = 1$ .*

**Lemma C.** *Let  $P$  be a graph polynomial, and  $\mathcal{C}_k$  the property of graphs with genus at most  $k$ . If there is a fixed planar graph  $H$  such that every graph  $G$  with a pendant appearance of  $H$  has a  $P$  mate  $G' \in \mathcal{C}_k$ , then  $P$  is weakly distinguishing on  $\mathcal{C}_k$ .*

*Proof.* By Corollary 3.9, almost all graphs  $G \in \mathcal{C}_k$  have a pendant appearance of  $H$ , and hence almost all graphs  $G \in \mathcal{C}_k$  have a  $P$ -mate, so  $P$  is weakly distinguishing on  $\mathcal{C}_k$ . □

### 4. The characteristic polynomial

**Definition 4.1.** Let  $G$  be a graph, and denote by  $A_G$  its adjacency matrix. The characteristic polynomial of  $G$ , denoted  $P_A(G)$  is defined as the characteristic polynomial of  $A_G$ . The set of roots of  $P_A(G)$  is referred to as the spectrum of  $G$ .

A  $P_A$ -unique graph is usually referred to as a graph determined by its spectrum, and if  $G$  and  $H$  are two non isomorphic graphs such that  $P_A(G) = P_A(H)$ ,  $G$  and  $H$  are said to be cospectral.

The characteristic polynomial and particularly its roots are widely studied; see [Brouwer and Haemers 2012], for example, for an introduction. A classic result of Schwenk states that almost all trees are not determined by their spectrum (see [Schwenk 1973] for details). Using Corollary 3.7, we can extend this result to all small addable graph properties.

We use the following recurrence relation for the characteristic polynomial:


**Lemma 4.2** (see [Clarke 1970]). *Let  $G_1$  and  $G_2$  be two graphs, and let  $v_1 \in V(G_1)$ ,  $v_2 \in V(G_2)$ . Denote by  $H$  the graph obtained from the disjoint union of  $G_1$  and  $G_2$  by adding an edge from  $v_1$  to  $v_2$ . Then*

$$P_A(H, x) = P_A(G_1, x)P_A(G_2, x) - P_A(G_1 - v_1, x)P_A(G_2 - v_2, x) \tag{4-1}$$

We now take two graphs considered in [Schwenk 1973], labeled  $R$  and  $S$  in Figure 4. They are isomorphic, and we can check that

$$P_A(S - v, x) = P_A(R - u, x) = x^8 - 6x^6 + 10x^4 - 4x^2.$$

**Lemma 4.3.** *Let  $G_S$  be a graph with a pendant appearance of  $S$  rooted at  $v$ . Denote by  $G_R$  the graph obtained from  $G$  by replacing the pendant appearance of  $S$  in  $G$  with a pendant appearance of  $R$ , rooted at  $u$ . Then  $G_S$  and  $G_R$  are cospectral mates.*



**Figure 4.** Cospectral graphs.

*Proof.* Denote by  $w \in V(G_S)$  the vertex adjacent to  $v$  that is not in the pendant appearance of  $S$ , and by  $H$  the graph obtained from  $G_S$  by deleting all the vertices in the pendant appearance of  $S$ . By applying relation (4-1) to  $G_S$  with the edge  $wv$ , we get

$$P_A(G_S, x) = P_A(S, x)P_A(H, x) - P_A(S - v, x)P_A(H - w, x)$$

By applying relation (4-1) to  $G_R$  with the edge  $wu$  we get

$$P_A(G_R, x) = P_A(R, x)P_A(H, x) - P_A(R - u, x)P_A(H - w, x)$$

But since  $P_A(R, x) = P_A(S, x)$  and  $P_A(S - v, x) = P_A(R - u, x)$ , we get that  $P_A(G_S, x) = P_A(G_R, x)$ , so  $G_S$  and  $G_R$  are cospectral mates.  $\square$

As unrooted graphs,  $S$  and  $R$  are isomorphic. Thus, we have:

**Theorem 4.4.** *Let  $\mathcal{A}$  be a small addable graph property such that  $S \in \mathcal{A}$ . Then  $P_A$  is weakly distinguishing on  $\mathcal{A}$ .*

*Proof.* From lemmas B and 4.3 we get that  $P_A$  is weakly distinguishing on  $\mathcal{A}$ .  $\square$

**Corollary 4.5.** *Let  $\mathcal{A}$  be a proper minor closed addable graph property. Then  $P_A$  is weakly distinguishing on  $\mathcal{A}$ .*

*Proof.* Since all forests are in  $\mathcal{A}$ , and  $S$  is a tree, by Theorems 4.4 and 3.5,  $P_A$  is weakly distinguishing on  $\mathcal{A}$ .  $\square$

Since all the properties in list 3.2 are addable and minor closed, we have:

**Corollary 4.6.**  *$P_A$  is weakly distinguishing on all the properties listed in Example 3.2.*

Similarly, we get:

**Theorem 4.7.** *Denote by  $\mathcal{C}_k$  the class of graphs of genus at most  $k$ . Then  $P_A$  is weakly distinguishing on  $\mathcal{C}_k$  for all  $k \in \mathbb{N}$ .*

*Proof.* Since  $S$  is a tree, from lemmas C and 4.3 we get that  $P_A$  is weakly distinguishing on  $\mathcal{C}_k$ .  $\square$

Instead of the characteristic polynomial  $P_A$ , which is based on the adjacency matrix, one can also look at the analogue graph polynomial  $P_L$  which is based on the Laplacian matrix  $L_G = D_G - A_G$ , where  $D_G$  is the diagonal matrix with diagonal elements  $d_{v,v} = \deg(v)$  for each  $v \in V(G)$ . Now  $P_L(G; x)$  is the characteristic polynomial of  $L_G$ . An early survey about the Laplacian polynomial may be found in [Mohar 1991]. The graph polynomials are *d.p.*-equivalent on regular graphs, but *d.p.*-incomparable on all finite graphs; see [Brouwer and Haemers 2012] and [Makowsky et al. 2019]. Our proofs of Theorems 4.4 and 4.7 do not work for  $P_L(G; x)$ .

**Problem 4.1.** Find two graphs  $R$  and  $S$  that can be used to prove analogues of Theorems 4.4 and 4.7.



**Figure 5.** A path of length 5.

### 5. The domination polynomial

The *domination polynomial* is the generating function of dominating sets in a given graph  $G$ . More formally:

**Definition 5.1.** Let  $G$  be a graph. A set  $S \subseteq V(G)$  is called a dominating set if for every  $v \in V(G)$ , either  $v \in S$  or  $v$  has a neighbor  $u \in S$ . Define  $\text{Dom}(G; x) = \sum_{S \subseteq V(G)} x^{|S|}$ , where the sum is over all dominating sets of  $G$ .

The domination polynomial was extensively studied in recent years; see [Alikhani and Peng 2014] for a survey.

**Lemma 5.2** [Kotek et al. 2012]. *Let  $G$  be a graph. We say a vertex  $v \in V(G)$  is a stem if it has a neighbor  $u \in V(G)$  with  $\text{deg}(u) = 1$ . Assume  $G$  has two stems  $u, u' \in G$ ,  $(u, u') \in E(G)$ , and denote by  $G'$  the graph resulting from  $G$  by deleting the edge  $(u, u')$ . Then for every set  $S \subseteq V(G)$ ,  $S$  is a dominating set of  $G$  if and only if  $S$  is a dominating set of  $G'$ .*

*Proof.* If  $S$  is a dominating set of  $G'$ , it is clearly also a dominating set of  $G$ . On the other hand, let  $S$  be a dominating set of  $G$ . After deleting  $(u, u')$ , the only vertices that are perhaps not dominated by  $S$  are  $u, u'$ . But  $u$  has a vertex of degree 1 as a neighbor, so either it or  $u$  are in  $S$ , and in either case  $u$  is dominated. The same applies to  $u'$ . So  $S$  is a dominating set of  $G'$ . □

**Theorem 5.3.** *Every graph  $G$  that has two distinct stems has a Dom-mate.*

*Proof.* This is a direct consequence of the lemma. □

**Corollary 5.4.** *Let  $G$  be a graph that has a pendant appearance of  $P_5$  rooted at  $r$  (see Figure 5), and let  $\hat{G}$  be the graph obtained from  $G$  by replacing the pendant appearance of  $P_5$  with a pendant appearance of  $\hat{P}_5$  rooted at  $r$ . Then  $G$  and  $\hat{G}$  are Dom-mates.*

**Theorem 5.5.** *Let  $\mathcal{A}$  be a small addable graph property such that  $P_5, \hat{P}_5 \in \mathcal{A}$ . Then Dom is weakly distinguishing on  $\mathcal{A}$ .*

*Proof.* From Lemma B and Corollary 5.4 we get that Dom is weakly distinguishing on  $\mathcal{A}$ . □

**Corollary 5.6.** *Let  $\mathcal{A}$  be a proper minor closed addable graph property, such that  $\hat{P}_5 \in \mathcal{A}$ . Then Dom is weakly distinguishing on  $\mathcal{A}$*

*Proof.* Since all forests are in  $\mathcal{A}$ ,  $P_5 \in \mathcal{A}$ , and hence from Theorems 5.5 and 3.5 Dom is weakly distinguishing on  $\mathcal{A}$ . □

As in the previous sections, we have:

**Corollary 5.7.** *Dom is weakly distinguishing on all the properties listed in Example 3.2.*

Similarly, we get:

**Theorem 5.8.** Denote by  $\mathcal{C}_k$  the property of graphs of genus less than  $k$ . Then Dom is weakly distinguishing on  $\mathcal{C}_k$  for all  $k \in \mathbb{N}$ .

*Proof.* Since  $P_5$  and  $\hat{P}_5$  are planar, from Lemma C and Corollary 5.4 we get that Dom is weakly distinguishing on  $\mathcal{C}_k$ .  $\square$

## 6. The $\xi$ polynomial

The  $\xi$  polynomial, introduced in [Averbouch et al. 2008; 2010], generalizes the matching and the Tutte polynomial. It is defined via a recurrence relation:

Let  $G$  be a graph and  $e$  an edge of  $G$ . We denote by  $G - e$  the graph resulting from  $G$  by deleting the edge  $e$ , by  $G/e$  the graph resulting from  $G$  by contracting the edge  $e$ , and by  $G \dagger e$  the graph resulting from  $G$  by extracting the edge  $e$ , i.e., deleting  $e$  together with its adjacent vertices.

**Definition 6.1.** Let  $G$  be a graph. Define the trivariate polynomial  $\xi(G; x, y, z)$  using the recursive relation

$$\begin{aligned} \xi(G; x, y, z) &= \xi(G - e; x, y, z) + y\xi(G/e; x, y, z) + z\xi(G \dagger e; x, y, z), \\ \xi(G \sqcup H; x, y, z) &= \xi(G; x, y, z)\xi(H; x, y, z) \end{aligned} \quad (6-1)$$

with the base conditions  $\xi(K_1; x, y, z) = x$  and  $\xi(\emptyset; x, y, z) = 1$ .

An alternative representation of the  $\xi$  polynomial was introduced in [Trinks 2012]:

**Definition 6.2.** Let  $G$  be a graph. The covered components polynomial  $C(G; x, y, z)$  of  $G$  is defined as

$$C(G; x, y, z) = \sum_{A \subseteq E(G)} x^{k(G\langle A \rangle)} y^{|A|} z^{c(G\langle A \rangle)},$$

where  $G\langle A \rangle$  is the graph  $(V(G), A)$ ,  $k(G\langle A \rangle)$  is the number of connected components of  $G\langle A \rangle$  and  $c(G\langle A \rangle)$  is the number of covered connected components of  $G\langle A \rangle$ , that is connected components in  $G\langle A \rangle$  with at least one edge.

We sometimes write  $C(G)$  for  $C(G; x, y, z)$  to simplify long expressions. The two definitions are connected by the following result:

**Proposition 6.3** [Trinks 2012]. For all graphs  $G$ ,


$$C(G; x, y, z) = \xi(G; x, y, xyz - xy). \quad (6-2)$$

**Corollary 6.4.** Let  $G$  and  $H$  be graphs. Then  $C(G; x, y, z) = C(H; x, y, z)$  if and only if  $\xi(G; x, y, z) = \xi(H; x, y, z)$

We will use the following recurrence relation:

**Theorem 6.5** [Trinks 2012]. Let  $G_1, G_2$  be graphs, and  $v_1 \in V(G_1), v_2 \in V(G_2)$ . Let  $H$  be the graph obtained from  $G_1$  and  $G_2$  by identifying  $v_1$  with  $v_2$ . Then

$$\begin{aligned} C(G) &= \left(\frac{1}{xz} + \frac{2}{x}\right)C(G_1)C(G_2) + \left(-\frac{1}{z} - 1\right)(C(G_1)C(G_2 - v_2) + C(G_1 - v_1)C(G_2)) \\ &\quad + \left(\frac{x}{z} + x\right)C(G_1 - v_1)C(G_2 - v_2). \end{aligned} \quad (6-3)$$



**Figure 6.** A tree with pseudosimilar vertices.

We next apply Theorem 6.5 to exhibit a pair of  $C$  mates.

**Lemma 6.6.** *Consider the graph  $S$  in Figure 6. Let  $G_v$  be a graph with a pendent appearance of  $S$  rooted at  $v$ , and let  $G_u$  be the graph obtained from  $G_v$  by replacing the pendent appearance of  $S$  rooted at  $v$  by a pendent appearance of  $S$  rooted at  $u$ . Then  $G_v$  and  $G_u$  are  $C$  mates.*

*Proof.* We apply relation (6-3) to  $G_v$ , with  $G_1 = S$  and  $G_2$  being the graph  $H_1$  obtained from  $G_v$  by replacing the pendent appearance of  $S$  with a single vertex  $w$ . We get

$$C(G_v) = \left(\frac{1}{xz} + \frac{2}{x}\right)C(S)C(H_1) + \left(-\frac{1}{z} - 1\right)(C(S)C(H_1 - w) + C(S - v)C(H_1)) + \left(\frac{x}{z} + x\right)C(S - v)C(H_1 - w). \quad (6-4)$$

Similarly, we apply relation (6-3) to  $G_u$ , with  $G_1 = S$  and  $G_2$  being the graph  $H_2$  obtained from  $G_u$  by replacing the pendent appearance of  $S$  with a single vertex  $w$ . We get

$$C(G_u) = \left(\frac{1}{xz} + \frac{2}{x}\right)C(S)C(H_2) + \left(-\frac{1}{z} - 1\right)(C(S)C(H_2 - w) + C(S - u)C(H_2)) + \left(\frac{x}{z} + x\right)C(S - u)C(H_2 - w).$$

Note that  $H_1 \cong H_2$ ,  $H_1 - w \cong H_2 - w$ , and  $S - v \cong S - u$ , and hence  $C(G_v) = C(G_u)$ . □

As a direct consequence of Lemma 6.6 and Corollary 6.4, we have:

**Corollary 6.7.** *Let  $G_v$  be a graph with a pendent appearance of  $S$  (depicted in Figure 6) rooted at  $v$ , and let  $G_u$  be the graph obtained from  $G_v$  by replacing the pendent appearance of  $S$  rooted at  $v$  by a pendent appearance of  $S$  rooted at  $u$ . Then  $G_v$  and  $G_u$  are  $\xi$  mates.*

From Lemma B and Corollary 6.7 we get:

**Theorem 6.8.** *Let  $\mathcal{A}$  be a small addable graph property such that  $S \in \mathcal{A}$ . Then  $\xi$  is weakly distinguishing on  $\mathcal{A}$ .*

**Corollary 6.9.** *Let  $\mathcal{A}$  be a proper minor closed addable graph property. Then  $\xi$  is weakly distinguishing on  $\mathcal{A}$ .*

*Proof.* Since all forests are in  $\mathcal{A}$ ,  $S \in \mathcal{A}$ , and hence from Theorems 6.8 and 3.5,  $M$  is weakly distinguishing on  $\mathcal{A}$ . □

As in the previous section, we get:

**Corollary 6.10.**  *$\xi$  is weakly distinguishing on all the properties listed in Example 3.2.*

Similarly, we get:

**Theorem 6.11.** *Denote by  $\mathcal{C}_k$  the class of graphs of genus less than  $k$ . Then  $\xi$  is weakly distinguishing on  $\mathcal{C}_k$  for all  $k \in \mathbb{N}$ .*

*Proof.* Since  $S$  is a tree, from lemmas C and 6.7 we get that  $\xi$  is weakly distinguishing on  $\mathcal{C}_k$ .  $\square$

## 7. $\xi$ -invariants

We investigate consequences of Theorems 6.8 and 6.11 using Lemma A. From Theorem 6.8 we get:

**Corollary 7.1.** *Let  $\mathcal{A}$  be a small addable graph property such that  $S \in \mathcal{A}$ , and let  $P$  be a graph polynomial such that  $P <_{s.d.p}^C \xi$ . Then  $P$  is weakly distinguishing on  $\mathcal{A}$ .*

**Corollary 7.2.** *Let  $\mathcal{A}$  be a proper minor closed addable graph property, and let  $P$  be a graph polynomial such that  $P <_{s.d.p}^C \xi$ . Then  $P$  is weakly distinguishing on  $\mathcal{A}$ .*

From Theorem 6.11 we get:

**Corollary 7.3.** *Let  $\mathcal{C}_k$  the class of graphs of genus less than  $k$ , and let  $P$  be a graph polynomial such that  $P <_{s.d.p}^C \xi$ . Then  $P$  is weakly distinguishing on  $\mathcal{C}_k$  for all  $k \in \mathbb{N}$ .*

We next consider results from the literature showing that there are many graph polynomials  $P$  where we can apply these results. Among them we find

- (i) the generalized chromatic polynomial from [Dohmen et al. 2003], including the chromatic polynomial;
- (ii) the matching polynomials;
- (iii) the independence polynomial, including the vertex cover polynomial;
- (iv) the Tutte polynomial, including the flow polynomial, the reliability polynomial and the Euler polynomial.

**The generalized chromatic polynomial.** The generalized chromatic polynomial  $\text{GC}(G; x, y)$  was introduced in [Dohmen et al. 2003]. and  $Z$ . Let  $Y$  and  $Z$  be two disjoint sets of colors. A generalized coloring of a graph  $G = (V(G), E(G))$  is a map  $c : V(G) \rightarrow (Y \sqcup Z)$  such that for all  $(u, v) \in E$ , if  $c(u) \in Y$  and  $c(v) \in Y$ , then  $c(u) = c(v)$ .  $Y$  is called the set of proper colors. For two positive integers  $x \geq y$ , the value of the polynomial  $\text{GC}(G; x, y)$  is the number of generalized colorings by  $x$  colors, where  $y$  of them are proper. The chromatic polynomial  $\chi(G; x)$  is obtained for the case  $x = y$ .

**Theorem 7.4** [Averbouch et al. 2010, Proposition 22]. *For all graphs  $G$ ,*

$$\text{GC}(G; x, y) = \xi(G; x, -1, x - y).$$

*Therefore  $\chi <_{d.p}^C CG <_{d.p}^C \xi$ .*

**The matching polynomial.** Several variants of the matching polynomial are considered in the literature:

**Definition 7.5.** Let  $G$  be a graph with  $n$  vertices. A *matching* in  $G$  is a spanning subgraph of  $G$  in which every connected component is either an isolated vertex or two vertices connected by a single edge. We say a matching is of size  $k$  if it has exactly  $k$  edges. Denote by  $m_k(G)$  the number of  $k$  matchings in  $G$ .

The *matching acyclic polynomial* (also known as the matching defect polynomial) is defined as

$$\mu(G; x) = \sum_{k=0}^{\lfloor n/2 \rfloor} (-1)^k m_k(G) x^{n-2k}$$

The *matching generating polynomial* is defined as

$$g(G; x) = \sum_{k=0}^{\lfloor n/2 \rfloor} m_k(G) x^k$$

The *bivariate matching polynomial* of  $G$  is defined as

$$M(G; w_1, w_2) = \sum_{k=0}^{\lfloor n/2 \rfloor} m_k(G) w_1^{n-2k} w_2^k$$

For an introduction to the bivariate matching polynomial, see [Farrell 1979]. For a recent survey on the acyclic matching polynomial; see [Gutman 2016].

For our purposes, we note the following fact:

**Fact 7.1.** All of the polynomials in Definition 7.5 are of the same distinctive power on similar graphs:.

$$\mu \sim_{s.d.p} g \sim_{s.d.p} M.$$

In fact, we also have  $\mu \sim_{d.p} M$ , but for the edgeless graphs  $E_n$  of order  $n$  we have  $g(E_n, x) = g(E_m, x)$  for all  $m, n \geq 1$ , but  $\mu(E_n, x) \neq \mu(E_m, x)$  for  $n \neq m$ .

Thus we will only consider the bivariate matching polynomial.

**Theorem 7.6** [Averbouch et al. 2010, Proposition 20]. *For all graphs  $G$*

$$M(G; x, y) = \xi(G; x, 0, y);$$

therefore  $M <_{s.d.p} \xi$ .

**Independence and clique polynomial.** The independence polynomial is defined as

$$\text{In}(G; x) = \sum_{A \subseteq V(G)} x^{|A|}$$

where the graph induced graph  $G[A]$  is edgless.

The vertex cover polynomial  $\text{VC}(G; x)$  is defined as


$$\text{VC}(G; x) = \sum_{A \subseteq V(G)} x^{|A|}$$

where  $A$  is a vertex cover of  $G$ .

The following is taken from (but possibly not originally due to) [Trinks 2012].

**Proposition 7.7.** *For all graphs  $G$  we have:*

- (i)  $\text{In}(G; x) = \text{GC}(G; x + 1, 1)$ .
- (ii)  $\text{VC}(G; x) = x^n \text{In}(G; \frac{1}{x})$ .



**Figure 7.** A path of length 3.

Hence,  $\text{VC} \sim_{s.d.p} \text{In} <_{s.d.p} \text{GC}$ .

The clique polynomial is defined as

$$\text{Cl}(G; x) = \sum_{A \subseteq V(G)} x^{|A|},$$

where the graph induced graph  $G[A]$  is a complete graph.

We note that  $\text{In}(G; x) = \text{Cl}(\bar{G})$  for simple graphs. Therefore  $\text{In} \sim_{d.p} \text{Cl}$  by Example 2.2(iii).

Both  $\text{In}(G; x)$  and  $\text{Cl}(G; x)$  were shown in [Makowsky and Rakita 2019] to be weakly distinguishing on all finite graphs. In the light of the above,  $\text{In}(G; x)$  is also weakly distinguishing on small addable graph properties, and on graphs of genus at most  $k$ .

This also holds if  $\mathcal{C}$  is addable, small, and closed under complements. If  $\mathcal{C}$  is addable and small, but not closed under complements we can use the following lemma:

**Lemma 7.8.** *Let  $G_r$  be a graph such that  $G$  has a pendant appearance of  $P_3$  (see Figure 7) rooted at  $r$ , and  $G_u$  the graph obtained from  $G_r$  by replacing the pendant appearance of  $P_3$  by a pendant appearance of  $P_3$  rooted at  $u$ . Then  $G_r$  and  $G_u$  are clique mates.*

*Proof.* For all  $k > 2$ ,  $G_r$  and  $G_u$  have the same number of  $k$  cliques (since no  $k$  clique can include vertices from the pendant appearance), and  $G$  and  $G'$  have the same number of vertices and edges, and hence the same number of 1 and 2 cliques. So  $\text{Cl}(G_r; x) = \text{Cl}(G_u; x)$ . □

From Lemmas B and 7.8 we get:

**Theorem 7.9.** *Let  $\mathcal{A}$  be a small addable graph property such that  $P_3 \in \mathcal{A}$ . Then  $\text{Cl}$  is weakly distinguishing on  $\mathcal{A}$ .*

**The Tutte polynomial.** Let  $G = (V(G), E(G))$  be a graph. The Tutte polynomial  $T(G; x, y)$  is defined by

$$T(G; x, y) = \sum_{A \subseteq E(G)} (x - 1)^{k(A) - k(E)} (y - 1)^{k(A) + |A| - |V(G)|}.$$

The partition function  $Z(G; q, w)$  is defined by

$$Z(G; q, w) = \sum_{A \subseteq E(G)} q^{k(F)} w^{|F|}.$$

They are related by the equation

$$T(G; x, y) = (x - 1)^{-k(E)} (y - 1)^{-|V(G)|} Z(G; (x - 1)(y - 1), (y - 1)).$$

The chromatic polynomial  $\chi(G; x)$  can be obtained from the Tutte polynomial by

$$\chi(G; x) = (-1)^{|V(G)| - k(G)} x^{k(G)} T(G; (1 - x), 0)$$

From this we get:

**Proposition 7.10.**  $\chi <_{s.d.p} T \sim_{s.d.p} Z <_{d.p} C \sim_{d.p} \xi$ .

A graph is *Eulerian* if all its vertices have even degree. It does not have to be connected. The Euler polynomial  $\mathcal{E}(G; x)$  of a graph is defined by

$$\mathcal{E}(G; x) = \sum_{\substack{A \subseteq E(G) \\ (V, A) \text{ is Eulerian}}} x^{|A|};$$

it is related to the Tutte polynomial as follows [Aigner 2007, Chapter 10, p. 468]:

$$\mathcal{E}(G; x) = (1-x)^{|E(G)|-|V(G)|+k(G)} x^{|V(G)|-k(G)} T\left(G; \frac{1}{x}, \frac{1+x}{1-x}\right).$$

Hence we get:

**Proposition 7.11.**  $\mathcal{E} <_{s.d.p} T$ .

The flow polynomial  $\text{Fl}(G, x)$  and the reliability polynomial  $R(G; p)$  are related to the Tutte polynomial by

$$\begin{aligned} \text{Fl}(G; x) &= (-1)^{|E(G)|-|V(G)|+k(G)} T(G; 0, 1-x) \\ R(G; p) &= (p)^{|E(G)|-|V(G)|+k(G)} (1-p)^{|V(G)|-k(G)} T\left(G; 1, \frac{1}{p}\right). \end{aligned}$$

Hence we get:

**Proposition 7.12.**  $\text{Fl} <_{s.d.p} T$  and  $R <_{s.d.p} T$ .

## 8. Conclusion

We have shown that many graph polynomials are weakly distinguishing on all proper minor closed addable graph properties, including many interesting properties such as planar graphs, graphs with tree width at most  $k$ , and  $K_k$  free graphs (for  $4 < k \in \mathbb{N}$ ). In addition, we proved that the domination, characteristic, and the edge elimination polynomial  $\xi(G; x, y, z)$  are weakly distinguishing on the properties of graphs with genus less than  $k$  for all  $k \in \mathbb{N}$ . This also applies to graph polynomials derivable from  $\xi$ , such as the generalized chromatic polynomial, Tutte polynomial and its variants, the matching, the independence, and the clique polynomials.

Our results relied on the fact that proper minor closed addable properties are, in a sense, small, and so they do not settle the question whether the domination, characteristic or Tutte polynomials are weakly distinguishing, almost complete, or otherwise on all graphs.

We have shown that for the above graph polynomials  $P$ , the sequence  $\alpha_P^{\mathcal{C}}(n) = \frac{|U_P^{\mathcal{C}}(n)|}{|\mathcal{C}(n)|}$  tends to 0 as  $n$  tends to infinity.

The following questions are natural extensions of the work in this paper:

**Problem 8.1.** What can be said about  $\alpha_P^{\mathcal{C}}(n)$  when  $\mathcal{C}$  is assumed to be a hereditary or monotone graph property?

**Problem 8.2.** Can we find a graph polynomial  $P$  and a graph property  $\mathcal{C}$  such that we can prove that  $\alpha_P^{\mathcal{C}}(n) \geq \beta$  for all sufficiently large  $n$  for some fixed  $\beta \in (0, 1]$ ?

**Problem 8.3.** For  $P$  one of the above polynomials and  $\mathcal{A}$  a proper minor closed addable property, select a random graph  $G_n$  uniformly at random in  $\mathcal{A}_n$ , and denote  $[G_n] = \{H \in \mathcal{A}_n : P(G) = P(H)\}$ . What can be said about the limit distribution of the random variable  $|[G_n]|$ ?

### Acknowledgements

Preliminary versions of our results were presented at the 19th International Conference on Random Structures and Algorithms, 15 to 19 July 2019, Zürich, Switzerland. The results of Sections 6 and 7 were partially obtained during Seminar 19401 in Dagstuhl (Comparative Theory for Graph Polynomials) in October 2019. We want to thank the participants of the Dagstuhl Seminar Delia Garijo, Anna de Mier, Marc Noy, Elena Ravve and Peter Tittmann for valuable discussions and comments. Thanks go especially to Peter Tittmann for his guidance and comments about invariants derivable from  $\xi$ -polynomial, which helped us formulating Section 7.

### References

- [Aigner 2007] M. Aigner, *A course in enumeration*, Graduate Texts in Mathematics **238**, Springer, 2007. MR Zbl
- [Alikhani and Peng 2014] S. Alikhani and Y.-h. Peng, “Introduction to domination polynomial of a graph”, *Ars Combin.* **114** (2014), 257–266. MR Zbl
- [Averbouch et al. 2008] I. Averbouch, B. Godlin, and J. A. Makowsky, “A most general edge elimination polynomial”, pp. 31–42 in *Graph-theoretic concepts in computer science*, edited by H. Broersma et al., Lecture Notes in Comput. Sci. **5344**, Springer, 2008. MR Zbl
- [Averbouch et al. 2010] I. Averbouch, B. Godlin, and J. A. Makowsky, “An extension of the bivariate chromatic polynomial”, *European J. Combin.* **31**:1 (2010), 1–17. MR Zbl
- [Bollobás et al. 2000] B. Bollobás, L. Pebody, and O. Riordan, “Contraction-deletion invariants for graphs”, *J. Combin. Theory Ser. B* **80**:2 (2000), 320–345. MR Zbl
- [Brouwer and Haemers 2012] A. E. Brouwer and W. H. Haemers, *Spectra of graphs*, Springer, 2012. MR Zbl
- [Clarke 1970] F. H. Clarke, *The characteristic polynomial of a graph*, master’s thesis, McGill University, 1970, Available at <https://escholarship.mcgill.ca/concern/theses/gh93h0968?locale=en>.
- [Dohmen et al. 2003] K. Dohmen, A. Pönitz, and P. Tittmann, “A new two-variable generalization of the chromatic polynomial”, *Discrete Math. Theor. Comput. Sci.* **6**:1 (2003), 69–89. MR Zbl
- [Farrell 1979] E. J. Farrell, “An introduction to matching polynomials”, *J. Combin. Theory Ser. B* **27**:1 (1979), 75–86. MR Zbl
- [Godsil and Gutman 1981] C. D. Godsil and I. Gutman, “On the theory of the matching polynomial”, *J. Graph Theory* **5**:2 (1981), 137–144. MR Zbl
- [Gutman 2016] I. Gutman, “A survey on the matching polynomial”, pp. 77–99 in *Graph polynomials*, edited by M. Dehmer et al., CRC Press, Boca Raton, FL, 2016. MR
- [Kotek et al. 2012] T. Kotek, J. Preen, F. Simon, P. Tittmann, and M. Trinks, “Recurrence relations and splitting formulas for the domination polynomial”, *Electron. J. Combin.* **19**:3 (2012), art. no. 47, 27 pp. MR Zbl
- [Lando and Zvonkin 2004] S. K. Lando and A. K. Zvonkin, *Graphs on surfaces and their applications*, Encyclopaedia of Mathematical Sciences **141**, Springer, 2004. MR Zbl
- [Lovász 2006] L. Lovász, “Graph minor theory”, *Bull. Amer. Math. Soc. (N.S.)* **43**:1 (2006), 75–86. MR Zbl
- [Makowsky and Rakita 2019] J. A. Makowsky and V. Rakita, “On weakly distinguishing graph polynomials”, *Discrete Math. Theor. Comput. Sci.* **21**:1 (2019), art. no. 4, 10 pp. MR Zbl
- [Makowsky and Zhang 2019] J. A. Makowsky and R. X. Zhang, “On  $P$ -unique hypergraphs”, *Australas. J. Combin.* **73** (2019), 456–465. MR Zbl

- [Makowsky et al. 2019] J. A. Makowsky, E. V. Ravve, and T. Kotek, “A logician’s view of graph polynomials”, *Ann. Pure Appl. Logic* **170**:9 (2019), 1030–1069. MR Zbl
- [McDiarmid 2008] C. McDiarmid, “Random graphs on surfaces”, *J. Combin. Theory Ser. B* **98**:4 (2008), 778–797. MR Zbl
- [McDiarmid 2009] C. McDiarmid, “Random graphs from a minor-closed class”, *Combin. Probab. Comput.* **18**:4 (2009), 583–599. MR Zbl
- [McDiarmid, Steger, Welsh 2005] C. McDiarmid, A. Steger, and D. J. A. Welsh, “Random planar graphs”, *J. Combin. Theory Ser. B* **93**:2 (2005), 187–205. MR Zbl
- [Mohar 1991] B. Mohar, “The Laplacian spectrum of graphs”, pp. 871–898 in *Graph theory, combinatorics, and applications* (Kalamazoo, MI, 1988), vol. 2, edited by Y. Alavi et al., Wiley, New York, 1991. MR Zbl
- [Mohar and Thomassen 2001] B. Mohar and C. Thomassen, *Graphs on surfaces*, Johns Hopkins University Press, Baltimore, MD, 2001. MR Zbl
- [Norine et al. 2006] S. Norine, P. Seymour, R. Thomas, and P. Wollan, “Proper minor-closed families are small”, *J. Combin. Theory Ser. B* **96**:5 (2006), 754–757. MR Zbl
- [Noy 2003] M. Noy, “Graphs determined by polynomial invariants”, *Theoret. Comput. Sci.* **307**:2 (2003), 365–384. MR Zbl
- [Schwenk 1973] A. J. Schwenk, “Almost all trees are cospectral”, pp. 275–307 in *New directions in the theory of graphs* (Ann Arbor, MI, 1971), edited by F. Harary, Academic Press, New York, 1973. MR Zbl
- [Trinks 2012] M. Trinks, “The covered components polynomial: a new representation of the edge elimination polynomial”, *Electron. J. Combin.* **19**:1 (2012), art. no. 50, 31 pp. MR Zbl

Received 5 Dec 2019. Revised 1 Mar 2020.

JOHANN A. MAKOWSKY:

janos@cs.technion.ac.il

Department of Computer Science, Technion - IIT, Haifa, Israel

VSEVOLOD RAKITA:

vsevolod@campus.technion.ac.il

Department of Computer Science, Technion - IIT, Haifa, Israel



# Moscow Journal of Combinatorics and Number Theory

msp.org/moscow

## EDITORS-IN-CHIEF

- Yann Bugeaud    Université de Strasbourg (France)  
bugaud@math.unistra.fr
- Nikolay Moshchevitin    Lomonosov Moscow State University (Russia)  
moshchevitin@gmail.com
- Andrei Raigorodskii    Moscow Institute of Physics and Technology (Russia)  
mraigor@yandex.ru
- Ilya D. Shkredov    Steklov Mathematical Institute (Russia)  
ilya.shkredov@gmail.com

## EDITORIAL BOARD

- Iskander Aliev    Cardiff University (United Kingdom)
- Vladimir Dolnikov    Moscow Institute of Physics and Technology (Russia)
- Nikolay Dolbilin    Steklov Mathematical Institute (Russia)
- Oleg German    Moscow Lomonosov State University (Russia)
- Michael Hoffman    United States Naval Academy
- Grigory Kabatiansky    Russian Academy of Sciences (Russia)
- Roman Karasev    Moscow Institute of Physics and Technology (Russia)
- Gyula O. H. Katona    Hungarian Academy of Sciences (Hungary)
- Alex V. Kontorovich    Rutgers University (United States)
- Maxim Korolev    Steklov Mathematical Institute (Russia)
- Christian Krattenthaler    Universität Wien (Austria)
- Antanas Laurinčikas    Vilnius University (Lithuania)
- Vsevolod Lev    University of Haifa at Oranim (Israel)
- János Pach    EPFL Lausanne (Switzerland) and Rényi Institute (Hungary)
- Rom Pinchasi    Israel Institute of Technology – Technion (Israel)
- Alexander Razborov    Institut de Mathématiques de Luminy (France)
- Joël Rivat    Université d'Aix-Marseille (France)
- Tanguy Rivoal    Institut Fourier, CNRS (France)
- Damien Roy    University of Ottawa (Canada)
- Vladislav Salikhov    Bryansk State Technical University (Russia)
- Tom Sanders    University of Oxford (United Kingdom)
- Alexander A. Sapozhenko    Lomonosov Moscow State University (Russia)
- József Solymosi    University of British Columbia (Canada)
- Andreas Strömbergsson    Uppsala University (Sweden)
- Benjamin Sudakov    University of California, Los Angeles (United States)
- Jörg Thuswaldner    University of Leoben (Austria)
- Kai-Man Tsang    Hong Kong University (China)
- Maryna Viazovska    EPFL Lausanne (Switzerland)
- Barak Weiss    Tel Aviv University (Israel)

## PRODUCTION

- Silvio Levy    (Scientific Editor)  
production@msp.org

Cover design: Blake Knoll, Alex Scorpan and Silvio Levy

See inside back cover or [msp.org/moscow](http://msp.org/moscow) for submission instructions.

The subscription price for 2020 is US \$310/year for the electronic version, and \$365/year (+\$20, if shipping outside the US) for print and electronic. Subscriptions, requests for back issues and changes of subscriber address should be sent to MSP.

Moscow Journal of Combinatorics and Number Theory (ISSN 2640-7361 electronic, 2220-5438 printed) at Mathematical Sciences Publishers, 798 Evans Hall #3840, c/o University of California, Berkeley, CA 94720-3840 is published continuously online. Periodical rate postage paid at Berkeley, CA 94704, and additional mailing offices.

MJCNT peer review and production are managed by EditFlow® from MSP.

PUBLISHED BY  
 **mathematical sciences publishers**  
nonprofit scientific publishing  
<http://msp.org/>  
© 2020 Mathematical Sciences Publishers

Thematic Issue on  
Complexity

Provenance analysis for logic and games ERICH GRÄDEL and VAL TANNEN	203
Generalizations of $k$ -dimensional Weisfeiler–Leman stabilization ANUJ DAWAR and DANNY VAGNOZZI	229
On defining linear orders by automata BRUNO COURCELLE, IRÈNE DURAND and MICHAEL RASKIN	253
First-order definitions of subgraph isomorphism through the adjacency and order relations OLEG GRIGORYAN, MIKHAIL MAKAROV and MAKSIM ZHUKOVSKII	293
On limit points of spectra of first-order sentences with quantifier depth 4 YURY YAROVIKOV	303
Weakly distinguishing graph polynomials on addable properties JOHANN A. MAKOWSKY and VSEVOLOD RAKITA	333