

Model Theory

no. 2

vol. 2

2023

MAT

**An improved bound for regular decompositions of
3-uniform hypergraphs of bounded VC_2 -dimension**

Caroline Terry



An improved bound for regular decompositions of 3-uniform hypergraphs of bounded VC₂-dimension

Caroline Terry

A regular partition \mathcal{P} for a 3-uniform hypergraph $H = (V, E)$ consists of a partition $V = V_1 \cup \dots \cup V_t$ and for each $ij \in \binom{[t]}{2}$, a partition $K_2[V_i, V_j] = P_{ij}^1 \cup \dots \cup P_{ij}^\ell$ such that certain quasirandomness properties hold. The *complexity* of \mathcal{P} is the pair (t, ℓ) . In this paper we show that if a 3-uniform hypergraph H has VC₂-dimension at most k , then there is such a regular partition \mathcal{P} for H of complexity (t, ℓ) , where ℓ is bounded by a polynomial in the degree of regularity. This is a vast improvement on the bound arising from the proof of this regularity lemma in general, in which the bound generated for ℓ is of Wowzer type. This can be seen as a higher arity analogue of the efficient regularity lemmas for graphs and hypergraphs of bounded VC-dimension due to Alon–Fischer–Newman, Lovász–Szegedy, and Fox–Pach–Suk.

1. Introduction

Szemerédi’s regularity lemma is an important theorem with many applications in extremal combinatorics. The proof of the regularity lemma, which first appeared in the 70s [24], was well known to produce tower-type bounds in ϵ . The question of whether this type of bound is necessary was resolved in the late 90s by Gowers’ lower bound construction [10], which showed tower bounds are indeed required (see also [7; 17; 6]).

Hypergraph regularity was developed in the 2000s by Frankl, Gowers, Kohayakawa, Nagle, Rödl, Skokan, Schacht [9; 11; 12; 21; 20; 19], in order to prove a general counting lemma for hypergraphs. These types of regularity lemmas are substantially more complicated than prior regularity lemmas. In particular, a regular partition of a k -uniform hypergraph involves a sequence $\mathcal{P}_1, \dots, \mathcal{P}_{k-1}$, where \mathcal{P}_i is a collection of subsets $\binom{V}{i}$ such that certain quasirandomness properties hold for each \mathcal{P}_i relative to $\mathcal{P}_1, \dots, \mathcal{P}_{i-1}$. The proofs of these strong regularity lemmas produce Ackerman style bounds for the size of each \mathcal{P}_i . Given a function f , let $f^{(i)}$ denote the i -times iterate of f . We then define $\text{Ack}_1(x) = 2^x$, and for $i > 1$,

The author was partially supported by NSF grant DMS-2115518.

MSC2020: 03C45, 05D99.

Keywords: model theory, VC-dimension, regularity lemmas, hypergraph regularity.

$\text{Ack}_i(x) = \text{Ack}_{k-1}^{(i)}(x)$. The proofs of the strong regularity lemma for k -uniform hypergraphs produce bounds for the size of each \mathcal{P}_i of the form Ack_k . It was shown by Moshkovitz and Shapira [18] that this type of bound is indeed necessary for the size of \mathcal{P}_1 , which corresponds to the partition of the vertex set.

In the case of 3-uniform hypergraphs, a decomposition in this sense consists of a partition $\mathcal{P}_1 = \{V_1, \dots, V_t\}$ of V , and a set

$$\mathcal{P}_2 = \{P_{ij}^\alpha : ij \in \binom{[t]}{2}, \alpha \in [\ell]\},$$

where for each $ij \in \binom{[t]}{2}$, $P_{ij}^1 \cup \dots \cup P_{ij}^\ell$ is a partition of $K_2[V_i, V_j]$. The *complexity* of \mathcal{P} is the pair (t, ℓ) . We give a formal statement of the regularity lemma for 3-graphs here for reference, and refer the reader to [Section 2B](#) for the precise definitions involved. The version stated below is a refinement of a regularity lemma due to Gowers [12] (for more details see [Section 2B](#)).

Theorem 1.1 (strong regularity lemma for 3-graphs). *For all $\epsilon_1 > 0$, and every function $\epsilon_2 : \mathbb{N} \rightarrow (0, 1]$, there exist positive integers T_0, L_0 , and n_0 such that for any 3-graph $H = (V, E)$ on $n \geq n_0$ vertices, there exists a $\text{dev}_{2,3}(\epsilon_1, \epsilon_2(\ell))$ -regular, $(t, \ell, \epsilon_1, \epsilon_2(\ell))$ -decomposition \mathcal{P} for H with $t \leq T_0$ and $\ell \leq L_0$.*

In [Theorem 1.1](#), the parameter T_0 is the bound for t , the size of the vertex partition, and L_0 is the bound for ℓ , the size of the partition of $K_2[V_i, V_j]$, for each $ij \in \binom{[t]}{2}$. The proof of [Theorem 1.1](#) generates a Wowzer (i.e., Ack_3) type bound for both t and ℓ . Moshkovitz and Shapira showed in [18] that there exist 3-uniform hypergraphs requiring a Wowzer type bound for the size of t in [Theorem 1.1](#). Less attention has been paid to the form of the bound L_0 , and it remains open whether this is necessarily of Wowzer type. In recent work of the author and Wolf [26], the partition \mathcal{P}_2 plays a crucial role in the proof of a strong version of [Theorem 1.1](#) in a combinatorially tame setting. This work suggests that understanding the form of the bound for ℓ is also an interesting problem.

In the case of graphs, it was shown that dramatic improvements on the bounds in Szemerédi's regularity lemma can be obtained under the hypothesis of bounded VC-dimension. In particular, Alon, Fischer, and Newman [1] showed that if a bipartite graph G has VC-dimension less than k , then it has an ϵ -regular partition of size at most $(k/\epsilon)^{O(k)}$. Lovász and Szegedy [16] extended this to all graphs of VC-dimension less than k , with a bound of the form $\epsilon^{-O(k^2)}$. Fox, Pach, and Suck [8] strengthened the bound to one of the form $c(\epsilon^{2k-1})$, and extended these results to hypergraphs of bounded VC-dimension. Related results were obtained with weaker polynomial bounds by Chernikov and Starchenko [3].

In this paper we prove an analogous theorem in the context of strong regularity for 3-uniform hypergraphs, where VC-dimension is replaced by a higher arity analogue called VC₂-dimension.

Definition 1.2. Suppose $H = (V, E)$ is a 3-graph. The VC₂-dimension of H , $\text{VC}_2(H)$, is the largest integer k so that there exist $a_1, \dots, a_k, b_1, \dots, b_k \in V$ and $c_S \in V$ for each $S \subseteq [k]^2$, such that $a_i b_j c_S \in E$ if and only if $(i, j) \in S$.

The notion of VC₂-dimension was first introduced by Shelah [22], who also studied it in the context of groups [23]. It was later shown to have nice model-theoretic characterizations by Chernikov, Palacin, and Takeuchi [5] to have further natural connections to groups and fields by Hempel and Chernikov [15; 2], and to have applications in combinatorics by the author [25].

Using infinitary techniques, Chernikov and Towsner [4] proved a strong regularity lemma for 3-uniform hypergraphs of bounded VC₂-dimension without explicit bounds (in fact they proved results for k -uniform hypergraphs of bounded VC _{$k-1$} -dimension). Similar results were proved by the author and Wolf [26] in the 3-uniform case with Wowzer type bounds. In this paper, we show that 3-uniform hypergraphs of uniformly bounded VC₂-dimension have regular decompositions with vastly improved bounds on the size of ℓ ; in particular, ℓ can be guaranteed to be polynomial in size, rather than Wowzer. We include the formal statement of our main theorem below, and refer the reader to the next section for details on the definitions involved.

Theorem 1.3. *For all $k \geq 1$, there are $\epsilon_1^* > 0$ and $\epsilon_2^* : \mathbb{N} \rightarrow (0, 1]$ such that the following holds. Suppose $0 < \epsilon_1 < \epsilon_1^*$ and $\epsilon_2 : \mathbb{N} \rightarrow (0, 1]$ satisfies $0 < \epsilon_2(x) < \epsilon_2^*(x)$ for all $x \in \mathbb{N}$. There is $T = T(\epsilon_1, \epsilon_2)$ such that every sufficiently large 3-graph $H = (V, E)$ has a $\text{dev}_{2,3}(\epsilon_1, \epsilon_2(\ell))$ -regular $(t, \ell, \epsilon_1, \epsilon_2(\ell))$ -decomposition with $\ell \leq \epsilon_1^{-O_k(k)}$ and $t \leq T$.*

The bound T in Theorem 1.3 is generated from an application of Theorem 1.1, and is also of Wowzer type (see Theorem 3.1 for a more precise statement regarding this). The regular partition in Theorem 1.3 has the additional property that the regular triads have edge densities near 0 or 1, which also occurs in the results from [4; 26]. The ingredients in the proof of Theorem 1.3 include the improved regularity lemma for 3-graphs of bounded VC₂-dimension from [26], a method of producing quotient graphs from regular partitions of 3-graphs developed in [26], and ideas from [8] for producing weak regular partitions of hypergraphs of bounded VC-dimension.

The fact that the bound for ℓ can be brought all the way down to polynomial in Theorem 1.3 is somewhat surprising, given that the proof for arbitrary hypergraphs yields a Wowzer bound. This raises the question of what the correct form of the bound is, in general, for ℓ . The author conjectures it is at least a tower function (i.e., Ack₂).

It was conjectured in [4] that the bound for t can also be made sub-Wowzer under the assumption of bounded VC₂-dimension, however, the author has been unable to prove this is the case. This leaves the following open problem.

Problem 1.4. Given a fixed integer $k \geq 1$, are there arbitrarily large 3-uniform hypergraphs of VC₂-dimension at most k which require Wowzer type bounds for T_0 in [Theorem 1.1](#)?

2. Preliminaries

In this section we cover the requisite preliminaries, including graph and hypergraph regularity ([Section 2B](#)), VC and VC₂-dimension ([Sections 2C, 2E, and 2F](#)), auxiliary graphs defined from regular decompositions of 3-graphs ([Section 2D](#)), and basic lemmas around regularity and counting ([Section 2G](#)).

2A. Notation. We include here some basic notation needed for the other preliminary sections. Given a set V and $k \geq 1$, let

$$\binom{V}{k} = \{X \subseteq V : |X| = k\}.$$

A k -uniform hypergraph is a pair (V, E) where $E \subseteq \binom{V}{k}$. For a k -uniform hypergraph G , $V(G)$ denotes the vertex set of V and $E(G)$ denotes the edge set of G . Throughout the paper, all vertex sets are assumed to be finite.

When $k = 2$, we refer to a k -uniform hypergraph as simply a *graph*. When $k = 3$, we refer to a k -uniform hypergraph as a *3-graph*.

Given distinct elements x, y , we write xy for the set $\{x, y\}$. Similarly, for distinct x, y, z , we write xyz for the set $\{x, y, z\}$. Given sets X, Y, Z , we set

$$K_2[X, Y] = \{xy : x \in X, y \in Y, x \neq y\} \quad \text{and} \\ K_3[X, Y, Z] = \{xyz : x \in X, y \in Y, z \in Z, x \neq y, y \neq z, x \neq z\}.$$

If $G = (V, E)$ is a graph and $X, Y \subseteq V$ are disjoint, we let $G[X, Y]$ be the bipartite graph $(X \cup Y, E \cap K_2[X, Y])$.

Given a k -uniform hypergraph $G = (V, E)$, $1 \leq i < k$, and $e \in \binom{V}{i}$, set

$$N_E(e) = \left\{ e' \in \binom{V}{k-i} : e \cup e' \in E \right\}.$$

A *bipartite edge-colored graph* is a tuple $G = (A \cup B, E_0, E_1, \dots, E_i)$, where $i > 1$ and $K_2[A, B] = E_0 \sqcup E_1 \sqcup \dots \sqcup E_i$. In this case, given $u \in \{0, 1, \dots, i\}$ and $x \in A \cup B$, we let $N_{E_u}(x) = \{y \in A \cup B : xy \in E_u\}$. Similarly, a *tripartite edge-colored 3-graph* is a tuple $G = (A \cup B \cup C, E_0, E_1, \dots, E_i)$, where $i > 1$ and $K_3[A, B, C] = E_0 \sqcup E_1 \sqcup \dots \sqcup E_i$. In this case, given $u \in \{0, 1, \dots, i\}$ and $x, y \in V := A \cup B \cup C$, we let $N_{E_u}(x) = \{uv \in \binom{V}{2} : xuv \in E_u\}$ and $N_{E_u}(xy) = \{v \in V : xuv \in E_u\}$.

For two functions $f_1, f_2 : \mathbb{N} \rightarrow (0, 1]$, we write $f_1 < f_2$ to denote that $f_1(x) < f_2(x)$ for all $x \in \mathbb{N}$. For real numbers r_1, r_2 and $\epsilon > 0$, we write $r_1 = r_2 \pm \epsilon$ to denote

that $r_1 \in (r_2 - \epsilon, r_2 + \epsilon)$. Given a natural number $n \geq 1$, $[n] = \{1, \dots, n\}$. An equipartition of a set V is a partition $V = V_1 \cup \dots \cup V_t$ such that for each $1 \leq i, j \leq t$, we have $||V_i| - |V_j|| \leq 1$.

2B. Regularity. In this section we define graph regularity, as well as a strong notion of regularity for 3-graphs. We state our definitions in terms of the quasirandomness notion known as “dev”, which is one of three notions of quasirandomness which are now known to be equivalent, the other two being “oct” and “disc”. For more details on these and the equivalences, we refer the reader to [19].

We begin a notion of quasirandomness for graphs.

Definition 2.1. Suppose $B = (U \cup W, E)$ is a bipartite graph, and $|E| = d_B |U| |W|$. We say B has $\text{dev}_2(\epsilon, d)$ if $d_B = d \pm \epsilon$ and

$$\sum_{u_0, u_1 \in U} \sum_{w_0, w_1 \in W} \prod_{i \in \{0,1\}} \prod_{j \in \{0,1\}} g(u_i, w_j) \leq \epsilon |U|^2 |W|^2,$$

where $g(u, v) = 1 - d_B$ if $uv \in E$ and $g(u, v) = -d_B$ if $uv \notin E$.

We now define a generalization of Definition 2.1 to 3-graphs due to Gowers [11]. If $G = (V, E)$ is a graph, let $K_3^{(2)}(G)$ denote the set of triples from V forming a triangle in G , i.e.,

$$K_3^{(2)}(G) := \left\{ xyz \in \binom{V}{3} : xy, yz, xz \in E \right\}.$$

Now given a 3-graph $H = (V, R)$ on the same vertex set, we say that G *underlies* H if $R \subseteq K_3^{(2)}(G)$.

Definition 2.2. Assume $\epsilon_1, \epsilon_2 > 0$, $H = (V, E)$ is a 3-graph, $G = (U \cup W \cup Z, E)$ is a 3-partite graph underlying H , and $|E| = d_3 |K_3^{(2)}(G)|$. We say that (H, G) has $\text{dev}_{2,3}(\epsilon_1, \epsilon_2)$ if there is $d_2 \in (0, 1)$ such that $G[U, W]$, $G[U, Z]$, and $G[W, Z]$ each have $\text{dev}_2(\epsilon_2, d_2)$, and

$$\sum_{u_0, u_1 \in U} \sum_{w_0, w_1 \in W} \sum_{z_0, z_1 \in Z} \prod_{(i,j,k) \in \{0,1\}^3} h_{H,G}(u_i, w_j, z_k) \leq \epsilon_1 d_2^{12} |U|^2 |W|^2 |Z|^2,$$

where

$$h_{H,G}(x, y, z) = \begin{cases} 1 - d_3 & \text{if } xyz \in E \cap K_3^{(2)}(G), \\ -d_3 & \text{if } xyz \in K_3^{(2)}(G) \setminus E, \\ 0 & \text{if } xyz \notin K_3^{(2)}(G). \end{cases}$$

For the reader unfamiliar with hypergraph regularity, we note that in the notation of Definition 2.2, $d_2^{12} |U|^2 |W|^2 |Z|^2$ is approximately the number of tuples $(u_0, u_1, w_0, w_1, z_0, z_1) \in U^2 \times W^2 \times Z^2$ with $u_i w_j z_k \in K_3^{(2)}(G)$ for each $(i, j, k) \in \{0, 1\}^3$ (this is a consequence of the graph counting lemma and the

assumption that $G[U, W]$, $G[U, Z]$, and $G[W, Z]$ have $\text{dev}_2(\epsilon_2, d_2)$). Therefore, the first displayed equation in Definition 2.2 is bounding the quantity

$$\sum_{u_0, u_1 \in U} \sum_{w_0, w_1 \in W} \sum_{z_0, z_1 \in Z} \prod_{(i, j, k) \in \{0, 1\}^3} h_{H, G}(u_i, w_j, z_k)$$

in terms of total number of tuples $(u_0, u_1, w_0, w_1, z_0, z_1) \in U^2 \times W^2 \times Z^2$, where $\prod_{(i, j, k) \in \{0, 1\}^3} h_{H, G}(u_i, w_j, z_k)$ is nonzero.

We now define a (t, ℓ) -decomposition for a vertex set V , which partitions V , as well as pairs from V .

Definition 2.3. Let V be a vertex set and $t, \ell \in \mathbb{N}^{>0}$. A (t, ℓ) -decomposition \mathcal{P} for V consists of a partition $\mathcal{P}_1 = \{V_1 \cup \dots \cup V_t\}$ of V , and for each $1 \leq i \neq j \leq t$, a partition $K_2[V_i, V_j] = P_{ij}^1 \cup \dots \cup P_{ij}^\ell$. We let $\mathcal{P}_2 = \{P_{ij}^\alpha : ij \in \binom{[t]}{2}, \alpha \leq \ell\}$.

A triad of \mathcal{P} is a 3-partite graph of the form

$$G_{\alpha, \beta, \gamma}^{ijk} := (V_i \cup V_j \cup V_k, P_{ij}^\alpha \cup P_{ik}^\beta \cup P_{jk}^\gamma),$$

for some $ijk \in \binom{[t]}{3}$ and $\alpha, \beta, \gamma \leq \ell$. Let $\text{Triads}(\mathcal{P})$ denote the set of all triads of \mathcal{P} , and observe that $\{K_3^{(2)}(G) : G \in \text{Triads}(\mathcal{P})\}$ partitions the set of triples $xyz \in \binom{V}{3}$ which are in distinct elements of \mathcal{P}_1 .

For a 3-graph $H = (V, R)$, a decomposition \mathcal{P} of V , and $G \in \text{Triads}(\mathcal{P})$, define $H|G := (V(G), R \cap K_3^{(2)}(G))$. Note that G always underlies $H|G$.

Definition 2.4. Given a 3-graph $H = (V, R)$, a decomposition \mathcal{P} of V , and $G \in \text{Triads}(\mathcal{P})$, we say G has $\text{dev}_{2,3}(\epsilon_1, \epsilon_2)$ with respect to H if $(H|G, G)$ has $\text{dev}_{2,3}(\epsilon_1, \epsilon_2)$.

To define a regular decomposition for a 3-graph, we need one more notion, namely that of an “equitable” decomposition.

Definition 2.5. We say that \mathcal{P} is a $(t, \ell, \epsilon_1, \epsilon_2)$ -decomposition if $\mathcal{P}_1 = \{V_1, \dots, V_t\}$ is an equipartition and for at least $(1 - \epsilon_1) \binom{|V|}{2}$ many $xy \in \binom{V}{2}$, there is some $P_{ij}^\alpha \in \mathcal{P}_2$ containing xy such that $(V_i \cup V_j, P_{ij}^\alpha)$ has $\text{dev}_2(\epsilon_2, 1/\ell)$.

Definition 2.6. Suppose that $H = (V, E)$ is a 3-graph and \mathcal{P} is a $(t, \ell, \epsilon_1, \epsilon_2)$ -decomposition of V . We say that \mathcal{P} is $\text{dev}_{2,3}(\epsilon_1, \epsilon_2)$ -regular for H if for all but at most $\epsilon_1 n^3$ many triples $xyz \in \binom{V}{3}$, the unique $G \in \text{Triads}(\mathcal{P})$ with $xyz \in K_3^{(2)}(G)$ satisfies $\text{dev}_{2,3}(\epsilon_1, \epsilon_2)$ with respect to H .

We can now restate the regularity lemma for $\text{dev}_{2,3}$ -quasirandomness.

Theorem 2.7. For all $\epsilon_1 > 0$, every function $\epsilon_2 : \mathbb{N} \rightarrow (0, 1]$, and every $\ell_0, t_0 \geq 1$, there exist positive integers $T_0 = T_0(\epsilon_1, \epsilon_2, t_0, \ell_0)$ and $L_0 = L_0(\epsilon_1, \epsilon_2, t_0, \ell_0)$, such that for every sufficiently large 3-graph $H = (V, E)$, there exists a $\text{dev}_{2,3}(\epsilon_1, \epsilon_2(\ell))$ -regular, $(t, \ell, \epsilon_1, \epsilon_2(\ell))$ -decomposition \mathcal{P} for H with $t_0 \leq t \leq T_0$ and $\ell_0 \leq \ell \leq L_0$.

This theorem was first proved in a slightly different form by Gowers in [11]. In particular, in [11], the partition of the pairs \mathcal{P}_2 is not required to be equitable as it is in Theorem 2.7. Theorem 2.7 as stated appears in [19], where it is pointed out that the additional equitability requirement can be obtained using techniques from [9].

2C. VC-dimension. In this subsection we give some preliminaries around VC and VC₂-dimension. We begin by defining VC-dimension.

Given a set V , $\mathcal{F} \subseteq \mathcal{P}(V)$, and $X \subseteq V$, let $|\mathcal{F} \cap X| := \{F \cap X : F \in \mathcal{F}\}$. We say that X is *shattered* by \mathcal{F} if $|\mathcal{F} \cap X| = 2^{|X|}$. The *VC-dimension* of \mathcal{F} is then defined to be the size of the largest subset of V which is shattered by \mathcal{F} .

For a graph $G = (V, E)$, the *VC-dimension* of G is the VC-dimension of the set system $\{N_E(x) : x \in V\} \subseteq \mathcal{P}(V)$. We now give a simple recharacterization of this. Given $k \geq 1$, let $A_k = \{a_i : i \in [k]\}$, and $C_{\mathcal{P}([k])} = \{c_S : S \subseteq [k]\}$.

Definition 2.8. For $k \geq 1$, define $U(k)$ to be the bipartite graph $(A_k \cup C_{\mathcal{P}([k])}, E)$, where $E = \{a_i c_S : i \in S\}$.

Then it is well known that a graph G has VC-dimension at least k if and only if there is a map $f : V(U(k)) \rightarrow V(G)$ so that for all $a \in A_k$ and $c \in C_{\mathcal{P}([k])}$, $ab \in E(U(k))$ if and only if $f(a)f(b) \in E(G)$.

2D. Encodings. In this subsection, we define an auxiliary edge-colored graph associated to a regular decomposition of a 3-graph. We then state a result from [26] which shows that encodings of $U(k)$ cannot occur when the auxiliary edge-colored graph arises from a regular decomposition of a 3-graph with VC₂-dimension less than k .

Definition 2.9. Suppose $\epsilon_1, \epsilon_2 > 0$, $\ell, t \geq 1$, V is a set, and \mathcal{P} is a $(t, \ell, \epsilon_1, \epsilon_2)$ -decomposition for V consisting of $\mathcal{P}_1 = \{V_i : i \in [t]\}$ and $\mathcal{P}_2 = \{P_{ij}^\alpha : ij \in \binom{[t]}{2}, \alpha \leq \ell\}$. Define

$$\begin{aligned} \mathcal{P}_{\text{cnr}} &= \left\{ P_{ij}^\alpha P_{ik}^\beta : ijk \in \binom{[t]}{3}, \alpha, \beta \leq \ell, \text{ and } P_{ij}^\alpha, P_{ij}^\beta \text{ satisfy } \text{dev}_2(\epsilon_2, 1/\ell) \right\}, \\ \mathcal{P}_{\text{edge}} &= \{P_{ij}^\alpha \in \mathcal{P}_2 : P_{ij}^\alpha \text{ satisfies } \text{dev}_2(\epsilon_2, 1/\ell)\}. \end{aligned}$$

In the above, *cnr* stands for ‘‘corner’’. Observe that for each $P_{ij}^\alpha \in \mathcal{P}_{\text{edge}}$ and $P_{uv}^\beta P_{uw}^\gamma \in \mathcal{P}_{\text{cnr}}$, if $\{v, w\} = \{i, j\}$, then the pair $(P_{ij}^\alpha, P_{uv}^\beta P_{uw}^\gamma)$ corresponds to a triad from \mathcal{P} , namely G_{ijs}^{uvw} .

Definition 2.10. Suppose $\epsilon_1, \epsilon_2 > 0$, $\ell, t \geq 1$, $H = (V, E)$ is a 3-graph, and \mathcal{P} is a $(t, \ell, \epsilon_1, \epsilon_2)$ -decomposition for V . Define

$$\begin{aligned} \mathbf{E}_0 &= \{P_{ij}^\alpha (P_{jk}^\beta P_{ik}^\gamma) \in K_2[\mathcal{P}_{\text{edge}}, \mathcal{P}_{\text{cnr}}] : |E \cap K_3^{(2)}(G_{ijk}^{\alpha\beta\gamma})| < \frac{1}{2} |K_3^{(2)}(G_{ijk}^{\alpha\beta\gamma})|\}, \\ \mathbf{E}_1 &= \{P_{ij}^\alpha (P_{jk}^\beta P_{ik}^\gamma) \in K_2[\mathcal{P}_{\text{edge}}, \mathcal{P}_{\text{cnr}}] : |E \cap K_3^{(2)}(G_{ijk}^{\alpha\beta\gamma})| \geq \frac{1}{2} |K_3^{(2)}(G_{ijk}^{\alpha\beta\gamma})|\}, \text{ and} \\ \mathbf{E}_2 &= K_2[\mathcal{P}_{\text{edge}}, \mathcal{P}_{\text{cnr}}] \setminus (\mathbf{E}_1 \cup \mathbf{E}_0). \end{aligned}$$

Note that [Definition 2.10](#) gives us a natural bipartite edge-colored graph with vertex set $\mathcal{P}_{\text{edge}} \cup \mathcal{P}_{\text{cnr}}$ and edge sets given by $\mathbf{E}_0, \mathbf{E}_1, \mathbf{E}_2$. The author and Wolf showed in [\[26\]](#) that these auxiliary edge-colored graphs are useful for understanding 3-graphs of bounded VC₂-dimension. To explain why, we require the following notion of an “encoding”.

Definition 2.11. Let $\epsilon_1, \epsilon_2 > 0$ and $t, \ell \geq 1$. Suppose $R = (A \cup B, E_R)$ is a bipartite graph, $H = (V, E)$ is a 3-graph, and \mathcal{P} is a $(t, \ell, \epsilon_1, \epsilon_2)$ -decomposition of V . An (A, B) -encoding of R in (H, \mathcal{P}) consists of a pair of functions (g, f) , where $g : A \rightarrow \mathcal{P}_{\text{cnr}}$ and $f : B \rightarrow \mathcal{P}_{\text{edge}}$ are such that the following hold for some $j_0 k_0 \in \binom{[t]}{2}$:

- (1) $\text{Im}(f) \subseteq \{P_{j_0 k_0}^\alpha : \alpha \leq \ell\}$, and $\text{Im}(g) \subseteq \{P_{i j_0}^\beta P_{i k_0}^\gamma : i \in [t], \beta, \gamma \leq \ell\}$.
- (2) For all $a \in A$ and $b \in B$, if $ab \in E_R$, then $g(a)f(b) \in \mathbf{E}_1$, and if $ab \notin E_R$, then $g(a)f(b) \in \mathbf{E}_0$.

An *encoding of $U(k)$* will always mean an $(A_k, C_{\mathcal{P}([k])})$ -encoding of $U(k)$. In [\[26\]](#), we proved the following proposition connecting encodings of $U(k)$ and VC₂-dimension (see [Theorem 6.5\(2\)](#) in [\[26\]](#)).

Proposition 2.12. *For all $k \geq 1$, there are $\epsilon_1 > 0$ and $\epsilon_2 : \mathbb{N} \rightarrow (0, 1]$ such that for all $t, \ell \geq 1$, there is N such that the following hold. Suppose $H = (V, E)$ is a 3-graph with $|V| \geq N$, and \mathcal{P} is a $\text{dev}_{2,3}(\epsilon_2(\ell), \epsilon_1)$ -regular $(t, \ell, \epsilon_1, \epsilon_2(\ell))$ -decomposition of V . If there exists an encoding of $U(k)$ in (H, \mathcal{P}) , then H has k -IP₂.*

Moreover, there is a constant $C = C(k)$ so that $\epsilon_1 = (1/2)^C$.

We remark here that [Proposition 2.12](#) is actually proved in [\[26\]](#) for an equivalent notion of quasirandomness called $\text{disc}_{2,3}$, and without the final “Moreover” statement regarding the quantitative form for ϵ_1 (see [Proposition 5.6](#) in [\[26\]](#)). Tracing the bounds in the proof of [Proposition 5.6](#) in [\[26\]](#), one finds that ϵ_1 has the form $\mu = \mu(\epsilon_1, k)$, where μ comes from a version of the counting lemma (see [Theorem 3.1](#) in [\[26\]](#)). An explicit value for this μ is unclear, as the proof of the counting lemma for $\text{disc}_{2,3}$ passes through its equivalence with $\text{oct}_{2,3}$, and then the counting lemma for $\text{oct}_{2,3}$. The author has not found proofs of these results in the literature which are explicit in the parameters (see [Corollary 2.3](#) in [\[19\]](#)). It seems that one could produce such an explicit result from [\[19\]](#) and [\[14\]](#) with some effort, however, we have instead chosen to side-step the issue by working with the quasirandomness notion dev , rather than disc .

In particular, all the ingredients used to prove [Proposition 5.6](#) of [\[26\]](#) have well known analogues for dev . By running the same arguments as in [\[26\]](#) using dev rather than disc , one obtains [Proposition 2.12](#) as stated. The additional “Moreover” statement about the explicit form for ϵ_1 then arises from the fact that there is a

proof of the counting lemma for $\text{dev}_{2,3}$ which is explicit in the parameters (see [11, Theorem 6.8]).

2E. Haussler's packing lemma. We will be applying techniques for proving improved regularity lemmas for graphs and hypergraphs of bounded VC-dimension to the edge-colored auxiliary graphs defined in the previous subsection. In particular, we will use ideas from the proof of Theorem 1.3 in [8]. We begin by describing the relevant result from VC-theory, namely Haussler's packing lemma.

Suppose V is a set and $\mathcal{F} \subseteq \mathcal{P}(V)$. We say that a subset $\mathcal{X} \subseteq \mathcal{F}$ is δ -separated if for all distinct $X, X' \in \mathcal{X}$, $|X \Delta X'| > \delta$. The following packing lemma, due to Haussler, shows that if \mathcal{F} has bounded VC-dimension, the size of a δ -separated family cannot be too large [13].

Theorem 2.13 (Haussler's packing lemma). *Suppose $\mathcal{F} \subseteq \mathcal{P}(V)$, where $|V| = n$ and \mathcal{F} has VC-dimension at most k . Then the maximal size of a δ -separated subcollection of \mathcal{F} is at most $c_1(n/\delta)^k$, for some constant $c_1 = c_1(k)$.*

We will apply Theorem 2.13 in the setting of edge-colored graphs. This technique is inspired by the proof of Theorem 1.3 in [8].

Suppose $G = (A \cup B, E_0, E_1, E_2)$ is a bipartite edge-colored graph. We say that G has an E_0/E_1 -copy of $U(k)$ if there are $v_1, \dots, v_k \in A$ and for each $S \subseteq [k]$ a vertex $w_S \in B$ such that $i \in S$ implies $v_i w_S \in E_1$ and $i \notin S$ implies $v_i w_S \in E_0$. Given $a, a' \in A$ and $\delta > 0$, write $a \sim_\delta a'$ if for each $u \in \{0, 1, 2\}$, $|N_{E_u}(a) \Delta N_{E_u}(a')| \leq \delta |B|$. Our main application of Theorem 2.13 is the following lemma.

Lemma 2.14. *Suppose $k \geq 1$ and $c_1 = c_1(k)$ is as in Theorem 2.13. Suppose $d \geq 1$ and $\delta, \epsilon > 0$ satisfy $\epsilon \leq c_1^{-2}(\delta/8)^{2k+2}$. Assume $G = (A \cup B, E_0, E_1, E_2)$ is a bipartite edge-colored graph, and assume there is no E_0/E_1 -copy of $U(k)$ in G , and that $|E_2| \leq \epsilon |A| |B|$.*

Then there is an integer $m \leq 2c_1(\delta/8)^{-k}$, vertices $x_1, \dots, x_m \in A$, and a set $U \subseteq A$ with $|U| \leq \sqrt{\epsilon} |A|$, so that for all $a \in A \setminus U$, $|N_{E_2}(a)| \leq \sqrt{\epsilon} |B|$ and there is some $1 \leq i \leq m$ so that $a \sim_\delta x_i$.

Proof. Let $U = \{v \in A : |N_{E_2}(v)| \geq \sqrt{\epsilon} |B|\}$. Since $|E_2| \leq \epsilon |A| |B|$, we know that $|U| \leq \sqrt{\epsilon} |A|$. Let $A' = A \setminus U$. Let m be maximal such that there exist $x_1, \dots, x_m \in A'$, so that $\{N_{E_1}(x_i) : i \in [m]\}$ is a $\delta/2$ -separated family of sets on B . We show $m \leq 2c_1(\delta/8)^{-k}$.

Suppose towards a contradiction that $m \geq \lceil 2c_1(\delta/8)^{-k} \rceil$. Let

$$B' = B \setminus \left(\bigcup_{i=1}^m E_2(x_i) \right),$$

and let $\mathcal{F} := \{N_{E_1}(x_i) \cap B' : i \in [m]\}$. Notice $|B \setminus B'| \leq m\sqrt{\epsilon}|B|$. We claim that \mathcal{F} is $\delta/4$ -separated. Consider $1 \leq i \neq j \leq m$. Then we know that

$$\begin{aligned} |N_{E_1}(x_i) \Delta N_{E_1}(x_j) \cap B'| &\geq |N_{E_1}(x_i) \Delta N_{E_1}(x_j)| - m\sqrt{\epsilon}|B| \\ &\geq |B|(\delta/2 - m\sqrt{\epsilon}) \\ &\geq |B|\delta/4, \end{aligned}$$

where the last inequality is by our assumptions on δ, ϵ . By [Theorem 2.13](#), \mathcal{F} shatters a set of size k . By construction, for each $1 \leq i \leq m$, $B' \setminus N_{E_1}(x_i) \subseteq N_{E_0}(x_i)$. Consequently, we must have that there exists an E_0/E_1 -copy of $U(k)$ in G , a contradiction.

Thus, $m \leq 2c_1(\delta/8)^{-k}$. For all $a \in A \setminus U$, we know that $|N_{E_2}(a)| \leq \sqrt{\epsilon}|B|$, and there is some $1 \leq i \leq m$ so that $|N_{E_1}(a) \Delta N_{E_1}(x_i)| \leq \delta|B|/2$. We claim that $a \sim_\delta x_i$. We already know that $|N_{E_1}(a) \cap N_{E_1}(x_i)| \leq \delta|B|$. Since a, x_i are both in A' , we have

$$|N_{E_2}(a) \Delta N_{E_2}(x_i)| \leq |N_{E_2}(a)| + |N_{E_2}(x_i)| \leq 2\sqrt{\epsilon}|B| < \delta|B|/2.$$

Combining these facts, we have that

$$|N_{E_0}(a) \Delta N_{E_0}(x_i)| \leq |N_{E_2}(a)| + |N_{E_2}(x_i)| + |N_{E_1}(a) \Delta N_{E_1}(x_i)| \leq \delta|B|.$$

Thus $a \sim_\delta x_i$, as desired. \square

2F. Tame regularity for 3-graphs of bounded VC_2 -dimension. In this subsection we state the tame regularity lemma for 3-graphs of bounded VC_2 -dimension from [\[26\]](#).

Definition 2.15. Suppose $H = (V, E)$ is a 3-graph with $|V| = n$ and $\mu > 0$. Suppose $t, \ell \geq 1$ and \mathcal{P} is a (t, ℓ) -decomposition of V . We say that \mathcal{P} is μ -homogeneous with respect to H if at least $(1 - \mu)\binom{n}{3}$ triples $xyz \in \binom{V}{3}$ satisfy the following: there is some $G \in \text{Triads}(\mathcal{P})$ such that $xyz \in K_3^{(2)}(G)$ and either

$$|E \cap K_3^{(2)}(G)| \leq \mu|K_3^{(2)}(G)| \quad \text{or} \quad |E \cap K_3^{(2)}(G)| \geq (1 - \mu)|K_3^{(2)}(G)|.$$

Given a 3-graph $H = (V, E)$ and a $(t, \ell, \epsilon_1, \epsilon_2)$ -decomposition \mathcal{P} of V , we say that \mathcal{P} is μ -homogeneous with respect to H if at least $(1 - \mu)\binom{|V|}{3}$ triples $xyz \in \binom{V}{3}$ are in a μ -homogeneous triad of \mathcal{P} . We have the following theorem from [\[26\]](#).

Theorem 2.16. For all $k \geq 1$, there are $\epsilon_1^* > 0$, $\epsilon_2^* : \mathbb{N} \rightarrow (0, 1]$, and a function $f : (0, 1] \rightarrow (0, 1]$ with $\lim_{x \rightarrow 0} f(x) = 0$ such that the following hold.

Suppose $t_0, \ell_0 \geq 1$, $0 < \epsilon_1 < \epsilon_1^*$, and $\epsilon_2 : \mathbb{N} \rightarrow (0, 1]$ satisfies $\epsilon_2 < \epsilon_2^*$. Let N, T , and L be as in [Theorem 2.7](#) for $\epsilon_1, \epsilon_2, t_0, \ell_0$. Suppose $H = (V, E)$ is a 3-graph with $|V| \geq N$ and $\text{VC}_2(H) < k$. Then there exist $t_0 \leq t \leq T$, $\ell_0 \leq \ell \leq L$,

and a $(t, \ell, \epsilon_1, \epsilon_2(\ell))$ -decomposition of V which is $\text{dev}_{2,3}(\epsilon_1, \epsilon_2(\ell))$ -regular and $f(\epsilon_1)$ -homogeneous with respect to H .

Moreover, f may be taken to have the form $x^{1/D}$, where $D \geq 1$ depends only on k .

Since the bounds in [Theorem 2.16](#) come from [Theorem 2.7](#), they are of Wowzer type. We also note that the proof of [Theorem 2.16](#) in fact guarantees something slightly stronger, namely that every $\text{dev}_{2,3}(\epsilon_1, \epsilon_2(\ell))$ -regular triad of \mathcal{P} is $f(\epsilon_1)$ -homogeneous.

We remark here that [Theorem 2.16](#) was proved in [\[26\]](#) for the notion of $\text{disc}_{2,3}$ rather than $\text{dev}_{2,3}$, and without the “moreover” statement regarding the form of the function f (see [Proposition 3.2](#) in [\[26\]](#)). Examination of the proof of [Proposition 3.2](#) in [\[26\]](#) shows that the function f depends on k and a version of the counting lemma for 3-graphs (namely [Theorem 3.1](#) in [\[26\]](#)). An explicit expression for $f(x)$ in [Proposition 3.2](#) of [\[26\]](#) would thus require a version of the counting lemma for $\text{disc}_{2,3}$ which is explicit in the parameters. However, one can rerun all the arguments in [\[26\]](#) using the quasirandomness notion $\text{dev}_{2,3}$ in place of $\text{disc}_{2,3}$ to obtain [Theorem 2.16](#) as stated. In this case, an explicit expression for f can be obtained using the counting lemma for $\text{dev}_{2,3}$ (see also the discussion following [Proposition 2.12](#)).

2G. Other preliminaries. In this subsection we give several lemmas, most of which are basic facts about regularity and counting. First, we will use the following version of the triangle counting lemma.

Proposition 2.17 (counting lemma). *Suppose $\epsilon, d > 0$. Let $G = (A \cup B \cup C, E)$ be a 3-partite graph such that each of $G[A, B]$, $G[B, C]$, and $G[A, C]$ has $\text{dev}_2(\epsilon, d)$. Then*

$$||K_3^{(2)}(G) - d^3|A||B||C|| \leq 4\epsilon^{1/4}|A||B||C|.$$

For a proof, see [\[11, Lemma 3.4\]](#). The following symmetry lemma was proved in [\[26\]](#) (see [Lemma 4.9](#) there).

Lemma 2.18 (symmetry lemma). *For all $0 < \epsilon < \frac{1}{4}$ there is n such that the following holds. Suppose $G = (U \cup W, E)$ is a bipartite graph, $|U|, |W| \geq n$, and $U' \subseteq U$, $W' \subseteq W$ satisfy $|U'| \geq (1 - \epsilon)|U|$ and $|W'| \geq (1 - \epsilon)|W|$. Suppose that for all $u \in U'$,*

$$\max\{|N(u) \cap W|, |\neg N(u) \cap W|\} \geq (1 - \epsilon)|W|,$$

and for all $w \in W'$,

$$\max\{|N(w) \cap U|, |\neg N(w) \cap U|\} \geq (1 - \epsilon)|U|.$$

Then $|E|/|U||W| \in [0, 2\epsilon^{1/2}] \cup (1 - 2\epsilon^{1/2}, 1]$.

We will use the following immediate corollary of this.

Corollary 2.19. *For all $0 < \epsilon < \frac{1}{4}$ there is n such that the following holds. Suppose $G = (U \cup W, E)$ is a bipartite graph with $|U|, |W| \geq n$, and $|E|/|U||W| \in (2\epsilon^{1/2}, 1 - 2\epsilon^{1/2})$. Then one of the following hold.*

(1) *There is $U' \subseteq U$ with $|U'| \geq \epsilon|U|$, so that for all $u \in U$,*

$$\frac{|N_E(u) \cap W|}{|W|} \in (\epsilon, 1 - \epsilon).$$

(2) *There is $W' \subseteq W$ with $|W'| \geq \epsilon|W|$, so that for all $w \in W$,*

$$\frac{|N_E(w) \cap U|}{|U|} \in (\epsilon, 1 - \epsilon).$$

We will use a lemma which was originally proved by Frankl and Rödl (see [9, Lemma 3.8]) for another notion of quasirandomness for graphs, called disc_2 .

Definition 2.20. Suppose $B = (U \cup W, E)$ is a bipartite graph, and $|E| = d_B|U||W|$. We say B has $\text{disc}_2(\epsilon, d)$ if $d_B = d \pm \epsilon$ and for all $U' \subseteq U$ and $W' \subseteq W$,

$$||E \cap K_2[U', W']| - d|U'||W'| | \leq \epsilon|U||W|.$$

Gowers proved the following quantitative equivalence between disc_2 and dev_2 (see Theorem 3.1 in [11]).

Theorem 2.21. *Suppose $B = (U \cup W, E)$ is a bipartite graph. If B has $\text{disc}_2(\epsilon, d)$ then it has $\text{dev}_2(\epsilon, d)$. If B has $\text{dev}_2(\epsilon, d)$, then it has $\text{disc}_2(\epsilon^{1/4}, d)$.*

Combining Theorem 2.21 with Lemma 3.8 in [9], we obtain the following.

Lemma 2.22. *For all $\epsilon > 0$, $\rho \geq 2\epsilon$, $0 < p < \rho/2$, and $\delta > 0$, there is $m_0 = m_0(\epsilon, \rho, \delta)$ such that the following holds. Suppose $|U| = |V| = m \geq m_0$, and $G = (U \cup V, E)$ is a bipartite graph satisfying $\text{dev}_2(\epsilon)$ with density ρ . Then if $\ell = \lceil 1/p \rceil$ and $\epsilon \geq 10(1/\ell m)^{1/5}$, there is a partition $E = E_0 \cup E_1 \cup \dots \cup E_\ell$ such that*

- (1) *For each $1 \leq i \leq \ell$, $(U \cup V, E_i)$ has $\text{dev}_2(\epsilon^{1/4})$ with density $\rho p(1 \pm \delta)$, and*
- (2) $|E_0| \leq \rho p(1 + \delta)m^2$.

Further, if $1/p \in \mathbb{Z}$, then $E_0 = \emptyset$.

We will also use the following fact, which can be obtained from Fact 2.3 in [26] along with Theorem 2.21.

Fact 2.23. *Suppose E_1 and E_2 are disjoint subsets of $K_2[U, V]$. If $(U \cup V, E_1)$ has $\text{dev}_2(\epsilon_1, d_1)$, and $(U \cup V, E_2)$ has $\text{dev}_2(\epsilon_2, d_2)$, then $(U \cup V, E_1 \cup E_2)$ has $\text{dev}_2(\epsilon_1^{1/4} + \epsilon_2^{1/4}, d_2 + d_1)$.*

Finally, we will use the fact that triads with density near 0 or 1 are quasirandom. For completeness, we include a proof of this in the [Appendix](#).

Proposition 2.24. *For all $0 < \epsilon < \frac{1}{2}$, $d_2 > 0$, and $0 < \delta \leq (d_2/2)^{48}$, there is N such that the following holds. Suppose $H = (V_1 \cup V_2 \cup V_3, R)$ is a 3-partite 3-graph on $n \geq N$ vertices, and for each $i, j \in [3]$, $||V_i| - |V_j|| \leq \delta|V_i|$. Suppose $G = (V_1 \cup V_2 \cup V_3, E)$ is a 3-partite graph, where for each $1 \leq i < j \leq 3$, $G[V_i, V_j]$ has $\text{dev}_2(\delta, d_2)$, and assume*

$$|R \cap K_3^{(2)}(G)| \leq \epsilon |K_3^{(2)}(G)|.$$

Then $(H|G, G)$ has $\text{dev}_{2,3}(\delta, 6\epsilon)$.

3. Proof of main theorem

We first give a more precise statement of our main theorem.

Theorem 3.1. *For all $k \geq 1$, there are polynomials $p_1(x), p_2(x, y), p_3(x)$, a constant $\epsilon_1^* > 0$, and a function $\epsilon_2^* : \mathbb{N} \rightarrow (0, 1]$ such that the following holds, where $T_0(x, y, z, w)$ is as in [Theorem 2.7](#).*

For all $0 < \epsilon_1 < \epsilon_1^$ and $\epsilon_2 : \mathbb{N} \rightarrow (0, 1]$ satisfying $\epsilon_2 < \epsilon_2^*$, there is $L \leq \epsilon_1^{-O_k(k)}$ such that the following holds for $T = T_0(p_1(\epsilon_1), \epsilon_2 \circ q_2, p_3(\epsilon_1^{-1}), 1)$, where $q_2(y) = p_2(\epsilon_1, y)$.*

Every sufficiently large 3-graph $H = (V, E)$ such that $\text{VC}_2(H) < k$ has a $\text{dev}_{2,3}(\epsilon_1, \epsilon_2(\ell))$ -regular $(t, \ell, \epsilon_1, \epsilon_2(\ell))$ -decomposition with $\ell \leq L$ and $t \leq T$.

We now give a few remarks regarding the bounds. As can be seen above, the bound T in [Theorem 3.1](#) is obtained by composing the bound T_0 from [Theorem 2.7](#) with several polynomial functions. This does not change the fundamental shape of the bound in terms of the Ackerman hierarchy, and thus the bound for t in [Theorem 3.1](#) remains a Wowzer type function. On the other hand, we see that the bound for ℓ becomes polynomial in ϵ_1^{-1} .

The polynomial p_3 in [Theorem 3.1](#) depends on the f in [Theorem 2.16](#), which in turn depends on the hypergraph counting lemma for $\text{dev}_{2,3}$. One could therefore obtain a quantitative version of [Theorem 3.1](#) for the equivalent quasirandomness notions of $\text{disc}_{2,3}$ and $\text{oct}_{2,3}$ using the same arguments, given a quantitative version of their respective counting lemmas.

The general strategy for the proof of [Theorem 3.1](#) is as follows. Given a large 3-graph H of VC_2 -dimension less than k , we first apply [Theorem 2.16](#) to obtain a homogeneous, regular partition \mathcal{P} for H . We then consider the auxiliary edge-colored graphs associated to \mathcal{P} , as described in [Section 2D](#). These contain no copies of $U(k)$ by [Proposition 2.12](#), allowing us to apply [Lemma 2.14](#). This yields decompositions for the auxiliary edge-colored graphs, which we eventually use to define a new decomposition \mathcal{Q} for H which is still regular and homogeneous, but

which has a polynomial bound for the parameter ℓ . This last part requires the most work, as well as most of the lemmas from [Section 2G](#).

We have not sought to optimize constants which do not effect the overall form of the bounds involved.

Proof of [Theorem 1.3](#). Fix $k \geq 1$ and let $c_1 = c_1(k)$ be as in [Theorem 2.13](#). Let $\rho_1 > 0$, $\rho_2 : \mathbb{N} \rightarrow (0, 1]$, and f be as in [Theorem 2.16](#) for k , and let $D = D(k)$ be so that $f(x) = x^{1/D}$ (see [Theorem 2.16](#)). Let $\mu_1 > 0$, $\mu_2 : \mathbb{N} \rightarrow (0, 1]$ be as in [Proposition 2.12](#) for k . Set $\epsilon_1^* = \min\{\mu_1, \rho_1, (1/4)^D\}$ and define $\epsilon_2^* : \mathbb{N} \rightarrow (0, 1]$ by setting $\epsilon_2^*(x) = \min\{\mu_2(x), \rho_2(x), (1/2x)^{48}\}$, for each $x \in \mathbb{N}$.

Suppose $0 < \epsilon_1 < \epsilon_1^*$ and $\epsilon_2 : \mathbb{N} \rightarrow (0, 1]$ satisfies $\epsilon_2 < \epsilon_2^*$. We now choose a series of new constants. Set $\tau_1 = \epsilon_1^{4D}$ and note $\tau_1 < f(\epsilon_1)$. Set $\delta = \tau_1^{400}/1000$, $\epsilon_1' = (\delta/8c_1)^{2k+1000}$, $m = \lceil 2c_1(\delta/8)^{-2k-2} \rceil$, and $\epsilon_1'' = (\epsilon_1')^2/1000$. Next, define $\epsilon_2', \epsilon_2'' : \mathbb{N} \rightarrow (0, 1]$ by setting, for each $x \in \mathbb{N}$, $\epsilon_2'(x) = \epsilon_1''\epsilon_2(x)\epsilon_2(2^4\delta^{-8k-10})$ and $\epsilon_2''(x) = \epsilon_2(\delta^{-4}m^4)\epsilon_2'(x)^5/4$. Note there are polynomials $p_1(x)$, $p_2(x, y)$ depending only on k such that $\epsilon_1'' = p_1(\epsilon_1)$ and $\epsilon_2''(x) = p_2(\epsilon_1, x)$. To aid the reader in keeping track of the constants, we point out that the following inequalities hold:

$$\epsilon_1'' < \epsilon_1' < \delta < \tau_1 < \epsilon_1 < \epsilon_1^* \quad \text{and} \quad \epsilon_2'' < \epsilon_2' < \epsilon_2 < \epsilon_2^*.$$

Choose t_0 sufficiently large so that

$$\begin{aligned} \frac{t^3}{6} &\geq (1 - \epsilon_1'') \binom{t}{3}, \\ \frac{(1 - 3\epsilon_1'')t^3}{12} &\geq (1 - \epsilon_1') \binom{t}{3}, \quad \text{and} \\ \binom{t}{3}(1 - 6(\epsilon_1')^{1/4} - (\epsilon_1')^{3/8}) &\geq (1 - \epsilon_1')^{1/8} \binom{t}{3}. \end{aligned}$$

Note there is some polynomial $p(x)$ depending only on k so that we can take $t_0 = p(\epsilon_1^{-1})$. Finally, choose T_1 , L_1 , and N_1 as in [Theorem 2.7](#) for $\epsilon_1'', \epsilon_2'', t_0$ and $\ell_0 = 1$.

Set $L = \lceil \delta^{-4}m^4 \rceil$, $T = T_1$, and choose N sufficiently large compared to all the previously chosen constants. Notice that $L = O_k(\epsilon_1^{-O_k(1)})$ and

$$T = T_0(p_1(\epsilon_1), \epsilon_2 \circ q_2, p(\epsilon_1^{-1}), 1),$$

where $T_0(x, y, z, w)$ is as in [Theorem 2.7](#) and $q_2(y) = p_2(\epsilon_1, y)$.

Suppose $H = (V, E)$ is a 3-graph with $|V| \geq N$ satisfying $\text{VC}_2(H) < k$. [Theorem 2.16](#) implies there exist $1 \leq \ell \leq L_1$, $t_0 \leq t \leq T_1$, and \mathcal{P}_1 a $(t, \ell, \epsilon_1'', \epsilon_2''(\ell))$ -decomposition of V which is $\text{dev}_{2,3}(\epsilon_1'', \epsilon_2''(\ell_1))$ -regular and $f(\epsilon_1'')$ -homogeneous with respect to H . Say

$$\mathcal{P}_1 = \{V_1, \dots, V_t\} \quad \text{and} \quad \mathcal{P}_2 = \left\{ P_{ij}^\alpha : ij \in \binom{[t]}{2}, \alpha \in [\ell] \right\}.$$

Note that $f(\epsilon_1'') = (\epsilon_1'')^{1/D} < \frac{1}{4}$. Recall that as mentioned after [Theorem 2.16](#), we may assume that all $\text{dev}_{2,3}(\epsilon_1'', \epsilon_2''(\ell))$ -regular triads of \mathcal{P} are $f(\epsilon_1'')$ -homogeneous with respect to H .

Given $ij \in \binom{[t]}{2}$ and $\alpha \in [\ell]$, let $G_{ij}^\alpha = (V_i \cup V_j, P_{ij}^\alpha)$. Given $ijs \in \binom{[t]}{3}$ and $1 \leq \alpha, \beta, \gamma \leq \ell$, set

$$\begin{aligned} G_{ijs}^{\alpha,\beta,\gamma} &= (V_i \cup V_j \cup V_s, P_{ij}^\alpha \cup P_{js}^\beta \cup P_{is}^\gamma) \quad \text{and} \\ H_{ijs}^{\alpha,\beta,\gamma} &= (V_i \cup V_j \cup V_s, E \cap K_3^{(2)}(G_{ijs}^{\alpha,\beta,\gamma})). \end{aligned}$$

We will use throughout that since $\epsilon_2''(x) \leq \epsilon_2'(x)^5/4$, [Proposition 2.17](#) implies that for all $ijs \in \binom{[t]}{3}$ and $\alpha, \beta, \gamma \in [\ell]$,

$$|K_3^{(2)}(G_{ijs}^{\alpha,\beta,\gamma})| = (1 \pm \epsilon_2'(\ell)) \left(\frac{n}{\ell t}\right)^3. \quad (1)$$

We use \mathcal{P} to construct a different decomposition of V , which we call \mathcal{Q} , so that $\mathcal{Q}_1 = \mathcal{P}_1$ but $\mathcal{Q}_2 \neq \mathcal{P}_2$. Set

$$\begin{aligned} \mathbf{F}_{\text{err}} &= \{G_{ijs}^{\alpha,\beta,\gamma} \in \text{Triads}(\mathcal{P}) : (H_{ijs}^{\alpha,\beta,\gamma}, G_{ijs}^{\alpha,\beta,\gamma}) \text{ fails } \text{disc}_3(\epsilon_1'', \epsilon_2''(\ell))\}, \\ \mathbf{F}_1 &= \{G_{ijs}^{\alpha,\beta,\gamma} \in \text{Triads}(\mathcal{P}) \setminus \mathbf{F}_{\text{err}} : d_{ijs}^{\alpha,\beta,\gamma} \geq 1 - f(\epsilon_1'')\}, \text{ and} \\ \mathbf{F}_0 &= \{G_{ijs}^{\alpha,\beta,\gamma} \in \text{Triads}(\mathcal{P}) \setminus \mathbf{F}_{\text{err}} : d_{ijs}^{\alpha,\beta,\gamma} \leq f(\epsilon_1'')\}. \end{aligned}$$

By assumption, $\text{Triads}(\mathcal{P}) = \mathbf{F}_{\text{err}} \sqcup \mathbf{F}_1 \sqcup \mathbf{F}_0$, and at most $\epsilon_1'' n^3$ triples $xyz \in \binom{V}{3}$ are in $K_3^{(2)}(G)$ for some $G \in \mathbf{F}_{\text{err}}$. By (1), this implies

$$|\text{Triads}(\mathcal{P}) \setminus \mathbf{F}_{\text{err}}| \geq \left(\binom{n}{3} - \epsilon_1'' n^3\right) / \left(\frac{n^3}{t^3 \ell^3} (1 - \epsilon_2'(\ell))\right) \geq \binom{t}{3} \ell^3 (1 - \epsilon_1''),$$

where the last inequality uses that $t \geq t_0$ and n is large. Thus, $|\mathbf{F}_{\text{err}}| \leq \epsilon_1' t^3 \ell^3$. Let

$$\Psi = \{V_i V_j : |\{G_{ijs}^{\alpha\beta\gamma} \in \mathbf{F}_{\text{err}} \text{ some } s \in [t] \text{ and } \alpha, \beta, \gamma \in [\ell]\}| \geq (\epsilon_1')^{3/4} \ell^3 t\}.$$

Since $|\mathbf{F}_{\text{err}}| \leq \epsilon_1' t^3 \ell^3$, we have that $|\Psi| \leq (\epsilon_1')^{1/4} t^2$. Given $ij \in \binom{[t]}{2}$, let ℓ_{ij} be the number of $\alpha \in [\ell]$ such that G_{ij}^α has $\text{dev}_2(\epsilon_2''(\ell), 1/\ell)$. After relabeling, we may assume $G_{ij}^1, \dots, G_{ij}^{\ell_{ij}}$ each have $\text{dev}_2(\epsilon_2''(\ell), 1/\ell)$. We claim that for $V_i V_j \notin \Psi$,

$$\ell_{ij} \geq (1 - 2(\epsilon_1')^{3/4})\ell.$$

Indeed, given $V_i V_j \notin \Psi$, if it were the case that $\ell_{ij} < (1 - 2(\epsilon_1')^{3/4})\ell$, then we would have that

$$\begin{aligned} |\{G_{ijs}^{\alpha\beta\gamma} \in \mathbf{F}_{\text{err}} \text{ some } s \in [t] \text{ and } \alpha, \beta, \gamma \in [\ell]\}| &\geq (t-2)\ell^2(\ell - \ell_{ij}) \\ &> 2(\epsilon_1')^{3/4}(t-2)\ell^3 \geq (\epsilon_1')^{3/4}\ell^3 t, \end{aligned}$$

contradicting $V_i V_j \notin \Psi$. Thus we have that for all $V_i V_j \notin \Psi$, $\ell_{ij} \geq (1 - 2(\epsilon_1')^{3/4})\ell$.

For each $V_i V_j \notin \Psi$, let H_{ij} be the edge-colored graph $(U_{ij} \cup W_{ij}, E_{ij}^0, E_{ij}^1, E_{ij}^2)$, where

$$\begin{aligned} W_{ij} &= \{P_{ij}^\alpha : \alpha \leq \ell_{ij}\}, \\ U_{ij} &= \{P_{is}^\beta P_{js}^\gamma : s \in [t] \setminus \{i, j\}, \beta \leq \ell_{is}, \gamma \leq \ell_{js}\}, \\ E_{ij}^1 &= \{P_{ij}^\alpha (P_{is}^\beta P_{js}^\gamma) \in K_2[W_{ij}, U_{ij}] : G_{ijk}^{\alpha, \beta, \gamma} \in \mathbf{F}_1\}, \\ E_{ij}^0 &= \{P_{ij}^\alpha (P_{is}^\beta P_{js}^\gamma) \in K_2[W_{ij}, U_{ij}] : G_{ijk}^{\alpha, \beta, \gamma} \in \mathbf{F}_0\}, \\ E_{ij}^2 &= \{P_{ij}^\alpha (P_{is}^\beta P_{js}^\gamma) \in K_2[W_{ij}, U_{ij}] : G_{ijk}^{\alpha, \beta, \gamma} \in \mathbf{F}_{\text{err}}\}. \end{aligned}$$

By [Proposition 2.12](#), and since $f(\epsilon_1'') < \frac{1}{2}$, H_{ij} contains no E_{ij}^1/E_{ij}^0 copy of $U(k)$, and since $V_i V_j \notin \Psi$, $|E_{ij}^2| \leq (\epsilon_1')^{3/4} \ell^3 t$. We will later need the following size estimates for W_{ij} and U_{ij} . By the above, $|W_{ij}| = \ell_{ij} \geq (1 - 2(\epsilon_1')^{3/4})\ell$. We claim that $|U_{ij}| \geq (1 - 2(\epsilon_1')^{3/4})\ell^2 t$. Indeed, observe that $|U_{ij}| = \sum_{s \in [t] \setminus \{i, j\}} \ell_{is} \ell_{js}$ and

$$\begin{aligned} &|\{G_{ijs}^{\alpha\beta\gamma} \in \mathbf{F}_{\text{err}} \text{ some } s \in [t] \text{ and } \alpha, \beta, \gamma \in [\ell]\}| \\ &\geq \sum_{s \in [t] \setminus \{i, j\}} \ell^2 (\ell - \ell_{is}) + \ell_{is} \ell (\ell - \ell_{js}) \\ &= \sum_{s \in [t] \setminus \{i, j\}} \ell^3 - \ell \ell_{is} \ell_{js} = (t - 2)\ell^3 - \ell |U_{ij}|. \end{aligned}$$

Since $V_i V_j \notin \Psi$, this shows that

$$(\epsilon_1')^{3/4} \ell^3 t \geq (t - 2)\ell^3 - \ell |U_{ij}|.$$

Rearranging, this yields that

$$|U_{ij}| \geq (t - 2)\ell^2 - (\epsilon_1')^{3/4} \ell^2 t \geq t\ell^2 (1 - 2(\epsilon_1')^{3/4}),$$

where the last inequality is because $t \geq t_0$.

Given $v, v' \in W_{ij}$, write $v \sim v' \in W_{ij}$ if for each $w \in \{0, 1, 2\}$,

$$|E_{ij}^w(v) \Delta E_{ij}^w(v')| \leq \delta |U_{ij}|.$$

By [Lemma 2.14](#), there are $W_{ij}^0 \subseteq W_{ij}$ of size at most $(\epsilon_1')^{3/8} |W_{ij}|$, an integer $m_{ij} \leq m$, and $x_{ij}^1, \dots, x_{ij}^{m_{ij}} \in W_{ij}$ so that for all $v \in W_{ij} \setminus W_{ij}^0$, there is $1 \leq \alpha \leq m_{ij}$ so that $v \sim x_{ij}^\alpha$, and further, $|N_{E_{ij}^2}(v)| \leq (\epsilon_1')^{3/8} |U_{ij}|$. For each $1 \leq u \leq m_{ij}$, let

$$W_{ij}^u = \{v \in W_{ij} \setminus W_{ij}^0 : v \sim x_{ij}^u \text{ and for all } 1 \leq u' < u, v \not\sim x_{ij}^{u'}\}.$$

Note $W_{ij}^1 \cup \dots \cup W_{ij}^{m_{ij}}$ is a partition of $W_{ij} \setminus W_{ij}^0$.

We now define a series of sets to help us zero in on certain well behaved sets of triples. First, define

$$\Omega_0 = \left\{ ijs \in \binom{[t]}{3} : V_i V_j, V_j V_s, V_i V_s \notin \Psi \right\} \quad \text{and}$$

$$\Omega = \{ W_{ij}^u W_{is}^v W_{js}^w : ijs \in \Omega_0, 1 \leq u \leq m_{ij}, 1 \leq v \leq m_{is}, 1 \leq w \leq m_{js} \}.$$

Since $|\Psi| \leq (\epsilon'_1)^{1/4} t^2$, $|\Omega_0| \geq \binom{t}{3} - |\Psi|t \geq (1 - 6(\epsilon'_1)^{1/4}) \binom{t}{3}$. Let

$$Y_0 = \bigcup_{W_{ij}^u W_{is}^v W_{js}^w \in \Omega} K_3[W_{ij}^u, W_{is}^v, W_{js}^w].$$

We have that for all $ijs \in \Omega_0$, $|W_{ij}^0|, |W_{is}^0|, |W_{js}^0| \leq (\epsilon'_1)^{3/8} \ell$, and therefore $|Y_0|$ is at least the following:

$$\begin{aligned} |Y_0| &\geq \binom{t}{3} \ell^3 - \ell^3 \left| \binom{[t]}{3} \setminus \Omega_0 \right| - |\Omega_0| (\epsilon'_1)^{3/8} \ell^3 \\ &\geq \binom{t}{3} \ell^3 - 6(\epsilon'_1)^{1/4} \binom{t}{3} \ell^3 - \binom{t}{3} (\epsilon'_1)^{3/8} \ell^3 \\ &\geq \binom{t}{3} \ell^3 (1 - (\epsilon'_1)^{1/8}), \end{aligned}$$

where the last inequality is since $t \geq t_0$.

Given $ij \notin \Psi$, let us call W_{ij}^u *nontrivial* if it has size at least $\delta^{1/2} \ell / m_{ij}$. Define

$$\Omega_1 = \{ W_{ij}^u W_{js}^v W_{is}^w \in \Omega : \text{each of } W_{ij}^u, W_{js}^v, W_{is}^w \text{ are nontrivial} \},$$

and set $Y_1 = \bigcup_{W_{ij}^u W_{js}^v W_{is}^w \in \Omega_1} K_3[W_{ij}^u, W_{js}^v, W_{is}^w]$. Then we have that

$$\begin{aligned} |Y_1| &\geq |Y_0| - t \ell^2 \sum_{ij \in \binom{[t]}{2}} \sum_{\{u \in [m_{ij}] : W_{ij}^u \text{ trivial}\}} \delta^{1/2} (\ell / m_{ij}) \\ &\geq |Y_0| - t \ell^2 (t^2 \delta^{1/2} \ell) = |Y_0| - \delta^{1/2} t^3 \ell^3. \end{aligned}$$

Define

$$R_1 = \{ P_{ij}^\alpha P_{is}^\beta P_{js}^\gamma : G_{ijk}^{\alpha, \beta, \gamma} \in \mathbf{F}_1 \},$$

$$R_0 = \{ P_{ij}^\alpha P_{is}^\beta P_{js}^\gamma : G_{ijk}^{\alpha, \beta, \gamma} \in \mathbf{F}_0 \},$$

$$R_2 = \{ P_{ij}^\alpha P_{is}^\beta P_{js}^\gamma : G_{ijk}^{\alpha, \beta, \gamma} \in \mathbf{F}_{\text{err}} \}.$$

Note that $(\mathcal{P}_2 \cup \mathcal{P}_2 \cup \mathcal{P}_2, R_0, R_1, R_2)$ is a 3-partite edge-colored 3-graph, and $|R_2| \leq \epsilon'_1 t^3 \ell^3$. Now set

$$\Omega_2 = \{ W_{ij}^u W_{js}^v W_{is}^w \in \Omega_1 : |R_2 \cap K_3[W_{ij}^u, W_{is}^v, W_{js}^w]| \leq \sqrt{\epsilon'_1} |W_{ij}^u| |W_{is}^v| |W_{js}^w| \}$$

and $Y_2 = \bigcup_{W_{ij}^u W_{js}^v W_{is}^w \in \Omega_2} K_3[W_{ij}^u W_{js}^v W_{is}^w]$. Note that

$$\begin{aligned} |R_2| &\geq \sum_{W_{ij}^u W_{js}^v W_{is}^w \in \Omega_1 \setminus \Omega_2} \sqrt{\epsilon'_1} |W_{ij}^u| |W_{is}^v| |W_{js}^w| \\ &\geq \sqrt{\epsilon'_1} \sum_{W_{ij}^u W_{js}^v W_{is}^w \in \Omega_1 \setminus \Omega_2} |W_{ij}^u| |W_{is}^v| |W_{js}^w|. \end{aligned}$$

Therefore,

$$\sum_{W_{ij}^u W_{js}^v W_{is}^w \in \Omega_1 \setminus \Omega_2} |W_{ij}^u| |W_{is}^v| |W_{js}^w| \leq \sqrt{\epsilon'_1}^{-1} |R_2| < \sqrt{\epsilon'_1}^{-1} \epsilon'_1 t^3 \ell^3 \leq \sqrt{\epsilon'_1} t^3 \ell^3.$$

This implies that $|Y_2| \geq |Y_1| - \sqrt{\epsilon'_1} t^3 \ell^3$.

Given $ijs \in \Omega_0$, let us call a triple $P_{ij}^\alpha P_{is}^\beta P_{js}^\gamma$ *troublesome* if one of the following hold:

- For some $u \in [m_{ij}]$, $P_{ij}^\alpha \in W_{ij}^u$, and there are $\sigma_1 \neq \sigma_2 \in \{0, 1, 2\}$ such that $P_{is}^\beta P_{js}^\gamma P_{ij}^\alpha \in R^{\sigma_1}$ and $P_{is}^\beta P_{js}^\gamma x_{ij}^u \in R^{\sigma_2}$.
- For some $w \in [m_{js}]$, $P_{js}^\gamma \in W_{js}^w$, and there are $\sigma_1 \neq \sigma_2 \in \{0, 1, 2\}$ such that $P_{is}^\beta P_{ij}^\alpha P_{js}^\gamma \in R^{\sigma_1}$ and $P_{is}^\beta P_{ij}^\alpha x_{js}^w \in R^{\sigma_2}$.
- For some $v \in [m_{is}]$, $P_{is}^\beta \in W_{is}^v$, and there are $\sigma_1 \neq \sigma_2 \in \{0, 1, 2\}$ such that $P_{ij}^\alpha P_{js}^\gamma P_{is}^\beta \in R^{\sigma_1}$ and $P_{ij}^\alpha P_{js}^\gamma x_{is}^v \in R^{\sigma_2}$.

Let Tr be the set of troublesome triples. Define

$$\Omega_3 = \{W_{ij}^u W_{js}^v W_{is}^w \in \Omega_2 : |K_3[W_{ij}^u W_{js}^v W_{is}^w] \cap \text{Tr}| \leq \delta^{1/4} |W_{ij}^u| |W_{js}^v| |W_{is}^w|\},$$

and set $Y_3 = \bigcup_{W_{ij}^u W_{js}^v W_{is}^w \in \Omega_3} K_3[W_{ij}^u W_{js}^v W_{is}^w]$. We claim $|Y_3| \geq \binom{t}{3} \ell^3 (1 - 2\delta^{1/2})$.

Given $V_i V_j \notin \Psi$, $1 \leq u \leq m_{ij}$, and $P_{ij}^\alpha \in W_{ij}^u$, we know that $P_{ij}^\alpha \sim x_{ij}^u$, and therefore

$$\begin{aligned} &|\{P_{is}^\beta P_{js}^\gamma : s \in [t] \setminus \{i, j\}, \beta, \gamma \leq \ell, P_{is}^\beta P_{js}^\gamma P_{ij}^\alpha \in \text{Tr}\}| \\ &\leq (\ell^2(t-2) - |U_{ij}|) + \sum_{x=0}^2 |N_{E_{ij}^x}(P_{ij}^\alpha) \Delta N_{E_{ij}^x}(x_{ij}^u)| \\ &\leq 2(\epsilon'_1)^{3/4} \ell^2 t + 3\delta t \ell^2 \\ &\leq 4\delta t \ell^2. \end{aligned}$$

Thus, $|\text{Tr}| \leq 4\delta t \ell^2 (\sum_{V_i V_j \notin \Psi, u \in [m_{ij}]} |W_{ij}^u|) \leq 4\delta t \ell^2 (t^2 \ell) = 4\delta t^3 \ell^3$. Therefore

$$\begin{aligned} 4\delta t^3 \ell^3 \geq |\text{Tr}| &\geq \sum_{W_{ij}^u W_{js}^v W_{is}^w \in \Omega_2 \setminus \Omega_3} \delta^{1/4} |W_{ij}^u| |W_{js}^v| |W_{is}^w| \\ &= \delta^{1/4} \left| \bigcup_{W_{ij}^u W_{js}^v W_{is}^w \in \Omega_2 \setminus \Omega_3} K_3[W_{ij}^u W_{js}^v W_{is}^w] \right|. \end{aligned}$$

Rearranging, this yields that

$$\left| \bigcup_{W_{ij}^u W_{js}^v W_{is}^w \in \Omega_2 \setminus \Omega_3} K_3[W_{ij}^u W_{js}^v W_{is}^w] \right| \leq \delta^{-1/4} 4\delta t^3 \ell^3 = 4\delta^{3/4} t^3 \ell^3.$$

Thus

$$\begin{aligned} |Y_3| &\geq |Y_2| - \delta^{3/4} t^3 \ell^3 \\ &\geq |Y_1| - \sqrt{\epsilon'_1} t^3 \ell^3 - 4\delta^{3/4} t^3 \ell^3 \\ &\geq |Y_0| - \delta^{1/2} t^3 \ell^3 - \sqrt{\epsilon'_1} t^3 \ell^3 - 4\delta^{3/4} t^3 \ell^3 \\ &\geq \binom{t}{3} \ell^3 (1 - 7(\epsilon'_1)^{1/8}) - \delta^{1/2} t^3 \ell^3 - \sqrt{\epsilon'_1} t^3 \ell^3 - 4\delta^{3/4} t^3 \ell^3 \\ &\geq \binom{t}{3} \ell^3 (1 - 2\delta^{1/2}). \end{aligned}$$

Therefore, using (1), we have

$$\begin{aligned} \left| \bigcup_{P_{ij}^\alpha P_{is}^\beta P_{js}^\gamma \in Y_3} K_3^{(2)}(G_{ijs}^{\alpha, \beta, \gamma}) \right| &\geq \binom{t}{3} \ell^3 (1 - 2\delta^{1/2}) \left(\frac{n^3}{t^3 \ell^3} (1 - \epsilon'_2(\ell)) \right) \\ &\geq \binom{n}{3} (1 - 3\delta^{1/2}), \end{aligned}$$

where the last inequality is because n is large.

Our next goal is to prove Claim 3.2, which says that for each $W_{ij}^u W_{is}^v W_{js}^w \in \Omega_3$, $K_3[W_{ij}^u, W_{is}^v, W_{js}^w]$ is either mostly contained in R_1 or mostly contained in R_0 . For the proof of this claim, we will require the following notation. Given $ijs \in \binom{[t]}{3}$, $\alpha, \alpha' \leq \ell$, $1 \leq v \leq m_{is}$, and $1 \leq w \leq m_{js}$, we write $P_{ij}^\alpha \sim_{js, vw} P_{ij}^{\alpha'}$ if $P_{ij}^\alpha, P_{ij}^{\alpha'} \in W_{ij}^u$ for some $1 \leq u \leq m_{ij}$, and

$$\begin{aligned} &|\{(P_{is}^\beta, P_{js}^\gamma) \in W_{is}^v \times W_{js}^w : \text{for some } \sigma_1 \neq \sigma_2 \in \{0, 1, 2\}, P_{is}^\beta P_{js}^\gamma P_{ij}^\alpha \in R^{\sigma_1} \\ &\quad \text{and } P_{is}^\beta P_{js}^\gamma P_{ij}^{\alpha'} \in R^{\sigma_2}\}| \leq \delta^{1/8} |W_{is}^v| |W_{js}^w|. \end{aligned}$$

Claim 3.2. For any $W_{ij}^u W_{is}^v W_{js}^w \in \Omega_3$, there is $\sigma \in \{0, 1\}$ such that

$$\frac{|R_\sigma \cap K_3[W_{ij}^u, W_{is}^v, W_{js}^w]|}{|K_3[W_{ij}^u, W_{is}^v, W_{js}^w]|} \geq 1 - \delta^{1/100}.$$

Proof. Suppose towards a contradiction there is $W_{ij}^u W_{is}^v W_{js}^w \in \Omega_3$ such that for each $\sigma \in \{0, 1\}$,

$$\frac{|R_\sigma \cap K_3[W_{ij}^u, W_{is}^v, W_{js}^w]|}{|K_3[W_{ij}^u, W_{is}^v, W_{js}^w]|} < 1 - \delta^{1/100}.$$

To ease notation, let $A = W_{ij}^u$, $B = W_{is}^v$, and $C = W_{js}^w$.

We now define a series of subsets of A which will contain “well behaved” vertices. First, we set $A_1 = \{a \in A : a \sim_{js, vw} x_{ij}^u\}$. Since $W_{ij}^u W_{is}^v W_{js}^w \in \Omega_3$,

$$\delta^{1/4} |W_{ij}^u| |W_{is}^v| |W_{js}^w| \geq |\text{Tr} \cap K_3[W_{ij}^u W_{is}^v W_{js}^w]| \geq |A \setminus A_1| \delta^{1/8} |W_{is}^v| |W_{js}^w|.$$

Thus $|A \setminus A_1| \leq \delta^{-1/8} \delta^{1/4} |W_{ij}^u| = \delta^{1/8} |W_{ij}^u|$. Now set

$$A_2 = \{a \in A : |N_{R_2}(a)| \leq (\epsilon'_1)^{1/4} |B| |C|\}.$$

Because $W_{ij}^u W_{is}^v W_{js}^w \in \Omega_2$, we have that

$$(\epsilon'_1)^{1/2} |A| |B| |C| \geq |R_2 \cap K_3[A, B, C]| \geq |A \setminus A_2| (\epsilon'_1)^{1/4} |B| |C|.$$

Therefore, $|A \setminus A_2| \leq (\epsilon'_1)^{1/4} |A|$. Now set

$$A_3 = \{a \in A : |N_{R_1}(a)| / |B| |C| \in (\delta^{1/64}, 1 - \delta^{1/64})\} \quad \text{and}$$

$$A'_3 = \{a \in A : |N_{R_1}(a)| / |B| |C| \in (\delta^{1/128}, 1 - \delta^{1/128})\}.$$

We claim $x_{ij}^u \in A'_3$. Suppose towards a contradiction that $x_{ij}^u \notin A'_3$. Suppose first that $|N_{R_1}(x_{ij}^u)| \geq (1 - \delta^{1/128}) |B| |C|$. Then for all $a \in A_1$, since $a \sim_{js, vw} x_{ij}^u$, we have $|N_{R_1}(a)| \geq (1 - \delta^{1/128} - \delta^{1/8}) |B| |C|$, and thus,

$$\begin{aligned} |R_1 \cap K_3[W_{ij}^u W_{is}^v W_{js}^w]| &\geq (1 - \delta^{1/128} - \delta^{1/8}) |A_1| |B| |C| \\ &\geq (1 - \delta^{1/128} - \delta^{1/8}) (1 - \delta^{1/8}) |A| |B| |C| \\ &\geq (1 - \delta^{1/100}) |A| |B| |C|, \end{aligned}$$

a contradiction. So we must have $|N_{R_1}(x_{ij}^u)| \leq \delta^{1/128} |B| |C|$. Then for all $a \in A_1 \cap A_2$, $a \sim_{js, vw} x_{ij}^u$ and $|N_{R_2}(a)| \leq (\epsilon'_1)^{1/4} |B| |C|$ implies

$$|N_{R_0}(a)| \geq (1 - \delta^{1/128} - \delta^{1/8} - (\epsilon'_1)^{1/4}) |B| |C|.$$

Therefore

$$\begin{aligned} |R_0 \cap K_3[W_{ij}^u W_{is}^v W_{js}^w]| &\geq (1 - \delta^{1/128} - \delta^{1/8} - (\epsilon'_1)^{1/4}) |A_1 \cap A_2| |B| |C| \\ &\geq (1 - \delta^{1/128} - \delta^{1/8} - (\epsilon'_1)^{1/4}) (1 - \delta^{1/8} - (\epsilon'_1)^{1/4}) |A| |B| |C| \\ &\geq (1 - \delta^{1/100}) |A| |B| |C|, \end{aligned}$$

again a contradiction. Thus, we must have that $x_{ij}^u \in A'_3$. This implies that for all $a \in A_1 \cap A_2$,

$$\begin{aligned} |N_{R_1}(a)| &\geq |N_{R_1}(x_{ij}^u)| - |N_{R_1}(x_{ij}^u) \Delta N_{R_1}(a)| \\ &\geq \delta^{1/128} |B| |C| (1 - \delta^{1/8}) \geq \delta^{1/64} |B| |C| \end{aligned}$$

and

$$\begin{aligned} |N_{R_0}(a)| &\geq |N_{R_0}(x_{ij}^u)| - |N_{R_0}(a)| - |N_{R_0}(x_{ij}^u) \Delta N_{R_0}(a)| \\ &\geq \delta^{1/128} |B| |C| (1 - \delta^{1/8} - (\epsilon'_1)^{1/4}) \\ &\geq \delta^{1/64} |B| |C|. \end{aligned}$$

Thus $a \in A_3$. This shows that $A_1 \cap A_2 \subseteq A_3$, and therefore

$$|A_3| \geq |A| (1 - \delta^{1/8} - (\epsilon'_1)^{1/4}).$$

Now define

$$\begin{aligned} A_B &= \{a \in A : |\{b \in B : |N_{R_1}(ab) \Delta N_{R_1}(ax_{is}^v)| \leq \delta^{1/16} |C|\}| \geq (1 - \delta^{1/16}) |B|\}, \\ A_C &= \{a \in A : |\{c \in C : |N_{R_1}(ac) \Delta N_{R_1}(ax_{js}^w)| \leq \delta^{1/16} |B|\}| \geq (1 - \delta^{1/16}) |C|\}. \end{aligned}$$

Observe that $4\delta^{1/4} |A| |B| |C| \geq |\text{Tr} \cap K_3[A, B, C]| \geq \delta^{1/16} |A \setminus A_B| |B| |C|$, and therefore $|A \setminus A_B| \leq \delta^{-1/16} 4\delta^{1/4} |A| = 4\delta^{3/16} |A|$. A similar computation shows $|A \setminus A_C| \leq 4\delta^{3/16} |A|$. Consequently, setting $A_4 := A_3 \cap A_B \cap A_C$, we have that

$$|A_4| \geq |A_3| - |A \setminus A_C| - |A \setminus A_B| \geq |A| (1 - 8\delta^{3/16} - (\epsilon'_1)^{1/4} - \delta^{1/8}) > 0.$$

Fix some $a_* \in A_4$. We will use a_* to control the other edges in the triple. Let

$$S_1 = N_{R_1}(a_*), \quad S_0 = N_{R_0}(a_*), \quad \text{and} \quad S_2 = N_{R_2}(a_*).$$

Note $(B \cup C, S_0 \cup S_1 \cup S_2)$ is a 3-partite edge-colored 3-graph. Since $a_* \in A_3$, $|S_1|/|B||C| \in (\delta^{1/64}, 1 - \delta^{1/64})$. Therefore, [Corollary 2.19](#) implies that one of the following hold:

- (a) There is $B_1 \subseteq B$ such that $|B_1| \geq \delta^{1/32} |B|/2$ and for all $b \in B_1$,

$$\frac{|N_{S_1}(b)|}{|C|} \in \left(\frac{\delta^{1/32}}{2}, 1 - \frac{\delta^{1/32}}{2} \right).$$

- (b) There is $C_1 \subseteq C$ such that $|C_1| \geq \delta^{1/32} |C|/2$ and for all $c \in C_1$,

$$\frac{|N_{S_1}(c)|}{|B|} \in \left(\frac{\delta^{1/32}}{2}, 1 - \frac{\delta^{1/32}}{2} \right).$$

Without loss of generality, let us assume (a) holds (other case is symmetric). Define $B_2 = \{b \in B_1 : |N_{S_2}(b)| \leq (\epsilon'_1)^{1/16} |C|\}$. We claim $|B_2| \geq \delta^{1/32} |B|/4$. Indeed, we know that since $a_* \in A_2$,

$$(\epsilon'_1)^{1/4} |B| |C| \geq |S_2| \geq (\epsilon'_1)^{1/16} |B_1 \setminus B_2| |C|.$$

Thus, $|B_1 \setminus B_2| \leq (\epsilon'_1)^{-1/16}(\epsilon'_1)^{1/4}|B| = (\epsilon'_1)^{1/12}|B|$, so

$$|B_2| \geq |B_1| - (\epsilon'_1)^{1/12}|B| \geq \left(\frac{\delta^{1/32}}{2} - (\epsilon'_1)^{1/12} \right) |B| \geq \frac{\delta^{1/32}|B|}{4}.$$

Note that for all $b \in B_2$, we have that $|N_{S_1}(b)| \geq \delta^{1/32}|C|/2 \geq \delta^{1/32}|C|/4$ and

$$|N_{S_0}(b)| \geq |C \setminus N_{S_1}(b)| - |N_{S_2}(b)| \geq \left(\frac{\delta^{1/32}}{2} - (\epsilon'_1)^{1/16} \right) |C| \geq \frac{\delta^{1/32}|C|}{4}.$$

Now, let

$$B_3 = \{b \in B_2 : |N_{S_1}(b) \Delta N_{S_1}(x_{i_s}^v)| \leq \delta^{1/16}|C|\}.$$

Since $a_* \in A_B$,

$$|B_3| \geq |B_2| - \delta^{1/16}|B| \geq \left(\frac{\delta^{1/32}}{4} - \delta^{1/16} \right) |B| \geq \frac{\delta^{1/32}|B|}{8} > 0.$$

Fix some $b_* \in B_3$ and set $Q_0 = N_{S_0}(b_*)$ and $Q_1 = N_{S_1}(b_*)$. By above, since $b_* \in B_2$, $\min\{|Q_1|, |Q_0|\} \geq \delta^{1/32}|C|/4$.

We claim $|S_1 \cap K_2[B_3, Q_1]| \geq (1 - 10\delta^{1/32})|Q_1||B_3|$. Indeed, fix $b \in B_3$. Then we know $|N_{S_1}(b) \Delta N_{S_1}(x_{i_s}^v)| \leq \delta^{1/16}|C|$ and $|N_{S_1}(b_*) \Delta N_{S_1}(x_{i_s}^v)| \leq \delta^{1/16}|C|$, and therefore $|N_{S_1}(b) \Delta N_{S_1}(b_*)| \leq 2\delta^{1/16}|C|$. Consequently,

$$\begin{aligned} |N_{S_1}(b) \cap Q_1| &\geq |Q_1| - 2\delta^{1/16}|C| \geq |Q_1| \left(1 - 2\delta^{1/16} \frac{|C|}{|Q_1|} \right) \\ &\geq |Q_1|(1 - 2\delta^{1/16}(4\delta^{-1/32})) \geq |Q_1|(1 - 10\delta^{1/32}). \end{aligned}$$

This shows that $|S_1 \cap K_2[B_3, Q_1]| \geq (1 - 10\delta^{1/32})|Q_1||B_3|$.

Similarly, we claim $|S_0 \cap K_2[B_3, Q_0]| \geq (1 - 10\delta^{1/32})|B_3||Q_0|$. Indeed, for all $b \in B_3$, $|N_{S_2}(b)| \leq (\epsilon'_1)^{1/16}|C|$ and, as above, $|N_{S_1}(b) \Delta N_{S_1}(b_*)| \leq 2\delta^{1/16}|C|$. Thus $|N_{S_0}(b) \Delta N_{S_0}(b_*)| \leq ((\epsilon'_1)^{1/16} + 2\delta^{1/16})|C|$. Therefore,

$$\begin{aligned} |N_{S_0}(b) \cap Q_0| &\geq |Q_0| - ((\epsilon'_1)^{1/4} + 2\delta^{1/16})|C| \\ &\geq |Q_0| \left(1 - ((\epsilon'_1)^{1/4} + 2\delta^{1/16}) \frac{|C|}{|Q_0|} \right) \\ &\geq |Q_0|(1 - ((\epsilon'_1)^{1/4} + 2\delta^{1/16})4\delta^{-1/32}) \\ &\geq |Q_0|(1 - 10\delta^{1/32}), \end{aligned}$$

where the last inequality uses the definition of ϵ'_1 . This shows

$$|S_0 \cap K_2[B_3, Q_0]| \geq (1 - 10\delta^{1/32})|B_3||Q_0|.$$

Now let

$$\begin{aligned} Q'_1 &= \{c \in Q_1 : |N_{S_1}(c) \cap B_3| \geq (1 - \sqrt{10}\delta^{1/64})|B_3|\} \quad \text{and} \\ Q'_0 &= \{c \in Q_0 : |N_{S_0}(c) \cap B_3| \geq (1 - \sqrt{10}\delta^{1/64})|B_3|\}. \end{aligned}$$

Since both

$$|S_1 \cap K_2[B_3, Q_1]| \geq (1 - 10\delta^{1/32})|Q_1||B_3| \text{ and}$$

$$|S_0 \cap K_2[B_3, Q_0]| \geq (1 - 10\delta^{1/32})|B_3||Q_0|,$$

we have that $|Q'_1| \geq (1 - \sqrt{10}\delta^{1/64})|Q_1|$ and $|Q'_0| \geq (1 - \sqrt{10}\delta^{1/64})|Q_0|$. Finally, let

$$C^* = \{c \in C : |N_{S_1}(c) \Delta N_{S_1}(x_{j_s}^w)| \leq \delta^{1/16}|B|\}.$$

Since $a_* \in A_C$, $|C^*| \geq (1 - \delta^{1/16})|C|$. Thus,

$$\begin{aligned} |Q'_1 \cap C^*| &\geq (1 - \sqrt{10}\delta^{1/64})|Q_1| - \delta^{1/16}|C| \\ &\geq (1 - \sqrt{10}\delta^{1/64})\frac{\delta^{1/32}}{4} - \delta^{1/16}|C| \geq \frac{\delta^{1/32}|C|}{10}. \end{aligned}$$

Similarly,

$$\begin{aligned} |Q'_0 \cap C^*| &\geq (1 - \sqrt{10}\delta^{1/64})|Q_0| - \delta^{1/16}|C| \\ &\geq (1 - \sqrt{10}\delta^{1/64})\frac{\delta^{1/32}}{4} - \delta^{1/16}|C| \geq \frac{\delta^{1/32}|C|}{10}. \end{aligned}$$

Consequently, there are $c_1 \in Q'_1 \cap C^*$ and $c_0 \in Q'_0 \cap C^*$. Since $c_0, c_1 \in C^*$, we can see that $|N_{S_1}(c_1) \Delta N_{S_1}(c_0)| \leq 2\delta^{1/16}|B|$. However, we also have that

$$\begin{aligned} |N_{S_1}(c_1) \cap N_{S_0}(c_0) \cap B_3| &\geq (1 - 2\sqrt{10}\delta^{1/64})|B_3| \\ &\geq (1 - 2\sqrt{10}\delta^{1/64})\delta^{1/32}\frac{|B|}{8} > 2\delta^{1/16}|B|. \end{aligned}$$

But this is a contradiction, since $N_{S_1}(c) \cap N_{S_0}(c_0) \cap B_3 \subseteq N_{S_1}(c_1) \Delta N_{S_1}(c_0)$. \square

Let $\ell_1 = \lceil \delta^{-4}m^4 \rceil$. Suppose $V_i V_j \notin \Psi$ and $1 \leq u \leq \ell_{ij}$ is such that W_{ij}^u is nontrivial. Define $\mathbf{W}_{ij}^u = \bigcup_{P_{ij}^\alpha \in W_{ij}^u} P_{ij}^\alpha$, let \mathbf{G}_{ij}^u be the bipartite graph $(V_i \cup V_j, \mathbf{W}_{ij}^u)$, and define

$$\rho_{ij}(u) = \frac{|\mathbf{W}_{ij}^u|}{|V_i||V_j|}.$$

By [Fact 2.23](#), \mathbf{G}_{ij}^u has $\text{dev}_2(\ell(\epsilon_2''(\ell))^{1/4})$ and

$$|\mathbf{W}_{ij}^u| = (1 \pm \ell(\epsilon_2''(\ell))^{1/4})\frac{|W_{ij}^u||V_i||V_j|}{\ell}.$$

Using the size estimate above and the fact that W_{ij}^u is nontrivial, we have

$$\rho_{ij}(u) = (1 \pm \ell(\epsilon_2''(\ell))^{1/4})\frac{|W_{ij}^u|}{\ell} \geq (1 \pm \ell(\epsilon_2''(\ell))^{1/4})\frac{\delta^{1/2}}{\ell} \geq 2\ell(\epsilon_2''(\ell))^{1/4},$$

where the last inequality is by choice of $\epsilon_2''(\ell)$. Set $p_{ij}(u) = \rho_{ij}(u)^{-1}/\ell_1$, and let $s_{ij}(u) = \lceil 1/p_{ij}(u) \rceil$. Observe that $\rho_{ij} p_{ij} = 1/\ell_1$. Note $(\epsilon_2''(\ell))^{1/4} \geq 10(1/s|V_i|)^{1/5}$

(since n is very large), and since W_{ij}^u is nontrivial and $\ell(\epsilon_2''(\ell))^{1/4} < \frac{1}{4}$,

$$\rho_{ij}(u) \geq (1 \pm \ell(\epsilon_2''(\ell))^{1/4}) |W_{ij}^u| / \ell \geq \delta^{1/2} \frac{(\ell/m_{ij})}{\ell} = \delta^{1/2} / m_{ij} \geq \frac{\delta^{1/2}}{m}.$$

Further, $0 < p_{ij}(u) < \rho_{ij}(u)/2$ since

$$p_{ij}(u) \leq (1 \pm \ell(\epsilon_2''(\ell))^{1/4})^{-1} m \delta^{-1/2} \ell_1^{-1} \leq m \delta^{-1/2} \frac{\delta^4}{m^4} \leq \frac{\delta^{3/2}}{m^3} < \frac{\rho_{ij}(u)}{2},$$

where the last inequality uses that $\rho_{ij}(u) \geq \delta^{1/2}/m$. Thus by [Lemma 2.22](#), there is a partition

$$\mathbf{W}_{ij}^u = \mathbf{W}_{ij}^u(0) \cup \dots \cup \mathbf{W}_{ij}^u(s_{ij}(u)),$$

so that $|\mathbf{W}_{ij}^u(0)| \leq \rho_{ij} p_{ij} (1 + \epsilon_1') |V_i| |V_j|$ and for each $1 \leq x \leq s_{ij}(u)$, the bipartite graph $\mathbf{G}_{ij}^u(x) := (V_i \cup V_j, \mathbf{W}_{ij}^u(x))$ has $\text{dev}_2(\ell(\epsilon_2''(\ell))^{1/4}, \rho_{ij} p_{ij})$, i.e., $\text{dev}_2(\ell(\epsilon_2''(\ell))^{1/4}, 1/\ell_1)$. Since $(\epsilon_2''(\ell))^{1/4} m < \epsilon_2(\ell_1)$, and by definition of ϵ_2'' , we have that for each $1 \leq x \leq s_{ij}(u)$, $\mathbf{G}_{ij}^u(x)$ has $\text{dev}_2(\epsilon_2(\ell_1), 1/\ell)$. Let

$$s_{ij} = \sum_{1 \leq u \leq m_{ij}} s_{ij}(u).$$

Give a reenumeration

$$\{X_{ij}^1, \dots, X_{ij}^{s_{ij}}\} = \{\mathbf{W}_{ij}^u(v) : 1 \leq v \leq s_{ij}(u), 1 \leq u \leq m_{ij}\}.$$

Then let $X_{ij}^{s_{ij}+1}, \dots, X_{ij}^{\ell_1}$ be any partition of $K_2[V_i, V_j] \setminus \bigcup_{x=1}^{s_{ij}} X_{ij}^x$.

For $V_i V_j \in \Psi$ choose a partition $K_2[V_i, V_j] = X_{ij}^1 \cup \dots \cup X_{ij}^{\ell_1}$ such that for each $1 \leq x \leq \ell_1$, X_{ij}^x has $\text{dev}_2(\epsilon_2(\ell_1), 1/\ell_1)$ (such a partition exists by [Lemma 2.22](#)). Now define \mathcal{Q} to be the decomposition of V with

$$\mathcal{Q}_1 = \{V_i : i \in [t]\} \quad \text{and} \quad \mathcal{Q}_2 = \left\{ X_{ij}^v : v \leq \ell_1, ij \in \binom{[t]}{2} \right\}.$$

We claim this is a $(t, \ell_1, \epsilon_1, \epsilon_2(\ell_1))$ -decomposition of V . Indeed, by construction, any $xy \in \binom{V}{2}$ which is not in an element of \mathcal{Q}_2 satisfying $\text{disc}_2(\epsilon_2(\ell_1), 1/\ell_1)$ is in the set

$$\Gamma := \bigcup_{V_i V_j \notin \Psi} X_{ij}^{s_{ij}+1} \cup \dots \cup X_{ij}^{\ell_1}.$$

Observe that

$$|\Gamma| \leq \sum_{V_i V_j \notin \Psi} \sum_{u=1}^{m_{ij}} |\mathbf{W}_{ij}^u(0)| + \left| K_2[V_i, V_j] \setminus \left(\bigcup_{P_{ij}^\alpha \in W_{ij}} P_{ij}^\alpha \right) \right|. \tag{2}$$

We have that

$$\begin{aligned}
 \sum_{V_i V_j \notin \Psi} \sum_{u=1}^{m_{ij}} |\mathbf{W}_{ij}^u(0)| &\leq \sum_{V_i V_j \notin \Psi} m_{ij} (1 + \epsilon'_1) \rho_{ij} p_{ij} |V_i| |V_j| \\
 &\leq \binom{t}{2} m (1 + 2\epsilon'_1) \frac{(n/t)^2}{\ell_1} \\
 &= \delta^4 \binom{t}{2} (1 + 2\epsilon'_1) \frac{(n/t)^2}{m^3} \leq 2\delta^2 \frac{\binom{n}{2}}{m^3},
 \end{aligned}$$

where the last inequality is because n is large. Then, by definition of δ and m , this shows that $\sum_{V_i V_j \notin \Psi} \sum_{u=1}^{m_{ij}} |\mathbf{W}_{ij}^u(0)| \leq \epsilon_1 \binom{n}{2} / 2$. We also have that

$$\begin{aligned}
 \sum_{V_i V_j \notin \Psi} \left| K_2[V_i, V_j] \setminus \left(\bigcup_{P_{ij}^\alpha \in W_{ij}} P_{ij}^\alpha \right) \right| &\leq \sum_{V_i V_j \notin \Psi} \left(|V_i| |V_j| - |W_{ij}| (1 + \epsilon'_2(\ell)) \frac{|V_i| |V_j|}{\ell} \right) \\
 &= \sum_{V_i V_j \notin \Psi} |V_i| |V_j| \left(1 - \ell_{ij} (1 + \epsilon'_2(\ell)) \frac{1}{\ell} \right) \\
 &\leq \sum_{V_i V_j \notin \Psi} |V_i| |V_j| (1 - (1 - 2(\epsilon'_1)^{3/4})(1 + \epsilon'_2(\ell))) \\
 &\leq \sum_{V_i V_j \notin \Psi} |V_i| |V_j| (\epsilon'_1)^{1/8} \leq (\epsilon'_1)^{1/8} \binom{n}{2}.
 \end{aligned}$$

Combining these with (2) yields that $|\Gamma| \leq \epsilon_1 \binom{n}{2} / 2 + (\epsilon'_1)^{1/8} \binom{n}{2} \leq \epsilon_1 \binom{n}{2}$, and therefore, \mathcal{Q} is a $(t, \ell, \epsilon_1, \epsilon_2(\ell))$ -decomposition of V .

We now show that \mathcal{Q} is $\epsilon_1/6$ -homogeneous with respect to H . We show first that for any $W_{ij}^u W_{is}^v W_{js}^w \in \Omega_3$, $\mathbf{G}_{ijs}^{uvw} := (V_i \cup V_j \cup V_s, \mathbf{W}_{ij}^u \cup \mathbf{W}_{is}^v \cup \mathbf{W}_{js}^w)$ is $2\delta^{1/100}$ -homogeneous with respect to H , and second that almost all $xyz \in K_3^{(2)}(\mathbf{G}_{ijs}^{uvw})$ are in an $\epsilon_1/6$ -homogenous triad of \mathcal{Q} .

Fix $W_{ij}^u W_{is}^v W_{js}^w \in \Omega_3$. We know by Claim 3.2, that there is $\sigma \in \{0, 1\}$ such that

$$|R_\sigma \cap K_3[W_{ij}^u, W_{is}^v, W_{js}^w]| \geq (1 - \delta^{1/100}) |K_3[W_{ij}^u, W_{is}^v, W_{js}^w]|.$$

This implies, by (1) and definition of R_σ , that the following holds, where $E^1 = E$ and $E^0 = \binom{V}{3} \setminus E^1$ (recall $E = E(H)$):

$$\begin{aligned}
 |E^\sigma \cap K_3^{(2)}(\mathbf{G}_{ijs}^{uvw})| &\geq (1 - \delta^{1/100}) (1 - \epsilon''_1) |K_3[W_{ij}^u, W_{is}^v, W_{js}^w]| (1 - \ell^3 \epsilon'_2(\ell)) |V_i| |V_j| |V_s| \frac{1}{\ell^3} \\
 &= (1 - \delta^{1/100}) (1 - \epsilon''_1) (1 - \ell^3 \epsilon'_2(\ell)) \frac{|W_{ij}^u| |W_{is}^v| |W_{js}^w|}{\ell^3}
 \end{aligned}$$

On the other hand, note that by (1),

$$|K_3^{(2)}(\mathbf{G}_{ijs}^{uvw})| = |W_{ij}^u| |W_{is}^v| |W_{js}^w| (1 \pm \ell^3 \epsilon'_2(\ell)) \frac{|V_i| |V_j| |V_s|}{\ell^3}.$$

Combining this with the above, we see that

$$\begin{aligned} |E^\sigma \cap K_3^{(2)}(\mathbf{G}_{ijs}^{uvw})| &\geq (1 - \delta^{1/100})(1 - \epsilon''_1)(1 - \ell^3 \epsilon'_2(\ell))(1 + \ell^3 \epsilon'_2(\ell))^{-1} |K_3^{(2)}(\mathbf{G}_{ijs}^{uvw})| \\ &\geq (1 - 2\delta^{1/100}) |K_3^{(2)}(\mathbf{G}_{ijs}^{uvw})|, \end{aligned}$$

where the last inequality is by definition of ϵ'_2 and ϵ''_1 . This shows \mathbf{G}_{ijs}^{uvw} is $2\delta^{1/100}$ -homogeneous. We now show that almost all $xyz \in K_3^{(2)}(\mathbf{G}_{ijs}^{uvw})$ are in an $\epsilon_1/6$ -homogeneous triad of \mathcal{Q} . Set

$$\Sigma_0(ijs, uvw) = \{0, \dots, s_{ij}(u)\} \times \{0, \dots, s_{is}(v)\} \times \{0, \dots, s_{js}(w)\}.$$

Given $(x, y, z) \in \Sigma_0$, set

$$\mathbf{G}_{ijs}^{uvw}(x, y, z) = (V_i \cup V_j \cup V_s; \mathbf{W}_{ij}^u(x) \cup \mathbf{W}_{is}^v(y) \cup \mathbf{W}_{js}^w(z)).$$

Note that $K_3^{(2)}(\mathbf{G}_{ijs}^{uvw}) = \bigcup_{(x,y,z) \in \Sigma_0(ijs, uvw)} K_3^{(2)}(\mathbf{G}_{ijs}^{uvw}(x, y, z))$. Define

$$\begin{aligned} \Sigma_1(ijs, uvw) &= \{(x, y, z) \in \{0, \dots, s_{ij}(u)\} \times \{0, \dots, s_{is}(v)\} \times \{0, \dots, s_{js}(w)\} : x, y \text{ or } z \text{ is } 0\}, \end{aligned}$$

and set $\Sigma_2(ijs, uvw) = \Sigma_0(ijs, uvw) \setminus \Sigma_1(ijs, uvw)$. Note that by construction, for all $(x, y, z) \in \Sigma_2(ijs, uvw)$, $\mathbf{G}_{ijs}^{uvw}(x, y, z) \in \text{Triads}(\mathcal{Q})$. Observe that

$$\begin{aligned} \sum_{(x,y,z) \in \Sigma_1(ijs, uvw)} |K_3^{(2)}(\mathbf{G}_{ijs}^{uvw}(x, y, z))| &\leq |\mathbf{W}_{ij}^u(0)| |V_s| + |\mathbf{W}_{is}^v(0)| |V_j| + |\mathbf{W}_{js}^w(0)| |V_i| \\ &\leq (1 + \epsilon'_1) |V_i| |V_j| |V_s| (\rho_{ij} p_{ij} + \rho_{is} p_{is} + \rho_{js} p_{js}) \\ &= 3(1 + \epsilon'_1) |V_i| |V_j| |V_s| \frac{1}{\ell_1} \\ &\leq 3(1 + \epsilon'_1) \delta^4 |V_i| |V_j| |V_s| m^{-4} \\ &\leq 3(1 + \epsilon'_1) \delta^4 \left(|W_{ij}^u| |W_{is}^v| |W_{js}^w| \frac{1}{\ell^3} \right)^{-1} m^{-4} |K_3^{(2)}(\mathbf{G}_{ijs}^{uvw})| \\ &\leq 3(1 + \epsilon'_1) \delta^4 \left(\frac{\delta^{1/2}}{m} \right)^{-3} m^{-4} |K_3^{(2)}(\mathbf{G}_{ijs}^{uvw})| \\ &= 3(1 + \epsilon'_1) \delta^{1/2} m^{-1} |K_3^{(2)}(\mathbf{G}_{ijs}^{uvw})| \\ &< \delta |K_3^{(2)}(\mathbf{G}_{ijs}^{uvw})|, \end{aligned}$$

where the last inequality uses the definition of m . Let $\Sigma_3(ijs, uvw)$ be the set of $(x, y, z) \in \Sigma_2(ijs, uvw)$ such that

$$|E^\sigma \cap K_3^{(2)}(\mathbf{G}_{ijs}^{uvw}(x, y, z))| < (1 - \delta^{1/200})|K_3^{(2)}(\mathbf{G}_{ijs}^{uvw}(x, y, z))|,$$

and set

$$\Sigma_4(ijs, uvw) = \Sigma_2(ijs, uvw) \setminus \Sigma_3(ijs, uvw).$$

By definition, and since $\delta^{1/200} < \epsilon_1/6$, every triad of the form $K_3^{(2)}(\mathbf{G}_{ijs}^{uvw}(x, y, z))$ for $(x, y, z) \in \Sigma_4(ijs, uvw)$ is in an $\epsilon_1/6$ -homogeneous triad of \mathcal{Q} . We now show that $\bigcup_{(x,y,z) \in \Sigma_4(ijs, uvw)} K_3^{(2)}(\mathbf{G}_{ijs}^{uvw})$ is most of $K_3^{(2)}(\mathbf{G}_{ijs}^{uvw})$. Observe

$$\begin{aligned} |E^\sigma \cap K_3^{(2)}(\mathbf{G}_{ijs}^{uvw})| &\leq \sum_{(x,y,z) \in \Sigma_1(ijs, uvw)} |K_3^{(2)}(\mathbf{G}_{ijs}^{uvw}(x, y, z))| \\ &\quad + (1 - \delta^{1/200}) \sum_{(x,y,z) \in \Sigma_3(ijs, uvw)} |K_3^{(2)}(\mathbf{G}_{ijs}^{uvw}(x, y, z))| \\ &\quad + \sum_{(x,y,z) \in \Sigma_4(ijs, uvw)} |K_3^{(2)}(\mathbf{G}_{ijs}^{uvw}(x, y, z))| \\ &\leq \delta |K_3^{(2)}(\mathbf{G}_{ijs}^{uvw})| \\ &\quad + (1 - \delta^{1/200}) \sum_{(x,y,z) \in \Sigma_3(ijs, uvw)} |K_3^{(2)}(\mathbf{G}_{ijs}^{uvw}(x, y, z))| \\ &\quad + \sum_{(x,y,z) \in \Sigma_4(ijs, uvw)} |K_3^{(2)}(\mathbf{G}_{ijs}^{uvw}(x, y, z))|. \end{aligned}$$

Thus, since $|E^\sigma \cap K_3^{(2)}(\mathbf{G}_{ijs}^{uvw})| \geq (1 - 2\delta^{1/100})|K_3^{(2)}(\mathbf{G}_{ijs}^{uvw})|$,

$$\begin{aligned} (1 - 2\delta^{1/100} - \delta)|K_3^{(2)}(\mathbf{G}_{ijs}^{uvw})| &\leq (1 - \delta^{1/200}) \sum_{(x,y,z) \in \Sigma_3(ijs, uvw)} |K_3^{(2)}(\mathbf{G}_{ijs}^{uvw}(x, y, z))| \\ &\quad + \sum_{(x,y,z) \in \Sigma_4(ijs, uvw)} |K_3^{(2)}(\mathbf{G}_{ijs}^{uvw}(x, y, z))| \\ &= \sum_{(x,y,z) \in \Sigma_2(ijs, uvw)} |K_3^{(2)}(\mathbf{G}_{ijs}^{uvw}(x, y, z))| \\ &\quad - \delta^{1/200} \sum_{(x,y,z) \in \Sigma_3(ijs, uvw)} |K_3^{(2)}(\mathbf{G}_{ijs}^{uvw}(x, y, z))|. \end{aligned}$$

Rearranging this inequality, we have the following upper bound for the sum $\sum_{(x,y,z) \in \Sigma_3(ijs, uvw)} |K_3^{(2)}(\mathbf{G}_{ijs}^{uvw}(x, y, z))|$:

$$\begin{aligned}
\delta^{-1/200} \left(\sum_{(x,y,z) \in \Sigma_2(ijs,uvw)} |K_3^{(2)}(\mathbf{G}_{ijs}^{uvw}(x,y,z))| - (1-2\delta^{1/100}-\delta) |K_3^{(2)}(\mathbf{G}_{ijs}^{uvw})| \right) \\
\leq \delta^{-1/200} |K_3^{(2)}(\mathbf{G}_{ijs}^{uvw})| 3\delta^{1/100} \\
\leq 3\delta^{1/200} |K_3^{(2)}(\mathbf{G}_{ijs}^{uvw})|.
\end{aligned}$$

Consequently,

$$\sum_{(x,y,z) \in \Sigma_4(ijs,uvw)} |K_3^{(2)}(\mathbf{G}_{ijs}^{uvw}(x,y,z))| \geq |K_3^{(2)}(\mathbf{G}_{ijs}^{uvw})| (1 - 3\delta^{1/200}).$$

We have now established that $\bigcup_{(x,y,z) \in \Sigma_4(ijs,uvw)} K_3^{(2)}(\mathbf{G}_{ijs}^{uvw})$ covers most of $K_3^{(2)}(\mathbf{G}_{ijs}^{uvw})$, and for all $(x,y,z) \in \Sigma_4(ijs,uvw)$, $\mathbf{G}_{ijs}^{uvw}(x,y,z)$ is an $\epsilon_1/6$ -homogeneous triad of \mathcal{Q} . For all $(x,y,z) \in \Sigma_4(ijs,uvw)$, $W_{ij}^u(x)$, $W_{is}^v(y)$, $W_{js}^w(z)$ all have $\text{dev}_2(\epsilon_2(\ell_1), 1/\ell_1)$, and thus, by [Proposition 2.24](#), $\mathbf{G}_{ijs}^{uvw}(x,y,z)$ has $\text{dev}_{2,3}(\epsilon_1, \epsilon_2(\ell_1))$ with respect to H .

Using this and our lower bound on the size of Y_3 , we can now give the following lower bound on the number of triples $xyz \in \binom{V}{3}$ in a $\text{dev}_{2,3}(\epsilon_2(\ell_1), \epsilon_1)$ -regular triad of \mathcal{P} :

$$\begin{aligned}
\sum_{W_{ij}^u W_{is}^v W_{js}^w \in \Omega_3} \sum_{(x,y,z) \in \Sigma_4(ijs,uvw)} |K_3^{(2)}(\mathbf{G}_{ijs}^{uvw}(x,y,z))| \\
\geq \sum_{W_{ij}^u W_{is}^v W_{js}^w \in \Omega_3} (1 - 3\delta^{1/200}) |K_3^{(2)}(\mathbf{G}_{ijs}^{uvw})| \\
= (1 - 3\delta^{1/200}) \sum_{W_{ij}^u W_{is}^v W_{js}^w \in \Omega_3} |K_3^{(2)}(\mathbf{G}_{ijs}^{uvw})| \\
\geq (1 - 3\delta^{1/200})(1 - 3\delta^{1/2}) \binom{n}{3} \\
\geq (1 - \epsilon_1) \binom{n}{3},
\end{aligned}$$

where the last inequality is by definition of δ . This finishes the proof. \square

Appendix: Proof of [Proposition 2.24](#)

We will use the following fact:

Lemma A.1. *For all $\delta, r, \mu \in (0, 1]$ satisfying $2^{12}\delta < \mu^2 r^{12}$, the following holds. Suppose $G = (V_1 \cup V_2 \cup V_3, E)$ is a 3-partite graph such that for each $ij \in \binom{[3]}{2}$, $\| |V_i| - |V_j| \| \leq \delta |V_i|$ and $G[V_i, V_j]$ has $\text{dev}_2(\delta, r)$. Given $u_0 v_0 w_0 \in K_3^{(2)}(G)$, define*

$$\begin{aligned}
& K_{2,2,2}[u_0, v_0, w_0] \\
&= \{u_1 v_1 w_1 \in K_3[V_1, V_2, V_3] : \text{for each } \epsilon \in \{0, 1\}^3, (u_{\epsilon_1}, v_{\epsilon_2}, w_{\epsilon_3}) \in K_3^{(2)}(G)\}.
\end{aligned}$$

Then if $J := \{uvw \in K_3^{(2)}(G) : |K_{2,2,2}[u_0, v_0, w_0]| \leq (1 + \mu)r^9|V_1||V_2||V_3|\}$, we have that $|J| \geq (1 - \mu)r^3|V_1||V_2||V_3|$.

Proof. Let $K_{2,2,2}^G[V_1, V_2, V_3]$ be the set

$$\{(u_0, u_1, w_0, w_1, z_0, z_1) \in V_1^2 \times V_2^2 \times V_3^3 \\ : \text{for each } \epsilon \in \{0, 1\}^3, u_{\epsilon_1} w_{\epsilon_2} z_{\epsilon_3} \in R \cap K_3^{(2)}(G)\}.$$

By Theorem 3.5 in [11],

$$|K_{2,2,2}[V_1, V_2, V_3]| \leq r^{12}|V_1|^2|V_2|^2|V_3|^2 + 2^{12}\delta^{1/4}|V_1|^2|V_2|^2|V_3|^2.$$

Suppose towards a contradiction that $|J| > (1 - \mu)r^3|V_1||V_2||V_3|$. Then

$$|K_{2,2,2}[V_1, V_2, V_3]| \geq |J|(1 + \mu)r^9|V_1||V_2||V_3| > (1 - \mu^2)r^{12}|V_1|^2|V_2|^2|V_3|^2.$$

Combining with the above, this implies $r^{12} + 2^{12}\delta^{1/4} > (1 - \mu^2)r^{12}$, which implies $\mu^2 r^{12} < 2^{12}\delta^{1/4}$, a contradiction. \square

Proof of Proposition 2.24. Fix $0 < \epsilon < \frac{1}{2}$, $0 < d_2 < \frac{1}{2}$, and $0 < \delta \leq (d_2/2)^{48}$, and choose N sufficiently large.

Suppose $H = (V_1 \cup V_2 \cup V_3, R)$ is a 3-partite 3-graph on $n \geq N$ vertices and for each $i, j \in [3]$, $||V_i| - |V_j|| \leq \delta|V_i|$. Suppose $G = (V_1 \cup V_2 \cup V_3, E)$ is a 3-partite graph, where for each $1 \leq i < j \leq 3$, $G[V_i, V_j]$ has $\text{dev}_2(\delta, d_2)$, and assume $|R \cap K_3^{(2)}(G)| \leq \epsilon|K_3^{(2)}(G)|$. Let d be such that $|R \cap K_3^{(2)}(G)| = d|K_3^{(2)}(G)|$. By assumption $d \leq \epsilon$. Define $g(x, y, z) : \binom{V}{3} \rightarrow [0, 1]$ by

$$g(x, y, z) = \begin{cases} 1 - d & \text{if } xyz \in R \cap K_3^{(2)}(G), \\ -d & \text{if } xyz \in K_3^{(2)}(G) \setminus R, \\ 0 & \text{otherwise.} \end{cases}$$

Given $u_0 v_0 w_0 \in K_3^{(2)}(G)$, define

$$K_{2,2,2}[u_0, v_0, w_0] = \{u_1 v_1 w_1 \in K_3[V_1, V_2, V_3] \\ : \text{for each } (i, j, k) \in \{0, 1\}^3, (u_i, v_j, w_k) \in K_3^{(2)}(G)\}.$$

Let $\mu = d_2^{12}$. Note that $2^{12}\delta < (d_2/2)^{36} < d_2^{36} = \mu^2 d_2^{12}$. Set

$$J := \{uvw \in K_3^{(2)}(G) : |K_{2,2,2}[u_0, v_0, w_0]| \leq (1 + \mu)d_2^9|V_1||V_2||V_3|\}.$$

By Lemma A.1, we have that $|J| \geq (1 - \mu)d_2^3|V_1||V_2||V_3|$. Now set

$$I_1 = \{(u_0, u_1, w_0, w_1, z_0, z_1) \in V_1^2 \times V_2^2 \times V_3^3 \\ : \text{for each } (i, j, k) \in \{0, 1\}^3, u_i w_j z_k \in R \cap K_3^{(2)}(G)\}$$

and let

$$I_2 = \{(u_0, u_1, w_0, w_1, z_0, z_1) \in (V_1^2 \times V_2^2 \times V_3^3) \setminus I_1$$

$$: \text{for each } (i, j, k) \in \{0, 1\}^3, u_i w_j z_k \in K_3^{(2)}(G)\}.$$

Then

$$\begin{aligned} & \sum_{u_0, u_1 \in V_1} \sum_{w_0, w_1 \in V_2} \sum_{z_0, z_1 \in V_3} \prod_{(i, j, k) \in \{0, 1\}^3} g(u_i, w_j, z_k) \\ & \leq \left| \sum_{u_0, u_1 \in V_1} \sum_{w_0, w_1 \in V_2} \sum_{z_0, z_1 \in V_3} \prod_{(i, j, k) \in \{0, 1\}^3} g(u_i, w_j, z_k) \right| \\ & \leq \sum_{u_0, u_1 \in V_1} \sum_{w_0, w_1 \in V_2} \sum_{z_0, z_1 \in V_3} \left| \prod_{(i, j, k) \in \{0, 1\}^3} g(u_i, w_j, z_k) \right| \\ & = \sum_{(u_0, u_1, w_0, w_1, z_0, z_1) \in I_1} (1-d)^9 \\ & \quad + \sum_{(u_0, u_1, w_0, w_1, z_0, z_1) \in I_2} \left| \prod_{(i, j, k) \in \{0, 1\}^3} g(u_i, w_j, z_k) \right|. \end{aligned}$$

For each $(u_0, u_1, w_0, w_1, z_0, z_1) \in I_2$,

$$\left| \prod_{(i, j, k) \in \{0, 1\}^3} g(u_i, w_j, z_k) \right| \leq d(1-d)^8,$$

since at least one of the $g(u_i, w_j, z_k)$ is equal to $-d$, and $|-d| < |1-d|$ (since $d \leq \epsilon < \frac{1}{2}$). Thus we have, by above, that

$$\sum_{u_0, u_1 \in V_1} \sum_{w_0, w_1 \in V_2} \sum_{z_0, z_1 \in V_3} \prod_{(i, j, k) \in \{0, 1\}^3} g(u_i, w_j, z_k) \leq (1-d)^9 |I_1| + d(1-d)^8 |I_2|.$$

Note

$$\begin{aligned} |I_1| & \leq \sum_{u_0 w_0 z_0 \in J} |K_{2,2,2}(u_0, w_0, z_0)| + \sum_{u_0 w_0 z_0 \in R \setminus J} |K_{2,2,2}(u_0, w_0, z_0)| \\ & \leq |J|(1+\mu)d_2^9 |V_1||V_2||V_3| + |R \setminus J||R| \\ & \leq |R|(1+\mu)d_2^9 |V_1||V_2||V_3| + \mu d_2^3 |V_1||V_2||V_3| d |K_3^{(2)}(G)| \\ & \leq d |K_3^{(2)}(G)| (1+\mu)d_2^9 |V_1||V_2||V_3| + \mu d_2^3 |V_1||V_2||V_3| d |K_3^{(2)}(G)| \\ & \leq |V_1||V_2||V_3| |K_3^{(2)}(G)| (d(1+\mu)d_2^{12} + dd_2^{12}), \end{aligned}$$

where the last inequality is by definition of μ . By the counting lemma [11, Theorem 3.5], $|K_3^{(2)}(G)| \leq (1 + 2^3\delta^{1/4})|V_1||V_2||V_3|$. Therefore, we have that

$$|I_1| \leq |V_1|^2|V_2|^2|V_3|^2(1 + 2^3\delta^{1/4})(d(1 + \mu)d_2^{12} + dd_2^{12}) \leq 3dd_2^{12}|V_1|^2|V_2|^2|V_3|^2.$$

On the other hand, $|I_2| \leq |K_{2,2,2}[V_2, V_2, V_3]|$, which, by [11, Theorem 3.5], has size at most $(d_2^{12} + 2^{12}\delta^{1/4})|V_1|^2|V_2|^2|V_3|^2$. Combining the bounds above with the fact that $d \leq \epsilon$, we have that

$$\begin{aligned} \sum_{u_0, u_1 \in V_1} \sum_{w_0, w_1 \in V_2} \sum_{z_0, z_1 \in V_3} \prod_{(i, j, k) \in \{0, 1\}^3} g(u_i, w_j, z_k) \\ \leq (1 - d)^9 |I_1| + d(1 - d)^8 |I_2| \\ \leq |V_1|^2|V_2|^2|V_3|^2(3\epsilon d_2^{12} + \epsilon(d_2^{12} + 2^{12}\delta^{1/4})) \\ \leq 6\epsilon d_2^{12}|V_1|^2|V_2|^2|V_3|^2, \end{aligned}$$

where the last inequality is due to $\delta < (d_2/2)^{48}$. This shows that (H, G) has $\text{dev}_{2,3}(\delta, 6\epsilon)$, as required. □

References

[1] N. Alon, E. Fischer, and I. Newman, “Efficient testing of bipartite graphs for forbidden induced subgraphs”, *SIAM J. Comput.* **37**:3 (2007), 959–976. [MR](#) [Zbl](#)

[2] A. Chernikov and N. Hempel, “On n -dependent groups and fields, II”, *Forum Math. Sigma* **9** (2021), art. id. e38. [MR](#) [Zbl](#)

[3] A. Chernikov and S. Starchenko, “Definable regularity lemmas for NIP hypergraphs”, *Q. J. Math.* **72**:4 (2021), 1401–1433. [MR](#) [Zbl](#)

[4] A. Chernikov and H. Towsner, “Hypergraph regularity and higher arity VC-dimension”, preprint, 2020. [arXiv 2010.00726](#)

[5] A. Chernikov, D. Palacin, and K. Takeuchi, “On n -dependence”, *Notre Dame J. Form. Log.* **60**:2 (2019), 195–214. [MR](#) [Zbl](#)

[6] D. Conlon and J. Fox, “Bounds for graph regularity and removal lemmas”, *Geom. Funct. Anal.* **22**:5 (2012), 1191–1256. [MR](#) [Zbl](#)

[7] J. Fox and L. M. Lovász, “A tight lower bound for Szemerédi’s regularity lemma”, *Combinatorica* **37**:5 (2017), 911–951. [MR](#) [Zbl](#)

[8] J. Fox, J. Pach, and A. Suk, “Erdős–Hajnal conjecture for graphs with bounded VC-dimension”, *Discrete Comput. Geom.* **61**:4 (2019), 809–829. [MR](#) [Zbl](#)

[9] P. Frankl and V. Rödl, “Extremal problems on set systems”, *Random Structures Algorithms* **20**:2 (2002), 131–164. [MR](#) [Zbl](#)

[10] W. T. Gowers, “Lower bounds of tower type for Szemerédi’s uniformity lemma”, *Geom. Funct. Anal.* **7**:2 (1997), 322–337. [MR](#) [Zbl](#)

[11] W. T. Gowers, “Quasirandomness, counting and regularity for 3-uniform hypergraphs”, *Combin. Probab. Comput.* **15**:1-2 (2006), 143–184. [MR](#) [Zbl](#)

[12] W. T. Gowers, “Hypergraph regularity and the multidimensional Szemerédi theorem”, *Ann. of Math. (2)* **166**:3 (2007), 897–946. [MR](#) [Zbl](#)

- [13] D. Haussler, “Sphere packing numbers for subsets of the Boolean n -cube with bounded Vapnik–Chervonenkis dimension”, *J. Combin. Theory Ser. A* **69**:2 (1995), 217–232. [MR](#) [Zbl](#)
- [14] P. E. Haxell, B. Nagle, and V. Rödl, “An algorithmic version of the hypergraph regularity method”, pp. 439–448 in *Proceedings of the 46th Annual IEEE Symposium on Foundations of Computer Science* (Pittsburgh, PA, 2005), IEEE Computer Society Press, Washington, DC, 2005.
- [15] N. Hempel, “On n -dependent groups and fields”, *MLQ Math. Log. Q.* **62**:3 (2016), 215–224. [MR](#) [Zbl](#)
- [16] L. Lovász and B. Szegedy, “Regularity partitions and the topology of graphons”, pp. 415–446 in *An irregular mind*, edited by I. Bárány and J. Solymosi, Bolyai Soc. Math. Stud. **21**, János Bolyai Math. Soc., Budapest, 2010. [MR](#) [Zbl](#)
- [17] G. Moshkovitz and A. Shapira, “A short proof of Gowers’ lower bound for the regularity lemma”, *Combinatorica* **36**:2 (2016), 187–194. [MR](#) [Zbl](#)
- [18] G. Moshkovitz and A. Shapira, “A tight bound for hypergraph regularity”, *Geom. Funct. Anal.* **29**:5 (2019), 1531–1578. [MR](#) [Zbl](#)
- [19] B. Nagle, A. Poerschke, V. Rödl, and M. Schacht, “Hypergraph regularity and quasi-randomness”, pp. 227–235 in *Proceedings of the Twentieth Annual ACM-SIAM Symposium on Discrete Algorithms*, edited by C. Mathieu, SIAM, Philadelphia, PA, 2009. [MR](#) [Zbl](#)
- [20] V. Rödl and J. Skokan, “Regularity lemma for k -uniform hypergraphs”, *Random Structures Algorithms* **25**:1 (2004), 1–42. [MR](#) [Zbl](#)
- [21] V. Rödl and J. Skokan, “Counting subgraphs in quasi-random 4-uniform hypergraphs”, *Random Structures Algorithms* **26**:1-2 (2005), 160–203. [MR](#) [Zbl](#)
- [22] S. Shelah, “Strongly dependent theories”, *Israel J. Math.* **204**:1 (2014), 1–83. [MR](#) [Zbl](#)
- [23] S. Shelah, “Definable groups for dependent and 2-dependent theories”, *Sarajevo J. Math.* **13(25)**:1 (2017), 3–25. [MR](#) [Zbl](#)
- [24] E. Szemerédi, “Regular partitions of graphs”, pp. 399–401 in *Problèmes combinatoires et théorie des graphes* (Orsay, France, 1976), Colloq. Internat. CNRS **260**, CNRS, Paris, 1978. [MR](#) [Zbl](#)
- [25] C. Terry, “ VC_ℓ -dimension and the jump to the fastest speed of a hereditary \mathcal{L} -property”, *Proc. Amer. Math. Soc.* **146**:7 (2018), 3111–3126. [MR](#) [Zbl](#)
- [26] C. Terry and J. Wolf, “Irregular triads in 3-uniform hypergraphs”, preprint, 2021. [arXiv 2111.01737](#)

Received 11 Apr 2022. Revised 1 Nov 2022.

CAROLINE TERRY:

terry.376@osu.edu

Department of Mathematics, The Ohio State University, Columbus, OH, United States

Model Theory

msp.org/mt

EDITORS-IN-CHIEF

- Martin Hils Westfälische Wilhelms-Universität Münster (Germany)
hils@uni-muenster.de
- Rahim Moosa University of Waterloo (Canada)
rmoosa@uwaterloo.ca

EDITORIAL BOARD

- Sylvy Anscombe Université Paris Cité (France)
sylvy.anscombe@imj-prg.fr
- Alessandro Berarducci Università di Pisa (Italy)
berardu@dm.unipi.it
- Emmanuel Breuillard University of Oxford (UK)
emmanuel.breuillard@gmail.com
- Artem Chernikov University of California, Los Angeles (USA)
chernikov@math.ucla.edu
- Charlotte Hardouin Université Paul Sabatier (France)
hardouin@math.univ-toulouse.fr
- François Loeser Sorbonne Université (France)
francois.loeser@imj-prg.fr
- Dugald Macpherson University of Leeds (UK)
h.d.macpherson@leeds.ac.uk
- Alf Onshuus Universidad de los Andes (Colombia)
aonshuus@uniandes.edu.co
- Chloé Perin The Hebrew University of Jerusalem (Israel)
perin@math.huji.ac.il

PRODUCTION

- Silvio Levy (Scientific Editor)
production@msp.org

See inside back cover or msp.org/mt for submission instructions.

Model Theory (ISSN 2832-904X electronic, 2832-9058 printed) at Mathematical Sciences Publishers, 798 Evans Hall #3840, c/o University of California, Berkeley, CA 94720-3840 is published continuously online.

MT peer review and production are managed by EditFlow® from MSP.

PUBLISHED BY
 **mathematical sciences publishers**
nonprofit scientific publishing
<https://msp.org/>

© 2023 Mathematical Sciences Publishers

Model Theory

no. 2 vol. 2 2023

Celebratory issue on the occasion of
Ehud Hrushovski's 60th Birthday

Introduction	133
ASSAF HASSON, H. DUGALD MACPHERSON and SILVAIN RIDEAU-KIKUCHI	
Mock hyperbolic reflection spaces and Frobenius groups of finite Morley rank	137
TIM CLAUSEN and KATRIN TENT	
Rigid differentially closed fields	177
DAVID MARKER	
Definable convolution and idempotent Keisler measures, II	185
ARTEM CHERNIKOV and KYLE GANNON	
Higher amalgamation properties in measured structures	233
DAVID M. EVANS	
Residue field domination in some henselian valued fields	255
CLIFTON EALY, DEIRDRE HASKELL and PIERRE SIMON	
Star sorts, Lelek fans, and the reconstruction of non- \aleph_0 -categorical theories in continuous logic	285
ITAÏ BEN YAACOV	
An improved bound for regular decompositions of 3-uniform hypergraphs of bounded VC ₂ -dimension	325
CAROLINE TERRY	
Galois groups of large simple fields	357
ANAND PILLAY and ERIK WALSBERG	
Additive reducts of real closed fields and strongly bounded structures	381
HIND ABU SALEH and YA'ACOV PETERZIL	
Remarks around the nonexistence of difference closure	405
ZOÉ CHATZIDAKIS	
An exposition of Jordan's original proof of his theorem on finite subgroups of $GL_n(\mathbb{C})$	429
EMMANUEL BREUILLARD	
Higher internal covers	449
MOSHE KAMENSKY	