Model Theory

no. 2 vol. 3 2024

**Zilber's skew-field lemma**

Adrien Deloro

msp

**Model Theory**
Vol. 3, No. 2, 2024
https://doi.org/10.2140/mt.2024.3.571

# Zilber's skew-field lemma

Adrien Deloro

We revisit one of Zilber's early results in model-theoretic algebra, viz., definability in Schur's lemma. This takes place in a broader context than the original version from the seventies.

*La droite laisse couler du sable.*
*Toutes les transformations sont possibles.*
Paul Éluard

The present contribution discusses and proves a linearisation result originating in Zilber's early work. Let us note to begin:

(1) *o*-minimal dimension and Borovik–Morley–Poizat rank are examples of finite dimensions.

(2) All necessary definitions are in Section 2.1.

(3) I have preferred not to conflate $T$ with $\mathbb{K}$ in the statement.

(4) There are classical corollaries in Section 2.4.

(5) The result bears no relationship to indecomposable generation discussed in Section 2.5.

**Theorem** (Zilber's skew-field lemma). *Work in a finite-dimensional theory. Let $V$ be a definable, connected, abelian group and $S, T \leq \mathrm{DefEnd}(V)$ be two invariant rings of definable endomorphisms such that*

- *$V$ is irreducible as an $S$-module*;
- *$C(S) = T$ and $C(T) = S$, with centralisers taken in* $\mathrm{DefEnd}(V)$;
- *$S$ and $T$ are infinite*;
- *$S$ **or** $T$ is unbounded.*

*Then there is a definable skew-field $\mathbb{K}$ such that $V \in \mathbb{K}$-**Vect**$_{<\aleph_0}$; moreover, $S \simeq \mathrm{End}(V : \mathbb{K}$-**Vect**$)$ and $T \simeq \mathbb{K}\,\mathrm{Id}_V$ are definable.*

---

The present exposition contains results stemming from more general research pursued with Frank O. Wagner [Deloro and Wagner ≥ 2024].

*MSC2020:* primary 03C60, 16B70; secondary 16G99.
*Keywords:* Schur's lemma, Zilber's field theorem, definability, linearisation.

© 2024 The Author, under license to MSP (Mathematical Sciences Publishers).

Section 1 provides context. Section 2 discusses the statement, and gives all definitions. The proof is in Section 3.

## 1. Introduction

Section 1.1 explains the relation to Schur's Lemma. Section 1.2 makes some historical remarks. Section 1.3 discusses a more famous corollary on fields in abstract groups.

**1.1.** ***Schur's lemma.*** Among the early work of Zilber are a couple of gems in model-theoretic algebra. (More on Zilber's early work is in [Hodges 2024] in the present volume.) This article deals with one of the phenomena he discovered: *many* $\aleph_1$*-categorical groups interpret infinite fields*. The result, or the method, or the general line of thought, is often called *Zilber's field theorem*. It stems from Schur's lemma in representation theory:

**Lemma** (Schur's lemma). *Let $R$ be a ring and $V$ be a simple $R$-module. Then the covariance ring $\mathbb{F} = C_{\mathrm{End}(V)}(R)$ is a skew-field, $V$ is a vector space over $\mathbb{F}$, and $R \hookrightarrow \mathrm{End}_R(V)$.*

Zilber's deep observation is simple:

*in many model-theoretically relevant cases, $\mathbb{F}$ is definable.*

A precise and modern form of the latter statement, given as Corollary 1 in Section 2.4, is a straightforward consequence of the main theorem above. (One should remember that every module is actually a bimodule by introducing Schur's covariance ring.) I shall henceforth call it (in long form) the *Schur–Zilber skew-field lemma*, hoping that Boris will not mind being in good company. Far be it from me to minimise its significance by dubbing it a lemma instead of a theorem; quite the opposite as lemmas are versatile devices — methods.

**1.2.** ***Editorial fortune of the lemma.*** This subsection is a layman's attempt at providing historical remarks. I apologise for misconceptions.

• As one learns from [Curtis 1999, p. 139], Schur's lemma itself appears in [Schur 1904, §2, I.] with comment: "*der auch in der Burnside'schen Darstellung der Theorie eine wichtige Rolle spielt*".

• Before Zilber's result was known, Cherlin [1979, §4.2, Theorem 1] found a definable field independently. There interpretation is obtained by hand (and seemingly by miracle), without a general method. Cherlin heard about Zilber's work after completing his own; [Cherlin 1979, §1.4] is very informative.

• The lemma itself seems not to have drawn as much attention as its corollary on soluble groups (Section 1.3). There are few traces of the lemma as a stand-alone statement.

• All sources discussing the topic [Zilber 1977; 1984; Thomas 1983; Nesin 1989a; 1989b; Poizat 1987; Loveys and Wagner 1993; Borovik and Nesin 1994; Macpherson and Pillay 1995] rely on indecomposable generation (however, see Section 2.5).

• This is different in the *o*-minimal context, but [Peterzil et al. 2000, Theorem 2.6] has its own techniques. (The earlier [Nesin et al. 1991, Proposition 2.4], which bears no reference to Zilber, resembles the coordinatisation by hand of [Cherlin 1979].) This and the above item may have given the impression that the Schur–Zilber lemma is a finite Morley rank gadget; *the present contribution shows that it isn't.*

• Most sources focus on the ring *generated* by the action instead of going to the centraliser; exceptions are [Nesin 1989a; Macpherson and Pillay 1995]. Only the under-cited [Nesin 1989a] discusses rings and makes the connection with Schur's lemma, while [Macpherson and Pillay 1995, p. 487] notices resemblances between various linearisation results but concludes:

> *There appear to be no immediate implications between this and the results recorded here, though it looks similar to Theorem 1.2.*

The present contribution elucidates the desired relations.

• My own interest in the topic started when I read [Nesin 1989a] while preparing [Deloro 2016]. This resulted in a very partial version of the theorem, in finite Morley rank and using indecomposability. After I gave a talk on generalising "Zilber's field theorem" in Lyon in January 2016, Wagner shared numerous ideas, which will bear all their fruits in the collaboration [Deloro and Wagner $\geq$ 2024].

**1.3.** ***Fields in soluble groups.*** To some extent, the Schur–Zilber lemma is the poor relation of the following theorem [Zilber 1984, Corollary, p. 175] (currently undergoing generalisation by Wagner):

> *connected*, *nonnilpotent*, *soluble groups of finite Morley rank interpret infinite fields.*

I believe the significance of the latter principle has been exaggerated for three reasons.

(1) In the local analysis of simple groups of finite Morley rank, different soluble subquotients may interpret nonisomorphic fields. Since there are strongly minimal structures interpreting *different* infinite fields [Hrushovski 1992], any field structure could be a false lead. (For more on how experts approach the algebraicity conjecture on simple groups of finite Morley rank, and the influence of finite group theory instead of pure model theory, see [Cherlin 2024; Poizat 2024].)

(2) Fields obtained by this method can have "bad" properties, typically nonminimal multiplicative group [Baudisch et al. 2009].

(3) The corollary focused on abstract groups and distracted us from doing representation theory (see the remarkable [Borovik 2024]).

## 2. The theorem

Section 2.1 contains all necessary definitions. Section 2.2 justifies the structure of the statement. Section 2.3 discusses optimality, Section 2.4 gives corollaries, and Section 2.5 considers the relation to "indecomposable generation".

The general version of the skew-field lemma is a double-centraliser theorem, repeated below. Alternative names could have been "bimodule theorem" or "double-centraliser linearisation".

**Theorem.** *Work in a finite-dimensional theory. Let $V$ be a definable, connected, abelian group and $S, T \leq \mathrm{DefEnd}(V)$ be two invariant rings of definable endomorphisms such that*

- *$V$ is irreducible as an $S$-module (viz., in the definable, connected category);*
- *$C(S) = T$ and $C(T) = S$, with centralisers taken in* $\mathrm{DefEnd}(V)$;
- *$S$ and $T$ are infinite;*
- *$S$ **or** $T$ is unbounded.*

*Then there is a definable skew-field $\mathbb{K}$ such that $V \in \mathbb{K}$-**Vect**$_{<\aleph_0}$; moreover, $S \simeq \mathrm{End}(V : \mathbb{K}\text{-}\mathbf{Vect})$ and $T \simeq \mathbb{K}\,\mathrm{Id}_V$ are definable.*

It would be interesting to recast this kind of double-centraliser result in the abstract ring $S \otimes T$, with no reference to $V$. (This is not planned in [Deloro and Wagner $\geq$ 2024].)

### 2.1. *Definitions.*

• Connected: with no definable proper subgroup of finite index. (Since the context does not provide a DCC, not all definable groups have a connected component.)

• Bounded: which does not grow larger when taking larger models. (The algebraist may fix a saturated model with inaccessible cardinality and argue there; bounded then means small. Also see [Halevi and Kaplan 2023].)

• Type-definable: a bounded intersection of definable sets.

• Invariant: a bounded union of type-definable sets. (The name comes from the action of the Galois group of a "large" model. Section 2.2 gives reasons for considering the invariant category instead of the definable one.)

• Irreducible: no nontrivial proper submodule — a submodule being definable *and connected*. (This is weaker than usual algebraic simplicity, which would also exclude finite submodules. Model theory will handle those in its own way.)

• Finite-dimensional: which bears a reasonable dimension on interpretable sets. Here [Wagner 2020] would say *fine*, *integer-valued*, *finite-dimensional*. The definition is as follows.

**Definition** [Wagner 2020]. A theory $T$ is [fine, integer-valued] finite-dimensional if there is a dimension function dim from the collection of all interpretable sets in models of $T$ to $\mathbb{N} \cup \{-\infty\}$, satisfying the following for a formula $\varphi(x, y)$ and interpretable sets $X$ and $Y$:

- Invariance: If $a \equiv a'$ then $\dim(\varphi(x, a)) = \dim(\varphi(x, a'))$.
- Algebraicity: $X$ is finite nonempty if and only if $\dim(X)=0$, and $\dim(\varnothing)=-\infty$.
- Union: $\dim(X \cup Y) = \max\{\dim(X), \dim(Y)\}$.
- Fibration: If $f : X \to Y$ is an interpretable map such that $\dim(f^{-1}(y)) \geq d$ for all $y \in Y$, then $\dim(X) \geq \dim(Y) + d$.

The dimension extends to type-definable, and then to invariant sets; of course one should no longer expect nice additivity properties.

Except for a key "field definability lemma" (Section 2.5) we shall use little from [Wagner 2020]. There is an ACC and a DCC on definable, *connected* subgroups.

**2.2.** ***Explaining the statement.*** Our statement deviates from traditional versions in several respects, and we make three cases for three notions.

*Skew-fields rather than fields.* Schur's lemma produces a skew-field, and so does Zilber's model-theoretic version.

• This went first unnoticed since $\aleph_1$-categorical skew-fields are commutative (answering a question of Macintyre's, proved by Cherlin and Shelah — see note on [Borovik and Nesin 1994, p. 139] — and independently by Zilber [1977].)

• It is easy to construct, in tame geometry, so-called "quaternionic representations", where the Schur field is the skew-field of quaternions.

• Also, the subring $\langle A \rangle \leq \mathrm{End}(V)$ generated by a *commutative* group action can be smaller than its Schur skew-field $C_{\mathrm{End}(V)}(A)$: classical focus on the former (as in most sources) captures only partial geometric information.

So skew-fields are naturally unavoidable. (There remains the question of which skew-fields can arise in a finite-dimensional theory. Skew-fields abound in number theory, but arguably number theory is far from tame. One can also doubt that the more exotic objects constructed in [Cohn 1995] will be finite-dimensional. The bold would conjecture that infinite skew-fields in finite-dimensional theories are

commutative and real closed, commutative and algebraically closed, or quaternionic over a commutative real closed field. The more reasonable may be content with conjecturing that they are finite extensions of their centres. Either of these claims, if true, would have an impact on their stability-theoretic properties.)

*Rings rather than groups.* Let $V$ be an abelian group; then $\mathrm{End}(V)$ is a ring. This accounts for studying representations of *rings*.

- If $G \leq \mathrm{Aut}(V)$ is a definable acting group, the subring of $\mathrm{End}(V)$ it generates need not be definable (see "invariance" below). This may have baffled pioneers in the topic.
- Rings were long neglected after the seminal [Zilber 1977] (a remarkable exception being [Nesin 1989a]). Going to the enveloping ring, however, gives powerful results, inaccessible to group-theoretic reasoning; see [Borovik 2024].

*Invariance rather than definability.* Leaving definability may have stopped first investigators of the matter; it is however salutary.

- If $G \leq \mathrm{Aut}(V)$ is a definable group, then the generated subring $\langle G \rangle \leq \mathrm{End}(V)$ is $\bigvee$-definable; this is closer to definability than invariance is. However (see "skew-fields" above), $\langle G \rangle$ does not capture enough geometric information. The double-centraliser $C(C(G)) \geq \langle G \rangle$ is more adapted to Schur-style arguments.
- So let $R \leq \mathrm{End}(V)$ be a definable ring. Then Schur's covariance ring $C_{\mathrm{DefEnd}(V)}(R)$ need not be definable, but it is invariant. And if $R$ itself is invariant, $C_{\mathrm{DefEnd}(V)}(R)$ is too.

So model-theoretic invariance arises as naturally as centralisers do.

### 2.3. *Optimality.*

- Both $S$ and $T$ must be infinite.

Let $\mathbb{K}$ be a pure algebraically closed field of positive characteristic $p$ and $V = \mathbb{K}_+$, which is definably minimal. Now $\mathrm{DefEnd}(V)$ consists of quasi-$p$-polynomials, viz., of all maps $x \mapsto \sum_{k=-n}^{n} a_{p^k} \mathrm{Fr}_{p^k}$, where Fr is the Frobenius automorphism of relevant power, and $a_{p^k} \in \mathbb{K}$; there is no bound on $n$. Only the action of $\mathbb{F}_p$ commutes to all these. We then let $S = \mathrm{DefEnd}(V)$ and $T = \mathbb{F}_p$ (or vice-versa). The first is not definable.

- At least one must be unbounded.

For the same $V$, now let $S$ be the ring of all quasi-$p$-polynomials *with coefficients in* $\mathbb{F}_p$, viz., the subring of $\mathrm{DefEnd}(V)$ generated by $\mathrm{Fr}_p$ and its inverse. Then one easily sees that $C(S) = S$ is countable, and not definable.

On the other hand, it so happens that $S$-irreducibility can be relaxed to irreducibility as an $(S, T)$-bimodule [Deloro and Wagner $\geq$ 2024]. So in retrospect, the main theorem can be retrieved as a corollary to [Deloro and Wagner $\geq$ 2024, Theorem 2].

**2.4.** ***Corollaries.*** I give three corollaries, proved in Section 3.5. The first relates the main, "double-centraliser" theorem to Schur's lemma. The second retrieves what is called "Zilber's field theorem" in sources such as [Borovik and Nesin 1994]. The third is a variation coming from Nesin's work and isolated by Poizat.

**Corollary 1** (Schur–Zilber, one-sided form). *Work in a finite-dimensional theory. Let $V$ be a definable, connected, abelian group and $S \leq \mathrm{DefEnd}(V)$ be an invariant, unbounded ring of definable endomorphisms. Suppose that $V$ is irreducible as an $S$-module. Then $C_{\mathrm{DefEnd}(V)}(S)$ is a definable skew-field.*

Corollary 1 is, however, not equivalent to our main result, which also covers the case of unbounded $T$ and infinite $S$.

**Corollary 2** (see [Deloro 2016, Théorème IV.1]). *Work in a finite-dimensional theory. Let $V$ be a definable, connected, abelian group and $G \leq \mathrm{DefAut}(V)$ be a definable group such that $V$ is irreducible as a $G$-module* ***and*** *$C_{\mathrm{DefEnd}(V)}(G)$ is infinite. Then $T = C_{\mathrm{DefEnd}(V)}(G)$ is a definable skew-field (so the action of $G$ is linear).*

Corollary 2 (or a minor variation) unifies and should replace various results such as [Zilber 1984, Lemma 2; Loveys and Wagner 1993, Theorem 4; Nesin 1989a, Lemma 12; Macpherson and Pillay 1995, Theorem 1.2(b); Deloro 2016, Théorème IV.1; Peterzil et al. 2000, Theorem 2.6; Macpherson et al. 2000, Proposition 4.1]. However, there are no claims on finite generation.

**Corollary 3** (after Nesin and Poizat). *Work in a finite-dimensional theory. Let $V$ be a definable, connected, abelian group and $R \leq \mathrm{DefEnd}(V)$ be an invariant, unbounded, commutative ring of definable endomorphisms. Suppose there is an invariant group $G \leq \mathrm{DefAut}(V)$ such that*

- *$V$ is irreducible as a $G$-module;*
- *$G$ normalises $R$;*
- *$G$ is connected.*

*Then there is a definable skew-field $\mathbb{K}$ such that $V \in \mathbb{K}\text{-}\mathbf{Vect}_{<\aleph_0}$; moreover, $R \hookrightarrow \mathbb{K}\,\mathrm{Id}_V$ and $G \hookrightarrow \mathrm{GL}(V : \mathbb{K}\text{-}\mathbf{Vect})$.*

It would be interesting to relax the assumption on commutativity of $R$. Further generalisations are expected using endogenies instead of endomorphisms [Deloro and Wagner $\geq$ 2024].

**2.5.** ***Indecomposable generation (and how to avoid it).*** Contrary to widespread belief, the Schur–Zilber lemma has nothing to do with another celebrated result from Boris' early work: the "indecomposability theorem" [Zilber 1977, Theorem 3.3], which by analogy with the algebraic case I prefer to call the *Chevalley–Zilber*

*generation lemma* (again with hope that Boris will not mind being in good company). For more on the topic, see [Poizat 2024, §8].

Both results are often presented jointly, which serves neither clarity nor purity of methods. In contrast, the proof given here relies on another phenomenon.

**Lemma** (field definability; extracted from [Wagner 2020, Proposition 3.6]). *Work in a finite-dimensional theory. Let $\mathbb{K}$ be an invariant skew-field such that*

- *there is an upper bound on dimensions of type-definable subsets of $\mathbb{K}$;*
- *$\mathbb{K}$ contains an invariant, unbounded subset.*

*Then $\mathbb{K}$ is definable.*

The first clause is satisfied if there is a definable $\mathbb{K}$-vector space of finite $\mathbb{K}$-linear dimension.

## 3. The proofs

The corollaries are derived in Section 3.5. Let $V, S, T$ be as in the theorem. The proof is a series of claims arranged in propositions.

*Proof of Zilber's skew-field lemma.* It is convenient to let $T$ act from the right and treat $V$ as an $(S, T)$-bimodule.

**Proposition.** (i) *$T$ is a domain acting by surjections with finite kernels*; *for $t \in T \setminus \{0\}$ one has $Vt = V$.*

This will later be reinforced in (x).

*Proof.* (i) Let $t \in T \setminus \{0\}$. Then $0 < Vt$ is $S$-invariant, definable, and connected; by $S$-irreducibility $Vt = V$, so $t$ is onto. In particular, $T$ is a domain. Finally, $\dim \ker t = \dim V - \dim Vt = 0$, so $\ker t$ is finite. □

The global behaviour is difficult to control, so we go down to a more "local" scale with a suitable notion of lines.

### 3.1. *Lines.*

**Notation.** Let $\delta = \min\{\dim sV : s \in S \setminus \{0\}\}$ and $\Lambda = \{sV : \dim sV = \delta\}$ be the set of *lines*.

**Proposition.** (ii) *Every line is $T$-invariant.*

(iii) *If $L \in \Lambda$ and $s \in S$ are such that $sL \neq 0$, then $sL \in \Lambda$; in particular, $L \cap \ker s$ is finite.*

(iv) *$V$ is a finite sum of lines.*

(v) *$S$ is transitive on $\Lambda$.*

Items (iii) and (iv) will later be reinforced in (vi) and (ix), respectively.

*Proof.* (ii) This is obvious since $S$ and $T$ commute.

(iii) Say $L = s_0 V$. If $sL \neq 0$, then $0 < \dim sL = \dim((ss_0)V) \leq \dim(s_0 V) = \delta$, so by minimality of $\delta$ one has $sL \in \Lambda$. This also implies $\dim(L \cap \ker s) = \dim \ker s_{|L} = \dim L - \dim sL = 0$, and $L \cap \ker s$ is finite.

(iv) The subgroup $0 < \sum \Lambda \leq V$ is definable, connected, and $S$-invariant; by $S$-irreducibility, it equals $V$. Since dimension is finite, it is a finite sum.

(v) Let $L_1, L_2 \in \Lambda$, say $L_i = s_i V$. Now as above, $V = \sum_S sL_1 \not\leq \ker s_2$, so there is $s \in S$ such that $s_2 sL_1 \neq 0$. But then $0 < s_2 sL_1 = s_2 ss_1 V \leq s_2 V = L_2$, and equality holds. □

### 3.2. *Linearising lines.*

**Proposition.** (vi) *If $L \in \Lambda$ and $s \in S$ are such that $sL \neq 0$, then $L \cap \ker s = 0$.*

(vii) *$T$ acts by automorphisms on every line.*

The proof is different depending on whether $S$ or $T$ is unbounded.

*Proof if $T$ is unbounded.* (vi) Suppose $sL \neq 0$; we show $L \cap \ker s = 0$. By (v), $S$ is transitive on $\Lambda$, so there is $s' \in S$ with $s'sL = L$. Now $L \cap \ker s \leq L \cap \ker(s's)$, so we may assume that $sL = L$. Recall that $\ker s_{|L} = L \cap \ker s$ is finite by (iii). Considering $s_{|L}^2 : L \to L$, which is onto, we inductively find $|\ker s_{|L}^n| = |\ker s_{|L}|^n$, so $K = \sum_{n \in \mathbb{N}} \ker s_{|L}^n$ is either trivial or countably infinite. *Since $T$ is unbounded*, there is $t \in T \setminus \{0\}$ annihilating $K$. But $t$ has a finite kernel by (i), so $K = 0$, as desired.

(vii) Let $t \in T$. Then $\ker t$ is finite and $S$-invariant, while $S$ is infinite; so there is $s_0 \in S \setminus \{0\}$ with $s_0(\ker t) = 0$.

Since $s_0 \neq 0$ and $V = \sum \Lambda$ by (iv), there is $L_0 \in \Lambda$ such that $s_0 L_0 \neq 0$. Then $s_0(L_0 \cap \ker t) = 0$ so $L_0 \cap \ker t \leq L_0 \cap \ker s_0$ by (vi).

Now if $L$ is any other line, then there is $s \in S$ with $sL = L_0$ by (v). Therefore $s(L \cap \ker t) \leq L_0 \cap \ker t = 0$, and $L \cap \ker t \leq L \cap \ker s = 0$ by (vi) again.

So $\ker t$ intersects each line trivially. □

*Proof if $S$ is unbounded.* The strategy is different here and we first prove weakened versions in reverse order.

Weak (vii)′: We first prove that $T$ acts by automorphisms on *some* line. By (iv), $V = \sum \Lambda$ is a finite sum, so there are $L_1, \ldots, L_n$ such that $\bigcap_{i=1}^n \mathrm{Ann}_S(L_i) = 0$. In particular $(S, +) \hookrightarrow \prod_i S/\mathrm{Ann}_S(L_i)$ as abelian groups. *Since $S$ is unbounded*, there exists some line $L$ such that the quotient group $\Sigma = S/\mathrm{Ann}_S(L)$ is unbounded. Let $t \in T \setminus \{0\}$. Then $K = \sum_{n \in \mathbb{N}} \ker t_{|L}^n$ is either trivial or countably infinite. Since $\Sigma$ is unbounded, there is $\sigma \in \Sigma \setminus \{0\}$ annihilating $K$, i.e., there is $s \in S$ annihilating $K$ but not $L$. By (iii) this shows $K = 0$, as desired.

Weak (vi)′: We next prove: *if $T$ acts by automorphisms on $L$*, then for $s \in S$ with $sL \neq 0$ one has $L \cap \ker s = 0$. Indeed, $L \cap \ker s$ is finite by (iii). Since $T$ is infinite there is $t \in T \setminus \{0\}$ with $(L \cap \ker s)t = 0$, but $t$ induces an automorphism of $L$. This proves (vi), but only for lines on which $T$ acts by automorphisms.

(vii) and (vi): By (vii)′, let $L$ be a line on which $T$ acts by automorphisms and $L'$ be another line. Then by transitivity (v), there is $s \in S$ with $sL = L'$. Suppose $w \in L' \cap \ker t$. Then there is $v \in L$ with $sv = w$. Now $s(vt) = (sv)t = wt = 0$, so $vt \in L \cap \ker s = 0$. Since $T$ acts by automorphisms on $L$, (vi)′ implies $v = 0$ and $w = 0$, as desired. □

Since it is unclear at this stage whether every element belongs to a line, we cannot immediately conclude that $T$ acts by automorphisms; this requires writing $V$ as a direct sum.

**3.3.** ***Globalising local geometries.*** Instead of *morphism of $T$-modules*, we simply say *$T$-covariant* map. We tend to reserve it for definable maps, even implicitly.

**Proposition.** (viii) *Lines are complemented as $T$-modules, viz., for $L \in \Lambda$ there is a definable, connected, $T$-invariant $H \leq V$ with $V = L \oplus H$.*

(ix) *$V$ is a finite, **direct** sum of lines.*

(x) *$T$ is a skew-field acting by automorphisms.*

*Proof.* (viii) Say $L = s_0 V$. Since $V = \sum_S sL$ by (iv) and (v), there is $s \in S$ with $s_0 sL \neq 0$, so $0 < s_0 sL = s_0 s s_0 V \leq L$. Let $s_1 = s_0 s$, so that $L = s_1 V = s_1 L$. Then for $v \in V$ there is $\ell \in L$ with $s_1 v = s_1 \ell$; in particular, $v = \ell + (v - \ell)$ with $\ell \in L$ and $v - \ell \in \ker s_1$. Therefore $H = \ker s_1$ is such that $V = L + H$; it also is $T$-invariant as $S$ and $T$ commute. Now $L \cap H = L \cap \ker s_1 = 0$ by (vi), so actually $V = L \oplus H$. Connectedness of $H$ follows.

Since $V = L \oplus H$ is a direct decomposition as a $T$-module, the associated projections are $T$-covariant (viz., morphisms of $T$-modules).

(ix) As long as possible, we recursively construct lines $L_1, \ldots, L_i$ with direct complements $H_j$ (as definable, connected $T$-modules) satisfying

> for $j \leq i$, one has $L_j \leq \bigcap_{k<j} H_k$ (viz., each new line is contained in all previous complements).

The construction starts by (viii). Now suppose $L_1, \ldots, L_i$ and $H_1, \ldots, H_i$ are as claimed. A quick induction yields:

$$V = \left( \bigoplus_{j=1}^{i} L_j \right) \oplus \left( \bigcap_{j=1}^{i} H_j \right).$$

Let $q$ project $V$ onto $\bigcap_{j=1}^{i} H_i$ with kernel $\bigoplus_{j=1}^{i} L_j$. Then $q$ is $T$-covariant, so $q \in C(T) = S$. If $\bigoplus_{j=1}^{i} L_j < V$, then $q \neq 0$. Now $V = \sum \Lambda$ so there is $L' \in \Lambda$

such that $qL' \neq 0$. Then let $L_{i+1} = qL' \in \Lambda$; it satisfies $L_{i+1} \leq \bigcap_{j=1}^{i} H_j$. Picking a complement as in (viii), we have reached stage $i+1$.

However the process must terminate because $\dim \bigoplus_{j=1}^{i} L_j = \delta \cdot i$ remains bounded by $\dim V$. So at some stage one obtains $\bigoplus_{j=1}^{i} L_j = V$, as wanted.

(x) Say $V = \bigoplus_{i=1}^{n} L_i$ by (ix). Then for $t \in T$ one has $\ker t = \bigoplus_{i=1}^{n} (L_i \cap \ker t) = 0$ by (vii). □

Hence $T$ is a skew-field and $V \in T$-**Vect**, but we still fall short of definability.

**3.4.** ***Definability.*** We return to lines. The next result is of a purely auxiliary nature.

**Proposition.** (xi) *Let $L_1, L_2 \in \Lambda$. If $\sigma : L_1 \simeq L_2$ is definable and $T$-covariant, then there is an* ***invertible*** *$s \in S^{\times}$ inducing $\sigma$.*

*Proof.* (xi) Using (viii), write $V = L_1 \oplus H_1$ for some $\pi_1 \in S$ with $L_1 = \operatorname{im} \pi_1$ and $H_1 = \ker \pi_1$.

If $L_2 \cap H_1 = 0$, then $H_1$ is a common direct complement for $L_1$ and $L_2$. Glue $\sigma : L_1 \to L_2$ with $\mathrm{Id}_H$ to produce a $T$-covariant map, viz., an element of $C_{\mathrm{DefEnd}(V)}(T) = S$, inducing $\sigma$. It clearly is invertible.

If $L_2 \leq H_1$, then the process proving (ix) enables us to take $L_1$ and $L_2$ as the first two lines in a direct sum decomposition. Consider the map given on $L_1$ by $\sigma$, on $L_2$ by $\sigma^{-1}$, and on the remaining sum by 1. It is $T$-covariant and bijective, hence invertible in $S$; it induces $\sigma$.

The case $0 < L_2 \cap H_1 < L_2$ cannot happen, for then $\ker \pi_{1|L_2} \geq L_2 \cap H_1 > 0$ so by definition of lines, $\pi_1 L_2 = 0$ and $L_2 \leq H_1$. □

**Notation.** For $L \in \Lambda$, by (viii) there exists a definable, connected, $T$-invariant $H$ such that $V = L \oplus H$.

- Let $\pi_L$ be the relevant projection and $S_L = \pi_L S \pi_L$.
- Also let $T_L \leq \mathrm{DefEnd}(L)$ be the image of $T$.

In full rigour, $S_L$ also depends on the complement chosen; we omit it from the notation. This will not create difficulties.

**Proposition.** (xii) *$S_L$ and $T_L$ are skew-fields contained in* $\mathrm{DefEnd}(L)$.

(xiii) *Inside* $\mathrm{DefEnd}(L)$ *one has $C(S_L) = T_L$ and $C(T_L) = S_L$.*

(xiv) *$T$ is definable.*

*Proof.* In case $T$ is unbounded, one may directly jump to (xiv).

(xii) Keep in mind that $S_L$ is an additive subgroup of $S$ closed under multiplication but it need not contain 1. (Sometimes $S_L$ is called a *subrng*, for "subring without identity".) However, $S_L$ per se is a ring with identity $\pi_L$, as the latter acts on $L$ as $\mathrm{Id}_L$. Moreover, if $\pi_L s \pi_L$ annihilates $L$, then since it annihilates the chosen direct

complement, it is 0 as an endomorphism of $V$, viz., $\pi_L s \pi_L = 0$ in $S$. So $S_L$ can be viewed as a subring of $\mathrm{DefEnd}(L)$, and it is exactly the subring of restrictions-corestrictions $\left\{ s_{|L}^{|L} : s \in \mathrm{Stab}_S(L) \right\}$. (This explains why the complement plays no role in our construction. It is however useful to have both points of view on $S_L$.)

Let $s \in S_L \setminus \{0\}$. Then $sL = L$, so by (vi) and since $S$ and $T$ commute, it induces some $T$-covariant automorphism $\sigma$ of $L$; by (xi) there is $s' \in S^\times$ inducing $\sigma$. Now $\pi_L s'^{-1} \pi_L$ is a two-sided inverse of $s$ in $S_L$. This proves that $S_L$ is a skew-field. So is $T$ by (x); now the restriction map $T \to T_L$, which is onto by definition, is injective since $T$ acts by automorphisms. Therefore $T_L$ is a skew-field as well.

(xiii) One of them is easy. Let $f : L \to L$ be a definable, $T_L$-covariant morphism, viz., $f \in C_{\mathrm{DefEnd}(L)}(T_L)$. By definition, $f$ commutes with the action of $T$. Take any $T$-invariant direct complement $H$ and set $\hat{f} = 0$ on $H$. Then $\hat{f} : V \to V$ is $T$-covariant. Hence $\hat{f} \in C(T) = S$ and $\pi_L \hat{f} \pi_L = f \in S_L$.

Now let $g : L \to L$ be definable and $S_L$-covariant, viz., $g \in C_{\mathrm{DefEnd}(L)}(S_L)$. We aim at extending $g$ to an $S$-covariant endomorphism of $V$.

For $M \in \Lambda$ first use transitivity (v) to choose $s \in S$ with $sL = M$. By (xi) we may assume $s \in S^\times$. Notice that $sgs^{-1}$ leaves $M$ invariant, and let $g_M \in \mathrm{DefEnd}(M)$ be the induced map. We claim that this does not depend on the choice of $s$. Indeed let $s'$ be another invertible choice, giving rise to $g'_M$. Then $s^{-1}s'$ induces an element of $S_L$, so $g$ commutes with it and we find $g_M = g'_M$.

We deduce as follows that $g_M \in C(S_M)$. For if $\eta \in S_M$ then we may assume $\eta \neq 0$ so by (xi) it is induced by an invertible element $h \in S^\times$ normalising $M$. Then $s' = hs$ is another invertible element taking $L$ to $M$. By the preceding paragraph, $s' g s'^{-1} = h g_M h^{-1}$ and $sgs^{-1} = g_M$ agree on $M$, so $g_M$ commutes with $\eta$ in the ring $S_M$.

We even prove: if $s \in S$ induces $\sigma : M \simeq N$, then $g_N \sigma = \sigma g_M$. Both are maps from $M$ to $N$. By (xi), we freely suppose $s$ invertible and pick invertible $s_M$, $s_N$ inducing $L \simeq M, N$. Then $s'_M = s^{-1} s_N \in S$ takes $L$ to $M$, so $s'_M g s'^{-1}_M$ agrees with $s_M g s_M^{-1} = g_M$ on $M$. Thus for arbitrary $m \in M$ we find

$$\begin{aligned} g_N \sigma(m) &= s s^{-1} \cdot s_N g s_N^{-1} \cdot s(m) \\ &= s \cdot (s^{-1} s_N) g (s_N^{-1} s)(m) = s g_M(m) = \sigma g_M(m). \end{aligned}$$

Therefore $g_N \sigma = \sigma g_M$, as claimed.

Finally take a direct sum $V = \bigoplus L_i$ as in (ix) and let $\hat{g}\left(\sum \ell_i\right) = \sum g_{L_i}(\ell_i)$, which is definable, well-defined, and extends $g$. We want to show $\hat{g} \in C(S)$. Let $s \in S$; also let $s_i = \pi_i s$. It is enough to show that $\hat{g}$ commutes with each $s_i$, and it is enough to show that they commute on each $L_j$. We have thus reduced to checking that $\hat{g}$ and $\sigma : L_j \simeq L_i$ induced by an element of $S$ commute. But this is the previous paragraph.

Hence $\hat{g} \in C(S) = T$ and therefore $g = \hat{g}_{|L} \in T_L$.

(xiv) Recall that $T$ is a skew-field by (x). If $T$ is unbounded we directly apply the field definability lemma from Section 2.5 (in that case, (xii) and (xiii) are not necessary). So we suppose that $S$ is unbounded.

We first prove that there is $L$ such that $S_L$ is unbounded. By (ix) take any decomposition $V = \bigoplus_{i=1}^{n} L_i$ and form projections $\pi_i$ onto $L_i$ with kernels $\bigoplus_{j \neq i} L_j$. Let $S_{i,j} = \pi_i S \pi_j$, an additive subgroup of $S$. We contend that one of them is unbounded. Indeed, the additive group homomorphism

$$S \to \prod_{i,j} S_{i,j}, \quad s \mapsto (\pi_i s \pi_j)_{i,j},$$

is injective since $\sum_k \pi_k = 1$. Now if $S_{L,M}$ and $S_{L',M'}$ are defined as the $S_{i,j}$, one easily sees $S_{L,M} \simeq S_{L',M'}$ definably; so all rings $S_L$ are unbounded.

A caveat: because $S_L$ and $T_L$ are mutual centralisers only in $\mathrm{DefEnd}(L)$ and not in $\mathrm{End}(L)$, the following paragraph cannot be made more trivial.

Therefore $S_L$ is an unbounded skew-field by (xii). By field definability of Section 2.5, $S_L$ is definable; now $\dim S_L > 0$ and $\dim L$ is finite, so $L \in S_L\text{-}\mathbf{Vect}_{<\aleph_0}$. In particular, all $S_L$-endomorphisms of $L$ are definable, so by (xiii) one has $T_L = \mathrm{End}(L : S_L\text{-}\mathbf{Vect})$. This is a skew-field by (xii), so the *linear* dimension over $S_L$ is 1 and $T \simeq T_L \simeq S_L^{\mathrm{op}}$ is unbounded as well. □

By field definability, the skew-field $T$ is definable and infinite, so $\dim T > 0$; now $\dim V$ is finite so $V \in T\text{-}\mathbf{Vect}_{<\aleph_0}$. Finally $S = C(T) = \mathrm{End}(V : T\text{-}\mathbf{Vect})$. Lines in our sense now coincide with 1-dimensional $T$-subspaces of $V$. This completes the proof of Zilber's skew-field lemma. □

**3.5.** ***Proofs of corollaries.*** We repeat the statements already given in Section 2.4.

**Corollary 1** (Schur–Zilber, one-sided form). *Work in a finite-dimensional theory. Let $V$ be a definable, connected, abelian group and $S \leq \mathrm{DefEnd}(V)$ be an invariant, unbounded ring of definable endomorphisms. Suppose that $V$ is irreducible as an $S$-module. Then $C_{\mathrm{DefEnd}(V)}(S)$ is a definable skew-field.*

*Proof.* Let $T = C_{\mathrm{DefEnd}(V)}(S)$. Notice that $T$ acts by surjective endomorphisms, so it is a domain. If it is finite, then it is a definable field. Otherwise we wish to apply our theorem, but it is unclear whether $S = C_{\mathrm{DefEnd}(V)}(T)$. It actually does not matter. Let $\hat{S} = C_{\mathrm{DefEnd}(V)}(T) \geq S$, which is invariant and unbounded. Moreover, $C_{\mathrm{DefEnd}(V)}(\hat{S}) = T$ as a "triple centraliser", and $V$ remains $\hat{S}$-minimal. So we apply the theorem with $(\hat{S}, T)$ and get definability of the skew-field $C_{\mathrm{DefEnd}(V)}(\hat{S}) = T$. □

**Corollary 2.** *Work in a finite-dimensional theory. Let $V$ be a definable, connected, abelian group and $G \leq \mathrm{DefAut}(V)$ be a definable group such that $V$ is irreducible*

*as a $G$-module* ***and*** *$C_{\mathrm{DefEnd}(V)}(G)$ is infinite. Then $T = C_{\mathrm{DefEnd}(V)}(G)$ is a definable skew-field* (*so the action of $G$ is linear*).

*Proof.* Let $T = C_{\mathrm{DefEnd}(V)}(G)$ and $S = C_{\mathrm{DefEnd}(V)}(T) \supseteq G$. Apply the theorem. □

**Corollary 3** (after Nesin and Poizat). *Work in a finite-dimensional theory. Let $V$ be a definable, connected, abelian group and $R \leq \mathrm{DefEnd}(V)$ be an invariant, unbounded, commutative ring of definable endomorphisms. Suppose there is an invariant group $G \leq \mathrm{DefAut}(V)$ such that*

- *$V$ is irreducible as a $G$-module*;
- *$G$ normalises $R$*;
- *$G$ is connected.*

*Then there is a definable skew-field $\mathbb{K}$ such that $V \in \mathbb{K}$-**Vect**$_{<\aleph_0}$; moreover, $R \hookrightarrow \mathbb{K}\,\mathrm{Id}_V$ and $G \hookrightarrow \mathrm{GL}(V : \mathbb{K}$-**Vect**$)$.*

*Proof.* Let $V$, $R$, $G$ be as in the statement. The proof follows that of [Poizat 1987, Théorème 3.8] closely. Let $W \leq V$ be $R$-irreducible, viz., minimal as a definable, connected, $R$-submodule; this exists by the DCC on definable, connected subgroups. Let $\mathfrak{p} = \mathrm{Ann}_R(W)$, a relatively definable ideal of $R$.

For $g \in G$, the definable, connected subgroup $gW \leq V$ is $R$-invariant, and hence an $R$-submodule. Clearly $\mathrm{Ann}_R(gW) = g\mathfrak{p}g^{-1}$. Moreover, $R/\mathfrak{p} \simeq R/(g\mathfrak{p}g^{-1})$.

Now, by $G$-irreducibility, $V = \sum_G gW$. So there are $g_1, \ldots, g_n \in G$ such that $V = \sum_{i=1}^n g_i W$. In particular, $\bigcap_{i=1}^n \mathrm{Ann}_R(g_i W) = 0$, and $R \hookrightarrow \prod R/(g_i \mathfrak{p} g_i^{-1})$. We just saw that all terms have the same cardinality. They are therefore unbounded.

Hence, the unbounded, commutative ring $R/\mathfrak{p}$ acts faithfully on the $R/\mathfrak{p}$-irreducible module $W$. Notice that $R/\mathfrak{p} \leq C_{\mathrm{DefEnd}(W)}(R/\mathfrak{p})$. By the theorem, the action of $R/\mathfrak{p}$ on $W$ is linearisable, and $R/\mathfrak{p}$ acts by scalars. The problem is to make this linear structure global without losing the action of $G$. But we know that $\mathfrak{p}$ is a prime ideal of $R$.

Now consider the set of prime ideals $P = \{h\mathfrak{p}h^{-1} : h \in G\}$. Suppose $\mathfrak{p}_1, \ldots, \mathfrak{p}_k \in P$ are distinct, say $\mathfrak{p}_i = h_i \mathfrak{p} h_i^{-1}$. By prime avoidance, there are elements $r_i \in \mathfrak{p}_i \setminus \bigcup_{j \neq i} \mathfrak{p}_j$. Then taking products, there are elements $r_i' \in \bigcap_{j \neq i} \mathfrak{p}_j \setminus \mathfrak{p}_i$. These are used to show that the sum $\sum_{i=1}^k h_i W$ is direct. In particular, $k \leq \dim V$ and $P$ is finite.

Since $G$ is connected and transitive on the finite set $P$, the latter is a singleton, namely $P = \{\mathfrak{p}\}$. But by faithfulness one had $\bigcap P = 0$, so $\mathfrak{p} = 0$.

Now let $r \in R \setminus \{0\}$. Then $r \notin \mathfrak{p}$ acts on $W$ as a nonzero scalar, so $W \leq \mathrm{im}\, r$. Since $r$ was arbitrary, for any $g \in G$, one has $gW \leq \mathrm{im}\, r$. Summing, $\mathrm{im}\, r = V$; this implies that $\ker r$ is finite. Then $K = \sum_{n \in \mathbb{N}} \ker r^n$ is either trivial or countably infinite. But by commutativity, it is $R$-invariant. Since $R$ is unbounded, there is $r_0 \in R \setminus \{0\}$ annihilating $K$. Since $r_0$ has a finite kernel in $V$, we see $K = 0$. Thus the domain $R$ acts by automorphisms on $V$.

Hence $\mathbb{F} = \mathrm{Frac}(R)$ is naturally a subring of $\mathrm{DefEnd}(V)$. By field definability, it is definable. Now $G$ normalises $\mathbb{F}$ and centralises it [Wagner 2020, §3.3]. In particular, $G$ centralises $R$. Therefore, $S = C_{\mathrm{DefEnd}(V)}(R)$, which contains $R$ by commutativity, also contains $G$. It follows that $V$ is $S$-irreducible and we apply the theorem globally to conclude. □

## Acknowledgements

It is a pleasure to thank, alphabetically: Tuna Altınel, who first taught me Zilber's field theorem; Alexandre Borovik and Gregory Cherlin, who took me further into model-theoretic algebra; Ali Nesin and Bruno Poizat, for sharing their culture and passion; Frank Wagner, to whom I have a great technical debt; and last but not least, for a long-lasting vision which inspired so many of us, Boris Zilber.

## References

[Baudisch et al. 2009] A. Baudisch, M. Hils, A. Martin-Pizarro, and F. O. Wagner, "Die böse Farbe", *J. Inst. Math. Jussieu* **8**:3 (2009), 415–443. MR Zbl

[Borovik 2024] A. Borovik, "Finite group actions on abelian groups of finite Morley rank", *Model Theory* **3**:2 (2024), 539–569.

[Borovik and Nesin 1994] A. Borovik and A. Nesin, *Groups of finite Morley rank*, Oxford Logic Guides **26**, Oxford Univ. Press, 1994. MR Zbl

[Cherlin 1979] G. Cherlin, "Groups of small Morley rank", *Ann. Math. Logic* **17**:1-2 (1979), 1–28. MR Zbl

[Cherlin 2024] G. Cherlin, "Around the algebraicity problem in odd type", *Model Theory* **3**:2 (2024), 505–538.

[Cohn 1995] P. M. Cohn, *Skew fields*, Encyclopedia of Mathematics and its Applications **57**, Cambridge Univ. Press, 1995. MR Zbl

[Curtis 1999] C. W. Curtis, *Pioneers of representation theory: Frobenius, Burnside, Schur, and Brauer*, History of Mathematics **15**, Amer. Math. Soc., Providence, RI, 1999. MR Zbl

[Deloro 2016] A. Deloro, *Un regard élémentaire sur les groupes algébriques*, mémoire d'habilitation, Université Pierre et Marie Curie, 2016, available at https://hal.science/tel-01989901.

[Deloro and Wagner ≥ 2024] A. Deloro and F. O. Wagner, "Endogenies and linearisation (working title)", in preparation.

[Halevi and Kaplan 2023] Y. Halevi and I. Kaplan, "Saturated models for the working model theorist", *Bull. Symb. Log.* **29**:2 (2023), 163–169. MR Zbl

[Hodges 2024] W. Hodges, "Meeting Boris Zilber", *Model Theory* **3**:2 (2024), 203–211.

[Hrushovski 1992] E. Hrushovski, "Strongly minimal expansions of algebraically closed fields", *Israel J. Math.* **79**:2-3 (1992), 129–151. MR Zbl

[Loveys and Wagner 1993] J. G. Loveys and F. O. Wagner, "Le Canada semi-dry", *Proc. Amer. Math. Soc.* **118**:1 (1993), 217–221. MR Zbl

[Macpherson and Pillay 1995] D. Macpherson and A. Pillay, "Primitive permutation groups of finite Morley rank", *Proc. London Math. Soc.* (3) **70**:3 (1995), 481–504. MR Zbl

[Macpherson et al. 2000] D. Macpherson, A. Mosley, and K. Tent, "Permutation groups in *o*-minimal structures", *J. London Math. Soc.* (2) **62**:3 (2000), 650–670. MR Zbl

[Nesin 1989a] A. Nesin, "Nonassociative rings of finite Morley rank", pp. 117–137 in *The model theory of groups* (Notre Dame, IN, 1985–1987), edited by A. Nesin and A. Pillay, Notre Dame Math. Lectures **11**, Univ. Notre Dame Press, 1989. MR Zbl

[Nesin 1989b] A. Nesin, "Solvable groups of finite Morley rank", *J. Algebra* **121**:1 (1989), 26–39. MR Zbl

[Nesin et al. 1991] A. Nesin, A. Pillay, and V. Razenj, "Groups of dimension two and three over *o*-minimal structures", *Ann. Pure Appl. Logic* **53**:3 (1991), 279–296. MR Zbl

[Peterzil et al. 2000] Y. Peterzil, A. Pillay, and S. Starchenko, "Simple algebraic and semialgebraic groups over real closed fields", *Trans. Amer. Math. Soc.* **352**:10 (2000), 4421–4450. MR Zbl

[Poizat 1987] B. Poizat, *Groupes stables*, Nur al-Mantiq wal-Ma'rifah **2**, Bruno Poizat, Villeurbanne, 1987. MR Zbl

[Poizat 2024] B. Poizat, "La conjecture d'algébricité, dans une perspective historique, et surtout modèle-théorique", *Model Theory* **3**:2 (2024), 479–504.

[Schur 1904] I. Schur, "Neue Begründung der Theorie der Gruppencharaktere", *Berl. Ber.* (1904), 406–432. Zbl

[Thomas 1983] S. R. Thomas, *Classification theory of simple locally finite groups*, Ph.D. thesis, Bedford College, University of London, 1983, available at https://www.proquest.com/docview/1791501281. MR

[Wagner 2020] F. O. Wagner, "Dimensional groups and fields", *J. Symb. Log.* **85**:3 (2020), 918–936. MR Zbl

[Zilber 1977] B. I. Zilber, "Группы и кольца, теория которых категорична", *Fund. Math.* **95**:3 (1977), 173–188. MR Zbl

[Zilber 1984] B. I. Zilber, "Some model theory of simple algebraic groups over algebraically closed fields", *Colloq. Math.* **48**:2 (1984), 173–180. MR Zbl

Received 14 Dec 2022. Revised 15 Oct 2023.

ADRIEN DELORO:

adrien.deloro@imj-prg.fr
Institut de Mathématiques de Jussieu, Sorbonne Université, Paris, France

# Model Theory

msp.org/mt


**EDITORS-IN-CHIEF**

Martin Hils — Westfälische Wilhelms-Universität Münster (Germany)
hils@uni-muenster.de

Rahim Moosa — University of Waterloo (Canada)
rmoosa@uwaterloo.ca

**EDITORIAL BOARD**

Sylvy Anscombe — Université Paris Cité (France)
sylvy.anscombe@imj-prg.fr

Alessandro Berarducci — Università di Pisa (Italy)
berardu@dm.unipi.it

Emmanuel Breuillard — University of Oxford (UK)
emmanuel.breuillard@gmail.com

Artem Chernikov — University of California, Los Angeles (USA)
chernikov@math.ucla.edu

Charlotte Hardouin — Université Paul Sabatier (France)
hardouin@math.univ-toulouse.fr

François Loeser — Sorbonne Université (France)
francois.loeser@imj-prg.fr

Dugald Macpherson — University of Leeds (UK)
h.d.macpherson@leeds.ac.uk

Alf Onshuus — Universidad de los Andes (Colombia)
aonshuus@uniandes.edu.co

Chloé Perin — The Hebrew University of Jerusalem (Israel)
perin@math.huji.ac.il

**PRODUCTION**

Silvio Levy — (Scientific Editor)
production@msp.org


See inside back cover or msp.org/mt for submission instructions.

Model Theory (ISSN 2832-904X electronic, 2832-9058 printed) at Mathematical Sciences Publishers, 798 Evans Hall #3840, c/o University of California, Berkeley, CA 94720-3840 is published continuously online.

MT peer review and production are managed by EditFlow® from MSP.

PUBLISHED BY
mathematical sciences publishers
nonprofit scientific publishing
https://msp.org/
© 2024 Mathematical Sciences Publishers

# Model Theory

no. 2 vol. 3 2024

Special issue on the occasion of the 75th birthday of
Boris Zilber

Introduction 199
MARTIN BAYS, MISHA GAVRILOVICH and JONATHAN KIRBY
Meeting Boris Zilber 203
WILFRID HODGES
Very ampleness in strongly minimal sets 213
BENJAMIN CASTLE and ASSAF HASSON
A model theory for meromorphic vector fields 259
RAHIM MOOSA
Revisiting virtual difference ideals 285
ZOÉ CHATZIDAKIS and EHUD HRUSHOVSKI
Boris Zilber and the model-theoretic sublime 305
JULIETTE KENNEDY
Approximate equivalence relations 317
EHUD HRUSHOVSKI
Independence and bases: theme and variations 417
PETER J. CAMERON
On the model theory of open generalized polygons 433
ANNA-MARIA AMMER and KATRIN TENT
New simple theories from hypergraph sequences 449
MARYANTHE MALLIARIS and SAHARON SHELAH
How I got to like graph polynomials 465
JOHANN A. MAKOWSKY
La conjecture d'algébricité, dans une perspective historique, et surtout modèle-théorique 479
BRUNO POIZAT
Around the algebraicity problem in odd type 505
GREGORY CHERLIN
Finite group actions on abelian groups of finite Morley rank 539
ALEXANDRE BOROVIK
Zilber's skew-field lemma 571
ADRIEN DELORO
Zilber–Pink, smooth parametrization, and some old stories 587
YOSEF YOMDIN
The existential closedness and Zilber–Pink conjectures 599
VAHAGN ASLANYAN
Zilber–Pink for raising to the power $i$ 625
JONATHAN PILA
Zilber's notion of logically perfect structure: universal covers 647
JOHN T. BALDWIN and ANDRÉS VILLAVECES
Positive characteristic Ax–Schanuel 685
PIOTR KOWALSKI
Analytic continuation and Zilber's quasiminimality conjecture 701
ALEX J. WILKIE
Logic Tea in Oxford 721
MARTIN BAYS and JONATHAN KIRBY