Model Theory

no. 2 vol. 4 2025

Definable compactness in o-minimal structures

Pablo Andújar Guerrero





Definable compactness in o-minimal structures

Pablo Andújar Guerrero

We characterize the notion of definable compactness for topological spaces definable in o-minimal structures, answering questions posed by Peterzil and Steinhorn (*J. London Math. Soc.* (2) **59**:3 (1999), 769–786) and Johnson (*J. Symb. Log.* **83**:4 (2018), 1477–1500). Specifically, we prove the equivalence of various definitions of definable compactness in the literature, including those in terms of definable curves, definable types and definable downward directed families of closed sets.

1. Introduction

In the study of first-order topological theories various definable notions of topological compactness have been helpful tools in tame settings by isolating classes of topological objects with desirable properties. The first of such notions was introduced in *o-minimal* theories for definable manifold spaces in [Peterzil and Steinhorn 1999], and corresponds to the property that every definable curve converges (here curve-compactness). This property was crucial in formulating Pillay's conjecture [2004, Conjecture 1.1] about o-minimal definably compact groups and their relationship with compact Lie groups. The research that led to the solution of this conjecture provided a deeper understanding of the relationship between neostability and tame topology. In particular results in o-minimal forking were used to reach another reasonable notion of o-minimal definable compactness [Peterzil and Pillay 2007]: for every definable family of closed sets with the finite intersection property there exists a finite set that intersects each set in the family (here transversalcompactness). On the other hand, Thomas [2012] and Walsberg [2015] generalized and applied curve-compactness to study topologies arising from o-minimal definable norms and metrics respectively. In collaboration with the author [Andújar Guerrero et al. 2021], they also explored a third notion of definable compactness within o-minimality: every downward directed definable family of nonempty closed sets has nonempty intersection (here filter-compactness). This definition has been independently studied by Johnson [2018] in the context of o-minimal quotient spaces, and in a general model-theoretic setting by Fornasiero [2015]. The o-minimal exploration of definable compactness (through the various notions mentioned above)

MSC2020: primary 03C64; secondary 54A05, 54D30.

Keywords: o-minimality, types, definable compactness, definable topological spaces.

has yielded in particular that, in many cases, definably compact spaces are definably homeomorphic to a set with the canonical o-minimal "Euclidean" topology (see Chapter 7 in [Andújar Guerrero 2021a]). Hrushovski and Loeser [2016] explored the tame topology of valued fields, including the introduction of yet another notion of definable compactness: every definable type has a limit (here type-compactness), where a limit is a point in every closed set in the type. Recently, some of these notions have also been approached in the p-adic setting in [Andújar Guerrero and Johnson 2024], and in the local o-minimal setting in [Fujita 2024].

In the present paper we prove the equivalence of all the above notions of definable compactness in the setting of Hausdorff definable topological spaces (Definition 5.1) in o-minimal structures. We also show that, if we drop the Hausdorffness assumption, curve-compactness is strictly weaker than all the other properties. Our main result is the following (see Sections 2.2 and 5.1 for definitions).

Theorem A. Fix an o-minimal structure $\mathcal{M} = (M, <, ...)$. Let (X, τ) be a definable topological space in \mathcal{M} . The following are equivalent:

- (1) Every downward directed definable family of nonempty τ -closed sets has nonempty intersection (filter-compactness).
- (2) Every definable type $p \in S_X(M)$ has a limit; i.e., there is a point in the intersection of every τ -closed set in p (type-compactness).
- (3) Every definable family of τ -closed sets that extends to a definable type in $S_X(M)$ has nonempty intersection.
- (4) Every consistent definable family of τ-closed sets admits a finite transversal; i.e., there exists a finite set that intersects every set in the family (transversalcompactness).
- (5) Every definable family C of τ -closed sets with the (m, n)-property, where $m \ge n > \dim \bigcup C$, has a finite transversal.
- (6) Every definable family C of τ -closed sets with the (m, n)-property, where $m \ge n$ and n is greater than the VC-codensity of C, has a finite transversal.

Moreover all the above imply and, if τ is Hausdorff or \mathcal{M} has definable choice, are equivalent to:

(7) Every definable curve in X is τ -completable (curve-compactness).

Theorem A and Remark 5.17 provide a positive answer to [Johnson 2018, Question 4.14], which asks whether curve-compactness and filter-compactness are equivalent for o-minimal definable manifold spaces.

In light of Theorem A we may present the following definition.

Definition 1.1. Let (X, τ) be a definable topological space in an o-minimal structure. We say that (X, τ) is *definably compact* if it satisfies any (all) of the conditions (1)–(6) in Theorem A.

We also prove that, if \mathcal{M} is an o-minimal expansion of the real line (\mathbb{R} , <), then every definable topological space in \mathcal{M} is definably compact if and only if it is compact in the classical sense (Corollary 5.19), which provides a positive answer (Remark 5.20) to part of [Peterzil and Steinhorn 1999, Question 2.5]. Furthermore, we show that definable compactness is definable uniformly in families (Proposition 5.22). Additionally, throughout the paper we comment on other reasonable notions of definable compactness, including definitions in terms of externally definable sets (Remark 5.9), chains (paragraph above Lemma 5.14) and nets (Remark 5.18).

We prove Theorem A using o-minimal combinatorial and geometrical facts which are either known to hold in more general settings or can be conjectured to do so. These facts include known characterizations of o-minimal nonforking formulas, the Alon–Kleitman–Matoušek (p, q)-theorem for VC classes, and two geometrical facts of independent interest about o-minimal types (Propositions 4.1 and 4.3), the first of which can be understood as a strong form of distal cell decomposition. Hence this paper can be seen as a road map to characterizing definable compactness in various NIP settings.

The structure of the paper is as follows. In Section 2 we include preliminaries. In Section 3 we gather the necessary literature results on Vapnik–Chervonenkis theory and on forking, extracting some easy corollaries. In Section 4 we prove our two main results about o-minimal types. In Section 5 we introduce our topological framework and prove Theorem A through a series of propositions, as well our other results on definable compactness.

This paper has been largely extracted from [Andújar Guerrero 2021b], which includes independent proofs within o-minimality of Fact 3.6 and of a version of Corollary 3.7 where $m \ge n > vc^*(S)$ is substituted by $m \ge n > \dim \bigcup S$, thus avoiding largely the use of forking or VC literature.

2. Preliminaries

2.1. *Conventions.* We fix a language $\mathcal{L} = \{<, ...\}$ and a first-order \mathcal{L} -structure $\mathcal{M} = (M, <, ...)$ expanding a dense linear order without endpoints. For a set of parameters *A* we denote by $\mathcal{L}(A)$ the expansion of \mathcal{L} by symbols for elements in *A*. Throughout unless otherwise specified "definable" means " $\mathcal{L}(M)$ -definable in \mathcal{M} ". All variables and parameters *x*, *a*, *u*... are *n*-tuples for some $n < \omega$. We denote the length of a variable or parameter *x* by |x|. We denote ordered pairs of tuples by $\langle x, y \rangle$. We use *n*, *m*, *k* and *l* to denote natural numbers.

Unless stated otherwise, any formula we consider is in $\mathcal{L}(M)$. For any formula $\varphi(x)$ and set $A \subseteq M^{|x|}$, let $\varphi(A) = \{a \in A : \mathcal{M} \models \varphi(a)\}$. For simplicity we write $\varphi(M)$ to mean $\varphi(M^{|x|})$. A (uniformly) definable family of sets is a family of the form $\{\varphi(M, b) : b \in \psi(M)\}$ for some formulas $\varphi(x, y)$ and $\psi(y)$, where we may always assume that $\varphi(x, y) \in \mathcal{L}$ (i.e., a formula without parameters). For any two formulas $\varphi(x)$ and $\psi(x)$, we write $\varphi(x) \vdash \psi(x)$ to mean $\mathcal{M} \models \forall x(\varphi(x) \rightarrow \psi(x))$). For sets of formulas p(x) and q(x) on free variables x, we write $p(x) \vdash q(x)$ to mean that, for every formula $\varphi(x) \in q(x)$, there is a finite subset $p'(x) \subseteq p(x)$ such that $\wedge p'(x) \vdash \varphi(x)$.

For a given *n*, let π denote the projection $M^{n+1} \to M^n$ onto the first *n* coordinates, where *n* will often be omitted and clear from context. For a family S of subsets of M^{n+1} let $\pi(S) = {\pi(S) : S \in S}$.

Recall that \mathcal{M} is o-minimal if every definable subset of M is a finite union of points and intervals with endpoints in $M \cup \{-\infty, +\infty\}$. For background in o-minimality we direct the reader to [van den Dries 1998]. We will use, in particular, the existence of uniform cell decompositions found in Chapter 3, Proposition 3.5 of that work. We use the following notation related to o-minimal cells: given two partial functions $f, g: M^n \to M \cup \{-\infty, +\infty\}$, with domains dom(f) and dom(g) respectively, let $(f, g) = \{\langle x, t \rangle : x \in \text{dom}(f) \cap \text{dom}(g), f(x) < t < g(x)\}$ (we relax thus the classical notation throughout by allowing that f and g have different domains). Whenever \mathcal{M} is o-minimal, we refer jointly to the order topology on Mand induced product topology on M^n as the *Euclidean topology*. Given a definable set $X \subseteq M^n$, we denote its closure in the Euclidean topology by cl(X), and its frontier by $\partial(X) = cl(X) \setminus X$. We also denote the o-minimal dimension of X by dim X.

2.2. *Intersecting families of sets and refinements.* We say that a family of sets S is *n*-consistent if every subfamily of cardinality at most *n* has nonempty intersection. A family is *consistent* if it is *n*-consistent for every *n*. We say that S is *n*-inconsistent if every subfamily of cardinality *n* has empty intersection.

A family of sets S has the (p, q)-property, for cardinals $p \ge q > 0$, if the sets in S are nonempty and, for every p distinct sets in S, there exist q among them with nonempty intersection. Note that S does not have the (p, q)-property if and only if it either contains the empty set or there exists a subfamily of S of size p that is q-inconsistent.

A family of sets S is *downward directed* if, for every $F_0, F_1 \in S$, there exists $F_2 \in S$ such that $F_2 \subseteq F_0 \cap F_1$. Equivalently if for every finite $\mathcal{F} \subseteq S$ there exists $F \in S$ with $F \subseteq \bigcap \mathcal{F}$.

Given a family of sets S and a set X let $X \cap S = S \cap X = \{S \cap X : S \in S\}$. Observe that, if S is downward directed, then, for every set X, it holds that $X \cap S$ is downward directed too. Given two families of sets S and F, we say that F is a refinement of S, or that F refines S if, for every $S \in S$, there exists $F \in F$ with $F \subseteq S$. Observe that, if F is a downward directed refinement of S, then, for every finite subfamily $\mathcal{G} \subseteq S$, there exists some $F \in F$ with $F \subseteq \bigcap \mathcal{G}$.

Given a family of sets S and a set X we say that X is a *transversal of* S if it intersects every set in S (i.e., $\emptyset \notin X \cap S$). In this paper we are interested in the property that a definable family of sets has a finite transversal, as a weakening of the property of having nonempty intersection (i.e., having a transversal of size 1).

The following lemma will be used throughout the paper. We leave the easy proof to the reader.

Lemma 2.1. Let S be a downward directed family of sets and X be a finite covering of a set X. If $S \cap X \neq \emptyset$ for every $S \in S$, then there exists some $Y \in X$ such that $S \cap Y \neq \emptyset$ for every $S \in S$. In particular, if S has a finite transversal, then $\bigcap S \neq \emptyset$.

2.3. *Type preliminaries.* All the types that we consider are consistent and, unless otherwise specified, complete over M. We denote the set of these types by S(M). We denote by $S_n(M)$ the set of *n*-types in S(M). We resort often and without warning to the common model-theoretic convention of identifying types with the family of sets defined by formulas in it. For a definable set $X \subseteq M^n$, we denote by $S_X(M)$ the family of all types $p \in S_n(M)$ with $X \in p$ (namely types that concentrate on X). We will investigate partial types which are downward directed,¹ and the refinement relation between partial types.

Recall that a type $p(x) \in S(M)$ is definable if, for every formula $\varphi(x, y) \in \mathcal{L}$, there is another formula $\psi(y) \in \mathcal{L}(M)$ such that $\psi(M) = \{b \in M^{|y|} : \psi(x, b) \in p(x)\}$. It is definable over $A \subseteq M$ if these formulas $\psi(y)$ can be chosen in $\mathcal{L}(A)$. Note that, if a type is definable, then its projection $\pi(p)$ is definable too.

Given a formula $\varphi(x)$ let $\varphi^1(x) = \varphi(x)$ and $\varphi^0(x) = \neg \varphi(x)$. Given a type p(x) and a formula $\varphi(x, y)$, recall that the restriction of p(x) to $\varphi(x, y)$ is the subtype $p|_{\varphi}(x) = \{\varphi^i(x, b) \in p(x) : i \in \{0, 1\}, b \in M^{|y|}\}$. We denote by $p|_{\varphi}^1(x)$ the restriction of p(x) to "positive" instances of $\varphi(x, y)$, i.e., $p|_{\varphi}^1(x) = \{\varphi(x, b) \in p(x) : b \in M^{|y|}\}$.

3. O-minimal VC theory and forking

3.1. VC *theory.* The following is an ad hoc presentation of the notion of VC-codensity and related results, with applications in Sections 3.2 and 5. For a more standard treatment of Vapnik–Chervonenkis (VC) theory in a model-theoretic context see [Aschenbrenner et al. 2016].

¹In the literature this property among types is also denoted by 1-*compressible*.

A pair (X, S), where X is a set and S is a family of subsets of X, is called a *set system*. For a subfamily $\mathcal{F} \subseteq S$, let BA (\mathcal{F}) denote the collection of *Boolean atoms* of \mathcal{F} , by which we mean the family of all maximal nonempty intersections of sets in $\mathcal{F} \cup \{X \setminus S : S \in \mathcal{F}\}$. The *dual shatter function* of S is the function $\pi_S^* : \omega \to \omega$ given by

$$\pi_{\mathcal{S}}^*(n) = \max_{\mathcal{F} \subseteq \mathcal{S}, \, |\mathcal{F}|=n} |\mathrm{BA}(\mathcal{F})|.$$

The VC-codensity of S, denoted by vc*(S), is the infimum over all real numbers $r \ge 0$ such that $\pi_S^*(n) = O(n^r)$ (that is, $\pi_S^*(n)/n^r$ is bounded at infinity). Observe that vc*(S) is independent of the ambient set X, and so throughout we omit it from our terminology. A theory T is *NIP* (not the independence property) if every definable family of sets in every model of T has finite VC-codensity. Every o-minimal theory is NIP [van den Dries 1998, Chapter 5].

For convenience we state the Alon–Kleitman–Matoušek (p, q)-theorem in terms of VC-codensity. For a finer statement see [Matoušek 2004, Theorem 4].

Fact 3.1 (Alon–Kleitman–Matoušek (p, q)-theorem [Matoušek 2004]). Let $p \ge q > 0$ be natural numbers and let S be a set system such that $vc^*(S) < q$. Then there is $n < \omega$ such that, for every finite subfamily $\mathcal{F} \subseteq S$, if \mathcal{F} has the (p, q)-property, then it has a transversal of size at most n.

The following easy corollary will be used in the proof of Corollary 3.7.

Corollary 3.2. Let $p \ge q > 0$ be natural numbers and let S be a set system such that $vc^*(S) < q$. If S has the (p, q)-property, then, for every $0 < q' < \omega$, there exists some natural number $p' = p'(q') \ge q'$ such that S has the (p', q')-property. In particular, S has the (ω, q') -property for every $0 < q' < \omega$.

Proof. Let S be as in the corollary, satisfying the (p, q)-property. Let n be as described by Fact 3.1. For any given q' > 0, let p' = n(q' - 1) + 1. Consider an arbitrary subfamily \mathcal{F} of S of size p'. By Fact 3.1, \mathcal{F} has a transversal A of size at most n. By the definition of p', there must exist some $a \in A$ such that $|F \in \mathcal{F} : a \in F| \ge q'$. It follows that S has the (p', q')-property. \Box

The following fact is a reformulation of the main result for weakly o-minimal structures (a class which contains o-minimal structures) in [Aschenbrenner et al. 2016] by Aschenbrenner, Dolich, Haskell, Macpherson and Starchenko. It was previously proved for o-minimal structures by Wilkie (unpublished) and Johnson and Laskowski [2010], and for o-minimal expansions of the field of reals by Karpinski and Macintyre [2000].

Fact 3.3 [Aschenbrenner et al. 2016, Theorem 6.1]. Let \mathcal{M} be an o-minimal structure and let \mathcal{S} be a definable family of subsets of M^n . Then vc^{*}(\mathcal{S}) $\leq n$.

We will apply Fact 3.3 in subsequent sections through the slight improvement given by the next corollary.

Corollary 3.4. Let \mathcal{M} be an o-minimal structure and let S be a definable family of sets with $n = \dim \bigcup S$. Then $vc^*(S) \le n$.

The proof of Corollary 3.4 follows immediately from the following lemma and o-minimal cell decomposition, the latter implying that, if X is a definable set in an o-minimal structure \mathcal{M} with dim $X \leq n$, then X admits a finite partition into definable subsets, each of which is in definable bijection with a subset of M^n .

Lemma 3.5. Let S be a set system and let X_1, \ldots, X_m be sets such that $\bigcup S \subseteq \bigcup_{i \le m} X_i$. Then

$$\operatorname{vc}^*(\mathcal{S}) = \max_{1 \le i \le m} \operatorname{vc}^*(X_i \cap \mathcal{S}).$$

Proof. First note that, for every $i \leq m$ and finite subfamily $\mathcal{F} \subseteq S$, we have $BA(X_i \cap \mathcal{F}) \leq BA(\mathcal{F}) + 1$, meaning that $\pi^*_{X_i \cap S}(n) \leq \pi^*_S(n) + 1$ for every *n*, and consequently $vc^*(X_i \cap S) \leq vc^*(S)$.

For the opposite inequality, let \mathcal{F} be a finite subfamily of \mathcal{S} . Observe that

$$BA(\mathcal{F}) \leq BA(X_1 \cap \mathcal{F}) + \dots + BA(X_m \cap \mathcal{F}).$$

Consequently

$$\pi_{\mathcal{S}}^*(n) \leq \pi_{X_1 \cap \mathcal{S}}^*(n) + \dots + \pi_{X_m \cap \mathcal{S}}^*(n)$$

for every *n*. It follows that, for any real number $r \ge 0$, if $\pi^*_{X_i \cap S}(n) = O(n^r)$ for all $i \le m$, then $\pi^*_S(n) = O(n^r)$. Hence there must exist some $i \le m$ such that $vc^*(S) \le vc^*(X_i \cap S)$.

Since throughout this paper p and q are employed as terminology for types, in subsequent sections we address the (p, q)-property in terms of m and n, e.g., the (m, n)-property.

3.2. Forking, dividing and definable types. In this section we recall some facts about nonforking formulas in o-minimal theories, and derive some consequences which we will need in Section 5. This is the subject of ongoing research among NIP theories [Simon 2015]. Throughout we fix an $|M|^+$ -saturated elementary extension $\mathcal{U} = (U, ...)$ of \mathcal{M} .

Recall that a formula $\varphi(x, b) \in \mathcal{L}(U)$ is said to *n*-divide over $A \subseteq U$, for some $n \ge 1$, if there exists a sequence of elements $(b_i)_{i < \omega}$ in $U^{|b|}$, with $\operatorname{tp}(b_i/A) = \operatorname{tp}(b/A)$ for every *i*, such that $\{\varphi(x, b_i) : i < \omega\}$ is *n*-inconsistent. Equivalently, $\varphi(x, b)$ is said to *n*-divide over *A* if the family $\{\varphi(U, b') : \operatorname{tp}(b'/A) = \operatorname{tp}(b/A)\}$ does not have the (ω, n) -property. A formula $\varphi(x, b)$ divides if it *n*-divides for some *n*. Conversely, a formula $\varphi(x, b)$ does not divide over *A* if and only if the family $\{\varphi(U, b') : \operatorname{tp}(b'/A) = \operatorname{tp}(b/A)\}$ has the (ω, n) -property for every *n*. Hence, not dividing is an intersection property.

A formula *forks over* A if it implies a finite disjunction of formulas that divide each over A. In NTP₂ theories (a class which includes NIP and simple theories) forking and dividing over a model are equivalent notions [Chernikov and Kaplan 2012, Theorem 1.1].

The next equivalence was proved first for o-minimal expansions of ordered fields² by Dolich [2004] (where he considered forking over small sets and not just models) and for unpackable VC-minimal theories, a class which includes o-minimal theories, by Cotter and Starchenko [2012]. The best generalization up to date is due to Simon and Starchenko [2014], and applies to a large class of dp-minimal theories (for details and precise definitions of unpackable VC-minimal and dp-minimal theory see [Cotter and Starchenko 2012; Simon and Starchenko 2014] respectively). We state the result for o-minimal theories.

Fact 3.6. Let *T* be an o-minimal \mathcal{L} -theory with monster model \mathcal{U} . Let $\mathcal{M} \models T$ and $\varphi(x, b) \in \mathcal{L}(U)$. The following are equivalent:

- (i) $\varphi(x, b)$ does not fork (equivalently, by [Chernikov and Kaplan 2012], does not divide) over *M*.
- (ii) $\varphi(x, b)$ extends to an *M*-definable type in $S_{|x|}(U)$.

In Section 5.2 we will apply Fact 3.6 in the form of the following corollary.

Corollary 3.7. Let \mathcal{M} be an o-minimal structure and S be a definable family of nonempty subsets of M^k . If there exist natural numbers $m \ge n > vc^*(S)$ such that S has the (m, n)-property, then there exists a finite covering $\{S_1, \ldots, S_l\}$ of S by definable subfamilies such that, for every $i \le l$, the family S_i extends to a definable type in $S_k(M)$.

Proof. Let $\varphi(x, y) \in \mathcal{L}$ and $\psi(y) \in \mathcal{L}(M)$ be formulas such that $S = \{\varphi(M, b) : b \in \psi(M)\}$. If *S* does not admit a covering as described in the corollary then, by model-theoretic compactness, there exists some $b \in \psi(U)$ such that $\varphi(U, b)$ does not extend to an *M*-definable type in $S_{|x|}(U)$. On the other hand, by Corollary 3.2, the family *S* has the (ω, n) -property for every n > 0, and consequently the formula $\varphi(x, b)$ does not divide over *M*. So, by Fact 3.6, $\varphi(x, b)$ extends to an *M*-definable type in $S_{|x|}(U)$, contradiction.

Remark 3.8. By [Andújar Guerrero 2021b, Theorem 3.21; Cotter and Starchenko 2012, Corollary 5.6], Fact 3.6 still holds if we substitute M with any (small) set $A \subseteq U$. It follows that, in Corollary 3.7, if S is A-definable for some $A \subseteq M$, then the finite covering $\{S_1, \ldots, S_l\}$ can be chosen so that each S_i extends to an A-definable type in $S_k(M)$.

²Dolich specifically works with "nice" o-minimal theories, a certain class of theories which includes o-minimal expansions of ordered fields.

Remark 3.9. There is a close relation between (p, q)-theorems and so-called fractional Helly theorems (see [Matoušek 2004]), both of which branched from the classical Helly theorem. In its infinite version, this classical theorem states that every family of closed and bounded convex subsets of \mathbb{R}^n that is (n+1)-consistent has nonempty intersection. Aschenbrenner and Fischer [2011, Theorem B] proved a definable version of Helly's theorem (i.e., for definable families of closed and bounded convex sets) in definably complete expansions of real closed fields.

Our Theorem A and the arguments in Section 3.2 of [Aschenbrenner and Fischer 2011] allow an obvious generalization of the o-minimal part of Aschenbrenner and Fischer's definable Helly theorem, by asking that the sets be definably compact and closed in some (any) definable topology, instead of closed and bounded in the Euclidean sense. Perhaps more interestingly, by using Corollary 3.7 to adapt the second proof of Theorem B in [loc. cit.] (the one right below Theorem 3.7), one may show that, in an o-minimal expansion \mathcal{M} of an ordered field, every definable family of convex subsets of M^n that is (n+1)-consistent extends to a definable type in $S_n(M)$.

4. O-minimal types

Throughout this section we assume that our structure \mathcal{M} is o-minimal. Our aim is to investigate the relationship between definable types and definable downward directed families of sets, in order to apply the results in Section 5. Our two main results, Propositions 4.1 and 4.3, are of independent interest.

Proposition 4.1 below can be seen as a strong nonparameter form of distal cell decomposition within o-minimality (see Theorem 21(2) in [Chernikov and Simon 2015]). It implies that every definable family of sets that extends to a definable type admits a refinement given by a definable downward directed family.

Proposition 4.1. Let $p(x) \in S(M)$ be a type and $\varphi(x, y)$ be a formula. There exists another formula $\psi(x, z)$ such that $p|_{\psi}^{1}(x)$ is downward directed and

$$p|_{\psi}^{1} \vdash p|_{\varphi}.$$

In particular, for every finite subtype $q \subseteq p|_{\varphi}$, there exists $c \in M^{|z|}$ such that $\psi(x, c) \in p(x)$ and $\psi(x, c) \vdash q(x)$.

To prove the above proposition we will use the following easy lemma, whose proof we leave to the reader.

Lemma 4.2. Let p(x) be a type and $q_1(x), \ldots, q_k(x)$ be finitely many partial subtypes of p(x). Suppose that, for every $i \le k$, there exists a formula $\psi_i(x, z_i)$ such that $p|_{\psi_i}^1$ is downward directed and $p|_{\psi_i}^1 \vdash q_i$. Then the conjunction

$$\psi(x, z_1, \ldots, z_k) = \bigwedge_{i \le k} \psi_i(x, z_i)'$$

satisfies that $p|_{\psi}^{1}$ is downward directed and

$$p|_{\psi}^1 \vdash \bigcup_{i \leq k} q_i.$$

We now present the proof of the proposition.

Proof of Proposition 4.1. We proceed by induction on |x|. We may assume throughout that p(x) is not realized, since otherwise it suffices to have $\psi(x, z)$ be the formula x = z, where |x| = |z|.

<u>Case |x| = 1</u>: By o-minimality it suffices to have $\psi(x, z_1, z_2)$, with $|z_1| = |z_2| = 1$, be one of the three formulas

$$(z_1 < x) \land (x < z_2), \quad z_1 < x, \quad x < z_1.$$

<u>Case |x| > 1</u>: Throughout let x = (u, t), where |t| = 1. Recall that $\pi(p) \in S_{|u|}(M)$ denotes the projection of the type p to the first |u| coordinates; i.e., $\pi(p)(u)$ is the family of all formulas $\lambda(u)$ such that $\lambda(u) \wedge (t = t)$ is in p(x).

Suppose that there exists a definable partial function $f: M^{|x|-1} \to M$ whose graph is contained in p. By extending f if necessary to a constant function outside its domain we may assume that the domain of f is in fact $M^{|x|-1}$. We may apply the induction hypothesis to the type $\pi(p)$ and formula

$$\varphi_f(u, y) = \exists t ((t = f(u)) \land \varphi(u, t, y)),$$

and obtain a formula $\psi_f(u, z_f)$ as described in the proposition. This allows us to construct our desired formula ψ as follows:

$$\psi(x, z_f) = \psi(u, t, z_f) = "(t = f(u)) \land \psi_f(u, z_f)".$$

We show that $\psi(x, z_f)$ has the desired properties. Observe that, since the graph of f is contained in p, for every $b \in M^{|y|}$ and $i \in \{0, 1\}$, the formula $\varphi^i(x, b)$ belongs in p if and only if $\varphi^i_f(u, b)$ belongs in $\pi(p)$. An analogous statement holds for $\psi(x, z_f)$ and $\psi_f(u, z_f)$. In particular, we may define

$$C = \{ c \in M^{|z_f|} : \psi_f(u, c) \in \pi(p) \} = \{ c \in M^{|z_f|} : \psi(x, c) \in p \}.$$

Since, by the induction hypothesis, the family of formulas $\{\psi_f(u, c) : c \in C\}$ is downward directed, the same clearly holds for $p|_{\psi}^1 = \{\psi(x, c) : c \in C\}$. Moreover, for any formula of the form $\varphi^i(x, b)$ in p(x), where $b \in M^{|y|}$ and $i \in \{0, 1\}$, there exists $c \in C$ such that $\psi_f(u, c) \vdash \varphi^i_f(u, b)$, and so $\psi(x, c) \in p$ and $\psi(x, c) \vdash \varphi^i(x, b)$. Hence $p|_{\psi}^1 \vdash p|_{\varphi}$.

Hence onwards we assume that there does not exist a definable partial function $f: M^{|x|-1} \to M$ whose graph is contained in p.

In the next paragraphs we reduce the remainder of the proof to the case where, for every $b \in M^{|y|}$, if the formula $\varphi(x, b)$ is in p, then it defines a set of the form $(f_b, +\infty)$ for some partial function $M^{|x|-1} \to M \cup \{-\infty\}$.

By o-minimal uniform cell decomposition [van den Dries 1998, Chapter 3, Proposition 3.5], there exist finitely many formulas $\sigma_1(x, y), \ldots, \sigma_k(x, y)$ such that, for every $b \in M^{|y|}$, the family { $\sigma_1(M, b), \ldots, \sigma_k(M, b)$ } is an o-minimal cell decomposition of $M^{|x|}$ compatible with $\varphi(M, b)$. Observe that

$$\bigcup_{i\leq k}p|_{\sigma_i}^1\vdash p|_{\varphi}.$$

By Lemma 4.2, it suffices to pass to an arbitrary $i \le k$ and prove the proposition for $p|_{\sigma_i}^1$ in place of $p|_{\varphi}$. Hence onwards let us assume that, for every $b \in M^{|y|}$, the formula $\varphi(x, b)$ defines a cell and, moreover, if $\varphi(x, b) \in p$, then, by the assumption on p, this cell is of the form (f_b, g_b) for f_b and g_b partial functions $M^{|x|-1} \to M \cup \{-\infty, +\infty\}$ with the same domain and with $f_b < g_b$. Additionally, to prove the proposition it suffices to find $\psi(x, z)$ such that $p|_{\psi}^1$ is downward directed and $p|_{\psi}^1(x) \vdash p|_{\varphi}^1(x)$.

Recall the notation x = (u, t), with |t| = 1. Let $B = \{b \in M^{|y|} : \varphi(x, b) \in p\}$. Let $\varphi_0(x, y) (= \varphi_0(u, t, y))$ denote the formula $\exists s (s \leq t) \land \varphi(u, s, y)$, and similarly let $\varphi_1(x, y)$ be the formula $\exists s (s \geq t) \land \varphi(u, s, y)$. That is, for every $b \in B$, the formulas $\varphi_0(x, b)$ and $\varphi_1(x, b)$ define the sets $(f_b, +\infty)$ and $(-\infty, g_b)$ respectively. In particular, when $b \in B$, the formula $\varphi(x, b)$ is equivalent to the conjunction $\varphi_0(x, b) \land \varphi_1(x, b)$. So $p|_{\varphi_0}^1 \cup p|_{\varphi_1}^1 \vdash p|_{\varphi}^1$. By Lemma 4.2, to prove the proposition it suffices to find formulas $\psi_0(x, z_0)$ and $\psi_1(x, z_1)$ such that, for every $j \in \{0, 1\}$, the restriction $p|_{\psi_j}^1$ is downward directed and $p|_{\psi_j}^1 \vdash p|_{\varphi_j}^1$. We prove this for j = 0, since the remaining case is analogous. For simplicity of notation we also assume that φ is equivalent to φ_0 .

Consider the formula

$$\theta(u, y, y') = ``\exists s \, \varphi(u, s, y) \land \exists t \, \varphi(u, t, y') \land \forall t \, (\varphi(u, t, y') \to \varphi(u, t, y))``$$

For every $b, b' \in M^{|y|}$ note that it holds that

$$\theta(u, b, b') \wedge \varphi(x, b') \vdash \varphi(x, b). \tag{1}$$

In particular, if *b* and *b'* are in *B*, then $\theta(u, b, b')$ defines the set of all *u* such that $f_b(u) \leq f_{b'}(u)$.

Recall the notation $\pi(p)$ for the projection of p to the first |u| = |x| - 1 coordinates. By the induction hypothesis on the formula $\theta(u, y, y')$ and the type $\pi(p)$, there exists a formula $\xi(u, z_{\xi})$ such that $\pi(p)|_{\xi}^{1}$ is downward directed and $\pi(p)|_{\xi}^{1} \vdash \pi(p)|_{\theta}$.

Finally, let $z = (z_{\xi}, y)$ and

$$\psi(x, z) = \psi(u, t, z_{\xi}, y) = ``\xi(u, z_{\xi}) \land \varphi(x, y)`'.$$

Clearly by construction $p|_{\psi}^1 \vdash p|_{\varphi}^1$. We show that $p|_{\psi}^1$ is downward directed.

Let $D = \{d \in M^{|z_{\xi}|} : \xi(u, d) \in \pi(p)\}$. Note that $\xi(u, d) \land \varphi(x, b)$ belongs in p if and only if $b \in B$ and $d \in D$. Let us fix $b, b' \in B$ and $d, d' \in D$. Recall that $\varphi(M, b) = (f_b, +\infty)$ and $\varphi(M, b') = (f_{b'}, +\infty)$. Consider the formula $\zeta(u, b, b') =$ " $\exists s \varphi(u, s, b) \land \exists t \varphi(u, t, b')$ ", which defines the intersection of the domains of f_b and $f_{b'}$. Clearly $\zeta(u, b, b') \in \pi(p)$. Observe that the sets $\theta(M, b, b')$ and $\theta(M, b', b)$ cover $\zeta(M, b, b')$, and so at least one of them belongs in $\pi(p)$. Without loss of generality we assume that $\theta(u, b, b') \in \pi(p)$.

Let $d'' \in D$ be such that $\xi(u, d'') \vdash \theta(u, b, b')$. By (1) we have that

 $\xi(u, d'') \land \varphi(x, b') \vdash \varphi(x, b).$

By downward directedness let $d''' \in D$ be such that

 $\xi(u, d''') \vdash \xi(u, d) \land \xi(u, d') \land \xi(u, d'').$

We conclude that

 $\xi(u, d''') \land \varphi(x, b') \vdash \xi(u, d) \land \varphi(x, b) \land \xi(u, d') \land \varphi(x, b'),$

or equivalently

$$\psi(x, d''', b') \vdash \psi(x, d, b) \land \psi(x, d', b').$$

So $p|_{\psi}^{1}$ is downward directed.

It seems likely that Proposition 4.1 is also true in weakly o-minimal structures. As far as the author knows, it is open among distal dp-minimal structures.

The following proposition shows that every definable downward directed family of nonempty sets extends to a definable type $p(x) \in S(M)$, and furthermore that p(x) can be chosen so that, for some formula $\psi(x, z)$, the restriction $p|_{\psi}^{1}(x)$ is a basis (in the sense of filter basis) of cells for p(x). We present a shorter proof than the one in [Andújar Guerrero 2021b, Lemma 2.7], applying ideas communicated to the author by Will Johnson.

Proposition 4.3. Let $\varphi(x, y)$ be a formula and $B \subseteq M^{|y|}$ be such that the family $\{\varphi(x, b) : b \in B\}$ is consistent and downward directed. Then there exists a type $p(x) \in S(M)$ with $\{\varphi(x, b) : b \in B\} \subseteq p(x)$, and a formula $\psi(x, z)$ such that $p|_{\psi}^{1}$ defines a family of cells, is downward directed, and $p|_{\psi}^{1} \vdash p$. Furthermore, if B is definable, then p(x) can be chosen definable too.

In particular, for every definable downward directed family of nonempty sets S, there exists a definable downward directed family of cells \mathcal{F} which refines S and furthermore \mathcal{F} generates a definable type in S(M).

Proof. We devote most of the proof to showing the existence of p(x) and $\psi(x, z)$ as described in the proposition except for the condition that $p|_{\psi}^{1}$ defines a family of cells. In the next two paragraphs we describe how, once we have these, by passing

if necessary to a formula in a cell decomposition of $\psi(x, z)$ we may assume that $p|_{\psi}^{1}$ defines a family of cells, completing the proof.

Applying uniform cell decomposition [van den Dries 1998, Chapter 3, Proposition 3.5] to the formula $\psi(x, z)$, let $\sigma_i(x, z)$, for $i \le k$, denote formulas such that, for every $c \in M^{|z|}$, the sets $\sigma_i(M, c)$, for $i \le k$, are a cell partition of $\psi(M, c)$. We claim that there exists some $i \le k$ such that the family $p|_{\sigma_i}^1$ is downward directed and $p|_{\sigma_i}^1 \vdash p|_{\psi}^1$ (hence $p|_{\sigma_i}^1 \vdash p)$. To see this let $p|_{\psi}^1 = \{\psi(x, c) : c \in C\}$ and, for every $i \le k$, let $p|_{\sigma_i}^1 = \{\sigma_i(x, c) : c \in C(i)\}$. We show that there exists $i \le k$ such that, for every $c \in C$, there exists some $c' \in C(i)$ with $\sigma_i(x, c') \vdash \psi(x, c)$ (i.e., $p|_{\sigma_i}^1$ refines $p|_{\psi}^1$); hence $p|_{\sigma_i}^1 \vdash p|_{\psi}^1$ and, using the facts that $p|_{\psi}^1 \vdash p$ and $p|_{\psi}^1$ is downward directed, it is also easy to derive that $p|_{\sigma_i}^1$ is downward directed.

Towards a contradiction suppose that, for every $i \leq k$, there exists some $c_i \in C$ such that $\sigma_i(x, c) \nvDash \psi(x, c_i)$ for every $c \in C(i)$. By downward directedness of $p|_{\psi}^1$, let $c_{k+1} \in C$ be such that $\psi(x, c_{k+1}) \vdash \bigwedge_{i \leq k} \psi(x, c_i)$. It follows that $\sigma_i(x, c) \nvDash$ $\psi(x, c_{k+1})$ for every $i \leq k$ and $c \in C(i)$. However this contradicts the facts that $\psi(x, c_{k+1}) \in p(x)$ and $\vdash \psi(x, c_{k+1}) \leftrightarrow \bigvee_{i \leq k} \sigma_i(x, c_{k+1})$, which imply that there exists some $i \leq k$ with $\sigma_i(x, c_{k+1}) \in p(x)$ (i.e., $c_{k+1} \in C(i)$) and $\sigma_i(x, c_{k+1}) \vdash$ $\psi(x, c_{k+1})$.

We now begin the proof of the existence of a type $p(x) \in S(M)$ extending $\{\varphi(x, b) : b \in B\}$ and a formula $\psi(x, z)$ satisfying that $p|_{\psi}^1$ is downward directed and $p|_{\psi}^1 \vdash p$ (i.e., $p|_{\psi}^1$ is a basis for p). We prove the case where B is definable. In the general case the same proof applies by considering throughout, instead of definable families of sets, subfamilies of fibers of definable sets in general. To make the presentation more succinct, we work explicitly with set notation rather than formulas.

We introduce some useful terminology. For a definable family of nonempty sets \mathcal{F} , let $d(\mathcal{F})$ denote the smallest $n \ge 0$ such that, for every set $F \in \mathcal{F}$, there exists $G \in \mathcal{F}$ with $G \subseteq F$ and dim(G) = n. Let $c(\mathcal{F})$ denote the smallest $m \ge 1$ such that, for every set $F \in \mathcal{F}$, there exists $G \in \mathcal{F}$ with $G \subseteq F$ such that G has exactly m definably connected components.

Let $S = \{\varphi(M, b) : b \in B\}$. Recall that a family of sets \mathcal{F} is a refinement of S if, for every $S \in S$, there exists $F \in \mathcal{F}$ with $F \subseteq S$. Let \mathcal{DR} denote the collection of all definable downward directed refinements of S which do not contain the empty set. Throughout we fix $n = \min\{d(\mathcal{F}) : \mathcal{F} \in \mathcal{DR}\}$ and $m = \min\{c(\mathcal{F}) : \mathcal{F} \in \mathcal{DR}, d(\mathcal{F}) = n\}$. We also fix $\mathcal{F} \in \mathcal{DR}$ with $d(\mathcal{F}) = n$ and $c(\mathcal{F}) = m$. We show that \mathcal{F} generates a (clearly definable) type in $S_{|x|}(M)$.

Towards a contradiction we assume that \mathcal{F} does not generate a type in $S_{|x|}(M)$, meaning that there exists a definable set $X \subseteq M^{|x|}$ satisfying that, for every $F \in \mathcal{F}$, $F \cap X \neq \emptyset$ and $F \setminus X \neq \emptyset$. Let us fix some $F_0 \in \mathcal{F}$ with dim $F_0 = n$. Consider the boundary of $F_0 \cap X$ in F_0 , i.e., the set

$$Z = F_0 \cap (\partial(F_0 \cap X) \cup \partial(F_0 \setminus X)).$$

Since dim $F_0 = n$, by o-minimality we have that dim Z < n. It follows that $\mathcal{F} \cap Z$ is a downward directed refinement of S composed of sets of dimension lower than n. By the definition of n, there must exist a set $F_1 \in \mathcal{F}$ with $F_1 \cap Z = \emptyset$. Now let $\mathcal{G} = \{F \cap X : F \in \mathcal{F}, F \subseteq F_0 \cap F_1\}$. By the downward directedness of \mathcal{F} and the definition of X, the definable family \mathcal{G} is a downward directed refinement of S that does not contain the empty set (i.e., $\mathcal{G} \in D\mathcal{R}$). By the definition of n it follows that $d(\mathcal{G}) = n$. We show that $c(\mathcal{G}) < m$, contradicting the definition of m.

We show that, for every $F \in \mathcal{F}$ with $F \subseteq F_0 \cap F_1$, the intersection $F \cap X \in \mathcal{G}$ has strictly fewer definably connected components than F. In particular, this implies that, for every set $F \in \mathcal{F}$ with $F \subseteq F_0 \cap F_1$, if $G \in \mathcal{F}$ is a subset of F with exactly m definably connected components, then $G \cap X \in \mathcal{G}$ has less than m definably connected components, and so $c(\mathcal{G}) < m$ as desired.

Let *Y* denote the interior of $F_0 \cap X$ in F_0 , i.e., $Y = F_0 \setminus cl(F_0 \setminus X)$. Let $C \subseteq F_0$ be a definably connected set. If $C \cap Z = \emptyset$, then by the definition of *Z* clearly *C* must be a subset of either *Y* or $F_0 \setminus (Y \cup Z)$. Since $Y \subseteq F_0 \cap X \subseteq Y \cup Z$, we know *C* must be a subset of either $F_0 \cap X$ or $F_0 \setminus X$. Now let us fix a set $F \in \mathcal{F}$ with $F \subseteq F_0 \cap F_1$. Since $F_1 \cap Z = \emptyset$, we have that $F \cap Z = \emptyset$, and so every definably connected component *C* of *F* is a subset of either $F_0 \cap X$ or $F_0 \setminus X$. Finally recall that, by the definition of *X*, the sets $F \cap X$ and $F \setminus X$ are both nonempty. Consequently we conclude that the set $F \cap X$ (as well as $F \setminus X$) has a positive number of definably connected components that is fewer than the number of definably connected components of *F*.

In Proposition 4.3, whenever *B* is definable, one may wonder if p(x) can always be chosen definable over the same parameters as *B*. This was proved to be false in general by Johnson in [Andújar Guerrero 2021b, Appendix B]. Nevertheless, by [loc. cit., Proposition 2.17] it does hold that every definable downward directed family { $\varphi(x, b) : b \in B$ } extends to a type in $S_{|x|}(M)$ definable over the same parameters as *B*. This can also be proved using Corollary 3.7 and Remark 3.8. For a similar result see [Hrushovski and Loeser 2016, Lemma 4.2.18].

Remark 4.4. Observe that Propositions 4.1 and 4.3 together yield a strong density result for types p(x) satisfying that there is a formula $\varphi(x, y)$ such that $p|_{\varphi}$ is downward directed and $p|_{\varphi} \vdash p$, namely types which have a basis (in the sense of filter basis) given by their restriction to a single formula. This is discussed in [Andújar Guerrero 2021b, Remarks 2.13 and 2.22]. In any o-minimal structure every 1-type is of this kind (it is either realized or has a basis of open intervals). On the other hand, it was shown in [Andújar Guerrero et al. 2021, Corollary 32]

that, in an o-minimal expansion of an ordered group, every definable type of this kind contains at least one set of dimension at most 2 (and of dimension at most 1 in o-minimal expansions of ordered fields). Using the Marker–Steinhorn theorem [1994, Theorem 2.1] one derives that, in any o-minimal expansion of the group of reals, there are *n*-types that do not have a basis given by their restriction to a single formula for every n > 2 (n > 1 in o-minimal expansions of the field of reals).

5. O-minimal definable compactness

5.1. *Topological preliminaries.* We introduce definable (explicitly in the sense of [Flum and Ziegler 1980]) topological spaces and various related definitions.

Definition 5.1. A *definable topological space* (X, τ) , with $X \subseteq M^n$, is a topological space such that there exists a definable family of subsets of X which is a basis for τ .

Any definable set in an o-minimal structure with its induced Euclidean topology is a definable topological space. For other examples within o-minimality, see the definable manifold spaces studied in [Pillay 1988] and [van den Dries 1998, Chapter 10], the definable Euclidean quotient spaces of the latter work and [Johnson 2018], the definable normed spaces of [Thomas 2012], and the definable metric spaces of [Walsberg 2015]. See moreover the author's doctoral dissertation [Andújar Guerrero 2021a] for an exhaustive exploration of o-minimal definable topological spaces. For a foundational treatment of definable tame topology generalizing ominimality, see [Pillay 1987]. For an exploration of dp-minimal tame topology, see the more recent work [Simon and Walsberg 2019], and related independent work [Dolich and Goodrick 2022].

Onwards we contextualize topological notions related to a given topological space (X, τ) by adding the prefix τ , e.g., τ -open, τ -closure etc. We recall some standard definitions.

Definition 5.2. Let (X, τ) be a definable topological space. A *definable curve in* X is a definable map $\gamma : (a, b) \to X$ for some $-\infty \le a < b \le +\infty$. We say that it τ -converges to $x \in X$ (i.e., x is a τ -limit of γ) as $t \to a$ if, for every τ -neighborhood A of x, there exists $t_A \in (a, b)$ such that $\gamma(s) \in A$ whenever $s \in (a, t_A)$. The notion of τ -convergence as $t \to b$ is defined analogously. We denote by $\lim_{t\to a}^{\tau} \gamma(t)$ (respectively $\lim_{t\to b}^{\tau} \gamma(t)$) the set of τ -limit points of x as $t \to a$ (respectively $t \to b$).

We say that γ is τ -completable if it τ -converges as $t \to a$ and as $t \to b$.

Given a definable topological space (X, τ) and a set $Y \subseteq X$ we denote the τ -closure of Y by $cl_{\tau}(Y)$. It is easy to check that a τ -limit of a definable curve $\gamma : (a, b) \to Y \subseteq X$ is always contained in $cl_{\tau}(Y)$. Furthermore, if τ is Hausdorff, then the sets $\lim_{t \to a}^{\tau} \gamma(t)$ and $\lim_{t \to b}^{\tau} \gamma(t)$ are always either empty or a singleton, and in the latter case we abuse terminology by identifying them with their single point.

We will use these facts in Section 5.2 without explanation. To erase ambiguity, at times we also use side convergence notation $t \to a^+$ and $t \to b^-$ (e.g., $\lim_{t \to a^+}^{\tau} \gamma(t)$), with the standard meaning.

The following definition is borrowed from [Hrushovski and Loeser 2016].

Definition 5.3. Let (X, τ) be a definable topological space and p be a (possibly partial) type with $X \in p$. We say that $x \in X$ is a τ -*limit*³ of p if x is contained in the τ -closure of every subset of X in p. If $p(x) \in S_X(M)$, then this is equivalent to saying that x is contained in every τ -closed set in p.

We now present various definitions extracted from the literature (for references see Section 1) which seek to capture the notion of definable compactness. We mostly maintain consistency with [Andújar Guerrero and Johnson 2024] in the names. (In particular we avoid using the adjective "definable" in our terminology to enable an easier read.) A more general approach to definable compactness, including more definitions than the ones in this paper, can be found in [Fornasiero 2015].

Definition 5.4. Let (X, τ) be a definable topological space. Then (X, τ) is

- (1) *curve-compact* if every definable curve in X is τ -completable,
- (2) *filter-compact* if every downward directed definable family of nonempty τ -closed subsets of X has nonempty intersection,
- (3) *type-compact* if every definable type $p(x) \in S_X(M)$ has a τ -limit in X,
- (4) *transversal-compact* if every consistent definable family of τ -closed subsets of X has a finite transversal.

The equivalence between curve-compactness and filter-compactness was proved for definable topological spaces in o-minimal expansions of ordered fields in [Andújar Guerrero et al. 2021, Corollary 44]. In this paper we present a deeper characterization in the general o-minimal setting.

5.2. Characterizing definable compactness. In this section we prove our results on definable compactness for definable topological spaces in o-minimal structures. Throughout we assume that our underlying structure \mathcal{M} is o-minimal.

We devote most of the section to proving the characterization of definable compactness given by Theorem A, which we divide into three propositions. Proposition 5.5 provides the equivalence $(2) \Leftrightarrow (3)$ in the theorem. In Proposition 5.7 we prove, using results from previous sections, the equivalence between (1), (3), (4), (5) and (6). Finally, in Proposition 5.12 we prove the implication $(1) \Rightarrow (7)$, and the reverse implication when τ is Hausdorff or \mathcal{M} has definable choice. We follow

³Fornasiero [2015], as well as Thomas, Walsberg and the author [Andújar Guerrero et al. 2021], use the word "specialization" (borrowed from real algebraic geometry) to refer to limits of types. Here we use instead the terminology from [Hrushovski and Loeser 2016, Chapter 4].

it with an example (Example 5.16) showing that the implication $(1) \Rightarrow (7)$ is strict in general. Throughout we also discuss other notions of definable compactness, and end the section with two additional results: definable compactness is equivalent to classical compactness in o-minimal expansions of (\mathbb{R} , <) (Corollary 5.19), and definable compactness is definable in families (Proposition 5.22).

The equivalence (1) \Leftrightarrow (2) in Proposition 5.5 below corresponds to the equivalence (2) \Leftrightarrow (3) in Theorem A. Note that the proof of this equivalence does not use o-minimality. Hence this characterization of type-compactness holds in any model-theoretic structure. Furthermore, the equivalence of type-compactness with classical compactness always holds whenever the underlying structure \mathcal{M} satisfies that every type in $S(\mathcal{M})$ is definable, as we point out in Remark 5.6 below.

Proposition 5.5. Let (X, τ) be a definable topological space. The following are equivalent:

- (1) (X, τ) is type-compact.
- (2) Every definable family of τ -closed sets that extends to a definable type in $S_X(M)$ has nonempty intersection.

If \mathcal{M} expands $(\mathbb{R}, <)$, then (1) and (2) are also equivalent to

(3) (X, τ) is compact.

Proof. To prove $(1) \Rightarrow (2)$, suppose that (X, τ) is type-compact and let C be a definable family of τ -closed sets that extends to a definable type $p \in S_X(M)$. Let $x \in X$ be a τ -limit of p. Then clearly $x \in \bigcap C$.

The key element to the rest of the proof is the fact that every closed set in a topological space is an intersection of basic closed sets.

To prove (2) \Rightarrow (1), let $p \in S_X(M)$ be a definable type. Let \mathcal{B} denote a definable basis (of opens) for the topology τ . Now let \mathcal{C} denote the definable family of basic τ -closed sets in p, i.e., the family of sets C in p of the form $X \setminus B$ for some $B \in \mathcal{B}$. If (2) holds, then there exists some $x \in X$ with $x \in \bigcap \mathcal{C}$. In this case it follows that x is a τ -limit of p.

Finally, suppose that \mathcal{M} expands $(\mathbb{R}, <)$. Clearly, if (X, τ) is compact, then it is type-compact. Conversely, suppose that (X, τ) is type-compact and let \mathcal{C} be a consistent family of τ -closed sets. The intersection $\bigcap \mathcal{C}$ can be rewritten as an intersection of basic closed sets. In particular, we may assume that \mathcal{C} contains only definable sets. Now, by the Marker–Steinhorn theorem [1994, Theorem 2.1], every type over $M = \mathbb{R}$ is definable. Consequently \mathcal{C} extends to a definable type $p \in S_X(M)$. Let x be a τ -limit of p. Then $x \in \bigcap \mathcal{C}$. So (X, τ) is compact. \Box

Remark 5.6. Note that the equivalence between type-compactness and classical topological compactness shown in Proposition 5.5 holds in every structure satisfying

that all types are definable. For example it remains true in the field of p-adic numbers $(\mathbb{Q}_p, +, \cdot)$, as observed in [Andújar Guerrero and Johnson 2024, Theorem 8.15].

More specifically, if $\varphi(x, y)$ defines a basis \mathcal{B} for the topology τ , i.e., $\mathcal{B} = \{\varphi(M, b) : b \in M^{|y|}\}$, then to have the equivalence between type-compactness and classical compactness it suffices to have that every maximal consistent subfamily of $\{X \setminus B : B \in \mathcal{B}\}$ is definable, which occurs in particular whenever every φ -type (i.e., restrictions of types in $S_{|x|}(M)$ to $\varphi(x, y)$) is definable. Observe that the latter always holds whenever $\varphi(x, y)$ is stable although, as already noted in the proof of [Pillay 1987, Proposition 1.2], every infinite T_1 topological space that has a basis defined by a stable formula must be discrete, and consequently not compact.

Proposition 5.7 below corresponds to the equivalence between (1), (3), (4), (5) and (6) in Theorem A. Its proof relies on Proposition 4.1 and Corollaries 3.4 and 3.7.

Proposition 5.7. Let (X, τ) be a definable topological space. The following are equivalent:

- (1) (X, τ) is filter-compact.
- (2) Every definable family of τ -closed sets that extends to a definable type in $S_X(M)$ has nonempty intersection.
- (3) Every definable family C of τ -closed sets with the (m, n)-property, where $m \ge n > vc^*(S)$, has a finite transversal.
- (4) Every definable family C of τ -closed sets with the (m, n)-property, where $m \ge n > \dim \bigcup C$, has a finite transversal.
- (5) (X, τ) is transversal-compact.

Proof. Note that, if a downward directed family of sets has a finite transversal, then, by Lemma 2.1, it has nonempty intersection. Hence (3), (4) and (5) each imply (1). We prove $(1) \Rightarrow (2)$ and $(2) \Rightarrow (3)$. Observe that implication $(3) \Rightarrow (4)$ follows from Corollary 3.4, and implication $(4) \Rightarrow (5)$ is trivial, completing the proof.

<u>Proof of (1) \Rightarrow (2)</u>. Suppose that (X, τ) is filter-compact and let $C = \{\varphi(M, b) : b \in B\}$ be a definable family of τ -closed sets that extends to a definable type $p(x) \in S_X(M)$. Let $\psi(x, z)$ be as given by Proposition 4.1 for $\varphi(x, y)$ and p(x). Let $\mathcal{F} = \{\psi(X, c) : \psi(x, c) \in p(x), c \in M^{|z|}\}$. By Proposition 4.1, \mathcal{F} is a definable downward directed family of subsets of X which refines C. Let $\mathcal{D} = \{cl_{\tau}(F) : F \in \mathcal{F}\}$. Clearly \mathcal{D} is a definable downward directed family of subsets of X which refines C, Let $\mathcal{D} = \{cl_{\tau}(F) : F \in \mathcal{F}\}$. Clearly \mathcal{D} is a definable downward directed family of τ -closed sets, so by filter-compactness there exists $a \in \bigcap \mathcal{D}$. Moreover observe that, since the sets in C are closed, \mathcal{D} is still a refinement of C, implying that $\bigcap \mathcal{D} \subseteq \bigcap C$, and so $a \in \bigcap C$.

<u>Proof of (2) \Rightarrow (3).</u> Let *C* be a definable family of τ -closed subsets of *X* with the (m, n)-property, where $m \ge n > vc^*(C)$. By Corollary 3.7 there exists a finite covering $\{C_1, \ldots, C_k\}$ of *C* by definable subfamilies, each of which extends to a

definable type in $S_X(M)$. If property (2) holds, then, for each $i \le k$, there exists some $a_i \in \bigcap C_i$ in X. The family $\{a_1, \ldots, a_k\}$ is clearly a transversal of C. \Box

Remark 5.8. We remark that, although omitted from the proof above, the implication $(2) \Rightarrow (1)$ in Proposition 5.7 (i.e., by Proposition 5.5 the implication *type-compactness* \Rightarrow *filter-compactness*) can be shown to follow easily from Proposition 4.3. In fact we claim that this implication, as well as $(2) \Rightarrow (5)$, hold in a more general dp-minimal setting by [Simon and Starchenko 2014, Theorem 5] (see the discussion above Fact 3.6). Additionally, the equivalence $(3) \Leftrightarrow (5)$ holds in all in NIP structures by recent work [Kaplan 2024, Corollary 4.9].

Remark 5.9. It was shown in [Peterzil and Steinhorn 1999, Theorem 2.1] that a definable set with the o-minimal Euclidean topology is curve-compact if and only if it is closed and bounded. In [Peterzil and Pillay 2007, Theorem 2.1] the authors extracted from [Dolich 2004] the following. Suppose that our o-minimal structure \mathcal{M} has definable choice (e.g., expands an ordered group). Let $\mathcal{U} = (U, ...)$ be a monster model and $\varphi(x, b)$ be a formula in $\mathcal{L}(U)$ such that $\varphi(U, b)$ is closed and bounded (in the Euclidean topology in $U^{|x|}$). If the family { $\varphi(U, b') : \operatorname{tp}(b'/M) = \operatorname{tp}(b/M)$ } is consistent, then $\varphi(U, b)$ has a point in $M^{|x|}$. Using a straightforward model-theoretic compactness argument they derive from this that every closed and bounded Euclidean space is transversal-compact [Peterzil and Pillay 2007, Corollary 2.2(i)].

Let (X, τ) be a definable topological space (in \mathcal{M}), whose definition in \mathcal{U} we denote by $(X(\mathcal{U}), \tau(\mathcal{U}))$. The property that every formula $\varphi(x, b) \in \mathcal{L}(U)$, satisfying that $\varphi(U, b)$ is $\tau(\mathcal{U})$ -closed and the family { $\varphi(U, b') : \operatorname{tp}(b'/M) = \operatorname{tp}(b/M)$ } is consistent, satisfies that $\varphi(M, b) \neq \emptyset$, is labeled Dolich's property in [Fornasiero 2015]. As mentioned in the previous paragraph, this property implies transversal-compactness (without any assumption on \mathcal{M}), and furthermore one may show, using [Kaplan 2024, Theorem 1.5], that the converse implication (transversal-compactness \Rightarrow Dolich's property) holds in all NIP structures.

Theorem A completes the characterization of closed and bounded definable sets with the Euclidean topology. Furthermore, it generalizes the aforementioned result of [Peterzil and Pillay 2007] in three ways. First, we drop the assumption of having definable choice in \mathcal{M} . Second, we weaken the consistency assumption to having an appropriate (n, m)-property (in their work they actually observe that it suffices to have k-consistency for some k in terms of |x| and |b|). Third, we establish, by means of the equivalence with transversal-compactness mentioned in the paragraph above, the relationship between Dolich's property and the other compactness notions in the full generality of any o-minimal definable topological space.

We now prove the connection within o-minimality between filter-compactness and curve-compactness stated in Theorem A, that is, that filter-compactness implies curve-compactness, and that both notions are equivalent when the topology is Hausdorff or when the underlying o-minimal structure has definable choice. This is Proposition 5.12. We follow the proposition with an example of a non-Hausdorff topological space definable in the dense linear order without endpoints (M, <) that is curve-compact but not filter-compact.

The next lemma allows us to apply definable choice in certain instances even when the underlying structure \mathcal{M} may not have the property.

Lemma 5.10 (definable choice in compact Hausdorff spaces). Let *C* be a definable nonempty τ -closed set in a curve-compact Hausdorff definable topological space (X, τ) . Let $A \subseteq M$ be such that τ and *C* are *A*-definable. Then there exists a point $x \in C \cap dcl(A)$, where dcl(A) denotes the set of finite tuples of elements in the definable closure of *A*.

Consequently, for every A-definable family $\{\varphi(M, b) : b \in B\}$ of nonempty subsets of X there exists an A-definable choice function $h : B \to X$ such that $h(b) \in cl_{\tau}(\varphi(M, b))$ for every $b \in B$.

Proof. We prove the first paragraph of the lemma. The uniform result is derived in the usual way by the use of first-order logic compactness.

For this proof we adopt the convention of the one-point Euclidean space $M^0 = \{0\}$. In particular, any projection $M^k \to M^0$ is simply the constant function **0**, and any relation $E \subseteq M^0 \times M^k$ is definable if and only if its projection to M^k is.

Let *C*, (X, τ) and *A* be as in the lemma, with $X \subseteq M^m$. Let $n \leq m$ be such that there exists an *A*-definable function $f : D \subseteq M^n \to C$ for *D* a nonempty set. If *n* can be chosen to be zero, then the lemma follows. We prove that this is the case by backwards induction on *n*.

Note that *n* can always be chosen equal to *m*, by letting *f* be the identity on *C*. Consider a positive $n \le m$. For every $x \in M^{n-1}$, let D_x denote the fiber $\{t \in M : \langle x, t \rangle \in D\}$. For each $x \in \pi(D)$, let $s_x = \sup D_x$, and consider the *A*-definable set $F = \{x \in \pi(D) : s_x \in D_x\}$.

If $F \neq \emptyset$, then let g be the map $x \mapsto f(s_x) : F \to C$. If $F = \emptyset$, then let g be the map $x \mapsto \lim_{t \to s_x^-} f(x, t) : \pi(D) \to C$ which, by curve-compactness and Hausdorffness, is well-defined. In both cases g is an A-definable nonempty partial function $M^{n-1} \to C$.

Remark 5.11. Let $A \subseteq M$ and C be an A-definable family of nonempty τ -closed sets in a curve-compact Hausdorff A-definable topological space (X, τ) . Lemma 5.10 implies that, if C has a finite transversal, then it also has one of the same size in dcl(A). To prove this it suffices to note that, for every $k \ge 1$, the set of k-tuples of points corresponding to a transversal of C is A-definable and closed in the product topology, which can easily be shown to be A-definable and curve-compact. It follows that, whenever \mathcal{M} has definable choice or τ is Hausdorff, the finite transversals in Theorem A (statements (4), (5) and (6)) can always be assumed to be definable over the same parameters as the family of closed sets C and topology τ .

Proposition 5.12. Let (X, τ) be a definable topological space. If (X, τ) is filtercompact, then it is curve-compact.

Suppose that either τ is Hausdorff or \mathcal{M} has definable choice. Then (X, τ) is filter-compact if and only if it is curve-compact.

We prove the left-to-right direction through a short lemma.

Lemma 5.13. Let (X, τ) be a filter-compact definable topological space. Then (X, τ) is curve-compact.

Proof. Let $\gamma : (a, b) \to X$ be a definable curve in *X*. Consider the definable family of τ -closed nested sets $C_{\gamma} = \{cl_{\tau} \gamma[(a, t)] : a < t < b\}$. By filter-compactness, there exists $x \in \bigcap C_{\gamma}$. By o-minimality, observe that γ satisfies that it τ -converges to x as $t \to a$. Similarly one shows that γ also τ -converges as $t \to b$.

We now prove a simpler case of the left-to-right implication in Proposition 5.12. We do so by implicitly introducing a weakening of filter-compactness corresponding to the property that every definable family of nonempty closed sets that is nested has nonempty intersection (say chain-compactness). We show that, when \mathcal{M} has definable choice or the underlying topology is Hausdorff, curve-compactness implies chain-compactness (the reverse implication always holds within o-minimality by the proof of Lemma 5.13). On the other hand, Example 5.16 describes a (non-Hausdorff) definable topological space in (M, <) that is curve-compact but not chain-compact. It is unclear whether chain-compactness is equivalent to definable compactness (Definition 1.1) in the general setting of o-minimal definable topological spaces.

Lemma 5.14. Let (X, τ) be a definable topological space. Suppose that either τ is Hausdorff or \mathcal{M} has definable choice. Let \mathcal{C} be a nested definable family of nonempty τ -closed subsets of X. If (X, τ) is curve-compact, then $\bigcap \mathcal{C} \neq \emptyset$.

Proof. Let (X, τ) and $C = \{\varphi(M, b) : b \in B\}$, with $B \subseteq M^n$, be as in the lemma. We assume that (X, τ) is curve-compact and show that $\bigcap C \neq \emptyset$. We proceed by induction on *n*.

<u>Case n = 1</u>: Consider the definable total preorder \leq in *B* given by $b \leq c$ if and only if $\varphi(M, b) \subseteq \varphi(M, c)$. If *B* has a minimum **b** with respect to \leq , then $\varphi(M, b) \subseteq \varphi(M, c)$ for every $c \in B$, and the result follows. We suppose that (B, \leq) does not have a minimum and consider the definable nested family of (necessarily infinite) sets $\{(-\infty, b) \leq b \in B\}$, where $(-\infty, b) \leq c \leq B : c \leq b\}$ for every $b \in B$. Now let $a = \sup\{\inf(-\infty, b) \leq b \in B\}$, where the infimum and supremum are taken in $M \cup \{-\infty, +\infty\}$ with respect to the order < in *M*. We show that one of the families $\{(a, t) : t > a\}$ or $\{(t, a) : t < a\}$ is a refinement of $\{(-\infty, b) \le b \in B\}$. We prove the case where $a \in M$, since the case where $a \in \{-\infty, +\infty\}$ is similar but more straightforward.

Towards a contradiction suppose that $\{(-\infty, b)_{\leq} : b \in B\}$ does not have a refinement as described. Then, by o-minimality, there exist some $b_1 \in B$ and $t_1 > a$ such that $(-\infty, b_1)_{\leq} \cap (a, t_1) = \emptyset$, and similarly there exist $b_2 \in B$ and $t_2 < a$ such that $(-\infty, b_2)_{\leq} \cap (t_2, a) = \emptyset$. Additionally, by the definition of *a*, there exists $b_3 \in B$ such that $\inf(-\infty, b_3)_{\leq} > t_2$. Finally let $b_4 \in B$ be such that $a \notin (-\infty, b_4)_{\leq}$. Now let $i \leq 4$ be such that b_i is the minimum with respect to $\leq \inf\{b_1, b_2, b_3, b_4\}$. Then observe that $\inf(-\infty, b_i)_{\leq} \geq t_1$, contradicting the definition of *a*.

Hence onwards we assume that the family of intervals $\{(a, t) : t > a\}$ is a refinement of $\{(-\infty, b)_{\leq} : b \in B\}$; the case where the refinement is given by the family $\{(t, a) : t < a\}$ is analogous. This means that, for every $b \in B$, there exists an element t(b) > a such that $\varphi(M, c) \subset \varphi(M, b)$ for every $c \in (a, t(b))$.

Since either \mathcal{M} has definable choice or τ is Hausdorff, there exists by Lemma 5.10 a definable function $f: B \to \bigcup \mathcal{C}$ satisfying that $f(b) \in \varphi(M, b)$ for every $b \in B$. By the above paragraph it follows that, for every $b \in B$, if $c \in (a, t(b))$, then $f(c) \in \varphi(M, b)$. Let d > a be such that $(a, d) \subseteq B$ and γ be the restriction of f to (a, d). We derive that, for every $C \in \mathcal{C}$, it holds that $\lim_{t \to a}^{\tau} \gamma(t) \subseteq C$. Since (X, τ) is curve-compact, we conclude that $\bigcap \mathcal{C} \neq \emptyset$.

<u>Case n > 1</u>: For every $x \in \pi(B)$, let C_x denote the family $\{\varphi(M, x, t) : t \in B_x\}$, where $B_x = \{t \in M : \langle x, t \rangle \in B\}$, and set $C(x) := \bigcap C_x$. By the case n = 1 the definable family of τ -closed sets $\mathcal{D} = \{C(x) : x \in \pi(B)\}$ does not contain the empty set. Clearly $\bigcap \mathcal{D} = \bigcap C$. We observe that the family \mathcal{D} is nested and the result follows from the induction hypothesis.

Given $x, y \in \pi(B)$, if, for every $C \in C_x$, there is $C' \in C_y$ with $C' \subseteq C$, then $\bigcap C_y \subseteq \bigcap C_x$. Otherwise, there is $C \in C_x$ such that, for every $C' \in C_y$, it holds that $C \subseteq C'$, in which case $\bigcap C_x \subseteq C \subseteq \bigcap C_y$.

We may now prove Proposition 5.12.

Proof of Proposition 5.12. By Lemma 5.13 we must only prove the right-to-left implication in the second paragraph. Let (X, τ) , with $X \subseteq M^m$, be a curve-compact definable topological space, where either τ is Hausdorff or otherwise \mathcal{M} has definable choice. Let \mathcal{C} be a definable downward directed family of nonempty subsets of X, not necessarily τ -closed. We show that $\bigcap \{ cl_{\tau}(C) : C \in \mathcal{C} \} \neq \emptyset$.

We proceed by induction on $n = \min\{\dim C : C \in C\}$. Applying Proposition 4.3, after passing to a refinement of C if necessary, we may assume that C is a downward directed family of cells that generates a type in $S_m(M)$. If n = 0, then there exists a finite set in C and so (see Lemma 2.1) $\bigcap C \neq \emptyset$. Hence onwards we assume that n > 0. We begin by proving the case n = m. Hence suppose that every $C \in C$ is

an open cell $C = (f_C, g_C)$ for functions $f_C, g_C : \pi(C) \to M \cup \{-\infty, +\infty\}$ with $f_C < g_C$. Onwards recall the notation fixed in the last paragraph of Section 2.1. For every $C \in C$, consider the definable set $D(C) = \bigcap \{ cl_{\tau}(f_C, g_C') : C' \in C \}$.

Claim 5.15. For every $C \in C$ it holds that $D(C) \neq \emptyset$.

Proof. Let us fix C = (f, g) and, for every $x \in \pi(C)$, let c_x denote a point in $\lim_{t \to f(x)^+} \langle x, t \rangle$. If τ is Hausdorff then, for every $x \in \pi(C)$, there is a unique choice for c_x ; otherwise we use definable choice to pick c_x definably in x. The definable set $C^0 = \{c_x : x \in \pi(C)\}$ has dimension at most $\dim(\pi(C)) = \dim(C) - 1 = n - 1$. For every $C' = (f', g') \in C$, since $C \cap C' \neq \emptyset$, the definable set $\{x \in \pi(C) \cap \pi(C') : f(x) < g'(x)\} \supseteq \pi(C \cap C')$ is nonempty, and so $C^0 \cap cl_{\tau}(f, g') \neq \emptyset$.

Note that, because C is downward directed, the definable family of nonempty sets $\{C^0 \cap cl_\tau(f, g_{C'}) : C' \in C\}$ is downward directed. Since dim $C^0 \le n-1$, by inductive hypothesis there is a point that belongs in the τ -closure of $C^0 \cap cl_\tau(f, g_{C'})$ — in particular in $cl_\tau(f, g_{C'})$ — for all $C' \in C$. Hence $D(C) \ne \emptyset$.

Note that, for every $C \in C$, it holds that $D(C) \subseteq cl_{\tau}(C)$. We now show that the definable family of nonempty (by Claim 5.15) τ -closed sets $\{D(C) : C \in C\}$ is nested. Then, by Lemma 5.14, $\bigcap \{D(C) : C \in C\} \neq \emptyset$, and thus $\bigcap \{cl_{\tau}(C) : C \in C\} \neq \emptyset$.

Let us fix $C_1 = (f_1, g_1)$ and $C_2 = (f_2, g_2)$ in C. We may partition $B = \pi(C_1) \cap \pi(C_2)$ into the definable sets

$$B_1 = \{x \in B : f_1(x) \le f_2(x)\}$$
 and $B_2 = \{x \in B : f_1(x) > f_2(x)\}$

Since *C* is a basis (i.e., a downward directed generating family) for a type in $S_m(M)$, there exists some $i \in \{1, 2\}$ and $C \in C$ such that $\pi(C) \subseteq B_i$. Without loss of generality suppose that i = 1, and let us fix $C_3 \in C$ with $\pi(C_3) \subseteq B_1$. For an arbitrary set $C = (f, g) \in C$, let $C' = (f', g') \in C$ be such that $C' \subseteq C \cap C_3$. Then, clearly $(f_2, g') \subseteq (f_1, g') \subseteq (f_1, g)$. It follows that $D(C_2) \subseteq D(C_1)$. This completes the proof of the case n = m > 0.

Finally, we describe how the proof in the case 0 < n < m can be obtained by adapting the arguments above. Fix $\widehat{C} \in C$ with dim $\widehat{C} = n < m$ and a projection $\widehat{\pi} : M^m \to M^n$ such that $\widehat{\pi}|_{\widehat{C}} : \widehat{C} \to \widehat{\pi}(\widehat{C})$ is a bijection. By passing to a refining subfamily of C if necessary, we may assume that every set in C is contained in \widehat{C} . By the definition of n it follows that the definable downward directed family $\widehat{\pi}(C) = \{\widehat{\pi}(C) : C \in C\}$ contains only open cells in M^n . Note moreover that this family is a basis for a type in $S_n(M)$.

Set $h := (\hat{\pi}|_{\widehat{C}})^{-1}$ and, for every $C \in \mathcal{C}$, let $\hat{\pi}(C) = (\hat{f}_C, \hat{g}_C)$. Then the proof in the case n = m can be applied by letting D(C) be the intersection $\bigcap \{ cl_{\tau} h[(\hat{f}_C, \hat{g}_{C'})] : C' \in \mathcal{C} \}$, and letting C^0 be a set given by points c_x in $\lim_{t \to \hat{f}(x)^+} h(x, t)$ chosen definably in $x \in \pi(\hat{\pi}(C))$.



Figure 1. Depicting (in gray) the set A(x', x'', x''', y', y'', y''').

The following is an example of a non-Hausdorff definable topological space in the unbounded dense linear order (M, <) that is curve-compact but not definably compact. In particular, the space admits a definable nested family of nonempty closed sets with empty intersection (see the comments above Lemma 5.14).

Example 5.16. Let $X = \{\langle x, y \rangle \in M^2 : y < x\}$. Consider the family \mathcal{B} of subsets of *X* of the form

$$A(x', x'', x''', y', y'', y''') = \{\langle x, y \rangle : y < y', y < x\} \cup \{\langle x, y \rangle : y'' < y < y''' \land (y < x < y''' \lor x' < x < x'' \lor x''' < x)\}$$

definable uniformly in y' < y'' < y''' < x' < x'' < x'''.

Given any $A_0 = A(x'_0, x''_0, x'''_0, y''_0, y''_0)$ and $A_1 = A(x'_1, x''_1, x''_1, y'_1, y''_1, y''_1)$ in \mathcal{B} , and any $\langle x, y \rangle \in A_1 \cap A_2$, since every set in \mathcal{B} is open in the Euclidean topology, we may find y'' < y < y''' < x' < x < x'' such that the box $(x', x'') \times (y'', y''')$ is a subset of $A_1 \cap A_2$. Let $y' < \min\{y'', y'_0, y'_1\}$ and $x''' > \max\{x'', x''_0, x''_1\}$. Then $\langle x, y \rangle \in A(x', x'', x''', y', y'', y''') \subseteq A_1 \cap A_2$. Hence the family \mathcal{B} is a definable basis for a topology $\tilde{\tau}$ on X.

This topology is T_1 , i.e., every singleton is $\tilde{\tau}$ -closed. For every $y \in M$,

$$\lim_{t \to y^+} \langle t, y \rangle = \lim_{t \to +\infty} \langle t, y \rangle = (M \times \{y\}) \cap X,$$

and, for every $x \in M$,

$$\lim_{t \to x^{-}} \langle x, t \rangle = (M \times \{x\}) \cap X \text{ and } \lim_{t \to -\infty} \tilde{\langle} x, t \rangle = X.$$

In particular, $\tilde{\tau}$ is not Hausdorff.

Now suppose that $\mathcal{M} = (M, <)$. By quantifier elimination we know that in this structure any definable partial map $M \to M$ is piecewise either constant or the identity. Let $\gamma = (\gamma_0, \gamma_1) : (a, b) \to X$ be an injective definable curve in X, where γ_0 and γ_1 are the projections to the first and second coordinates respectively. Let

 $I = (a, c) \subseteq (a, b)$ be an interval where γ_0 and γ_1 are either constant or the identity. Since the graph of the identity is disjoint from X and γ is injective it must be that γ_i is constant and γ_{1-i} is the identity on I for some $i \in \{0, 1\}$.

Suppose that $\gamma_1|_I$ is constant with value y. Then, by the observations made above about the topology $\tilde{\tau}$, the curve γ satisfies that it $\tilde{\tau}$ -converges as $t \to a$ to either $\langle a, y \rangle$ (if y < a) or $(M \times \{y\}) \cap X$ (if a = y). On the other hand, if $\gamma_0|_I$ has a constant value x, then $\gamma \tilde{\tau}$ -converges as $t \to a$ to either $\langle x, a \rangle$ (if $a > -\infty$) or the whole space X (if $a = -\infty$). Analyzing the limit as $t \to b$ similarly allows us to conclude that γ is $\tilde{\tau}$ -completable. Hence the space $(X, \tilde{\tau})$ is curve-compact.

Meanwhile, the definable nested family of $\tilde{\tau}$ -closed sets { $X \cap (M \times [b, +\infty))$: $b \in M$ } has empty intersection. In particular, $(X, \tilde{\tau})$ is not definably compact.

Remark 5.17. Question 4.14 of [Johnson 2018] asks whether curve-compactness and filter-compactness are equivalent for o-minimal definable manifold spaces [van den Dries 1998, Chatper 10, Section 1]. While these spaces are not necessarily Hausdorff observe that, because they admit a covering by finitely many Hausdorff open subspaces, every definable curve in them converges to only finitely many points. It follows that the proofs of Lemmas 5.10 and 5.14 and Proposition 5.12 adapt to these spaces without any assumption that they are Hausdorff or that \mathcal{M} has definable choice. Hence we can answer Johnson's question in the affirmative. In fact, by Theorem A, every definable manifold space is curve-compact if and only if it is definably compact.

In the next remark we relate definable compactness to definable nets.

Remark 5.18. In [Andújar Guerrero et al. 2021, Section 6], Thomas, Walsberg and the author introduce the notion of *definable net* $\gamma : (B, \preceq) \rightarrow (X, \tau)$ to mean a definable map from a definable directed set (B, \preceq) into a definable topological space (X, τ) . Recall that a (*Kelley*) subnet of γ is a net $\gamma' : (B', \preceq') \rightarrow (X, \tau)$ such that $\gamma' = \gamma \circ \mu$, where $\mu : B' \rightarrow B$ satisfies that, for every $b \in B$, there exists $c \in B'$ satisfying that $\mu(d) \succeq b$ for every $d \succeq c$. We say that such a net γ' is definable if (B', \preceq') and μ are definable.

Classically, a topological space is compact if and only if every net in it has a converging subnet. Following the classical proof of this result one may show that, in any model-theoretic structure (regardless of the axiom of o-minimality), filter-compactness implies that every definable net has a definable converging subnet (say net-compactness). The reverse implication follows whenever the structure has definable choice. In the o-minimal case one may easily show that net-compactness implies curve-compactness and so, by Theorem A, net-compactness implies definable compactness whenever the topology is Hausdorff or \mathcal{M} has definable choice. See [Andújar Guerrero et al. 2021, Corollary 44] for a detailed proof of the equivalence between net-compactness, curve-compactness and filter-compactness in o-minimal

expansions of ordered groups. Furthermore, one may show that Example 5.16 above is not net-compact. The author is unaware of whether net-compactness implies definable compactness always within o-minimality.

In general topology there are further characterizations of compactness, such as the property that every net has a cluster point, which are not addressed in this paper.

In light of Definition 1.1, the following corollary is a direct consequence of Proposition 5.5.

Corollary 5.19. Suppose that \mathcal{M} is an o-minimal expansion of $(\mathbb{R}, <)$ and let (X, τ) be a definable topological space. Then (X, τ) is definably compact if and only if it is compact.

Remark 5.20. Part of [Peterzil and Steinhorn 1999, Question 2.5] asks whether curve-compactness is equivalent to classical compactness among definable manifold spaces in o-minimal expansions of (\mathbb{R} , <). Theorem A, Remark 5.17 and Corollary 5.19 above provide a positive answer to this question. The other part of [Peterzil and Steinhorn 1999, Question 2.5] asks whether curve-compactness for o-minimal definable manifold spaces is maintained after passing to an o-minimal expansion. Recall that Example 5.16 describes a (non-Hausdorff) definable topological space (X, $\tilde{\tau}$) in an arbitrary dense linear order (M, <) which is curve-compact but not definably compact. The linear order (M, <) could be chosen to have an o-minimal expansion \mathcal{N} with definable choice, in which case, by Theorem A, in \mathcal{N} the space (X, $\tilde{\tau}$) would lose the property of curve-compactness. Hence the above question has a negative answer when directed at all (non-Hausdorff) definable topological spaces. On the other hand, by Corollary 5.19, definable compactness is always maintained after passing to an expansion in o-minimal structures over (\mathbb{R} , <). It remains open however whether the same is true in arbitrary o-minimal structures.

Notice that all the characterizations of definable compactness in Theorem A are upfront expressible with infinitely many sentences (possibly with parameters) in the language of \mathcal{M} , i.e., you have to check all relevant definable families of closed sets or all definable curves. Recall that, by [Peterzil and Steinhorn 1999, Theorem 2.1] (or alternatively [Johnson 2018, Proposition 3.10]) and Theorem A, a definable set with the Euclidean topology is definably compact if and only if it is closed and bounded. Being closed and bounded is expressible by a single formula (in the same parameters used to define the set). Furthermore, given a definable family of sets with the Euclidean topology, the subfamily of those that are closed and bounded is definable.

We generalize this last observation to all definably compact spaces definable in o-minimal structures, showing along the way that definable compactness is expressible with a single formula (in the same parameters that define the topological space). We prove this for type-compactness and apply Theorem A. We use the following fact regarding strict prodefinability of the space of definable types from [Kovacsics et al. 2021], which can also be proved using Propositions 4.1 and 4.3, together with the Marker–Steinhorn theorem [1994, Theorem 2.1]. Recall that throughout this section our structure \mathcal{M} is o-minimal.

Fact 5.21 [Kovacsics et al. 2021, Theorems 2.4.8 and 3.2.1]. For every formula $\varphi(x, y)$, there exists a formula $\psi(y, z)$ and a definable (over \emptyset) set $Z \subseteq M^{|z|}$ with the following properties. For every $d \in Z$, there exists a definable type $p(x) \in S_{|x|}(M)$ such that

$$\{b \in M^{|y|} : \varphi(x, b) \in p(x)\} = \psi(M, d),$$
(2)

and, vice versa, for every definable type $p \in S_{|x|}(M)$ there exists some $d \in Z$ satisfying (2).

Proposition 5.22. Let $\{(X_c, \tau_c) : c \in C\}$, be a definable family of topological spaces, i.e., there exists a definable set $B \subseteq M^{n+m}$ and some formula $\sigma(x, u, v)$, with |u| = n and |v| = m, such that C is the projection of B to the last m coordinates and, for every $c \in C$, the family $\{\sigma(M, b, c) : b \in B_c\}$, where $B_c = \{b \in M^n : \langle b, c \rangle \in B\}$, is a basis for τ_c . Then the subfamily of all definably compact spaces is definable, i.e., there exists a definable set $D \subseteq C$ such that (X_c, τ_c) is definably compact if and only if $c \in D$.

In particular, given any definable family of subsets of a definable topological space, the subfamily of those that are definably compact is definable.

Proof. Let $\varphi(x, u, v)$ be a formula such that, for every $c \in C \subseteq M^{|v|}$, the family $\{\varphi(M, b, c) : b \in B_c\}$ is a basis of closed sets for τ_c . Recall that, as observed in the proof of Proposition 5.5, a type has a limit if and only if the intersection of all *basic* closed sets in it is nonempty.

Let $\psi(u, v, z)$ and $Z \subseteq M^{|z|}$ be as given by Fact 5.21 with y = (u, v). It follows that, for any fixed $c \in C$, the space (X_c, τ_c) is type-compact if and only if either X_c is empty or the following holds:

$$\forall z \in Z \ (\exists x \ \forall u \ (\psi(u, c, z) \land u \in B_c \to \varphi(x, u, c))).$$

Since the set Z is definable, this completes the proof.

In the case where \mathcal{M} is an o-minimal expansion of an ordered field, the completeness of the theory of tame pairs [van den Dries and Lewenberg 1995] can be used to circumvent some of the proofs in this paper. In particular Thomas, Walsberg and the author [Andújar Guerrero et al. 2021, Corollary 47] use it to prove the equivalence between curve-compactness and type-compactness among definable topological spaces, drawing inspiration from Walsberg's previous proof [2015, Proposition 6.6] of the equivalence between curve-compactness and sequential compactness among definable metric spaces in the case where \mathcal{M} expands the field of reals.

 \square

Acknowledgements

The author thanks Sergei Starchenko for the idea of a characterization of definable compactness in terms of VC theory and finite transversals, which motivated much of the work in this paper. He also thanks Margaret Thomas, Pantelis Eleftheriou, Matthias Aschenbrenner, and the anonymous referee for extensive feedback on various versions of the paper. In particular, he thanks Professor Aschenbrenner for suggesting the terminology "*n*-consistent" and "*n*-inconsistent". The author also thanks Antongiulio Fornasiero [2015] for sharing his unpublished notes on definable compactness. Finally, the author is deeply indebted to Will Johnson for some very helpful conversations on the contents of Section 4, which resulted in the current proof to Proposition 4.3, and in Johnson's authoring of [Andújar Guerrero 2021b, Appendix A], a counterexample to a parameter version of Proposition 4.3.

During the research into the contents of this paper the author was supported by the Institut Henri Poincaré (during the 2018 Program on Model Theory, Combinatorics and Valued Fields), the Canada Natural Sciences and Engineering Research Council (NSERC) Discovery Grant RGPIN-06555-2018, the Graduate School and Department of Mathematics at Purdue University, the Fields Institute for Research in Mathematical Sciences (during the 2021 Thematic Program on Trends in Pure and Applied Model Theory), and the Engineering and Physical Sciences Research Council (EPSRC) Grant EP/V003291/1.

References

- [Andújar Guerrero 2021a] P. Andújar Guerrero, *Definable topological spaces in o-minimal structures*, Ph.D. thesis, Purdue University, 2021, available at https://www.proquest.com/docview/2838331268. MR
- [Andújar Guerrero 2021b] P. Andújar Guerrero, "Types, transversals and definable compactness in o-minimal structures", preprint, 2021. arXiv 2111.03802
- [Andújar Guerrero and Johnson 2024] P. Andújar Guerrero and W. Johnson, "Around definable types in *p*-adically closed fields", *Ann. Pure Appl. Logic* **175**:10 (2024), art. id. 103484. MR Zbl
- [Andújar Guerrero et al. 2021] P. Andújar Guerrero, M. E. M. Thomas, and E. Walsberg, "Directed sets and topological spaces definable in o-minimal structures", *J. Lond. Math. Soc.* (2) **104**:3 (2021), 989–1010. MR Zbl
- [Aschenbrenner and Fischer 2011] M. Aschenbrenner and A. Fischer, "Definable versions of theorems by Kirszbraun and Helly", *Proc. Lond. Math. Soc.* (3) **102**:3 (2011), 468–502. MR Zbl
- [Aschenbrenner et al. 2016] M. Aschenbrenner, A. Dolich, D. Haskell, D. Macpherson, and S. Starchenko, "Vapnik-Chervonenkis density in some theories without the independence property, I", *Trans. Amer. Math. Soc.* **368**:8 (2016), 5889–5949. MR Zbl
- [Chernikov and Kaplan 2012] A. Chernikov and I. Kaplan, "Forking and dividing in NTP₂ theories", *J. Symbolic Logic* **77**:1 (2012), 1–20. MR Zbl
- [Chernikov and Simon 2015] A. Chernikov and P. Simon, "Externally definable sets and dependent pairs, II", *Trans. Amer. Math. Soc.* **367**:7 (2015), 5217–5235. MR Zbl

- [Cotter and Starchenko 2012] S. Cotter and S. Starchenko, "Forking in VC-minimal theories", *J. Symbolic Logic* **77**:4 (2012), 1257–1271. MR Zbl
- [Dolich 2004] A. Dolich, "Forking and independence in o-minimal theories", *J. Symbolic Logic* **69**:1 (2004), 215–240. MR Zbl
- [Dolich and Goodrick 2022] A. Dolich and J. Goodrick, "Tame topology over definable uniform structures", *Notre Dame J. Form. Log.* **63**:1 (2022), 51–79. MR Zbl
- [van den Dries 1998] L. van den Dries, *Tame topology and o-minimal structures*, London Mathematical Society Lecture Note Series **248**, Cambridge University Press, 1998. MR Zbl
- [van den Dries and Lewenberg 1995] L. van den Dries and A. H. Lewenberg, "*T*-convexity and tame extensions", *J. Symbolic Logic* **60**:1 (1995), 74–102. MR Zbl
- [Flum and Ziegler 1980] J. Flum and M. Ziegler, *Topological model theory*, Lecture Notes in Mathematics **769**, Springer, 1980. MR Zbl
- [Fornasiero 2015] A. Fornasiero, "Definable compactness for topological structures", unpublished notes, 2015.
- [Fujita 2024] M. Fujita, "Definable compactness in definably complete locally o-minimal structures", *Fund. Math.* **267**:2 (2024), 129–156. MR Zbl
- [Hrushovski and Loeser 2016] E. Hrushovski and F. Loeser, *Non-archimedean tame topology and stably dominated types*, Annals of Mathematics Studies **192**, Princeton University Press, 2016. MR Zbl
- [Johnson 2018] W. Johnson, "Interpretable sets in dense o-minimal structures", J. Symb. Log. 83:4 (2018), 1477–1500. MR Zbl
- [Johnson and Laskowski 2010] H. R. Johnson and M. C. Laskowski, "Compression schemes, stable definable families, and o-minimal structures", *Discrete Comput. Geom.* 43:4 (2010), 914–926. MR Zbl
- [Kaplan 2024] I. Kaplan, "A definable (p, q)-theorem for NIP theories", Adv. Math. 436 (2024), art. id. 109418. MR Zbl
- [Karpinski and Macintyre 2000] M. Karpinski and A. Macintyre, "Approximating volumes and integrals in o-minimal and *p*-minimal theories", pp. 149–177 in *Connections between model theory and algebraic and analytic geometry*, edited by A. Macintyre, Quad. Mat. **6**, Dept. Math., Seconda Univ. Napoli, Caserta, 2000. MR Zbl
- [Kovacsics et al. 2021] P. C. Kovacsics, M. Hils, and J. Ye, "Beautiful pairs", preprint, 2021. arXiv 2112.00651
- [Marker and Steinhorn 1994] D. Marker and C. I. Steinhorn, "Definable types in O-minimal theories", *J. Symbolic Logic* **59**:1 (1994), 185–198. MR Zbl
- [Matoušek 2004] J. Matoušek, "Bounded VC-dimension implies a fractional Helly theorem", *Discrete Comput. Geom.* **31**:2 (2004), 251–255. MR Zbl
- [Peterzil and Pillay 2007] Y. Peterzil and A. Pillay, "Generic sets in definably compact groups", *Fund. Math.* **193**:2 (2007), 153–170. MR Zbl
- [Peterzil and Steinhorn 1999] Y. Peterzil and C. Steinhorn, "Definable compactness and definable subgroups of o-minimal groups", *J. London Math. Soc.* (2) **59**:3 (1999), 769–786. MR Zbl
- [Pillay 1987] A. Pillay, "First order topological structures and theories", J. Symbolic Logic **52**:3 (1987), 763–778. MR Zbl
- [Pillay 1988] A. Pillay, "On groups and fields definable in *o*-minimal structures", *J. Pure Appl. Algebra* **53**:3 (1988), 239–255. MR Zbl

- [Pillay 2004] A. Pillay, "Type-definability, compact Lie groups, and o-minimality", *J. Math. Log.* **4**:2 (2004), 147–162. MR Zbl
- [Simon 2015] P. Simon, "Invariant types in NIP theories", J. Math. Log. 15:2 (2015), art. id. 1550006. MR Zbl
- [Simon and Starchenko 2014] P. Simon and S. Starchenko, "On forking and definability of types in some DP-minimal theories", *J. Symb. Log.* **79**:4 (2014), 1020–1024. MR Zbl
- [Simon and Walsberg 2019] P. Simon and E. Walsberg, "Tame topology over dp-minimal structures", *Notre Dame J. Form. Log.* **60**:1 (2019), 61–76. MR Zbl
- [Thomas 2012] M. E. M. Thomas, "Convergence results for function spaces over o-minimal structures", J. Log. Anal. 4 (2012), art. id. 1. MR Zbl
- [Walsberg 2015] E. Walsberg, "On the topology of metric spaces definable in o-minimal expansions of fields", preprint, 2015. arXiv 1510.07291
- Received 29 Feb 2024. Revised 10 Jan 2025.
- PABLO ANDÚJAR GUERRERO:

pa377@cantab.net

School of Mathematics, University of Leeds, Leeds, United Kingdom



Model Theory

msp.org/mt

EDITORS-IN-CHIEF

Martin Hils	Westfälische Wilhelms-Universität Münster (Germany) hils@uni-muenster.de
Rahim Moosa	University of Waterloo (Canada) rmoosa@uwaterloo.ca
EDITORIAL BOARD	

Sylvy Anscombe	Université Paris Cité (France) sylvy.anscombe@imj-prg.fr
Alessandro Berarducci	Università di Pisa (Italy) berardu@dm.unipi.it
Emmanuel Breuillard	University of Oxford (UK) emmanuel.breuillard@gmail.com
Artem Chernikov	University of California, Los Angeles (USA) chernikov@math.ucla.edu
Charlotte Hardouin	Université Paul Sabatier (France) hardouin@math.univ-toulouse.fr
François Loeser	Sorbonne Université (France) francois.loeser@imj-prg.fr
Dugald Macpherson	University of Leeds (UK) h.d.macpherson@leeds.ac.uk
Alf Onshuus	Universidad de los Andes (Colombia) aonshuus@uniandes.edu.co
Chloé Perin	The Hebrew University of Jerusalem (Israel) perin@math.huji.ac.il
PRODUCTION	
Silvio Levy	(Scientific Editor) production@msp.org

See inside back cover or msp.org/mt for submission instructions.

Model Theory (ISSN 2832-904X electronic, 2832-9058 printed) at Mathematical Sciences Publishers, 798 Evans Hall #3840, c/o University of California, Berkeley, CA 94720-3840 is published continuously online.

MT peer review and production are managed by EditFlow[®] from MSP.

PUBLISHED BY mathematical sciences publishers nonprofit scientific publishing https://msp.org/ © 2025 Mathematical Sciences Publishers

Model Theory

no. 2 vol. 4 2025

101
131
163
177