


Gauge Theory and Low-Dimensional Topology: Progress and Interaction

On the spectral sets of Inoue surfaces

Daniel Ruberman and Nikolai Saveliev



On the spectral sets of Inoue surfaces

Daniel Ruberman and Nikolai Saveliev

We study the Inoue surfaces S_M with the Tricerri metric and the canonical spin^c structure, and the corresponding chiral Dirac operators twisted by a flat \mathbb{C}^* -connection. The twisting connection is determined by $z \in \mathbb{C}^*$, and the points for which the twisted Dirac operators \mathcal{D}_z^\pm are not invertible are called spectral points. We show that there are no spectral points inside the annulus $\alpha^{-1/4} < |z| < \alpha^{1/4}$, where $\alpha > 1$ is the only real eigenvalue of the matrix M that determines S_M , and find the spectral points on its boundary. Via Taubes' theory of end-periodic operators, this implies that the corresponding Dirac operators are Fredholm on any end-periodic manifold whose end is modeled on S_M .

1. Introduction

Inoue surfaces are compact complex surfaces with zero second Betti number which are most remarkable in that they contain no holomorphic curves. These surfaces, constructed by Inoue [18], belong to the class VII_0 in Kodaira's classification [6], which is to say that they are minimal connected compact complex surfaces X with Kodaira dimension $\kappa(X) = -\infty$ and the first Betti number $b_1(X) = 1$. In fact, any class VII_0 surface with vanishing second Betti number and no holomorphic curves is biholomorphic to an Inoue surface; see Bogomolov [7; 8] and Teleman [28]. Inoue surfaces, which are not Kähler because their first Betti number is odd, have been extensively studied from the viewpoints of both algebraic and differential geometry.

In this paper, we restrict ourselves to the Inoue surfaces X of class S_M associated with certain integral matrices $M \in \mathrm{SL}(3, \mathbb{Z})$ with one real eigenvalue $\alpha > 1$ and two complex eigenvalues $\beta \neq \bar{\beta}$. These surfaces, described in detail in Section 2, are known to be diffeomorphic to the mapping torus of a self-diffeomorphism of the 3-torus induced by M . It is in this incarnation that the surfaces S_M are best known

Ruberman was partially supported by NSF Grants DMS-1811111 and DMS-1952790, and Saveliev was partially supported by NSF Grant DMS-1952762.

MSC2020: primary 32J15, 53C55; secondary 57R57, 58J50.

Keywords: Dirac operator, Seiberg–Witten, Inoue surface, Tricerri metric.

to topologists. In particular, Cappell and Shaneson [9; 10] independently used some of the matrices M to construct a fake \mathbb{RP}^4 and interesting fibered 2-spheres in a homotopy 4-sphere. From this point of view, the manifolds S_M are given by surgery on this homotopy 4-sphere along those knots. The question of when this homotopy 4-sphere is in fact diffeomorphic to S^4 has received considerable attention [2; 1; 3; 14].

Inoue surfaces are an intriguing class of examples to which to apply our work on the Seiberg–Witten invariants [24] and the end-periodic index theorem [25]. Spectral properties of chiral Dirac operators $\mathcal{D}^\pm(X)$ play an important role in determining the index of associated Dirac operators on end-periodic manifolds whose end is modeled on an infinite cyclic cover of X . In applications of those papers to date [21; 20; 22], the infinite cyclic cover was a Riemannian product of the real line and a 3-manifold. In the case of an Inoue surface, while this cover is topologically the product of the real line and a 3-torus, it is not a metric product. (This is related to the fact that the monodromy of the bundle $X \rightarrow S^1$ has infinite order.) Since the end-periodic index is metric dependent, this makes for an index problem that must be investigated analytically. We study this problem for the Tricerri metric on X , which makes it into a locally conformal Kähler manifold, and the canonical spin^c structure; see Section 2.

More specifically, we are interested in the spectral sets of the associated chiral Dirac operators $\mathcal{D}^\pm(X)$. Recall from [24] that $z \in \mathbb{C}^*$ is a spectral point of $\mathcal{D}^\pm(X)$ if and only if the operator

$$z^f \circ \mathcal{D}^\pm(X) \circ z^{-f} = \mathcal{D}^\pm(X) - \ln z \cdot df$$

has nonzero kernel, where $f : X \rightarrow S^1$ is a smooth function realizing a generator of $H^1(X; \mathbb{Z}) = \mathbb{Z}$, and df operates by Clifford multiplication. One can easily check that the spectral sets of $\mathcal{D}^+(X)$ and $\mathcal{D}^-(X)$ are obtained from each other by inversion $\tau(z) = 1/\bar{z}$ with respect to the unit circle. The following theorem, which was announced in [25, Section 6.4], is the main result of this paper.

Theorem 1.1. *The operators $\mathcal{D}^\pm(X)$ have no spectral points in the annulus $\alpha^{-1/4} < |z| < \alpha^{1/4}$. Furthermore, the only spectral points of $\mathcal{D}^+(X)$ on the circles $|z| = \alpha^{-1/4}$ and $|z| = \alpha^{1/4}$ are, respectively, $z = \alpha^{1/4}\beta$ and $z = \alpha^{1/4}$.*

Let Z_∞ be a spin^c end-periodic manifold whose end is modeled on the infinite cyclic cover of an Inoue surface X . According to Taubes [27, Lemma 4.3], the Dirac operators $\mathcal{D}^\pm(Z_\infty)$ are Fredholm in the usual Sobolev L^2 completion if and only if their spectral sets are disjoint from the unit circle $|z| = 1$.

Corollary 1.2. *The operators $\mathcal{D}^\pm(Z_\infty) : L^2_1(Z_\infty) \rightarrow L^2(Z_\infty)$ are Fredholm on any end-periodic spin^c manifold Z_∞ whose end is modeled on an Inoue surface X of type S_M .*

Remark 1.3. Inoue surfaces do not admit metrics of positive scalar metric, as was proved by Albanese [4, Theorem 4.5]. This also follows from Cecchini and Schick [11], making use of the fact that Inoue surfaces are solvmanifolds (see Wall [31; 32] and Hasegawa [16]) and hence are enlargeable in the sense of Gromov and Lawson [15]. In particular, one cannot prove that the operators $\mathcal{D}^\pm(Z_\infty)$ of Corollary 1.2 are Fredholm by using the (uniformly) positive scalar curvature at infinity condition as in [15].

Once we establish that the operators $\mathcal{D}^\pm(Z_\infty)$ are Fredholm, their index can in principle be calculated as in [25] in terms of an integral term and the periodic eta-invariant $\eta(X)$. The latter is a spectral invariant which generalizes the eta-invariant of Atiyah, Patodi, and Singer [5] and which can be viewed as a regularized count of points in the spectral set of $\mathcal{D}^\pm(X)$. The partial information about the spectral set we obtain in this paper is not sufficient to calculate $\eta(X)$ or the associated index of $\mathcal{D}^\pm(Z_\infty)$. However, even this modest attempt leads to some fascinating analysis which we felt was worth sharing.

It is worth mentioning that our original interest in end-periodic index theory grew out of our work [24] with Mrowka on Seiberg–Witten theory for 4-manifolds X with $b_2(X) = 0$ and $b_1(X) = 1$. In that paper, a Seiberg–Witten invariant $\lambda_{\text{SW}}(X)$ was defined as a sum of two metric dependent terms. One is a count of solutions to the Seiberg–Witten equations, and the other is an index-theoretic correction term, whose most important part is the index of the Dirac operator $\mathcal{D}^+(Z_\infty)$.

Evaluating $\lambda_{\text{SW}}(X)$ for an Inoue surface X presents quite a challenge. One can actually solve a modified version of the Seiberg–Witten equations for the Tricerri metric; see [26; 23]. However, the modification involves a certain twisting of the Dirac operator used in the formulation of the Seiberg–Witten equations. In order to turn this into a calculation of $\lambda_{\text{SW}}(X)$, one would have to first relate this modified Seiberg–Witten equation to the one used in [25]. The second step would be to evaluate the correction term; this is essentially the same as finding the invariant $\eta(X)$. As mentioned above, we are quite far from achieving this.

In conclusion, we mention that a recent paper of Holt and Zhang [17] uses related techniques to investigate $\bar{\partial}$ -harmonic forms on a different non-Kähler complex manifold, the Kodaira–Thurston surface [19; 29].

2. Inoue surfaces

The Inoue surfaces X we are interested in are all compact quotients of $\mathcal{H} \times \mathbb{C}$, where $\mathcal{H} = \{w = w_1 + iw_2 \in \mathbb{C} \mid w_2 > 0\}$ is the upper complex half-plane. To construct X , start with an integral matrix $M \in \text{SL}(3, \mathbb{Z})$ with one real eigenvalue $\alpha > 1$ (which must therefore be irrational) and two complex conjugate eigenvalues $\beta \neq \bar{\beta}$. For

example, the matrices

$$A_m = \begin{pmatrix} 0 & 1 & 0 \\ 0 & 1 & 1 \\ 1 & 0 & m+1 \end{pmatrix},$$

which are equivalent to the Cappell and Shaneson [10] family, will do as long as $-2 \leq m \leq 3$. Let $a = (a_1, a_2, a_3)$ be a real eigenvector corresponding to α , and $b = (b_1, b_2, b_3)$ a complex eigenvector corresponding to β . Let G_M be the group of complex analytic transformations of $\mathcal{H} \times \mathbb{C}$ generated by

$$\begin{aligned} g_0(w, z) &= (\alpha w, \beta z), \\ g_i(w, z) &= (w + a_i, z + b_i), \quad i = 1, 2, 3. \end{aligned}$$

The group G_M acts on $\mathcal{H} \times \mathbb{C}$ freely and properly discontinuously so that the quotient $X = (\mathcal{H} \times \mathbb{C})/G_M$ is a compact complex surface.

Inoue [18] showed that, as a smooth manifold, X is a 3-torus bundle over a circle whose monodromy is given by the matrix M , and that $b_1(X) = 1$ and $b_2(X) = 0$. One can check, for example, that $H_*(X) = H_*(S^1 \times S^3)$ for all manifolds X obtained from the Cappell–Shaneson matrices A_m . Define a function $f : \mathcal{H} \times \mathbb{C} \rightarrow \mathbb{R}$ by the formula $f(w, z) = \ln w_2 / \ln \alpha$. One can easily see that df is a well-defined 1-form on X , whose cohomology class generates $H^1(X; \mathbb{Z}) = \mathbb{Z}$.

The complex surface X admits no global Kähler metric. We will however consider the following Hermitian metric on $\mathcal{H} \times \mathbb{C}$, called the Tricerri metric:

$$g = \frac{dw \otimes d\bar{w}}{w_2^2} + w_2 dz \otimes d\bar{z},$$

see [30; 12]. Let ω be the Kähler form associated with this metric, then $d\omega = d \ln w_2 \wedge \omega$, with the torsion form $d \ln w_2 = \ln \alpha \cdot df$. The metric g is G_M -invariant; hence it defines a metric on X which makes X into a locally conformal Kähler manifold (or l.c.K. manifold, for short).

The complex surface X admits a canonical spin^c structure with respect to which

$$S^+ = \Lambda^{0,0}(X) \oplus \Lambda^{0,2}(X) \quad \text{and} \quad S^- = \Lambda^{0,1}(X).$$

Let $\mathcal{D}^\pm(X)$ be the chiral Dirac operators associated with the Tricerri metric and the canonical spin^c structure on X . These are the operators that Theorem 1.1 is concerned with. The proof of Theorem 1.1 will take up the rest of these notes.

3. Reduction to the Dirac–Dolbeault operator

Let $\mathcal{D}^-(X)$ be the negative chiral Dirac operator associated with the Tricerri metric and the canonical spin^c structure on X . According to Gauduchon [13, page 283],

there is an isomorphism

$$\mathcal{D}^-(X) + \frac{1}{4} \ln \alpha \cdot df = \sqrt{2}(\bar{\partial} \oplus \bar{\partial}^*), \tag{1}$$

where

$$\bar{\partial} \oplus \bar{\partial}^* : \Omega^{0,1}(X) \rightarrow \Omega^{0,2}(X) \oplus \Omega^{0,0}(X) \tag{2}$$

is the Dirac–Dolbeault operator on the complex surface X . To prove Theorem 1.1, it will suffice to compute the spectral set of (2). The spectral set of $\mathcal{D}^-(X)$ will be obtained from it via multiplication by $\alpha^{-1/4}$, and the spectral set of $\mathcal{D}^+(X)$ by further inversion.

4. The periodic boundary value problem

To compute the spectral set of (2), we will complete the operator (2) to an operator $L_1^2 \rightarrow L^2$ and look for $z = e^\mu \in \mathbb{C}^*$ such that the kernel of the operator

$$e^{\mu f} \circ (\bar{\partial} \oplus \bar{\partial}^*) \circ e^{-\mu f} = (\bar{\partial} \oplus \bar{\partial}^*) - \mu \cdot df$$

on X is nonzero. Equivalently, after passing to the universal covering space $\mathcal{H} \times \mathbb{C} \rightarrow X$, we will look for μ such that the following periodic boundary problem on $\mathcal{H} \times \mathbb{C}$ has a nonzero solution $\omega \in \Omega^{0,1}(\mathcal{H} \times \mathbb{C})$:

$$(\bar{\partial} \oplus \bar{\partial}^*)(\omega) = 0, \quad \text{where } g_i^* \omega = \omega \text{ for } i = 1, 2, 3, \text{ and } g_0^* \omega = e^{-\mu} \cdot \omega.$$

Let us restate this periodic boundary problem by writing $\omega = ad\bar{w} + bd\bar{z}$ on $\mathcal{H} \times \mathbb{C}$. The equation $(\bar{\partial} \oplus \bar{\partial}^*)(\omega) = 0$ turns into the system

$$\begin{cases} \frac{\partial a}{\partial \bar{z}} - \frac{\partial b}{\partial \bar{w}} = 0, \\ \frac{\partial(w_2 a)}{\partial w} + \frac{1}{w_2^2} \cdot \frac{\partial b}{\partial z} = 0, \end{cases}$$

and, after introducing the new function $c = w_2 a$ and the new variable $t = \ln w_2$ into the system,

$$\left(\frac{\partial}{\partial t} + iB_t \right) \begin{pmatrix} b \\ c \end{pmatrix} = 0 \tag{3}$$

with

$$B_t = \begin{pmatrix} -e^t \frac{\partial}{\partial w_1} & 2 \frac{\partial}{\partial \bar{z}} \\ 2e^{-t} \frac{\partial}{\partial z} & e^t \frac{\partial}{\partial w_1} \end{pmatrix}.$$

Taking into account the periodic boundary conditions $g_i^* \omega = \omega$ for $i = 1, 2, 3$, this can be viewed as a system on the product $\mathbb{R} \times T^3$, with the coordinates t on

the real line and (w_1, z_1, z_2) on the torus T^3 . The remaining periodic boundary condition $g_0^* \omega = e^{-\mu} \cdot \omega$ can be expressed in the language of $(0, 1)$ -forms as

$$g_0^*(a(w, z)d\bar{w} + b(w, z)d\bar{z}) = e^{-\mu} \cdot (a(w, z)d\bar{w} + b(w, z)d\bar{z}).$$

After switching to $c = w_2 \cdot a$, this turns into

$$\bar{\beta} \cdot b(\alpha w, \beta z) = e^{-\mu} \cdot b(w, z) \quad \text{and} \quad c(\alpha w, \beta z) = e^{-\mu} \cdot c(w, z). \quad (4)$$

It is the periodic boundary value problem (3), (4) on the manifold $\mathbb{R} \times T^3$ that we now wish to solve.

5. Fourier analysis

We will use Fourier analysis on the 3-torus to solve the system (3). First, consider the following basis in \mathbb{R}^3 :

$$\xi = (a_1, \operatorname{Re} b_1, \operatorname{Im} b_1),$$

$$\eta = (a_2, \operatorname{Re} b_2, \operatorname{Im} b_2),$$

$$\zeta = (a_3, \operatorname{Re} b_3, \operatorname{Im} b_3),$$

where $a = (a_1, a_2, a_3)$ and $b = (b_1, b_2, b_3)$ are, as before, the eigenvectors of M corresponding to the eigenvalues α and β . The quotient of \mathbb{R}^3 by the integer lattice spanned by the vectors ξ, η, ζ is our 3-torus. The matrix whose rows are the vectors ξ, η, ζ will be called Y so that

$$Y = \begin{pmatrix} \xi_1 & \xi_2 & \xi_3 \\ \eta_1 & \eta_2 & \eta_3 \\ \zeta_1 & \zeta_2 & \zeta_3 \end{pmatrix}.$$

Without loss of generality, we will assume that $\det Y = 1$. The columns of the matrix

$$Y^{-1} = \begin{pmatrix} \xi_1^* & \eta_1^* & \zeta_1^* \\ \xi_2^* & \eta_2^* & \zeta_2^* \\ \xi_3^* & \eta_3^* & \zeta_3^* \end{pmatrix}$$

form the dual basis ξ^*, η^*, ζ^* with respect to the usual dot product (\cdot, \cdot) on \mathbb{R}^3 . One can easily check that the functions $T^3 \rightarrow \mathbb{C}$ defined by

$$\theta \rightarrow \exp(2\pi i(\theta, k\xi^* + \ell\eta^* + m\zeta^*)) \quad \text{for all } (k, \ell, m) \in \mathbb{Z}^3, \quad (5)$$

where $\theta = (\theta_1, \theta_2, \theta_3) = (w_1, z_1, z_2)$, form an orthonormal basis in the L^2 -space of complex-valued functions on the 3-torus.

For each $t \in \mathbb{R}$, expand the functions $b(t, \theta)$ and $c(t, \theta) : T^3 \rightarrow \mathbb{C}$ into Fourier series,

$$b(t, \theta) = \sum_{k, \ell, m} b_{k\ell m}(t) \exp(2\pi i(\theta, k\xi^* + \ell\eta^* + m\zeta^*))$$

and

$$c(t, \theta) = \sum_{k, \ell, m} c_{k\ell m}(t) \exp(2\pi i(\theta, k\xi^* + \ell\eta^* + m\zeta^*)),$$

and plug them into (3). For each individual triple of integers (k, ℓ, m) , we obtain the system

$$\begin{pmatrix} b'_{k\ell m} \\ c'_{k\ell m} \end{pmatrix} = \begin{pmatrix} -e^t P_{k\ell m} & Q_{k\ell m} \\ e^{-t} \bar{Q}_{k\ell m} & e^t P_{k\ell m} \end{pmatrix} \begin{pmatrix} b_{k\ell m} \\ c_{k\ell m} \end{pmatrix}, \tag{6}$$

where the prime stands for the t -derivative,

$$P_{k\ell m} = 2\pi(k\xi_1^* + \ell\eta_1^* + m\zeta_1^*) \in \mathbb{R}, \quad \text{and}$$

$$Q_{k\ell m} = 2\pi(k\xi_2^* + \ell\eta_2^* + m\zeta_2^*) + 2\pi i(k\xi_3^* + \ell\eta_3^* + m\zeta_3^*) \in \mathbb{C}.$$

This is a linear system of ordinary differential equations with nonconstant coefficients. Note that $P_{k\ell m}$ and $Q_{k\ell m}$ are actually constants so the only dependence of the coefficients on t comes from the factors of e^t and e^{-t} . For future use, we make the following observation.

Lemma 5.1. *For no choice of $(k, \ell, m) \neq (0, 0, 0)$ can $Q_{k\ell m}$ be equal to zero.*

Proof. Observe that

$$Y \begin{pmatrix} P_{k\ell m} \\ \operatorname{Re} Q_{k\ell m} \\ \operatorname{Im} Q_{k\ell m} \end{pmatrix} = 2\pi \begin{pmatrix} k \\ \ell \\ m \end{pmatrix}.$$

If $Q_{k\ell m} = 0$, the first column of Y , which is an eigenvector of M with the eigenvalue α , is proportional to the vector with integral coordinates k, ℓ , and m . The latter vector is then also an eigenvector of $M \in \operatorname{SL}(3, \mathbb{Z})$ with the eigenvalue α , which contradicts the fact that α is irrational. \square

Next, we need to take care of the boundary conditions (4). In our θ -notation, we have $\beta z = (\beta_1 + i\beta_2)(z_1 + iz_2) = (\beta_1 + i\beta_2)(\theta_2 + i\theta_3) = (\beta_1\theta_2 - \beta_2\theta_3) + i(\beta_2\theta_2 + \beta_1\theta_3)$ and $\alpha w = \alpha(w_1 + iw_2) = \alpha\theta_1 + ie^{t+\ln\alpha}$. To simplify notation, introduce the matrix

$$A = \begin{pmatrix} \alpha & 0 & 0 \\ 0 & \beta_1 & -\beta_2 \\ 0 & \beta_2 & \beta_1 \end{pmatrix},$$

then the boundary conditions (4) become

$$\bar{\beta} \cdot b(t + \ln \alpha, A(\theta)) = e^{-\mu} \cdot b(t, \theta), \quad c(t + \ln \alpha, A(\theta)) = e^{-\mu} \cdot c(t, \theta).$$

In order to rewrite these in terms of the Fourier coefficients $b_{k\ell m}$ and $c_{k\ell m}$, we need the following technical result.

Lemma 5.2. *For any integers k, ℓ and m , we have $(A(\theta), k\xi^* + \ell\eta^* + m\zeta^*) = (\theta, k'\xi^* + \ell'\eta^* + m'\zeta^*)$, where*

$$\begin{pmatrix} k' \\ \ell' \\ m' \end{pmatrix} = M \begin{pmatrix} k \\ \ell \\ m \end{pmatrix}. \quad (7)$$

Proof. A straightforward calculation with matrices shows that $MY = YA^t$. Viewing θ as a column, we obtain

$$\begin{aligned} (A(\theta), k\xi^* + \ell\eta^* + m\zeta^*) &= \theta^t A^t Y^{-1} \begin{pmatrix} k \\ \ell \\ m \end{pmatrix} = \theta^t Y^{-1} M \begin{pmatrix} k \\ \ell \\ m \end{pmatrix} = \theta^t Y^{-1} \begin{pmatrix} k' \\ \ell' \\ m' \end{pmatrix} \\ &= (\theta, k'\xi^* + \ell'\eta^* + m'\zeta^*). \quad \square \end{aligned}$$

Now, substitute the Fourier expansions of $b(t, \theta)$ and $c(t, \theta)$ into the boundary conditions to obtain

$$\begin{aligned} \bar{\beta} \cdot b(t + \ln \alpha, A(\theta)) &= \bar{\beta} \sum_{k, \ell, m} b_{k\ell m}(t + \ln \alpha) \exp(2\pi i (A(\theta), k\xi^* + \ell\eta^* + m\zeta^*)) \\ &= \bar{\beta} \sum_{k, \ell, m} b_{k\ell m}(t + \ln \alpha) \exp(2\pi i (\theta, k'\xi^* + \ell'\eta^* + m'\zeta^*)) \\ &= e^{-\mu} \sum_{k', \ell', m'} b_{k'\ell'm'}(t) \exp(2\pi i (\theta, k'\xi^* + \ell'\eta^* + m'\zeta^*)), \end{aligned}$$

and similarly for c . A term-by-term comparison of the coefficients allows us to conclude that

$$\bar{\beta} \cdot b_{k\ell m}(t + \ln \alpha) = e^{-\mu} \cdot b_{k'\ell'm'}(t), \quad c_{k\ell m}(t + \ln \alpha) = e^{-\mu} \cdot c_{k'\ell'm'}(t), \quad (8)$$

where the triples (k, ℓ, m) and (k', ℓ', m') are related by (7). Therefore, to fit $b_{k\ell m}(t)$ and $c_{k\ell m}(t)$ together into a Fourier series solution, we need to know how M acts on the triples (k, ℓ, m) .

6. Finite orbits

The infinite cyclic subgroup of $\mathrm{SL}(3, \mathbb{Z})$ generated by the matrix M acts on the lattice \mathbb{Z}^3 . The only finite orbit of this action consists of the triple $(k, \ell, m) = (0, 0, 0)$. The solutions of (6) corresponding to this triple must be constant; we will denote them by b and c . The boundary conditions (8) then translate into $\bar{\beta}b = e^{-\mu}b$ and $c = e^{-\mu}c$, resulting in exactly two choices for the spectral point $z = e^{\mu}$ of the operator (2), namely, $z = 1$ and $z = 1/\bar{\beta} = \alpha\beta$. These correspond to the spectral points $z = \alpha^{1/4}$ and $z = \alpha^{1/4}\beta$ of the operator $\mathcal{D}^+(X)$ as claimed in Theorem 1.1.

7. Infinite orbits

For any fixed triple of integers $(k_0, \ell_0, m_0) \neq (0, 0, 0)$, the triples (k_n, ℓ_n, m_n) , $n \in \mathbb{Z}$, in its orbit can be found from the equation

$$\begin{pmatrix} k_n \\ \ell_n \\ m_n \end{pmatrix} = M^n \begin{pmatrix} k_0 \\ \ell_0 \\ m_0 \end{pmatrix}.$$

Denote $b_n(t) = b_{k_n \ell_n m_n}(t)$ and $c_n(t) = c_{k_n \ell_n m_n}(t)$. It follows from (8) that, once we know $b_0(t)$ and $c_0(t)$, the rest of $b_n(t)$ and $c_n(t)$ can be determined uniquely from the recursive relation

$$b_{n+1}(t) = \bar{\beta} \cdot e^\mu \cdot b_n(t + \ln \alpha), \quad c_{n+1}(t) = e^\mu \cdot c_n(t + \ln \alpha).$$

Therefore, each infinite orbit gives rise to the infinite series

$$\begin{aligned} b(t, \theta) &= \sum_{n \in \mathbb{Z}} \bar{\beta}^n \cdot e^{n\mu} \cdot b_0(t + n \ln \alpha) \cdot \exp(2\pi i(\theta, k_n \xi^* + \ell_n \eta^* + m_n \zeta^*)), \\ c(t, \theta) &= \sum_{n \in \mathbb{Z}} e^{n\mu} \cdot c_0(t + n \ln \alpha) \cdot \exp(2\pi i(\theta, k_n \xi^* + \ell_n \eta^* + m_n \zeta^*)). \end{aligned}$$

The question becomes whether these formal series solutions converge to a solution of (3). We will show that, for certain values of μ , the series cannot converge in L^2 norm unless $b_0(t) = c_0(t) = 0$; this will imply that the corresponding $z = e^\mu$ are not in the spectral set of the operator $\bar{\partial} \oplus \bar{\partial}^*$. To this end, denote by δ the real number

$$\delta = \operatorname{Re} \mu / \ln \alpha - 1/4$$

and introduce the notation

$$u(t) = b_0(t) \quad \text{and} \quad v(t) = e^{t/2} c_0(t).$$

Lemma 7.1. *The above Fourier series for $b(t, \theta)$ and $c(t, \theta)$ converge to L^2_1 sections on X if and only if both $u(t)$ and $v(t)$ belong to $L^2_{1, \delta-1/4}(\mathbb{R})$.*

Proof. Let $z = \bar{\beta} \cdot e^\mu$, then $z^{t/\ln \alpha} \cdot b(t, \theta)$ is the Fourier–Laplace transform [24] of the function $u(t) \exp(2\pi i(\theta, k_0 \xi^* + \ell_0 \eta^* + m_0 \zeta^*))$ on $\mathbb{R} \times T^3$ with respect to the covering translation $(t, \theta) \rightarrow (t + \ln \alpha, A(\theta))$. One can easily check that

$$|z^{1/\ln \alpha}| = e^{\delta-1/4}.$$

From this point on, we follow the proof of [24, Proposition 4.2] and use the fact that the functions $\exp(2\pi i(\theta, k_n \xi^* + \ell_n \eta^* + m_n \zeta^*))$ form an orthonormal basis on

the fibers $\{t\} \times T^3$. For example, it follows by direct calculation that

$$\begin{aligned} \|z^{t/\ln \alpha} \cdot b(t, \theta)\|_{L^2(X)}^2 &= \sum_{n \in \mathbb{Z}} \int_0^{\ln \alpha} |z|^{2(n+t/\ln \alpha)} \cdot |u(t + n \ln \alpha)|^2 dt \\ &= \int_{-\infty}^{\infty} |z|^{2t/\ln \alpha} \cdot |u(t)|^2 dt = \|u\|_{L^2_{\delta-1/4}(\mathbb{R})}^2. \end{aligned}$$

The proof for the function $c(t, \theta)$ is similar. □

One can easily check using (6) that the functions $u(t)$ and $v(t)$ solve the system of ordinary differential equations

$$\begin{pmatrix} u' \\ v' \end{pmatrix} = \begin{pmatrix} -Pe^t & Qe^{-t/2} \\ \bar{Q}e^{-t/2} & 1/2 + Pe^t \end{pmatrix} \begin{pmatrix} u \\ v \end{pmatrix}, \tag{9}$$

where $P = P_{k_0 \ell_0 m_0} \in \mathbb{R}$ and $Q = Q_{k_0 \ell_0 m_0} \in \mathbb{C}$. Because of Lemma 7.1, we are only interested in solutions $u(t)$ and $v(t)$ which belong to $L^2_{1, \delta-1/4}(\mathbb{R})$.

Proposition 7.2. *Suppose that $-1/4 \leq \delta \leq 1/4$, then all solutions $u(t), v(t)$ of the system (9) which belong to $L^2_{1, \delta-1/4}(\mathbb{R})$ are identically zero.*

Proof. Decoupling (9) turns it into the following pair of Sturm–Liouville problems:

$$-u'' + (Pe^t(Pe^t - 1) + |Q|^2e^{-t})u = 0 \quad \text{and} \tag{10}$$

$$-v'' + (Pe^t(Pe^t + 2) + |Q|^2e^{-t} + 1/4)v = 0. \tag{11}$$

Without loss of generality, we will assume that u and v are real-valued functions. We will separate our argument into three cases, depending on whether P is positive, negative, or zero.

If $P < 0$, introduce the positive real numbers $p = -P$ and $q = |Q|$ and rewrite (10) in the form $-u'' + U(t)u = 0$ with the everywhere-positive potential $U(t) = pe^t(pe^t + 1) + q^2e^{-t}$. For any choice of $a < b$, we then have

$$-\int_a^b u''(t)u(t) dt + \int_a^b U(t)u^2(t) dt = 0$$

and, after integration by parts,

$$\int_a^b u'(t)^2 dt + u(a)u'(a) - u(b)u'(b) + \int_a^b U(t)u(t)^2 dt = 0. \tag{12}$$

The first and the last terms in this formula are nonnegative for any choice of $a < b$. We will show that there exist a arbitrarily close to $-\infty$ and b arbitrarily close to $+\infty$ such that the other two terms in (12) are nonnegative as well. This will imply that $u(t) = 0$. Plugging $u(t) = 0$ back into (9) will then imply that $v(t) = 0$ because $Q \neq 0$ by Lemma 5.1.

We first show that for any a_0 there exists $a \leq a_0$ such that $u(a)u'(a) \geq 0$. If $u(a_0) = 0$, we are finished. Otherwise, suppose that $u(t)u'(t) < 0$ for all $t \leq a_0$. Then $(u^2(t))' = 2u(t)u'(t) < 0$ so that $u^2(t)$ is a decreasing function and hence $u^2(t) \geq u^2(a_0) > 0$ for all $t \leq a_0$. This contradicts the fact that $u \in L^2_{\delta-1/4}(\mathbb{R})$ with $\delta \leq 1/4$.

Next, we show that for any b_0 there exists $b \geq b_0$ such that $u(b)u'(b) \leq 0$. If $u(b_0) = 0$ we are finished. Otherwise, suppose that $u(t)u'(t) > 0$ for all $t \geq b_0$. Then $(u^2(t))' = 2u(t)u'(t) > 0$ so that $u^2(t)$ is an increasing function and hence $u^2(t) \geq u^2(b_0) > 0$ for all $t \geq b_0$. Using the formula (12) with $a = b_0$ we obtain the estimate

$$u(b)u'(b) \geq \int_{b_0}^b U(t)u^2(t) dt \geq u^2(b_0) \int_{b_0}^b U(t) dt,$$

and using the fact that $U(t) \geq p^2e^{2t}$ for all t , the estimate

$$u(b)u'(b) \geq \frac{1}{2}p^2u^2(b_0)(e^{2b} - e^{2b_0}) \quad \text{for all } b \geq b_0.$$

Since $u(t)$ and $u'(t)$ belong to $L^2_{\delta-1/4}(\mathbb{R})$, it follows from the Hölder inequality that $u(t)u'(t) \in L^1_{2(\delta-1/4)}(\mathbb{R})$. This contradicts the above estimate for $\delta \geq -1/4$.

If $P > 0$, essentially the same argument using (11) shows that $v(t) = 0$. After plugging $v(t) = 0$ back in (9), we see that $u(t) = 0$ as well.

In the remaining case of $P = 0$, both (10) and (11) admit explicit solutions in terms of Bessel functions. To be precise, the general solution of (11) is of the form

$$C_1 \cdot I_1(2qe^{-t/2}) + C_2 \cdot K_1(2qe^{-t/2}), \tag{13}$$

where $I_1(x)$ and $K_1(x)$ are the modified Bessel functions of the first and second kind, solving the equation $x^2y'' + xy' - (x^2 + 1)y = 0$. One can check that the zero function is the only function among (13) that belongs to $L^2_{\delta-1/4}(\mathbb{R})$ with $-1/4 \leq \delta \leq 1/4$. □

Proposition 7.2 together with the discussion in Section 6 completes the proof of Theorem 1.1.

Acknowledgments

We thank Tom Mrowka, Leonid Parnovski, and Andrei Teleman for generously sharing their expertise, and the anonymous referee for useful comments.

References

- [1] I. R. Aitchison and J. H. Rubinstein, “Fibered knots and involutions on homotopy spheres”, pp. 1–74 in *Four-manifold theory* (Durham, N.H., 1982), Contemp. Math. **35**, Amer. Math. Soc., Providence, RI, 1984. MR Zbl
- [2] S. Akbulut and R. Kirby, “An exotic involution of S^4 ”, *Topology* **18**:1 (1979), 75–81. MR Zbl

- [3] S. Akbulut and R. Kirby, “A potential smooth counterexample in dimension 4 to the Poincaré conjecture, the Schoenflies conjecture, and the Andrews–Curtis conjecture”, *Topology* **24**:4 (1985), 375–390. MR Zbl
- [4] M. Albanese, “The Yamabe invariants of Inoue surfaces, Kodaira surfaces, and their blowups”, *Ann. Global Anal. Geom.* **59**:2 (2021), 179–195. MR Zbl
- [5] M. F. Atiyah, V. K. Patodi, and I. M. Singer, “Spectral asymmetry and Riemannian geometry, I”, *Math. Proc. Cambridge Philos. Soc.* **77** (1975), 43–69. MR Zbl
- [6] W. P. Barth, K. Hulek, C. A. M. Peters, and A. Van de Ven, *Compact complex surfaces*, 2nd ed., *Ergeb. Math. Grenzgeb.* **4**, Springer, 2004. MR Zbl
- [7] F. A. Bogomolov, “Classification of surfaces of class VII_0 with $b_2 = 0$ ”, *Izv. Akad. Nauk SSSR Ser. Mat.* **40**:2 (1976), 273–288. MR
- [8] F. A. Bogomolov, “Surfaces of class VII_0 and affine geometry”, *Izv. Akad. Nauk SSSR Ser. Mat.* **46**:4 (1982), 710–761. MR
- [9] S. E. Cappell and J. L. Shaneson, “Some new four-manifolds”, *Ann. of Math. (2)* **104**:1 (1976), 61–72. MR Zbl
- [10] S. E. Cappell and J. L. Shaneson, “There exist inequivalent knots with the same complement”, *Ann. of Math. (2)* **103**:2 (1976), 349–353. MR Zbl
- [11] S. Cecchini and T. Schick, “Enlargeable metrics on nonspin manifolds”, *Proc. Amer. Math. Soc.* **149**:5 (2021), 2199–2211. MR Zbl
- [12] S. Dragomir and L. Ornea, *Locally conformal Kähler geometry*, *Progress in Mathematics* **155**, Birkhäuser Boston, Boston, 1998. MR Zbl
- [13] P. Gauduchon, “Hermitian connections and Dirac operators”, *Boll. Un. Mat. Ital. B (7)* **11**:2 (1997), 257–288. MR Zbl
- [14] R. E. Gompf, “Killing the Akbulut–Kirby 4-sphere, with relevance to the Andrews–Curtis and Schoenflies problems”, *Topology* **30**:1 (1991), 97–115. MR Zbl
- [15] M. Gromov and H. B. Lawson, Jr., “Spin and scalar curvature in the presence of a fundamental group, I”, *Ann. of Math. (2)* **111**:2 (1980), 209–230. MR Zbl
- [16] K. Hasegawa, “Complex and Kähler structures on compact solvmanifolds”, pp. 749–767 in *Conference on Symplectic Topology*, vol. 3, 2005. MR Zbl
- [17] T. Holt and W. Zhang, “Harmonic forms on the Kodaira–Thurston manifold”, *Adv. Math.* **400** (2022), art. id. 108277. MR
- [18] M. Inoue, “On surfaces of Class VII_0 ”, *Invent. Math.* **24** (1974), 269–310. MR Zbl
- [19] K. Kodaira, “On the structure of compact complex analytic surfaces, I”, *Amer. J. Math.* **86** (1964), 751–798. MR Zbl
- [20] J. Lin, D. Ruberman, and N. Saveliev, “On the Frøyshov invariant and monopole Lefschetz number”, preprint, 2018. To appear in *J. Differential Geom.* arXiv 1802.07704
- [21] J. Lin, D. Ruberman, and N. Saveliev, “A splitting theorem for the Seiberg–Witten invariant of a homology $S^1 \times S^3$ ”, *Geom. Topol.* **22**:5 (2018), 2865–2942. MR Zbl
- [22] J. Lin, D. Ruberman, and N. Saveliev, “On the monopole Lefschetz number of finite-order diffeomorphisms”, *Geom. Topol.* **25**:7 (2021), 3591–3628. MR Zbl
- [23] P. Lupascu, “The Seiberg–Witten equations on Hermitian surfaces”, *Math. Nachr.* **242** (2002), 132–147. MR Zbl
- [24] T. Mrowka, D. Ruberman, and N. Saveliev, “Seiberg–Witten equations, end-periodic Dirac operators, and a lift of Rohlin’s invariant”, *J. Differential Geom.* **88**:2 (2011), 333–377. MR Zbl

- [25] T. Mrowka, D. Ruberman, and N. Saveliev, “An index theorem for end-periodic operators”, *Compos. Math.* **152**:2 (2016), 399–444. MR Zbl
- [26] C. Okonek and A. Teleman, “Seiberg–Witten invariants for 4-manifolds with $b_+ = 0$ ”, pp. 347–357 in *Complex analysis and algebraic geometry*, de Gruyter, Berlin, 2000. MR Zbl
- [27] C. H. Taubes, “Gauge theory on asymptotically periodic 4-manifolds”, *J. Differential Geom.* **25**:3 (1987), 363–430. MR Zbl
- [28] A. D. Teleman, “Projectively flat surfaces and Bogomolov’s theorem on class VII₀ surfaces”, *Internat. J. Math.* **5**:2 (1994), 253–264. MR Zbl
- [29] W. P. Thurston, “Some simple examples of symplectic manifolds”, *Proc. Amer. Math. Soc.* **55**:2 (1976), 467–468. MR Zbl
- [30] F. Tricerri, “Some examples of locally conformal Kähler manifolds”, *Rend. Sem. Mat. Univ. Politec. Torino* **40**:1 (1982), 81–92. MR Zbl
- [31] C. T. C. Wall, “Geometries and geometric structures in real dimension 4 and complex dimension 2”, pp. 268–292 in *Geometry and Topology* (College Park, Md., 1983/84), Lecture Notes in Math. **1167**, Springer, 1985. MR Zbl
- [32] C. T. C. Wall, “Geometric structures on compact complex analytic surfaces”, *Topology* **25**:2 (1986), 119–153. MR Zbl

Received 23 Jan 2021. Revised 5 May 2021.

DANIEL RUBERMAN: ruberman@brandeis.edu

Department of Mathematics, Brandeis University, Waltham, MA, United States

NIKOLAI SAVELIEV: saveliev@math.miami.edu

Department of Mathematics, University of Miami, Coral Gables, FL, United States

Volume Editors:

John A. Baldwin
Boston College
Boston, MA
United States

Hans U. Boden
McMaster University
Hamilton, ON
Canada

John B. Etnyre
Georgia Institute of Technology
Atlanta, GA
United States

Liam Watson
University of British Columbia
Vancouver, BC
Canada

The cover image is based on an illustration from the article “Khovanov homology and strong inversions”, by Artem Kotelskiy, Liam Watson and Claudius Zibrowius (see p. 232).

The contents of this work are copyrighted by MSP or the respective authors.
All rights reserved.

Electronic copies can be obtained free of charge from <http://msp.org/obs/5> and printed copies can be ordered from MSP (contact@msp.org).

The Open Book Series is a trademark of Mathematical Sciences Publishers.

ISSN: 2329-9061 (print), 2329-907X (electronic)

ISBN: 978-1-935107-11-8 (print), 978-1-935107-10-1 (electronic)

First published 2022.



MATHEMATICAL SCIENCES PUBLISHERS

798 Evans Hall #3840, c/o University of California, Berkeley CA 94720-3840
contact@msp.org

<https://msp.org>

Gauge Theory and Low-Dimensional Topology: Progress and Interaction

This volume is a proceedings of the 2020 BIRS workshop *Interactions of gauge theory with contact and symplectic topology in dimensions 3 and 4*. This was the 6th iteration of a recurring workshop held in Banff. Regrettably, the workshop was not held onsite but was instead an online (Zoom) gathering as a result of the Covid-19 pandemic. However, one benefit of the online format was that the participant list could be expanded beyond the usual strict limit of 42 individuals. It seemed to be also fitting, given the altered circumstances and larger than usual list of participants, to take the opportunity to put together a conference proceedings.

The result is this volume, which features papers showcasing research from participants at the 6th (or earlier) *Interactions* workshops. As the title suggests, the emphasis is on research in gauge theory, contact and symplectic topology, and in low-dimensional topology. The volume contains 16 refereed papers, and it is representative of the many excellent talks and fascinating results presented at the *Interactions* workshops over the years since its inception in 2007.

TABLE OF CONTENTS

| | |
|--|-----|
| Preface — John A. Baldwin, Hans U. Boden, John B. Etnyre and Liam Watson | ix |
| A friendly introduction to the bordered contact invariant — Akram Alishahi, Joan E. Licata, Ina Petkova and Vera Vértesi | 1 |
| Branched covering simply connected 4-manifolds — David Auckly, R. İnanç Baykur, Roger Casals, Sudipta Kolay, Tye Lidman and Daniele Zuddas | 31 |
| Lifting Lagrangian immersions in $\mathbb{C}P^{n-1}$ to Lagrangian cones in \mathbb{C}^n — Scott Baldrige, Ben McCarty and David Vela-Vick | 43 |
| L-space knots are fibered and strongly quasipositive — John A. Baldwin and Steven Sivek | 81 |
| Tangles, relative character varieties, and holonomy perturbed traceless flat moduli spaces — Guillem Cazassus, Chris Herald and Paul Kirk | 95 |
| On naturality of the Ozsváth–Szabó contact invariant — Matthew Hedden and Lev Tovstopyat-Nelip | 123 |
| Dehn surgery and nonseparating two-spheres — Jennifer Hom and Tye Lidman | 145 |
| Broken Lefschetz fibrations, branched coverings, and braided surfaces — Mark C. Hughes | 155 |
| Small exotic 4-manifolds and symplectic Calabi–Yau surfaces via genus-3 pencils — R. İnanç Baykur | 185 |
| Khovanov homology and strong inversions — Artem Kotelskiy, Liam Watson and Claudius Zibrowius | 223 |
| Lecture notes on trisections and cohomology — Peter Lambert-Cole | 245 |
| A remark on quantum Hochschild homology — Robert Lipshitz | 265 |
| On uniqueness of symplectic fillings of links of some surface singularities — Olga Plamenevskaya | 269 |
| On the spectral sets of Inoue surfaces — Daniel Ruberman and Nikolai Saveliev | 285 |
| A note on thickness of knots — András I. Stipsicz and Zoltán Szabó | 299 |
| Morse foliated open books and right-veering monodromies — Vera Vértesi and Joan E. Licata | 309 |