

Motivic Geometry

Introduction to framed correspondences

Marc Hoyois and Nikolai Opdan



$$\begin{array}{ccc}
 X \xleftarrow{j} \circ \xrightarrow{j} \overline{X} \xleftarrow{j} \partial \overline{X} & & 0 \rightarrow K_{2-n}^M(k)/24 \rightarrow \pi_{n+1,n} \mathbf{1} \rightarrow \pi_{n+1,n} \mathbf{kq} \rightarrow 0 \\
 \downarrow f & \downarrow \bar{f} & \downarrow g \\
 \text{Spec}(k) & & \coprod_{\infty} X \simeq g_* j_* f^!(1)
 \end{array}$$

$\sum_{\text{lines}} \text{Tr}_{L/k}(\alpha) = 15\langle 1 \rangle + 12\langle -1 \rangle$

$u_*(\mathcal{X}(X/k)) = \pi_*(e^*(T_{X/k})) \in \mathcal{E}^{0,0}(k)$
 $\text{DM}^{\text{eff}}(k) \doteq \log \text{DM}^{\text{eff}}(k)$

Introduction to framed correspondences

Marc Hoyois and Nikolai Opdan

We give an overview of the theory of framed correspondences in motivic homotopy theory. Motivic spaces with framed transfers are the analogue in motivic homotopy theory of E_∞ -spaces in classical homotopy theory, and in particular they provide an algebraic description of infinite \mathbb{P}^1 -loop spaces. We will discuss the foundations of the theory (following Voevodsky, Garkusha, Panin, Ananyevskiy, and Neshitov), some applications such as the computations of the infinite loop spaces of the motivic sphere and of algebraic cobordism (following Elmanto, Hoyois, Khan, Sosnilo, and Yakerson), and some open problems.

1. Introduction	107
2. Transfers for motivic spectra	111
3. The ∞ -category of framed correspondences	114
4. The recognition principle	115
5. Algebraic cobordism	119
Acknowledgement	125
References	125

1. Introduction

Voevodsky introduced in an unpublished note [13] framed correspondences to get a more computation-friendly model for motivic homotopy theory. In Section 5 we will see how this can be used in practice to get results for algebraic cobordism. First, we introduce framed correspondences as a tool to answer two fundamental questions:

This text is based on a series of three lectures given by Hoyois in August 2020 during the Motivic Geometry program at the Centre for Advanced Study at the Norwegian Academy of Science and Letters in Oslo, Norway.

MSC2020: primary 14F42; secondary 55P47.

Keywords: framed correspondences, motivic homotopy theory, infinite loop space, algebraic cobordisms.

Question 1.1. *What kind of transfers do cohomology theories represented by motivic spectra have?*

Question 1.2. *Is every cohomology theory with these transfers represented by a motivic spectrum?*

These questions will be answered in Sections 3 and 4, respectively.

We begin with an analysis of these problems in classical topology.

Definition 1.3. Let \mathbf{Man} denote the category of smooth manifolds and \mathbf{Spc} the ∞ -category of spaces. A *cohomology theory* on \mathbf{Man} is a functor

$$F : \mathbf{Man}^{\text{op}} \rightarrow \mathbf{Spc}$$

satisfying

- (1) descent with respect to arbitrary open coverings,
- (2) homotopy invariance: $F(M) \xrightarrow{\simeq} F(M \times \mathbb{R})$ for all $M \in \mathbf{Man}$.

We have the following classification.

Theorem 1.4. *The evaluation functor*

$$\begin{aligned} \{\text{cohomology theories on } \mathbf{Man}\} &\xrightarrow{\simeq} \mathbf{Spc}, \\ F &\longmapsto F(*), \end{aligned}$$

is an equivalence.

Proof. The constant sheaf \underline{X} on \mathbf{Man} with fiber a space X is given by

$$M \mapsto \text{Map}(M, X).$$

(This is a nontrivial but well-known computation, which uses the fact that manifolds are sufficiently nice topological spaces, in particular locally contractible.) This computation shows that \underline{X} is homotopy invariant. In particular, $X \mapsto \underline{X}$ is the left adjoint to the given evaluation functor, and since $\underline{X}(*) = X$ it is fully faithful. To show that it is an equivalence of categories it remains to show that evaluation on the point is conservative.

Let f be a morphism between cohomology theories and suppose that it is an isomorphism on a point. We need to show that it is an isomorphism in general. By homotopy invariance, we see that f is an isomorphism on the Euclidean spaces \mathbb{R}^n for all n , and by using the descent property on good covers (covers are copies of \mathbb{R}^n such that all intersections are either empty or isomorphic to \mathbb{R}^m) we can deduce that it is indeed an isomorphism for all manifolds M . \square

In light of Theorem 1.4, cohomology theories can be described by giving a space X . If X happens to be the infinite loop space of a spectrum E , then the associated cohomology theory

$$M \mapsto \text{Map}(M, \Omega^\infty E)$$

acquires some extra structure.

Example 1.5 (Atiyah duality). If M is compact, then $\Sigma_+^\infty M$ has as dual the Thom spectrum M^{-T_M} , where T_M is the tangent bundle of M . This implies that for every morphism $f : M \rightarrow N$ between compact manifolds there is an induced morphism of Thom spectra $M^{-T_M} \leftarrow N^{-T_N}$. Mapping these spectra into another spectra E we get a covariant pushforward map

$$\text{Map}(M^{-T_M}, E) \rightarrow \text{Map}(N^{-T_N}, E).$$

If T_M and T_N are oriented with respect to E (e.g., if E is complex oriented and M and N are complex manifolds), then the pushforward involves ordinary shifts.

One can more generally define a cohomological pushforward along any proper morphism $f : M \rightarrow N$ between (not necessarily compact) smooth manifolds. This is a consequence of the “formalism of six operations”, which is a vast generalization of Poincaré/Atiyah duality introduced by Grothendieck. For any morphism $f : M \rightarrow N$ in **Man**, there is the usual pullback/pushforward adjunction

$$\text{Shv}(M, \mathbf{Spt}) \begin{array}{c} \xleftarrow{f^*} \\ \xrightarrow{f_*} \end{array} \text{Shv}(N, \mathbf{Spt}),$$

as well as the “exceptional” adjunction

$$\text{Shv}(M, \mathbf{Spt}) \begin{array}{c} \xrightarrow{f_!} \\ \xleftarrow{f^!} \end{array} \text{Shv}(N, \mathbf{Spt}).$$

Here, $f_!$ is the pushforward with compact support, which is essentially determined by the following two properties:

- If f is proper, then $f_! = f_*$.
- If f is étale (i.e., a local homeomorphism), then $f^! = f^*$.

Defining the virtual tangent bundle of f by

$$T_f := T_M - f^*T_N,$$

there exists then a canonical natural transformation

$$\mathfrak{p}_f : \Sigma^{T_f} f^* \rightarrow f^!,$$

where Σ^{T_f} denotes suspension by the virtual tangent bundle, i.e., smashing with its Thom spectrum. This functor is an automorphism of $\text{Shv}(M, \mathbf{Spt})$, and locally on M , it is the ordinary suspension by the rank of T_f . We do not explain here the construction of \mathfrak{p}_f . Suffice it to say, because any morphism factors as the composition of a submersion and a closed immersion and because of the tubular neighborhood theorem, the transformation \mathfrak{p}_f is essentially determined by the cases where f is a submersion (in which case it is an *equivalence* $\Sigma^{T_f} f^* \simeq f^!$), and where f is the zero section of a vector bundle. We will explain later the algebro-geometric analogue of \mathfrak{p}_f in more details.

If E_M denotes the constant sheaf on M with value E , we obtain morphisms

$$\Gamma(M, \Sigma^{T_f} E_M) \simeq \Gamma(M, \Sigma^{T_f} f^* E_N) \xrightarrow{\mathfrak{p}_f} \Gamma(M, f^! E_N) \simeq \text{Map}(\mathbb{S}_M, f^! E_N).$$

If f is proper we furthermore have morphisms

$$\text{Map}(\mathbb{S}_M, f^! E_N) \simeq \text{Map}(f_* f^*(\mathbb{S}_N), E_N) \rightarrow \text{Map}(\mathbb{S}_N, E_N) = \Gamma(N, E_N).$$

Composing these morphisms gives us a pushforward map

$$\Gamma(M, \Sigma^{T_f} E_M) \rightarrow \Gamma(N, E_N)$$

also known as the *transfer* along f . This provides an answer to Question 1.1 in the classical topological context.

For Question 1.2, the question becomes whether one can recover the spectrum structure on E from the infinite delooping $\Omega^\infty E$ by using these transfers. In particular, if $f : M \rightarrow N$ is a finite étale map (i.e., a finite covering map), then $T_f = 0$, in which case there is a canonical transfer

$$\text{Map}(M, \Omega^\infty E) \rightarrow \text{Map}(N, \Omega^\infty E).$$


By taking N to be a point and M to be two distinct points, the transfer provides an addition map

$$\Omega^\infty E \times \Omega^\infty E \xrightarrow{+} \Omega^\infty E.$$


We need to encode these transfers in a suitably coherent manner to recover the commutativity of $+$ up to coherent homotopy. Historically, this has been difficult because one lacked the tools to precisely express the required coherence. To achieve this we introduce the 2-category of correspondences:

Definition 1.6. The *category of finite étale correspondences* in \mathbf{Man} , denoted by $\mathbf{Cor}^{\text{fét}}(\mathbf{Man})$, is the 2-category where objects are smooth manifolds, and morphisms

from M to P are spans



such that the map $N \rightarrow M$ is finite étale. These spans are known as *correspondences*. Composition of two correspondences $(M \leftarrow N \rightarrow P)$ and $(P \leftarrow Q \rightarrow R)$ is defined as the pullback in the diagram



The pullback exists in \mathbf{Man} because the map $Q \rightarrow P$ is finite étale.

The following theorem then provides an answer to Question 1.2.

Theorem 1.7. *There exists a functor*

$$\begin{aligned}
 \mathbf{Spt} &\rightarrow \{\text{cohomology theories on } \mathbf{Cor}^{\text{fét}}(\mathbf{Man})\}, \\
 E &\mapsto \text{Map}(-, \Omega^\infty E),
 \end{aligned}$$

which restricts to an equivalence between connective spectra $\mathbf{Spt}_{\geq 0}$ and grouplike cohomology theories on $\mathbf{Cor}^{\text{fét}}(\mathbf{Man})$.

Such cohomology theories are automatically evaluated in the ∞ -category of E_∞ -spaces, hence the ‘‘grouplike’’ condition means that the addition structure

$$\Omega^\infty E \times \Omega^\infty E \rightarrow \Omega^\infty E$$

should be a group, that is, elements should have inverses up to homotopy. We refer to [2, Appendix C] for a proof of Theorem 1.7 given Theorem 1.4.

2. Transfers for motivic spectra

We will now focus on Question 1.1, i.e., transfers for motivic spectra.

Let $\mathcal{SH}(S)$ denote the ∞ -category of motivic spectra over S . There is a six-functor formalism on the system of categories $S \mapsto \mathcal{SH}(S)$, that is, functors

$$f^*, f_*, f^!, f_!, \underline{\text{Hom}}, \otimes,$$

with adjunctions $f^* \dashv f_*$ and $f_! \dashv f^!$, and equivalences $f^! \simeq f^*$ if f is étale and $f_! \simeq f_*$ if f is proper.

For the case of smooth manifolds, we saw in the previous section that there are transfers for any proper morphism. A key difference in the algebraic-geometric context is that we are now also willing to consider schemes which are not smooth. This changes the picture since morphisms of smooth manifolds are very far from being arbitrary morphisms, and in particular they are always of local complete intersections. The condition of local complete intersection turns out to be the most general condition for transfers to exist in this setting.

Definition 2.1. A morphism of schemes $f : X \rightarrow S$ is a *local complete intersection* (abbreviated “lci”) if locally on X it factors as

$$\begin{array}{ccc} X & \xrightarrow{i} & V \\ & \searrow f & \downarrow p \\ & & S, \end{array}$$

where i is a closed immersion which locally is cut out by a regular sequence (i.e., a *regular* closed immersion) and p is a smooth morphism.

For any local complete intersection morphism $f : X \rightarrow S$ with a global factorization as above, we can define the *virtual tangent bundle* of f by

$$T_f := i^*T_p - N_i \in K(X).$$

Here, T_p is the relative tangent bundle of p , N_i is the normal bundle of i , and $K(X)$ denotes the K -theory space of the scheme X . One can show that T_f does not depend on the factorization of f (it is in fact the image in $K(X)$ of the tangent complex of f , which is defined even if no factorization exists).

We will now sketch the fundamental construction of Déglise, Jin, and Khan [3], which is the source of the transfers in stable motivic homotopy theory. We recall that any element $\xi \in K(X)$ has an associated Thom spectrum in $\mathcal{SH}(X)$, and smashing with it defines a self-equivalence

$$\Sigma^\xi : \mathcal{SH}(X) \xrightarrow{\simeq} \mathcal{SH}(X).$$

Construction 2.2 [3]. *For an lci morphism $f : X \rightarrow S$ with a global factorization, we construct a canonical transformation*

$$\mathfrak{p}_f : \Sigma^{T_f} f^* \rightarrow f^!.$$

Sketch of proof. Because of the global lci factorization, it suffices to consider the case of smooth morphisms and regular closed immersions.

If f is smooth we let \mathfrak{p}_f be the well-known purity equivalence

$$\Sigma^{T_f} f^* \simeq f^!$$

due to Voevodsky and Ayoub.

If $f = i : X \hookrightarrow V$ is a regular closed immersion we consider the deformation of V to the normal cone, which is a scheme $D_X V$ that fits in a diagram

$$\begin{array}{ccccc}
 N_i & \xleftarrow{t} & D_X V & \xleftarrow{u} & \mathbb{G}_m \times V \\
 \swarrow \pi & \downarrow & \downarrow & & \parallel \\
 X & & & & \\
 \searrow i & \downarrow & \downarrow & & \\
 V & \xleftarrow{0} & \mathbb{A}_V^1 & \xleftarrow{\circ} & \mathbb{G}_m \times V \\
 & \parallel & \downarrow & \swarrow q & \\
 & & V, & &
 \end{array}$$

where $0 : V \hookrightarrow \mathbb{A}_V^1$ is the zero section, $\pi : N_i \rightarrow X$ is the normal bundle of i , $\mathbb{G}_m \times V \hookrightarrow D_X V$ and $\mathbb{G}_m \times V \hookrightarrow \mathbb{A}_V^1$ are the open complements, and both squares are Cartesian.¹ Let p denote the composition of the maps $D_X V \rightarrow \mathbb{A}_V^1 \rightarrow V$.

The top row gives rise to a localization triangle

$$t_* t^! \rightarrow \text{id}_{\mathcal{SH}(D_X V)} \rightarrow u_* u^! \xrightarrow{\partial} t_* t^![1],$$

and in $\mathcal{SH}(\mathbb{G}_m)$ there is a canonical (universal unit) map $S^0 \rightarrow \mathbb{G}_m$. Suspending this map with S^1 we get a new map

$$S^1 \xrightarrow{(*)} S^1 \wedge \mathbb{G}_m = T,$$

where $T = S^1 \wedge \mathbb{G}_m$ is the Thom space of the trivial line bundle. We want to define a natural transformation

$$p_i : i^* \rightarrow \Sigma^{N_i} i^!,$$

or equivalently, by adjunction, a map

$$\text{id} \rightarrow i_* \Sigma^{N_i} i^!.$$

This we obtain from the composition

$$\text{id}_{\mathcal{SH}(V)}[1] \xrightarrow{(*)} q_* \Sigma_T q^* \simeq q_* q^! \simeq p_* u_* u^! p^! \xrightarrow{\partial} p_* t_* t^! p^![1] \simeq i_* \pi_* \pi^! i^![1],$$

and using the purity equivalence $p_\pi : \pi^* \Sigma^{N_i} \simeq \pi^!$ and the fact that π^* is a fully faithful functor, we further get

$$i_* \pi_* \pi^! i^! \simeq i_* \Sigma^{N_i} i^!.$$

Composing these maps gives the desired natural transformation

$$\text{id}_{\mathcal{SH}(V)} \rightarrow i_* \Sigma^{N_i} i^!.$$

¹For an arbitrary closed immersion i , the fiber of $D_X V$ over 0 is the normal cone of i , which agrees with the normal bundle in case of a regular immersion.

It remains to prove that this procedure is independent of the factorization, composing various f 's, etc. We refer the curious reader to [3] for a thorough treatment of these facts. \square

Definition 2.3. Let $E \in \mathcal{SH}(S)$ be a motivic spectrum over S . For a morphism $f : X \rightarrow S$ and an element $\xi \in K(X)$ we define

$$E(X, \xi) := \text{Map}_{\mathcal{SH}(X)}(\mathbb{1}_X, \Sigma^\xi f^* E) \in \mathbf{Spc}.$$

Example 2.4. If $\xi = n$, then $\Sigma^\xi \cong \Sigma^{2n, n}$. Thus

$$\pi_i E(X, \xi) = E^{2n-i, n}(X).$$

Example 2.5. If $E = \text{KGL}$, then $E(X, \xi) \cong \text{KH}(X)$, where KH denotes the homotopy K -theory of X . Note that this does not depend on ξ .

Example 2.6. If $E = H\mathbb{Z}$ and X is smooth over a field or a discrete valuation ring, then $E(X, \xi) \cong z^r(X, \bullet)$ where $r = \text{rank}(\xi)$. Here $z^r(X, \bullet)$ denotes Bloch's cycle complex.

Theorem 2.7. *If $f : X \rightarrow S$ is proper and lci, then the natural transformation $\mathfrak{p}_f : \Sigma^{T_f} f^* \rightarrow f^!$ induces a transfer map*

$$E(X, T_f + f^* \xi) \rightarrow E(S, \xi),$$

where $\xi \in K(S)$.

Remark 2.8. More generally, if $f : X \rightarrow S$ is proper and X^{der} is a derived structure on X which is lci/quasismooth over S , a theorem by Adeel A. Khan [10] gives a transfer map

$$E(X, T_f^{\text{der}}) \rightarrow E(S).$$

3. The ∞ -category of framed correspondences

Following the approach to Question 1.1 in classical topology, we are interested in those transfers that do not shift the degree, i.e., that induce a covariant functoriality on the “unshifted” cohomology theory. For this to work we saw in the previous section that we need the virtual tangent bundle of a map f to vanish. We are thus led to the following definition:

Definition 3.1. Suppose $f : X \rightarrow S$ is a lci morphism. A (stable 0-dimensional) *framing* of f is a trivialization of the virtual tangent bundle T_f , that is, a path in $K(X)$ between T_f and 0.

If f is a framed proper lci morphism, we see by Theorem 2.7 that it induces a transfer map in cohomology that does not shift the degree. This provides an answer to Question 1.1.

Definition 4.1. Let $\mathcal{H}(S)$ denote the ∞ -category of \mathbb{A}^1 -invariant Nisnevich sheaves on Sm_S , and let $\mathcal{H}^{\mathrm{fr}}(S)$ be the ∞ -category of \mathbb{A}^1 -invariant Nisnevich sheaves on $\mathbf{Cor}^{\mathrm{fr}}(\mathrm{Sm}_S)$ (i.e., presheaves on $\mathbf{Cor}^{\mathrm{fr}}(\mathrm{Sm}_S)$ whose restriction to Sm_S are \mathbb{A}^1 -invariant Nisnevich sheaves). We define $\mathcal{SH}(S)$ and $\mathcal{SH}^{\mathrm{fr}}(S)$ as the ∞ -categories of T -spectra in $\mathcal{H}(S)$ and $\mathcal{H}^{\mathrm{fr}}(S)$, respectively.

The functor $\mathrm{Sm}_S \rightarrow \mathbf{Cor}^{\mathrm{fr}}(\mathrm{Sm}_S)$ induces adjoint functor pairs

$$\begin{array}{ccc} \mathcal{H}(S) & \begin{array}{c} \xrightarrow{\text{free}} \\ \xleftarrow{\text{forget}} \end{array} & \mathcal{H}^{\mathrm{fr}}(S) \\ \Sigma_+^\infty \updownarrow \Omega^\infty & & \Sigma_{\mathrm{fr}}^\infty \updownarrow \Omega_{\mathrm{fr}}^\infty \\ \mathcal{SH}(S) & \begin{array}{c} \xrightarrow{\text{free}} \\ \xleftarrow{\text{forget}} \end{array} & \mathcal{SH}^{\mathrm{fr}}(S). \end{array}$$

Theorem 4.2 (reconstruction theorem, [7, Theorem 16]). *For every scheme S , the canonical functor*

$$\mathcal{SH}(S) \xrightarrow{\sim} \mathcal{SH}^{\mathrm{fr}}(S)$$

is an equivalence of ∞ -categories.

Thus every motivic spectrum has a unique structure of framed transfers.

This result requires new ideas since the analogue statement is false for Voevodsky's correspondences because it would yield an equivalence $\mathcal{SH}^{\mathrm{fr}}(S) \simeq \mathbf{DM}(S)$.

We provide a brief sketch of the main ideas involved in the proof.

Sketch of proof. The key idea is to apply Voevodsky's lemma: that is, if X is a scheme, and $U \subset X$ is an open subscheme, we can identify $L_{\mathrm{Nis}}(X/U)(Y)$ with the set of pairs (Z, ϕ) such that $Z \subset Y$ is a closed subset and $\phi : Y_Z^h \rightarrow X$ is a map such that $\phi^{-1}(X \setminus U) = Z$.

Then consider, for $X, Y \in \mathrm{Sm}_S$, the mapping space

$$\mathrm{Map}((\mathbb{P}^1)^\wedge \wedge X_+, L_{\mathrm{Nis}}(\mathbb{A}^n / \mathbb{A}^n - 0) \wedge Y_+) \cong (\Omega_{\mathbb{P}^1}^n L_{\mathrm{Nis}} \Sigma_T^n Y_+)(X).$$

Applying Voevodsky's lemma we find that this equals the set of pairs (Z, ϕ) such that

- (1) $Z \subset (\mathbb{P}_X^1)^n$ is a closed subscheme,
- (2) $Z \cap \partial(\mathbb{P}_X^1)^n = \emptyset$,
- (3) $\phi : ((\mathbb{P}_X^1)^n)_Z^h \rightarrow \mathbb{A}_Y^1$ such that $\phi^{-1}(0) = Z$.

Thus Z is cut out by n equations with codimension n , and, in particular, it is framed. This shows that there is a forgetful map to $\mathbf{Cor}_S^{\mathrm{fr}}(X, Y)$. We conclude by [5, Corollary 2.3.27] which says that the forgetful map

$$\mathrm{colim}_{n \rightarrow \infty} \Omega_{\mathbb{P}^1}^n L_{\mathrm{Nis}} \Sigma_T^n Y_+ \rightarrow \mathbf{Cor}_S^{\mathrm{fr}}(-, Y)$$

is a motivic equivalence if $Y \in \mathrm{Sm}_S$. □

Theorem 4.3 (cancellation theorem, [5, Proposition 5.2.7]). *Let k be a perfect field. The functors*

$$S^1 \wedge (-), \mathbb{G}_m \wedge (-) : \mathcal{H}^{\text{fr}}(k)^{\text{gp}} \rightarrow \mathcal{H}^{\text{fr}}(k)^{\text{gp}}$$

are fully faithful, where $\mathcal{H}^{\text{fr}}(k)^{\text{gp}}$ is the full subcategory of $\mathcal{H}^{\text{fr}}(k)$ consisting of grouplike objects.

Let L_{Zar} denote the Zariski sheafification functor and $L_{\mathbb{A}^1}$ the functor defined by

$$(L_{\mathbb{A}^1} F)(U) = |F(U \times \mathbb{A}^{\bullet})|.$$

Theorem 4.4 (strict \mathbb{A}^1 -invariance theorem, [5, Theorem 3.4.11]). *Let k be a perfect field. If $F \in \mathcal{P}_{\Sigma}(\mathbf{Cor}^{\text{fr}}(\text{Sm}_k))^{\text{gp}}$, then $L_{\text{Zar}}L_{\mathbb{A}^1}F$ is an \mathbb{A}^1 -invariant Nisnevich sheaf.*

Theorems 4.3 and 4.4 are analogues of Voevodsky’s results for finite correspondences ([14] and [11, §3.2]) and the proofs are similar. For Theorem 4.3, Bachmann has a paper [1] where he proves a cancellation theorem for finite flat correspondences, which applies to these framed correspondences as well.

The next result is an analogue of Theorem 1.7 for framed motivic spectra and thus answers Question 1.2.

Corollary 4.5 (motivic recognition principle, [5, Theorem 3.5.16]). *Let k be a perfect field. There is an equivalence*

$$\mathcal{H}^{\text{fr}}(k)^{\text{gp}} \simeq \mathcal{SH}(k)^{\text{veff}},$$

where $\mathcal{SH}(k)^{\text{veff}} \subset \mathcal{SH}(k)$ denotes the full subcategory generated under colimits by suspension spectra $\Sigma_+^{\infty} X$.

Proof. Theorem 4.3 implies that the infinite suspension functor

$$\Sigma_{\text{fr}}^{\infty} : \mathcal{H}^{\text{fr}}(k)^{\text{gp}} \rightarrow \mathcal{SH}^{\text{fr}}(k)$$

is fully faithful. Using Theorem 4.2 we identify its essential image as $\mathcal{SH}(k)^{\text{veff}}$. \square

The main application of Theorem 4.2 is that the framed suspension functor $\Sigma_{\text{fr}}^{\infty}$ becomes a “machine” for producing motivic spectra: Let $F \in \mathcal{P}_{\Sigma}(\mathbf{Cor}^{\text{fr}}(\text{Sm}_S))$ be a presheaf with framed transfers. Then

$$\Sigma_{\text{fr}}^{\infty} F \in \mathcal{SH}^{\text{fr}}(S),$$

which by Theorem 4.2 can be identified with an object of $\mathcal{SH}(S)$. This allows one to build motivic spectra from presheaves with framed transfers.

In the special case of $S = \text{Spec } k$, where k is a perfect field, we can use Corollary 4.5 to compute the infinite loop space of such a spectrum by using the equivalence

$$\Omega_{\text{fr}}^{\infty} \Sigma_{\text{fr}}^{\infty} F \simeq L_{\text{Zar}}L_{\mathbb{A}^1}F^{\text{gp}}.$$

We have the following consequences of this in action:

Example 4.6. Applying the infinite framed suspension functor $\Sigma_{\text{fr}}^\infty$ to the right side of this diagram one obtains the left side:

$$\begin{array}{ccc}
 \mathbb{1} & & \text{FSyn}^{\text{fr}} \\
 \downarrow & & \downarrow \\
 \text{MSL} & \longrightarrow & \text{MGL} \\
 \downarrow & & \downarrow \\
 \text{ko} & \longrightarrow & \text{kgl} \\
 \parallel & & \parallel \\
 \text{ko} & \longrightarrow & \text{kgl} \\
 \downarrow & & \downarrow \\
 H\tilde{\mathbb{Z}} & \longrightarrow & H\mathbb{Z}
 \end{array}
 \qquad
 \begin{array}{ccc}
 \text{FSyn}^{\text{fr}} & & \text{FSyn} \\
 \downarrow & & \downarrow \\
 \text{FSyn}^{\text{or}} & \longrightarrow & \text{FSyn} \\
 \downarrow & & \downarrow \\
 \text{FGor}^{\text{or}} & \longrightarrow & \text{FFlat} \\
 \downarrow & & \downarrow \\
 \text{Vect}^{\text{sym}} & \longrightarrow & \text{Vect} \\
 \downarrow & & \downarrow \\
 \underline{\text{GW}} & \longrightarrow & \underline{\mathbb{Z}},
 \end{array}$$

In more details:

- FSyn^{fr} is the stack of framed finite syntomic schemes, which is the unit object in $\mathcal{P}_\Sigma(\mathbf{Cor}^{\text{fr}}(\text{Sm}_k))$. The claim that $\Sigma_{\text{fr}}^\infty \text{FSyn}^{\text{fr}} \simeq \mathbb{1}$ is therefore tautological.
- FSyn is the stack of finite syntomic schemes and FSyn^{or} is the stack of finite syntomic schemes with an orientation (i.e., a trivialization of the dualizing sheaf). That these give MGL and MSL over a general base is proved in [4, Theorems 3.4.1, 3.4.3].
- Vect is the stack of vector bundles and FFlat is the stack of finite flat schemes. That both give the effective K -theory spectrum kgl over a field is proved in [8, Corollary 5.2, Theorem 5.4].
- Vect^{sym} is the stack of vector bundles with a nondegenerate symmetric bilinear form, and FGor^{or} is the stack of oriented finite Gorenstein (dualizing sheaf is a line bundle) schemes. That both give the effective hermitian K -theory spectrum ko over a field of characteristic not 2 is proved in [9, Proposition 7.7, Theorem 7.12].
- $\underline{\mathbb{Z}}$ is the constant sheaf with fiber \mathbb{Z} . It is proved in [7, Theorem 21] that $\Sigma_{\text{fr}}^\infty \underline{\mathbb{Z}} \simeq H\mathbb{Z}$ over any Dedekind domain, where $H\mathbb{Z}$ is the motivic spectrum representing Bloch–Levine motivic cohomology.
- Finally, $\underline{\text{GW}}$ is the sheaf of unramified Grothendieck–Witt rings, that is, the Zariski sheafification of the usual Grothendieck–Witt groups. It is proved in [9, Theorem 8.3] that $\Sigma_{\text{fr}}^\infty \underline{\text{GW}} \simeq H\tilde{\mathbb{Z}}$ over any Dedekind domain in which 2 is invertible.

Here is one nice application of this picture: all morphisms on the right-hand side are easily seen to be E_∞ -ring maps, hence so are all morphisms on the left-hand side.

5. Algebraic cobordism

The topic for this section is algebraic cobordism, following [4]. Specifically, we will apply the above theory to provide a framed model for algebraic cobordism which allows for more friendly computations.

In particular the question about the convergence of the divisibility spectral sequence for algebraic cobordism would benefit greatly from knowing that elements of algebraic cobordism can be represented by some kind of geometric objects.

-V. Voevodsky, 2000, My view of the current state of motivic homotopy theory [12, p. 11].

We begin by considering the moduli stack PQSm of proper quasismooth derived schemes. It is natural to study this stack in connection with algebraic cobordism, because these are precisely the schemes that give rise to algebraic cobordism classes. Namely, a proper quasismooth derived scheme of virtual dimension $-n$ has a cobordism class in $\text{MGL}^{2n,n}$ by, for instance, using the transfers from Section 2. The \mathbb{A}^1 -localization

$$L_{\mathbb{A}^1}\text{PQSm} := |\text{PQSm}(- \times \mathbb{A}^\bullet)|$$

can be described informally as follows. Points in $L_{\mathbb{A}^1}\text{PQSm}$ are schemes, and a path between two points is a scheme over \mathbb{A}^1 , which may be viewed as a cobordism from the fiber over 0 to the fiber over 1 (see Figure 1). A path between two paths is then a cobordism of cobordisms, etc. The analogous construction in topology, with \mathbb{R}^1 in place of \mathbb{A}^1 , is one way to define the cobordism spaces that appear, for example, in the cobordism hypothesis. One may thus interpret the functor $L_{\mathbb{A}^1}$ as forming the cobordism spaces of moduli stacks of schemes.

Recall the forgetful functor

$$\mathbf{Cor}^{\text{fr}}(\text{Sm}_S) \rightarrow \mathbf{Cor}^{\text{fsyn}}(\text{Sm}_S)$$




Figure 1. A cobordism from two circles to one.

from the ∞ -category of framed correspondences of smooth schemes to the 2-category of finite syntomic correspondences of smooth schemes, given by forgetting the framing and using that the morphism in the left span is finite syntomic.

Let FQSm^n denote the moduli stack of finite quasismooth derived S -schemes of relative dimension $-n$. This is obviously a presheaf on $\mathbf{Cor}^{\mathrm{fsyn}}(\mathrm{Sms}_S)$, and hence also on $\mathbf{Cor}^{\mathrm{fr}}(\mathrm{Sms}_S)$, using the above forgetful functor.

Theorem 5.1. *For every scheme S and every integer $n \geq 0$ there is an equivalence*

$$\Sigma_T^n \mathrm{MGL} \simeq \Sigma_{\mathrm{fr}}^\infty \mathrm{FQSm}^n$$

in $\mathcal{SH}(S) \simeq \mathcal{SH}^{\mathrm{fr}}(S)$.

The main ingredient of the proof is to understand the framed models for Thom spectra of virtual vector bundles coming from Theorem 5.10. We therefore suspend the proof of this theorem until we have established a proof of the result.

Remark 5.2. There is an equivalence

$$\mathrm{FQSm}^0 \cong \mathrm{FSyn}.$$

The next theorem is a specialization of Theorem 5.1 to the case of perfect fields.

Theorem 5.3. *If k is a perfect field and $n \geq 0$, then*

$$\Omega_T^\infty \Sigma_T^n \mathrm{MGL} \simeq L_{\mathrm{Zar}} L_{\mathbb{A}^1} (\mathrm{FQSm}^n)^{\mathrm{gp}}.$$

In particular,

$$\Omega_T^\infty \mathrm{MGL} \simeq L_{\mathrm{Zar}} L_{\mathbb{A}^1} \mathrm{FSyn}^{\mathrm{gp}}.$$

If $n \geq 1$, then

$$\Omega^\infty \Sigma_T^n \mathrm{MGL} \simeq L_{\mathrm{Nis}} L_{\mathbb{A}^1} \mathrm{FQSm}^n.$$

Remark 5.4. The last point of Theorem 5.3 means that one can trade the group completion by instead taking Nisnevich sheafification. More precisely, the Nisnevich sheaf $L_{\mathrm{Nis}} L_{\mathbb{A}^1} \mathrm{FQSm}^n$ is connected, hence is already grouplike, so we do not need to take its group completion.

A posteriori, we know by Morel's connectivity theorem that $\Omega^\infty \Sigma_T^n \mathrm{MGL}$ is $(n-1)$ -connected.

Remark 5.5. We state Theorems 5.1 and 5.3 for MGL, but there are analogous statements for all Thom spectra that one would like to think about, for example, MSL, MSp, \mathbb{S} , etc.

Theorem 5.6. *For every scheme S there is an equivalence of symmetric monoidal ∞ -categories*

$$\mathrm{Mod}_{\mathrm{MGL}}(\mathcal{SH}(S)) \simeq \mathcal{SH}^{\mathrm{fsyn}}(S),$$

where $\mathcal{SH}^{\mathrm{fsyn}}(S)$ is analogous to $\mathcal{SH}^{\mathrm{fr}}(S)$ using $\mathbf{Cor}^{\mathrm{fsyn}}(\mathrm{Sms}_S)$ in lieu of $\mathbf{Cor}^{\mathrm{fr}}(\mathrm{Sms}_S)$.

Theorem 5.7. *If k is a perfect field, then*

$$\mathrm{Mod}_{\mathrm{MGL}}(\mathcal{SH}(S)^{\mathrm{veff}}) \simeq \mathcal{H}^{\mathrm{fsyn}}(k)^{\mathrm{gp}}.$$

This theorem is a combination of Theorem 5.6 with a cancellation theorem in $\mathcal{H}^{\mathrm{fsyn}}(k)^{\mathrm{gp}}$.

Framed models for effective Thom spectra.

Definition 5.8. Consider a smooth S -scheme X and $\xi \in K(X)_{\geq 0}$. We define a framed presheaf

$$(X, \xi)^{\mathrm{fr}} \in \mathcal{P}_{\Sigma}(\mathbf{Cor}^{\mathrm{fr}}(\mathrm{Sm}_S))$$

by sending a scheme U to the space of spans $(U \xleftarrow{f} Z \xrightarrow{g} X)$ where f is finite and quasismooth together with $\alpha : T_f \simeq -g^*(\xi) \in K(Z)$.

This definition is not enough to define a presheaf on $\mathbf{Cor}^{\mathrm{fr}}(\mathrm{Sm}_S)$; there is some work involved in making this precise which can be found in [4, Appendix B].

Remark 5.9. If $\mathrm{rank}(\xi) = 0$, then Z is underived. For example,

$$(X, 0)^{\mathrm{fr}} = \mathbf{Cor}_S^{\mathrm{fr}}(-, X).$$

These presheaves turn out to be framed models for effective Thom spectra.

Theorem 5.10 [4, Theorem 3.2.1]. *There is an equivalence*

$$\mathrm{Th}_X(\xi) \simeq \Sigma_{\mathrm{fr}}^{\infty}(X, \xi)^{\mathrm{fr}}$$

in $\mathcal{SH}(S) \simeq \mathcal{SH}^{\mathrm{fr}}(S)$.

The proof consists of two parts: first constructing a map between these two objects, and then showing that it is an equivalence.

Proof. Suppose that ξ is a vector bundle V on X , and let V^{\times} denote the complement of the zero section inside V . We define a map as follows:

Consider the diagram

$$\begin{array}{ccc} \emptyset & \xrightarrow{\quad} & X \\ \downarrow & & \swarrow z \\ V^{\times} & \hookrightarrow & V \end{array} \quad \begin{array}{c} \\ \\ \\ \parallel \\ \\ \end{array} \begin{array}{c} \\ \\ \\ X \\ \\ \end{array}$$

where z is the zero section. The canonical equivalence $T_z \simeq -V$ defines a map

$$V \rightarrow (X, V)^{\mathrm{fr}},$$

and we get a diagram

$$\begin{array}{ccc}
 (V^\times)^{\text{fr}} & & \\
 \downarrow & \searrow 0 & \\
 V^{\text{fr}} & \longrightarrow & (X, V)^{\text{fr}} \\
 \downarrow & & \nearrow \text{---} \\
 \text{Th}_X(V)^{\text{fr}}, & &
 \end{array} \tag{5-1}$$

where the dotted arrow is the comparison map we were looking for.

One can show that the map is a motivic equivalence by reducing to the case where the base is a field and using a moving lemma from [6].

The general case for virtual tangent bundles is now mostly formal. The main point is to use the identification $K(-) \simeq L_{\text{Zar}}\text{Vect}(-)^{\text{gp}}$. The Thom spectrum functor $\text{Th}_X : \text{Vect}(X) \rightarrow \mathcal{SH}(X)$ takes the direct sum of vector bundles to the tensor product of spectra and lands in the Picard ∞ -groupoid $\text{Pic}(\mathcal{SH}(X))$, and thus factors uniquely through the group completion of $\text{Vect}(X)$.

We can then formally extend (5-1) to a map

$$\text{Th}_X(\xi)^{\text{fr}} \rightarrow (X, \xi)^{\text{fr}},$$

which is natural in the virtual vector bundle ξ over X .

To see that this map is an equivalence we use that virtual vector bundles of rank ≥ 0 are actual vector bundles Zariski locally. □

We state a result about K -theory that is needed in the proof of Theorem 5.1.

Proposition 5.11 (Bhatt–Lurie, [4, A.0.6]). *Let R be a commutative ring. Then the K -theory functor*

$$K : \text{CAlg}_R^{\text{der}} \rightarrow \mathbf{Spc}$$

is left-Kan extended from $\text{CAlg}_R^{\text{Sm}}$.

Proof. The formula for left-Kan extension is

$$\text{LKE}(K \mid \text{CAlg}_R^{\text{Sm}})(T) = \text{colim}_{S \rightarrow T} K(S),$$

where $S \in \text{CAlg}_R^{\text{Sm}}$. The index category $\{S \rightarrow T\}$ has finite coproducts given by tensor products, which make it sifted.

By the identity

$$K = \text{Vect}^{\text{gp}}$$

and the fact that the group completion commutes with sifted colimits, we are reduced to show that $\text{Vect}(-)$ is left-Kan extended from $\text{CAlg}_R^{\text{Sm}}$. Using that

$\text{Vect} = \text{colim}_n \text{Vect}_{\leq n}$ we reduce to showing that each $\text{Vect}_{\leq n}$ is left-Kan extended. One way of showing this is by taking the Jouanolou device

$$U_n \rightarrow \text{Gr}_n,$$

which is an affine bundle whose total space is affine. Then there is a map

$$\coprod_{k \leq n} U_k \xrightarrow{f} \text{Vect}_{\leq n}.$$

The map f is a sectionwise surjection because we are dealing with affine schemes, so that vector bundles are generated by their global sections. Hence it may be lifted to a point in the Grassmannian, which further can be lifted to the Jouanolou device since it is an affine bundle over an affine scheme.

We therefore have

$$\text{Vect}_{\leq n} \simeq \text{colim } \check{C}_\bullet(f),$$

where $\check{C}_\bullet(f) = \Delta^{\text{op}} \rightarrow \text{Ind}(\text{smooth affine schemes})$. The same remains true of all the self-intersections of $\coprod_{k \leq n} U_k$, since $\text{Vect}_{\leq n}$ is a stack with smooth affine diagonals. Thus the iterated fiber product of $\coprod_{k \leq n} U_k$ is still a colimit of smooth and affine schemes. Combining the colimits we thereby find that the presheaf Vect is a colimit of smooth affine R -schemes, which is precisely what it means to be left-Kan extended from $\text{CAIg}_R^{\text{Sm}}$. \square

Now for the proof of Theorem 5.1.

Proof of Theorem 5.1. For $n = 0$ we can write

$$\text{MGL} \simeq \text{colim}_{X, \xi} \text{Th}_X(\xi),$$

by [2, Section 16], where the colimit is taken over $X \in \text{Sm}_S$ and $\xi \in K(X)$ with $\text{rank}(\xi) = 0$. Hence

$$\text{MGL} \simeq \Sigma_{\text{fr}}^\infty \text{colim}_{X, \xi} (X, \xi)^{\text{fr}}$$

by Theorem 4.2. Further, there is a forgetful map

$$\begin{array}{c} \text{colim}_{X, \xi} (X, \xi)^{\text{fr}} \\ \downarrow \\ \text{FSyn} \end{array}$$

by using that an object in $\text{colim}_{X, \xi} (X, \xi)^{\text{fr}}$ consists of a span $U \xleftarrow{f} Z \xrightarrow{g} X$ which we forgetfully map to $U \xleftarrow{f} Z$. We claim that this map is an equivalence.

By taking the fiber one obtains a Cartesian square

$$\begin{array}{ccc}
\operatorname{colim}_{X,\xi} (X, \xi)^{\text{fr}} & \longleftarrow & \operatorname{colim}_{X,\xi} \{Z \xrightarrow{g} X, T_f \simeq -g^*(\xi)\} \\
\downarrow & & \downarrow \\
\text{FSyn} & \longleftarrow & \{U \xleftarrow{f} Z\}.
\end{array}$$

Formally rewriting we find that

$$\begin{aligned}
\operatorname{colim}_{X,\xi} \{Z \xrightarrow{g} X, T_f \simeq -g^*(\xi)\} &\simeq \operatorname{colim}_{X,\xi,g} \{T_f \simeq -g^*(\xi)\} \\
&\simeq \operatorname{colim}_{X,g} \operatorname{fib}_{T_f}(K(X) \xrightarrow{g^*} K(Z)),
\end{aligned}$$

and exchanging the fiber and colimit we get

$$\operatorname{colim}_{X,g} \operatorname{fib}_{T_f}(K(X) \xrightarrow{g^*} K(Z)) \simeq \operatorname{fib}_{T_f}(\operatorname{colim}_{X,g} K(X) \rightarrow K(Z)).$$

This we recognize as the formula for the left-Kan extension, that is,

$$\operatorname{colim}_{X,g} K(X) \simeq \operatorname{LKE}(K | \operatorname{Sm}_S)(Z).$$

It remains to show that

$$\operatorname{colim}_{X,\xi} \{Z \xrightarrow{g} X, T_f \simeq -g^*(\xi)\}$$

is contractible, which we have reduced to showing that

$$\operatorname{fib}_{T_f}(\operatorname{LKE}(K | \operatorname{Sm}_S)(Z) \rightarrow K(Z))$$

is contractible. However, it follows from Proposition 5.11 that the map

$$\operatorname{LKE}(K | \operatorname{Sm}_S)(Z) \rightarrow K(Z)$$

is an equivalence, hence the fiber must be trivial.

The proof for $\Sigma_T^n \text{MGL}$ for $n > 0$ is similar, but with $\xi \in K(X)$ such that $\operatorname{rank}(\xi) = n$. \square

Question 5.12. *Is Theorem 5.1 still true if one relaxes the “finiteness” condition to a “properness” condition?*

Note that the finiteness condition was not necessary to get a cobordism class. If one drops this condition, then FSyn becomes equal to PQSm^0 , which is a much larger stack. To see this one can start with any smooth proper scheme of any dimension n and find a derived scheme structure on it which is quasismooth and of dimension 0; we may for instance take the vanishing locus of the zero function n times.

It turns out that the naive cobordisms space $L_{\mathbb{A}^1}\mathrm{PQSm}^0$ is a group under disjoint union, which is not true for $L_{\mathbb{A}^1}\mathrm{FSyn}$. There should then be maps

$$\mathrm{FSyn} \hookrightarrow \mathrm{PQSm}^0 \xrightarrow{\text{conjecturally}} \Omega_T^\infty \mathrm{MGL}, \quad (5-2)$$

where the last map uses the transfers in MGL-cohomology. This would imply that $\Sigma_{\mathrm{fr}}^\infty \mathrm{PQSm}^0$ has MGL as a retract. Over perfect fields one can therefore conclude that $\Omega^\infty \mathrm{MGL}$ is a direct factor of $L_{\mathrm{Zar}} L_{\mathbb{A}^1} \mathrm{PQSm}^0$. It is an open problem whether they in fact are equivalent.

Acknowledgement

The authors greatly acknowledge the support from the research project ‘‘Motivic Geometry’’ at the Centre for Advanced Study at the Norwegian Academy of Science and Letters in Oslo, Norway.

References

- [1] T. Bachmann, ‘‘Cancellation theorem for motivic spaces with finite flat transfers’’, *Doc. Math.* **26** (2021), 1121–1144. MR Zbl
- [2] T. Bachmann and M. Hoyois, *Norms in motivic homotopy theory*, Astérisque **425**, Société Mathématique de France, Paris, 2021. MR Zbl
- [3] F. Déglise, F. Jin, and A. A. Khan, ‘‘Fundamental classes in motivic homotopy theory’’, *J. Eur. Math. Soc. (JEMS)* **23**:12 (2021), 3935–3993. MR Zbl
- [4] E. Elmanto, M. Hoyois, A. A. Khan, V. Sosnilo, and M. Yakerson, ‘‘Modules over algebraic cobordism’’, *Forum Math. Pi* **8** (2020), art. id. e14. MR Zbl
- [5] E. Elmanto, M. Hoyois, A. A. Khan, V. Sosnilo, and M. Yakerson, ‘‘Motivic infinite loop spaces’’, *Camb. J. Math.* **9**:2 (2021), 431–549. MR Zbl
- [6] G. Garkusha, A. Neshitov, and I. Panin, ‘‘Framed motives of relative motivic spheres’’, *Trans. Amer. Math. Soc.* **374**:7 (2021), 5131–5161. MR Zbl
- [7] M. Hoyois, ‘‘The localization theorem for framed motivic spaces’’, *Compos. Math.* **157**:1 (2021), 1–11. MR Zbl
- [8] M. Hoyois, J. Jelisiejew, D. Nardin, B. Totaro, and M. Yakerson, ‘‘The Hilbert scheme of infinite affine space and algebraic K-theory’’, preprint, 2021. To appear in *J. Eur. Math. Soc.* arXiv 2002.11439v3
- [9] M. Hoyois, J. Jelisiejew, D. Nardin, and M. Yakerson, ‘‘Hermitian K-theory via oriented Gorenstein algebras’’, *J. Reine Angew. Math.* **793** (2022), 105–142. MR Zbl
- [10] A. A. Khan, ‘‘Algebraic K-theory of quasi-smooth blow-ups and CDH descent’’, *Ann. H. Lebesgue* **3** (2020), 1091–1116. MR Zbl
- [11] V. Voevodsky, ‘‘Cohomological theory of presheaves with transfers’’, pp. 87–137 in *Cycles, transfers, and motivic homology theories*, Ann. of Math. Stud. **143**, Princeton Univ. Press, 2000. MR Zbl
- [12] V. Voevodsky, ‘‘My view of the current state of motivic homotopy theory’’, notes, 2000, available at <http://tinyurl.com/Voevodsky-Note>.

- [13] V. Voevodsky, “Notes on framed correspondences”, unpublished notes, 2001, available at <https://www.math.ias.edu/vladimir/sites/math.ias.edu.vladimir/files/framed.pdf>.
- [14] V. Voevodsky, “Cancellation theorem”, *Doc. Math.* Suslin birthday volume (2010), 671–685.
MR Zbl

Received 26 Jan 2022. Revised 4 Jul 2022.

MARC HOYOIS: marc.hoyois@ur.de

Fakultät für Mathematik, Universität Regensburg, Regensburg, Germany

NIKOLAI OPDAN: ntmarti@math.uio.no

Department of Mathematics, University of Oslo, Oslo, Norway

Volume Editor:

Paul Arne Østvær
University of Oslo
Oslo, Norway

Cover courtesy of Paul Arne Østvær.

The contents of this work are copyrighted by MSP or the respective authors.
All rights reserved.

Electronic copies can be obtained free of charge from <http://msp.org/obs/6> and
printed copies can be ordered from MSP (contact@msp.org).

The Open Book Series is a trademark of Mathematical Sciences Publishers.

ISSN: 2329-9061 (print), 2329-907X (electronic)

ISBN: 978-1-935107-13-2 (print), 978-1-935107-14-9 (electronic)

First published 2025.



MATHEMATICAL SCIENCES PUBLISHERS

2000 Allston Way # 59
Berkeley CA 94701-4004

contact@msp.org
<https://msp.org>

Motivic Geometry

Based on lectures given at the Centre for Advanced Study (CAS) of the Norwegian Academy of Science and Letters, this book provides a panorama of developments in motivic homotopy theory and related fields.

A common goal of the research program underlying this volume is the understanding of the geometric nature of spaces, revealed through algebraic and homotopical invariants. The articles in this volume, contributed by leading experts, together touch on an extensive network of related topics in algebraic geometry, homotopy theory, K -theory and related areas.

The volume has a significant expository component, making it accessible to students, while also containing information and in-depth discussion of interest to all practitioners including specialists.

TABLE OF CONTENTS

Preface — Paul Arne Østvær	ix
Notes on motivic infinite loop space theory — Tom Bachmann and Elden Elmanto	1
An introduction to six-functor formalisms — Martin Gallauer	63
Introduction to framed correspondences — Marc Hoyois and Nikolai Opdan	107
Lectures on the cohomology of reciprocity sheaves — Nikolai Opdan and Kay Rülling	127
The Grothendieck ring of varieties and algebraic K -theory — Oliver Röndigs	165
Stable homotopy groups of motivic spheres — Oliver Röndigs and Markus Spitzweck	197