


PURE and APPLIED ANALYSIS

PAA

YUANLONG CHEN AND HART F. SMITH

**DISPERSIVE ESTIMATES FOR THE WAVE EQUATION ON
RIEMANNIAN MANIFOLDS OF BOUNDED CURVATURE**



vol. 1 no. 1 2019

DISPERSIVE ESTIMATES FOR THE WAVE EQUATION ON RIEMANNIAN MANIFOLDS OF BOUNDED CURVATURE

YUANLONG CHEN AND HART F. SMITH

We prove space-time dispersive estimates for solutions to the wave equation on compact Riemannian manifolds with bounded curvature tensor, where we assume that the metric tensor is of $W^{1,p}$ regularity for some $p > d$, which ensures that the curvature tensor is well-defined in the weak sense. The estimates are established for the same range of Lebesgue and Sobolev exponents that hold in the case of smooth metrics. Our results are for bounded time intervals, so by finite propagation velocity they hold also on noncompact manifolds under appropriate uniform geometry conditions.

1. Introduction

We assume throughout this paper that (M, g) is a d -dimensional Riemannian manifold of C^1 structure with the following property: there exists $r_0 > 0$, $C_0 < \infty$, and $p \in (d, \infty]$ such that for each $z \in M$ there is a C^1 coordinate chart $\Phi_z : B_{r_0} \rightarrow M$, with $\Phi_z(0) = z$, in which the induced metric g_{ij} on $B_{r_0} \subset \mathbb{R}^d$ satisfies

$$g_{ij}(0) = \delta_{ij}, \quad \sup_{ij} \|g_{ij}\|_{W^{1,p}} \leq C_0.$$

As shown in [Taylor 2000, Chapter 3 §9] or Section 2 of this paper, the Riemannian curvature tensor components R_{ijkl} are then well-defined as distributions in $W^{-1,p}(B_{r_0})$. We make the assumption that the R_{ijkl} are measurable functions, and that for some C_0 uniform over the coordinate charts,

$$\sup_{ijkl} \|R_{ijkl}\|_{L^\infty(B_{r_0})} \leq C_0.$$

In Theorem 2.2 we show that the Sobolev spaces $H^s(M)$ for $-2 \leq s \leq 2$ defined using local harmonic coordinates are equivalent to those defined using fractional powers of $-\Delta_g$ via the spectral calculus. For $-1 \leq s \leq 2$ the following Cauchy problem for the wave equation on (M, g) can then be solved using the spectral decomposition for Δ_g and Duhamel's formula:

$$\begin{aligned} (\partial_t^2 - \Delta_g)u(t, x) &= F(t, x) \in L^1([-T, T]; H^{-2}(M)), \\ u(0, x) &= f(x) \in H^s(M), \\ \partial_t u(0, x) &= g(x) \in H^{s-1}(M). \end{aligned} \tag{1-1}$$

This material is based upon work supported by the National Science Foundation under Grant DMS-1500098.
MSC2010: primary 58J45; secondary 35L15.

Keywords: wave equation, dispersive estimates.

In this paper we prove two types of dispersive estimates on the solution u , under the above assumptions on (M, g) . Recall that a triple (s, q, r) with $2 \leq q, r \leq \infty$ is said to be *admissible* for the wave equation if

$$\frac{1}{q} + \frac{d}{r} = \frac{d}{2} - s, \quad \frac{1}{q} \leq \frac{d-1}{2} \left(\frac{1}{2} - \frac{1}{r} \right).$$

Theorem 1.1 (Strichartz estimates). *If (s, q, r) and $(1-s, \tilde{q}, \tilde{r})$ are admissible, and $r, \tilde{r} < \infty$, then for a positive T depending only on (M, g) , solutions to (1-1) defined using the spectral decomposition of Δ_g satisfy*

$$\begin{aligned} \|u\|_{L^q([-T, T]; L^r(M))} + \|u\|_{L^\infty([-T, T]; H^s(M))} + \|\partial_t u\|_{L^\infty([-T, T]; H^{s-1}(M))} \\ \leq C (\|f\|_{H^s(M)} + \|g\|_{H^{s-1}(M)} + \|F\|_{L^{\tilde{q}'}([-T, T]; L^{\tilde{r}'}(M))}). \end{aligned}$$

Note that under these assumptions $0 \leq s \leq 1$, and since $\tilde{q} \geq 2$ we see that $H^{\frac{3}{2}}(M) \subset L^{\tilde{r}}(M)$; hence $F \in L^1([-T, T]; H^{-\frac{3}{2}}(M))$.

The next estimate is due in the smooth case to Mockenhaupt, Seeger and Sogge [1993]. Here we consider only the critical exponent q_d , but similar results with $s_d \leq s \leq 2$ hold by Sobolev embedding.

Theorem 1.2 (square function estimate). *Let $q_d = 2(d+1)/(d-1)$, and $s_d = 1/q_d$. Then for a positive T depending only on (M, g) , solutions to (1-1) satisfy*

$$\|u\|_{L^{q_d}(M; L^2([-T, T]))} \leq C (\|f\|_{H^{s_d}(M)} + \|g\|_{H^{s_d-1}(M)} + \|F\|_{L^1([-T, T]; H^{s_d-1}(M)}).$$

A straightforward consequence of the square function estimate are the following $L^2 \rightarrow L^q$ bounds for unit-width spectral projection operators, which were originally established for smooth metrics in [Sogge 1988].

Corollary 1.3. *Suppose that $\lambda \geq 0$, and let $\Pi_{[\lambda, \lambda+1]}$ denote the $L^2(M)$ projection onto the span of eigenfunctions $\{\phi_j\}$ such that $-\Delta_g \phi_j = \lambda_j^2 \phi_j$ with $\lambda_j \in [\lambda, \lambda+1]$. Then for some C depending only on (M, g) ,*

$$\|\Pi_{[\lambda, \lambda+1]} f\|_{L^q(M)} \leq C \lambda^{d(\frac{1}{2} - \frac{1}{q}) - \frac{1}{2}} \|f\|_{L^2(M)}, \quad q_d \leq q \leq \infty.$$

Corollary 1.3 is proven for $q = q_d$ from Theorem 1.2, and for $q > q_d$ it follows by Sobolev embedding. See [Smith 2006a] for details. It is shown there that the $q = \infty$ case, which is related to the spectral counting remainder estimates of Avakumović, Levitan and Hörmander, holds more generally on compact manifolds with metrics g of Lipschitz regularity.

The first version of Strichartz estimates was obtained globally on \mathbb{R}^{d+1} by Strichartz [1970; 1977] for $s = \frac{1}{2}$ and $q = r = 2(d+1)/(d-1)$. The results were subsequently extended to other values of the exponents, and to the setting of smooth Riemannian manifolds using a Fourier integral representation of the fundamental solution. More details can be found in [Sogge 1993; Ginibre and Velo 1997; Keel and Tao 1998; Lindblad and Sogge 1995]. Of particular interest are the critical indices, when equality holds in the second admissibility condition.

For a nonsmooth metric g , the standard constructions of the fundamental solution do not work. However, in [Smith 1998] one of us used paradifferential techniques and wave packet parametrices to

prove homogeneous Strichartz estimates in dimensions $d = 2, 3$ under the condition that the metric g is $C^{1,1}$. For all dimensions this is the minimal regularity condition on g in the context of Hölder spaces that implies the Strichartz estimates. Indeed, Smith and Sogge [1994] produced explicit examples of $C^{1,\alpha}$ metrics for which the homogeneous Strichartz estimates fail, for each $0 < \alpha < 1$.

The key idea in handling nonsmooth metrics is to introduce a paradifferential approximation P to $\sqrt{-\Delta_g}$, in that $P^2 + \Delta_g$ behaves as a first-order operator on a suitable range of Sobolev spaces. By energy estimates it then suffices to establish the bounds of Theorems 1.1 and 1.2 when Δ_g is replaced by $-P^2$ in (1-1). The operator P has symbol of class $S_{1,\frac{1}{2}}^1$ and is obtained by mollifying the coefficients of g over scale $2^{-\frac{k}{2}}$ when acting on functions at frequency scale 2^k .

One then seeks a construction of the evolution operator e^{-itP} for which the desired dispersive bounds can be proven. In [Smith 1998], an approximation $E(t)$ to e^{-itP} was obtained by working in a frame of dyadic-parabolic wave packets (curvelets). A key property of such wave packets is that the action of e^{-itP} on each element of the frame is well approximated by rigid translation of the packet along the Hamiltonian flow of P , and $E(t)$ was defined as this rigid motion. This operator $E(t)$ failed to satisfy the unitary group property $E(t)E(s)^* = E(t-s)$, however, which is a crucial requirement for the established proofs of dispersive bounds such as in [Keel and Tao 1998]. This limited the results of [Smith 1998] to low dimensions. The Strichartz estimates of Theorem 1.1 for $C^{1,1}$ metrics and general dimensions were subsequently established by Tataru [2000; 2001; 2002], where space-time bounds on the FBI transform were used. Smith [2006b] used a modified FBI transform to translate the problem to phase-space, and e^{-itP} was approximated on the transform side by the Hamiltonian flow map. This forms a unitary group, and the estimates in Theorems 1.1 and 1.2 (with $F = 0$ in Theorem 1.1) were established for $C^{1,1}$ metrics, in all dimensions.

For metrics of bounded curvature the paradifferential construction of the self-adjoint operator P goes through as above, provided one works in harmonic coordinates on (M, g) . In such coordinates the metric g has second derivatives belonging to BMO, which is sufficient to show that $P^2 + \Delta_g$ maps $H^s \rightarrow H^{s-1}$ for a range of s . The wave packet methods fail to give a useful construction of e^{-itP} , however, since the error estimates for the rigid translation or Hamiltonian flow approximations depend explicitly on pointwise bounds on $\partial_x^2 g^{ij}(x)$. On the other hand, by the Jacobi variation formula, L^∞ bounds on the Riemannian curvature tensor imply that the geodesic and Hamiltonian flows are bilipschitz. A consequence is that the solution to the eikonal equation in any local harmonic coordinate system has bounded second derivatives, the same regularity as for $C^{1,1}$ metrics.

This naturally leads us in this paper to imitate the Lax parametrix construction for e^{-itP} . It turns out that solving the transport equations for the amplitude produces no further improvement beyond setting the amplitude to be identically 1, as all terms in the expansion of the amplitude would be symbols of order 0, due to the fact that the symbol of P is of class $S_{1,\frac{1}{2}}^1$. On the other hand, to have a unitary group we need work with the exact operator e^{-itP} . We achieve this by producing e^{-itP} exactly as an iterative expansion of the Lax approximation, which we show converges uniformly on finite time intervals in the H^s operator norm for every $s \in \mathbb{R}$.

To prove the dispersive estimates of Theorems 1.1 and 1.2 we establish bounds on the integral kernel of e^{-itP} localized dyadically in frequency. These bounds capture the pointwise decay of the fundamental

solution away from the light cone, and are of the exact same form as for smooth metrics. An advantage of this proof is that we can obtain the inhomogeneous estimates stated in [Theorem 1.1](#). We establish the kernel bounds using a version of the wave packet frame of [\[Smith 1998\]](#) rescaled by time t . This method is well adapted to handle the multiple products arising in the iterative expression for e^{-itP} , since the bounds can be phrased in terms of operator bounds in certain weighted norm spaces.

The proof of [Theorems 1.1](#) and [1.2](#) is composed of multiple distinct steps, and we divide it up into sections as follows. A more detailed summary of each section is included at its beginning.

In [Section 2](#), we present the details of harmonic coordinates on (M, g) and the regularity results for g in such coordinates. The procedure is similar to that in [\[Taylor 2000, Chapter 3, §9\]](#). We then reduce matters to working with a compact perturbation of the Euclidean metric on \mathbb{R}^d . We introduce the paradifferential operator approximation P , and equate the estimates of [Theorems 1.1](#) and [1.2](#) to Lebesgue space mapping properties for e^{-itP} .

In [Section 3](#), we use the Jacobi variation formula to study the regularity of the geodesic flow for the metric g_k that is obtained by mollifying g at scale $2^{-\frac{k}{2}}$. The estimates on the derivatives of the geodesic flow are exactly those obtained in the case $g \in C^{1,1}$.

In [Section 4](#), we use the results derived in [Section 3](#) and a dilation argument to prove symbol-type estimates on the solution $\varphi_k(t, x, \eta)$ of the eikonal equation for g_k . A key result is obtaining better estimates for small t , which is crucial to proving the dispersive estimates on the kernel of e^{-itP} when $|t| \ll 1$.

In [Section 5](#), we introduce an approximation $W(t)$ to e^{-itP} , which is a sum over k of terms

$$(W_k(t)f)(x) = \frac{1}{(2\pi)^d} \int e^{i\varphi_k(t,x,\eta)} \psi_k(\eta) \hat{f}(\eta) d\eta,$$

where ψ_k is a Littlewood–Paley partition of unity. We show that

$$(\partial_t + iP_k)(W_k(t)f) = B_k(t)f,$$

where $B_k(t)$ is an oscillatory integral operator with phase φ_k , and symbol $b_k(t, x, \eta)$ of order 0 that satisfies derivative bounds similar to those for φ_k .

[Section 6](#) is concerned with energy flow properties of iterated compositions of $W(t)$ and $B(t)$, which arise in the expansion of e^{-itP} . In particular, we show that multiple compositions preserve dyadic localization in frequency up to smoothing errors. Thus, in proving dispersive estimates for e^{-itP} we need only handle the composition of terms W_k and B_k , all of which are localized at the same dyadic scale. We also prove “sideways” energy estimates that arise in the proof of [Theorem 1.2](#).

In [Section 7](#) we prove that, for small t , the kernel $K_k(t, x, y)$ of $e^{-itP} \psi_k(D)$ satisfies, modulo a smoothing operator, the same bounds as for smooth metrics:

$$|K_k(t, x, y)| \leq C_N 2^{kd} (1 + 2^k |t|)^{-\frac{d-1}{2}} (1 + 2^k \text{dist}(x, S_t(y)))^{-N},$$

where $S_t(y)$ is the geodesic sphere centered at y and $\text{dist}(\cdot, \cdot)$ the geodesic distance for g_k . Together with standard arguments, these estimates yield [Theorems 1.1](#) and [1.2](#). The proof of this estimate proceeds, for a given value of t , by representing $e^{-itP} \psi_k(D)$ in a wave packet frame that is obtained by scaling by $|t|$ the dyadic-parabolic frame from [\[Smith 1998\]](#). The kernel estimates follow by showing that the operator

$e^{-itP} \psi_k(D)$ maps a frame element at time 0 to a similar function translated along the Hamiltonian flow through its center. This fact is deduced from showing the same result for the terms $W_k(s)$ and $B_k(s)$ for $0 \leq s \leq t$ that arise in the iterative formula for e^{-itP} .

2. Preliminaries and reduction to the model operator

In this section we establish regularity estimates for the metric g in local harmonic coordinate charts. We then consider Sobolev spaces on M , and define the wave group for $\sqrt{-\Delta_g}$ using the orthonormal basis for $L^2(M)$ consisting of eigenfunctions of Δ_g . We conclude by reducing the proof of [Theorem 1.1](#) to estimates for the evolution group e^{-itP} of the self-adjoint first-order pseudodifferential operator P on \mathbb{R}^d , where P is an extension to \mathbb{R}^d of a paradifferential approximation to $\sqrt{-\Delta_g}$ in one of a finite cover by M of local harmonic coordinate charts.

Harmonic coordinates on (M, g) . We start with the assumption that (M, g) is a Riemannian manifold of C^1 structure with the following condition: there exists $r_0 > 0$, $C_0 < \infty$, and $p \in (d, \infty]$, and for each $z \in M$ a coordinate chart $\Phi_z : B_{r_0} \rightarrow M$, with $\Phi_z(0) = z$, so that the induced metric g on $B_{r_0} \subset \mathbb{R}^d$ satisfies

$$g_{ij}(0) = \delta_{ij}, \quad \sup_{ij} \|g_{ij}\|_{W^{1,p}} \leq C_0.$$

Since $W^{1,p}$ functions are of Hölder regularity $1 - \frac{d}{p} > 0$, by shrinking r_0 if needed we may additionally assume that, given $c_0 > 0$ to be determined,

$$\sup_{x \in B_{r_0}} |g_{ij}(x) - \delta_{ij}| \leq c_0.$$

Following [\[Taylor 2000, Chapter 3 §9\]](#), in particular Chapter 3, Proposition 9.1 and the comments following Chapter 3, (9.39) in that paper, after replacing r_0 by $\rho_0 = \rho_0(d, p, C_0, c_0)$, we may assume that the induced coordinate functions, $f_z^i : \Phi_z(B_{\rho_0}) \rightarrow \mathbb{R}$, are harmonic functions with respect to the Laplace–Beltrami operator of g , and that overlapping harmonic coordinate charts have transition functions of regularity $W^{2,p}$ on their overlaps. The harmonic coordinates are related to the coordinate functions of Φ_z by a $W^{2,p}$ change of coordinates over B_{ρ_0} , and it follows that the original coordinates were necessarily of regularity $W^{2,p} \subset C^{1,1-\frac{d}{p}}$ on their overlaps. Consequently, M is a manifold with $W^{2,p}$ structure. This is consistent with the fact that a metric g maintains its $W^{1,p}$ regularity under a $W^{2,p}$ change of coordinates, which can be seen by [\(2-1\)](#) below.

For every integer $m \geq 0$, there is a continuous linear extension operator of $W^{m,p}(B_{\rho_0})$ to $W^{m,p}(\mathbb{R}^d)$; see, e.g., [\[Stein 1970, Chapter VI, §3, Theorem 5\]](#). We may thus apply [\[Taylor 2000, Chapter 2, Proposition 1.1\]](#), together with the inclusions

$$W^{1,p}(\mathbb{R}^d) \subset L^\infty(\mathbb{R}^d), \quad H^1(\mathbb{R}^d) = W^{1,2}(\mathbb{R}^d) \subset L^{\frac{2p}{p-2}}(\mathbb{R}^d),$$

to see that the following hold, both on \mathbb{R}^d and B_{ρ_0} :

$$\|fg\|_{W^{1,p}} \leq C \|f\|_{W^{1,p}} \|g\|_{W^{1,p}}, \quad \|fg\|_{H^1} \leq C \|f\|_{W^{1,p}} \|g\|_{H^1}. \quad (2-1)$$

The Riemannian curvature tensor R for g is given in coordinates by

$$R_{ijkl} = \frac{1}{2} \left[\frac{\partial^2 g_{ik}}{\partial x_j \partial x_\ell} + \frac{\partial^2 g_{j\ell}}{\partial x_i \partial x_k} - \frac{\partial^2 g_{i\ell}}{\partial x_j \partial x_k} - \frac{\partial^2 g_{jk}}{\partial x_i \partial x_\ell} \right] + Q(g, \partial g),$$

where $Q(g, \partial g)$ is a quadratic form in first-order derivatives of g_{ij} , with coefficients given by a combination of coefficients of g ; hence $Q(g, \partial g) \in L^{\frac{d}{2}}$ when $g \in W^{1,p}$ with $p > d$. Then R is defined as a distribution, and our key assumption is that R_{ijkl} is a bounded measurable function such that uniformly in the local coordinates F_z ,

$$\sup_{ijkl} \|R_{ijkl}\|_{L^\infty(B_{\rho_0})} \leq C_0.$$

This is implied by assuming that R is a measurable function, together with the geometric condition that for all continuous vector fields v_j ,

$$\| \langle R(v_1, v_2)v_3, v_4 \rangle \|_{L^\infty(M)} \leq C_0 \quad \text{if } \|g(v_j)\|_{L^\infty(M)} \leq 1.$$

In harmonic coordinates, the Ricci tensor Ric can be written, see for example [DeTurck and Kazdan 1981], in the form

$$\text{Ric}_{ij} = \sum_{mn} \partial_{x_m} (g^{mn} \partial_{x_n} g_{ij}) + Q(g, \partial g).$$

Since $\text{Ric}_{ij} \in L^\infty(B_{\rho_0})$, following [Taylor 2000, Chapter 3, §10] we conclude $g_{ij} \in W^{2,q}(B_\rho)$ for all $\rho < \rho_0$ and all $q < \infty$; hence $g_{ij} \in \text{Lip}(B_{.9\rho_0})$.

Take $\phi \in C_c^\infty(B_{.8\rho_0})$ with $\phi = 1$ on $B_{.7\rho_0}$, and $\chi \in C_c^\infty(B_{.9\rho_0})$ with $\chi = 1$ on $B_{.8\rho_0}$, and assume ϕ and χ take values in $[0, 1]$.

We form a Riemannian metric $\tilde{g}_{ij} = \phi g_{ij} + (1 - \phi)\delta_{ij}$ on \mathbb{R}^d , and uniformly elliptic coefficients $a^{ij} = \chi g^{ij} + (1 - \chi)\delta^{ij}$ on \mathbb{R}^d . Note that $Q(g, \partial g) \in L^\infty(B_{.9\rho_0})$ since $g \in \text{Lip}(B_{.9\rho_0})$. Then the following holds globally on \mathbb{R}^d :

$$\sum_{m,n=1}^d \partial_{x_m} (a^{mn} \partial_{x_n} \tilde{g}_{ij}) \in L_c^\infty.$$

Since the a^{mn} are globally Lipschitz, from [Taylor 2000, Chapter 3, Proposition 10.3] we conclude that $\partial_x^2 \tilde{g}_{ij} \in \text{BMO}_c(\mathbb{R}^d)$; more precisely $\partial_x^2 \tilde{g}_{ij}$ belongs to $\text{BMO}(\mathbb{R}^d)$ and is supported in $B_{.8\rho_0}$.

Note that the Riemannian curvature tensor \tilde{R} of \tilde{g} belongs to $L_c^\infty(\mathbb{R}^d)$, where we use that \tilde{g} is Lipschitz, so $\tilde{R} = \phi R$ modulo products of g and $\partial_x g$ and functions in $C_c^\infty(B_{.9\rho_0})$. After shrinking ρ_0 by a factor of 2, we conclude:

Lemma 2.1. *Given $c_0 > 0$, there exists $\rho_0 > 0$ and $C_0 < \infty$ so that for each $z \in M$ there is a harmonic coordinate chart $\Phi_z : B_{\rho_0} \rightarrow M$, with $\Phi_z(0) = z$, such that the induced metric on B_{ρ_0} agrees with the restriction of a metric g defined on \mathbb{R}^d that satisfies $g_{ij} = \delta_{ij}$ if $|x| > 2\rho_0$, and*

$$\|\partial_x^2 g_{ij}\|_{\text{BMO}} + \|g_{ij}\|_{\text{Lip}} + \|R_{ijkl}\|_{L^\infty} \leq C_0, \quad \|g_{ij} - \delta_{ij}\|_{L^\infty} \leq c_0.$$

In particular, $g_{ij} - \delta_{ij}$ belongs to $W_c^{2,q}(\mathbb{R}^d)$ for all $q < \infty$.

We now cover M by a finite collection of harmonic coordinate charts $\Phi_j \equiv \Phi_{z_j} : B_{\rho_0} \rightarrow M$, each of which satisfies the conditions of [Lemma 2.1](#), such that there is a partition of unity χ_j on M with $\text{supp}(\chi_j) \subset \Phi_j(B_{\frac{1}{3}\rho_0})$ and $\chi_j \circ \Phi_i \in W^{2,p}(B_{\rho_0})$ for each i, j . In particular, $\chi_j \circ \Phi_j \in W_c^{2,p}(B_{\rho_0})$.

By (2-1), multiplication by $\chi_j \circ \Phi_j$ maps $H_{\text{loc}}^s(B_{\rho_0})$ into $H_c^s(B_{\rho_0})$ for $s = 0, 1, 2$. By interpolation this holds for $0 \leq s \leq 2$. We may then introduce Sobolev spaces $H^s(M) \subset L^2(M)$, for $0 \leq s \leq 2$, by the condition

$$\begin{aligned} f \in H^s(M) &\iff f \circ \Phi_j \in H_{\text{loc}}^s(B_{\rho_0}) \text{ for all } j, \\ \|f\|_{H^s(M)} &= \sum_j \|(\chi_j f) \circ \Phi_j\|_{H^s(\mathbb{R}^d)}. \end{aligned} \quad (2-2)$$

If $g \in H_c^s(B_{\rho_0})$ then

$$\|(\chi_j \cdot g \circ \Phi_i^{-1}) \circ \Phi_j\|_{H^s} \leq C \|g\|_{H^s}$$

for C depending on the support of g . This holds for $s = 0, 1$ since $\Phi_i^{-1} \circ \Phi_j$ is a C^1 diffeomorphism. It holds for $s = 2$ since $D(\Phi_i^{-1} \circ \Phi_j) \in W^{1,p}$ is a multiplier on H^1 by (2-1). It then holds by interpolation for $0 \leq s \leq 2$. Consequently, there are natural continuous inclusions $H_c^s(B_{\rho_0}) \rightarrow H^s(M)$ for $0 \leq s \leq 2$ given by $g \rightarrow g \circ \Phi_j^{-1}$, and one may identify $H^s(M)$ with a closed subspace of the finite direct sum over j of $H^s(B_{\rho_0})$.

An element of $(H^s)^*$ thus induces an element of $H_{\text{loc}}^{-s}(B_{\rho_0})$, and if we identify $H^{-s}(M)$ with $(H^s)^*$ for $0 \leq s \leq 2$, then the condition (2-2) holds for $-2 \leq s \leq 2$, with approximate equality for the norm.

We observe here the following regularity property for Δ_g in harmonic coordinates, which follows, for example, from [\[Gilbarg and Trudinger 1983, Theorem 8.9\]](#). Suppose that $u \in H^1(B_{\rho_0})$ is a weak solution to $\Delta_g u = f$, where $f \in L^2(B_{\rho_0})$. Then $u \in H^2(B_\rho)$ for all $\rho < \rho_0$, and

$$\|u\|_{H^2(B_\rho)} \leq C_\rho (\|u\|_{H^1(B_{\rho_0})} + \|f\|_{L^2(B_{\rho_0})}). \quad (2-3)$$

The Sobolev spaces for $|s| \leq 2$ can also be characterized using the spectral decomposition of Δ_g on $L^2(M)$. Consider the quadratic form on $H^1(M)$ given by

$$Q(u, v) = - \int \bar{u} (\Delta_g v) dm_g = \int g(d\bar{u}, dv) dm_g.$$

Then Q is symmetric, nonnegative, and coercive. By the Rellich compactness theorem there is a complete orthonormal basis $\{v_j\}$ of $L^2(M, dm_g)$ that diagonalizes Q , in that for $f, g \in H^1(M)$

$$Q(f, g) = \sum_j \lambda_j^2 \overline{c_j(f)} c_j(g), \quad c_j(f) = \int_M \bar{v}_j f dm_g,$$

and $0 = \lambda_0 \leq \lambda_1 \leq \dots$ is a sequence of real numbers converging to ∞ . The v_j are weak solutions in $H^1(M)$ to $-\Delta_g v_j = \lambda_j^2 v_j$; hence (2-3) gives $\|v_j\|_{H^2(M)} \leq C \lambda_j^2$. It follows that $c_j(f)$ can be defined for $f \in H^s(M)$ when $-2 \leq s \leq 0$ as the action of f on \bar{v}_j .

The operator $(1 - \Delta_g)$ is equivalent to multiplication by $(1 + \lambda_j^2)$ in the basis $\{v_j\}$, and the following theorem then gives a more natural definition of $H^s(M)$.

Theorem 2.2. For $-2 \leq s \leq 2$, the mapping $f \rightarrow \{c_j(f)\}_{j=0}^\infty$ defines a homeomorphism of $H^s(M)$ with the space $\ell^2(\mathbb{N}, (1 + \lambda_j^2)^s)$. In particular, uniformly over $-2 \leq s \leq 2$, we have

$$\|f\|_{H^s(M)}^2 \approx \sum_{j=0}^\infty (1 + \lambda_j^2)^s |c_j(f)|^2, \quad c_j(f) = \int_M f \bar{v}_j \, dm_g,$$

and $\sum_{j=0}^\infty c_j(f) v_j$ converges to f in the topology of $H^s(M)$.

Proof. The theorem holds for $s = 0$ by orthonormality, and for $s = 1$ since $\|f\|_{H^1}^2 \approx \|f\|_{L^2}^2 + Q(f, f)$. For $s = 2$, we note that the partial sums

$$\sum_{j=0}^N c_j((1 - \Delta_g)f) v_j = \sum_{j=0}^N (1 + \lambda_j^2) c_j(f) v_j = (1 - \Delta_g) \sum_{j=0}^N c_j(f) v_j$$

converge in $L^2(M)$ to $(1 - \Delta_g)f$ if $f \in H^2(M)$. It follows by elliptic regularity that $\sum_j c_j(f) v_j$ converges in $H^2(M)$ to f . Surjectivity onto $\ell^2(\mathbb{N}, (1 + \lambda_j^2)^2)$ follows similarly. The theorem follows for $0 \leq s \leq 2$ by interpolation, and for $-2 \leq s \leq 0$ by duality. \square

We note that the proof also shows that $-\Delta_g$ conjugates to multiplication by $\{\lambda_j^2\}$ in the basis $\{v_j\}$, as a map from $H^s(M) \rightarrow H^{s-2}(M)$, provided $0 \leq s \leq 2$.

The wave equation on (M, g) . For data $(f, g) \in L^2(M) \oplus H^{-1}(M)$ and $F \in L_t^1([-T, T]; H^{-2}(M))$ we define the solution of the Cauchy problem (1-1) to be

$$u(t, x) = \sum_{j=0}^\infty \left(\cos(t\lambda_j) c_j(f) + \lambda_j^{-1} \sin(t\lambda_j) c_j(g) + \int_0^t \lambda_j^{-1} \sin((t-s)\lambda_j) c_j(F(s, \cdot)) \right) v_j(x), \quad (2-4)$$

where we set $0^{-1} \sin(0t) = t$. We show here that [Theorem 1.1](#) can be deduced from the following assertion:

Assume that $u \in C^0(H^s(M)) \cap C^1(H^{s-1}(M))$, and that u is given by (2-4). Then for $s, q, \tilde{q}, r, \tilde{r}$ as in [Theorem 1.1](#), the following estimate holds:

$$\|u\|_{L_t^q([-T, T]; L^r(M))} \leq C \left(\|u\|_{L_t^\infty([-T, T]; H^s(M))} + \|\partial_t u\|_{L_t^\infty([-T, T]; H^{s-1}(M))} + \|F\|_{L_t^{\tilde{q}'}([-T, T]; L^{\tilde{r}'}(M))} \right).$$

To see that this result implies [Theorem 1.1](#), consider first the case $F = 0$. Then by the spectral representation of u we have

$$\|u\|_{L_t^\infty([-T, T]; H^s(M))} + \|\partial_t u\|_{L_t^\infty([-T, T]; H^{s-1}(M))} \approx \|f\|_{H^s(M)} + \|g\|_{H^{s-1}(M)},$$

and [Theorem 1.1](#) follows from the assertion. We apply this to the triple $(1-s, \tilde{q}, \tilde{r})$ and use duality to see that, when $f = g = 0$,

$$\|u\|_{L_t^\infty([-T, T]; H^s(M))} + \|\partial_t u\|_{L_t^\infty([-T, T]; H^{s-1}(M))} \leq C \|F\|_{L_t^{\tilde{q}'}([-T, T]; L^{\tilde{r}'}(M))}.$$

The continuity of u and $\partial_t u$ follows by translation continuity, and [Theorem 1.1](#) then follows from the assertion for the case $F \neq 0$.

As a result we may assume that

$$u \in C^0(H^s(M)) \cap C^1(H^{s-1}(M)) \cap C^2(H^{s-2}(M)),$$

and in particular, $\partial_t^2 u = \Delta_g u$ in the weak sense on B_{ρ_0} in each of the local harmonic coordinate charts Φ_j .

If the data (f, g, F) is localized in $\Phi_j(B_{\frac{1}{3}\rho_0})$, then finite propagation velocity shows that $u(t)$ is supported in $\Phi_j(B_{\frac{2}{3}\rho_0})$ if $|t| \leq \frac{1}{6}\rho_0$, where we use $W^{2,p}$ regularity of g for all $p < \infty$, and closeness of g_{ij} to δ_{ij} for c_0 small.

Using the partition of unity χ_j , we can reduce the proof of [Theorem 1.1](#) to the case that the Cauchy data is supported in $\Phi_j(B_{\frac{1}{3}\rho_0})$, and thus work on \mathbb{R}^d with a metric satisfying the conditions of [Lemma 2.1](#). After rescaling space and time by a factor $R \geq 1$, where $R^{-1}C_0 \leq c_d$, we can reduce [Theorems 1.1](#) and [1.2](#) with $T = \frac{1}{6}R^{-1}\rho_0$ to the following [Theorem 2.3](#). The constant c_d will be fixed depending only on the dimension, and in particular will be small enough to rule out conjugate points for $|t| \leq 1$.

Theorem 2.3. *Assume g is a Riemannian metric on \mathbb{R}^d such that for a prescribed constant c_d depending on the dimension d ,*

$$\|\mathbf{R}_{ijkl}\|_{L^\infty} + \|g_{ij} - \delta_{ij}\|_{\text{Lip}} + \|\partial_x^2 g_{ij}\|_{\text{BMO}} \leq c_d.$$

Assume that (s, q, r) and $(1-s, \tilde{q}, \tilde{r})$ are admissible with $r, \tilde{r} < \infty$, and let $u \in C^0([0, 1]; H^s(\mathbb{R}^d)) \cap C^1([0, 1]; H^{s-1}(\mathbb{R}^d))$ be a weak solution to

$$(\partial_t^2 - \Delta_g)u = F, \quad u(0, \cdot) = f, \quad \partial_t u(0, \cdot) = g.$$

Then there is a constant $C < \infty$, depending only on d , so that

$$\|u\|_{L^q([0,1]; L^r(\mathbb{R}^d))} \leq C \left(\|u\|_{L^\infty([0,1]; H^s(\mathbb{R}^d))} + \|\partial_t u\|_{L^\infty([0,1]; H^{s-1}(\mathbb{R}^d))} + \|F\|_{L^{\tilde{q}'}([0,1]; L^{\tilde{r}'}(\mathbb{R}^d))} \right).$$

If $q_d = 2(d+1)/(d-1)$ and $s = s_d = q_d^{-1}$, then

$$\|u\|_{L^{q_d}(\mathbb{R}^d; L^2([0,1]))} \leq C \left(\|f\|_{H^{s_d}(\mathbb{R}^d)} + \|g\|_{H^{s_d-1}(\mathbb{R}^d)} + \|F\|_{L^1([0,1]; H^{s_d-1}(\mathbb{R}^d))} \right).$$

The model operator P. We construct here the paradifferential approximation to $\sqrt{-\Delta_g}$, where we will assume that g is a metric on \mathbb{R}^d that satisfies the conditions of [Theorem 2.3](#).

We fix a family of dyadically supported functions $\beta_k(\xi)$ for $k \geq 0$ such that $\beta_k(\xi) = \beta_1(2^{1-k}\xi)$ if $k \geq 1$, and such that $\psi_k(\xi) = \beta_k(\xi)^2$ gives a Littlewood–Paley partition of unity. We will assume that

$$\text{supp}(\beta_1) \subset \left\{ \frac{9}{10} \leq |\xi| \leq \frac{20}{9} \right\}, \quad \beta_0(\xi)^2 + \sum_{k=1}^{\infty} \beta_k(\xi)^2 = 1.$$

We introduce a family of metrics $g_k(x)$ that are mollifications of $g(x)$ on spatial scale $2^{-\frac{k}{2}}$. Precisely, fix a radial function $\chi \in C_c^\infty(B_1)$ so that

$$\int \chi(x) dx = 1, \quad \int x^\alpha \chi(x) dx = 0 \quad \text{if } 1 \leq |\alpha| \leq 3.$$

For $k \geq 1$ define a smooth metric \mathfrak{g}_k on \mathbb{R}^d by

$$(\mathfrak{g}_k)^{ij}(x) = 2^{\frac{kd}{2}} \int \chi(2^{\frac{k}{2}}(x-y)) g^{ij}(y) dy.$$

From the conditions on g in [Theorem 2.3](#) it follows that $\|\mathfrak{g}_k - \mathbb{I}\|_{\text{Lip}} \leq c_d$. Also,

$$\|\partial_x^\beta \mathfrak{g}_k^{ij}\|_{L^\infty} \leq C_\alpha \begin{cases} 1 + \log(k), & |\beta| = 2, \\ 2^{\frac{k}{2}(|\beta|-2)}, & |\beta| \geq 3. \end{cases}$$

The estimate for $|\beta| = 2$ holds when $k = 1$ since $\partial_x^2(\chi * g) = (\partial_x \chi) * (\partial_x g)$. For $k \geq 2$ we use that $\chi(2^{\frac{1}{2}} \cdot) - \chi(\cdot)$ is an H^1 -atom, and $\partial_x^2 g \in \text{BMO}(\mathbb{R}^d)$. The estimate for $|\beta| \geq 3$ follows by writing

$$\partial_x^\beta (\mathfrak{g}_k)^{ij}(x) = 2^{\frac{k}{2}(d+|\beta|-2)} \int (\partial_x^{\beta-2} \chi)(2^{\frac{k}{2}}(x-y)) \partial_y^2 g^{ij}(y) dy,$$

and using that $\partial_x^\theta \chi$ is an H^1 -atom, with norm C_α , when $|\theta| \geq 1$.

We also note here the bound

$$\|\partial_x^\beta (\mathfrak{g}_k - \mathfrak{g}_{k-1})\|_{L^\infty} \leq C_\beta 2^{-k+\frac{1}{2}|\beta|k}. \quad (2-5)$$

For this, write

$$\chi(\xi) - \chi(2^{\frac{1}{2}}\xi) = |\xi|^2 \rho(\xi), \quad \rho \in \mathcal{S}(\mathbb{R}^d), \quad \rho(0) = 0.$$

Then, setting $\rho_k(\xi) = \rho(2^{-\frac{k}{2}}\xi)$, we have

$$\mathfrak{g}_k - \mathfrak{g}_{k-1} = 2^{-k} \hat{\rho}_k * (\Delta g).$$

The bound (2-5) then follows from [\[Stein 1993, IV.1.1.4\]](#) as above.

Many of the steps in subsequent estimates use only the weaker estimates that follow from the Lipschitz bounds on g ,

$$\|\partial_x^\beta \mathfrak{g}_k^{ij}\|_{L^\infty} \leq C_\alpha \begin{cases} 1, & |\beta| \leq 1, \\ 2^{\frac{k}{2}(|\beta|-1)}, & |\beta| \geq 2. \end{cases} \quad (2-6)$$

Define $p_k(x, \xi) = (\sum_{i,j=1}^d \mathfrak{g}_k^{ij}(x) \xi_i \xi_j)^{\frac{1}{2}}$, so that $p_k(x, \xi)$ is homogeneous of degree 1 in ξ . Then by (2-6) and the conditions of [Theorem 2.3](#)

$$\begin{aligned} |p_k(x, \xi) - |\xi|| + |\partial_x p_k(x, \xi)| &\leq c_d |\xi|, \\ |\partial_\xi^\alpha \partial_x^\beta p_k(x, \xi)| &\leq C_{\alpha,\beta} 2^{\frac{k}{2} \max(0, |\beta|-1)} |\xi|^{1-|\alpha|}. \end{aligned} \quad (2-7)$$

Hence, $\partial_x^\beta p_k(x, \xi) \psi_k(\xi) \in S_{1, \frac{1}{2}}^1$, uniformly over $k \geq 1$, if $|\beta| \leq 1$. Similarly, by (2-5) we see that

$$(p_{k\pm 1} - p_k) \psi_k \in S_{1, \frac{1}{2}}^0 \quad \text{uniformly over } k. \quad (2-8)$$

Define

$$P = \beta_0(D)^2 + \frac{1}{2} \sum_{k=1}^{\infty} \beta_k(D) (p_k(x, D) + p_k(x, D)^*) \beta_k(D),$$

and let $p(x, \xi)$ be the symbol of P . Then P is self-adjoint, and the $S_{1, \frac{1}{2}}$ pseudodifferential calculus shows that

$$p(x, \xi) - \sum_{k=1}^{\infty} p_k(x, \xi) \psi_k(\xi) \in S_{1, \frac{1}{2}}^0.$$

In particular,

$$\partial_x^\beta p \in S_{1, \frac{1}{2}}^1 \quad \text{for } |\beta| \leq 1.$$

We note for future use that the Gårding inequality for P follows easily. It can be verified by letting

$$b(x, \xi) = \left(\psi_0(\xi) + \sum_{k=1}^{\infty} p_k(x, \xi) \psi_k(\xi) \right)^{\frac{1}{2}}.$$

Then $b(x, D)^* b(x, D) - P \in \text{Op}(S_{1, \frac{1}{2}}^0)$; hence for $f \in H^{\frac{1}{2}}$, and some real C_1

$$\langle Pf, f \rangle \geq -C_1 \|f\|_{L^2}^2. \quad (2-9)$$

Lemma 2.4. *The following holds for $0 \leq s \leq 2$:*

$$\|P^2 u + \Delta_g u\|_{H^{s-1}(\mathbb{R}^d)} \leq C \|u\|_{H^s(\mathbb{R}^d)}.$$

Proof. By (2-7), we deduce that $\partial_x^\beta p_k(x, \xi) \beta_k(\xi) \in S_{1, \frac{1}{2}}^1$ for $|\beta| \leq 1$, with uniform bounds over k . Furthermore, β_k has disjoint support from β_j if $|j - k| > 1$. The composition calculus, together with (2-8), thus shows that

$$P^2 = \sum_{k=0}^{\infty} \left(\sum_{i,j=1}^d g_k^{ij}(x) D_i D_j \right) \psi_k(D) + r(x, D), \quad r(x, \xi) \in S_{1, \frac{1}{2}}^1,$$

and in particular $r(x, D) : H^s \rightarrow H^{s-1}$ for all s . We next write

$$-\Delta_g = \sum_{i,j=1}^d g^{ij}(x) D_i D_j + \det(g)^{-\frac{1}{2}} (D_i (\det(g)^{\frac{1}{2}} g^{ij})) D_j.$$

By (2-1) we see that $\det(g)^{-\frac{1}{2}} (D_i (\det(g)^{\frac{1}{2}} g^{ij})) \in W^{1,p}$ is a multiplier on H^s for $|s| \leq 1$, so the second term maps $H^s \rightarrow H^{s-1}$ for $0 \leq s \leq 2$.

We thus need to establish that, for each i, j , we have

$$\left\| \sum_{k=0}^{\infty} (g^{ij}(x) - g_k^{ij}(x)) \psi_k(D) D_i u \right\|_{H^s} \leq C \|u\|_{H^s} \quad \text{if } -1 \leq s \leq 1. \quad (2-10)$$

By the vanishing moment condition on the radial function $\chi \in C_c^\infty$, we can write

$$1 - \hat{\chi}(\xi) = |\xi|^2 h(\xi), \quad \text{where } |\partial^\alpha h(\xi)| \leq C_\alpha \begin{cases} \min(1, |\xi|^{2-|\alpha|}), & |\xi| \leq 1, \\ |\xi|^{-2-|\alpha|}, & |\xi| \geq 1. \end{cases}$$

For $j, k \geq 0$, we let $h_{j,k}(\xi) = \psi_j(\xi) h(2^{-\frac{k}{2}} \xi)$ and then have

$$|\partial_\xi^\alpha h_{j,k}(\xi)| \leq C_\alpha 2^{-|2j-k|} 2^{-j|\alpha|}. \quad (2-11)$$

That is, $\{2^{|2j-k|}h_{j,k}\}_{j=0}^{\infty}$ satisfies the derivative estimates and localization properties of a Littlewood–Paley partition of unity in j , uniformly over k . We then write

$$g - g_k = 2^{-k} \sum_{j=0}^{\infty} g_{j,k}, \quad \text{where } g_{j,k} = -(2\pi)^{-n} \hat{h}_{j,k} * (\Delta g).$$

We observe that

$$\text{supp}(\hat{g}_{j,k}) \subset \{2^{j-1} \leq |\xi| \leq 2^{j+2}\}, \quad \|g_{j,k}\|_{L^\infty} \leq 2^{-|2j-k|}.$$

For the second estimate we use that $\|\hat{h}_{j,k} * (\Delta g)\|_{L^\infty} \leq C 2^{-|2j-k|} \|\Delta g\|_{\text{BMO}}$. This follows for $j \neq 0$ from dilation invariance of BMO and the bound

$$\int h_{j,k}(x) dx = 0, \quad |h_{j,k}(x)| \leq C 2^{-|2j-k|} 2^{jn} (1 + 2^j |x|)^{-n-1}.$$

See for example [Stein 1993, IV.1.1.4]. For $j = 0$ we write $g_{0,k} = (\nabla \hat{h}_{0,k}) * (\nabla g)$.

If $j < k - 1$, the function $g_{j,k} \psi_k(D)u$ has Fourier transform supported in $\{2^{k-1} \leq |\xi| \leq 2^{k+2}\}$, so we can use orthogonality to estimate the corresponding terms in (2-10) over $j < k - 1$,

$$\begin{aligned} \left\| \sum_{k=0}^{\infty} \sum_{j=0}^{k-2} 2^{-k} g_{j,k} \psi_k(D)Du \right\|_{H^s}^2 &\leq C \sum_{k=0}^{\infty} \left\| \sum_{j=0}^{k-2} 2^{-k} g_{j,k} \psi_k(D)Du \right\|_{H^s}^2 \\ &\leq C \sum_{k=0}^{\infty} \left(\sum_{j=0}^{k-2} 2^{-|2j-k|} \|\psi_k(D)u\|_{H^s} \right)^2 \\ &\leq C \sum_{k=0}^{\infty} \|\psi_k(D)u\|_{H^s}^2 \leq C \|u\|_{H^s}^2. \end{aligned}$$

If $j > k + 1$, then $g_{j,k} \psi_k(D)u$ is frequency supported in $\{2^{j-1} \leq |\xi| \leq 2^{j+2}\}$, and we estimate the corresponding terms in (2-10) over $j > k + 1$,

$$\begin{aligned} \left\| \sum_{k=0}^{\infty} \sum_{j=k+2}^{\infty} 2^{-k} g_{j,k} \psi_k(D)Du \right\|_{H^s} &\leq \sum_{k=0}^{\infty} \sum_{j=k+2}^{\infty} 2^{-k} \|g_{j,k} \psi_k(D)Du\|_{H^s} \\ &\leq C \sum_{k=0}^{\infty} \sum_{j=k+2}^{\infty} 2^{-k+j s} \|g_{j,k} \psi_k(D)Du\|_{L^2} \\ &\leq C \sum_{k=0}^{\infty} \sum_{j=k+2}^{\infty} 2^{k(1-s)+j(s-2)} \|\psi_k(D)u\|_{H^s} \\ &\leq C \sum_{k=0}^{\infty} 2^{-k} \|\psi_k(D)u\|_{H^s} \leq C \|u\|_{H^s}. \end{aligned}$$

It remains to handle the case $|j - k| \leq 1$. For this, we note that, by (2-11), the function $a_k(\xi) := 2^k \sum_{|j-k| \leq 1} h_{j,k}(\xi)$ satisfies the properties of a Littlewood–Paley partition of unity, as does the function

$2^{-k} \psi_k(D)D := \tilde{\psi}_k(D)$. We then rewrite the remaining terms in (2-10) as

$$\left\| \sum_{k=0}^{\infty} 2^{-k} (a_k(D) \Delta g) (\tilde{\psi}_k(D) u) \right\|_{H^s}.$$

For $-1 \leq s \leq 0$, we can dominate this using the inequality

$$\left\| \sum_{k=0}^{\infty} (a_k(D) \Delta g) (2^{-k} \tilde{\psi}_k(D) u) \right\|_{L^2} \leq C \|\Delta g\|_{\text{BMO}} \|u\|_{H^{-1}}. \quad (2-12)$$

This inequality is a discrete version of Theorem 33 of [Coifman and Meyer 1978]; for completeness we sketch the proof here. The key estimate is that

$$d\mu = \sum_{k=0}^{\infty} |(a_k(D) \Delta g)(x)|^2 dx \delta_{2^{-k}}(t)$$

is a Carleson measure, and $\|d\mu\|_c \leq C \|\Delta g\|_{\text{BMO}}^2$. This follows from the proof of [Stein 1993, IV.4.3, (37)], which goes through using that $\{a_k(2^k \xi)\}_{k=0}^{\infty}$ is a uniformly bounded set in $C_c^\infty(\frac{1}{4} \leq |\xi| \leq 8)$. To verify (2-12) we test the left-hand side against $h \in L^2(\mathbb{R}^d)$. Fix a Schwartz function ϕ with $\phi(\xi) = 1$ for $|\xi| \leq 8$. Then since $(a_k(D) \Delta g)(\tilde{\psi}_k(D) u)$ is frequency supported in $|\xi| \leq 2^{k+3}$, we have

$$\begin{aligned} & \left| \int \bar{h} \sum_{k=0}^{\infty} (a_k(D) \Delta g) (2^{-k} \tilde{\psi}_k(D) u) dx \right| \\ &= \left| \int \sum_{k=0}^{\infty} \overline{(\phi(2^{-k} D) h)} (a_k(D) \Delta g) (2^{-k} \tilde{\psi}_k(D) u) dx \right| \\ &\leq \left(\int \sum_{k=0}^{\infty} |\phi(2^{-k} D) h|^2 |a_k(D) \Delta g|^2 dx \right)^{\frac{1}{2}} \left(\int \sum_{k=0}^{\infty} |2^{-k} \tilde{\psi}_k(D) u|^2 dx \right)^{\frac{1}{2}}. \end{aligned}$$

By a result of [Carleson 1962], see [Stein 1993, II.2.2, Theorem 2 and II.2.4, (24)], the penultimate term is dominated by

$$\|h\|_{L^2} \|d\mu\|_c^{\frac{1}{2}} \leq C \|h\|_{L^2} \|\Delta g\|_{\text{BMO}},$$

and by orthogonality the last term is dominated by $\|u\|_{H^{-1}}$.

For $s \geq 0$, we use the frequency support of $(a_k(D) \Delta g)(\tilde{\psi}_k(D) u)$ to bound

$$\begin{aligned} \left\| \sum_{k=0}^{\infty} 2^{-k} (a_k(D) \Delta g) (\tilde{\psi}_k(D) u) \right\|_{H^s} &\leq \sum_{k=0}^{\infty} 2^{k(s-1)} \|(a_k(D) \Delta g) (\tilde{\psi}_k(D) u)\|_{L^2} \\ &\leq C \sum_{k=0}^{\infty} 2^{k(s-1)} \|\Delta g\|_{\text{BMO}} \|\tilde{\psi}_k(D) u\|_{L^2} \\ &\leq C \sum_{k=0}^{\infty} 2^{-k} \|\Delta g\|_{\text{BMO}} \|\tilde{\psi}_k(D) u\|_{H^s} \\ &\leq C \|\Delta g\|_{\text{BMO}} \|u\|_{H^s}. \end{aligned}$$

□

Reduction to a first-order equation. Write $(\partial_t^2 + P^2)u = F + G$, where $G = (P^2 + \Delta_g)u$. By [Lemma 2.4](#),

$$\|G\|_{L_t^\infty([0,1]; H^{s-1}(\mathbb{R}^d))} \leq C \|u\|_{L_t^\infty([0,1]; H^s(\mathbb{R}^d))}.$$

If v solves $(\partial_t^2 + P^2)v = G$ with Cauchy data set to 0, then by the Duhamel formula and energy estimates we can deduce

$$\|v\|_{L_t^q L_x^r} + \|v\|_{L_t^\infty H_x^s} + \|\partial_t v\|_{L_t^\infty H_x^{s-1}} \leq C \|u\|_{L_t^\infty H_x^s},$$

provided that we prove homogeneous Strichartz estimates for $\partial_t^2 + P^2$. By splitting $u = v + (u - v)$, the Strichartz estimates of [Theorem 2.3](#) can thus be reduced to the same estimates with $-\Delta_g$ replaced by P^2 , that is, by proving that the following holds on $[0, 1] \times \mathbb{R}^d$, provided $u \in C^0 H^s \cap C^1 H^{s-1}$:

$$\|u\|_{L_t^q L_x^r} \leq C (\|u\|_{L_t^\infty H_x^s} + \|\partial_t u\|_{L_t^\infty H_x^{s-1}} + \|(\partial_t^2 + P^2)u\|_{L_t^{\bar{q}'} L_x^{\bar{r}'}}). \quad (2-13)$$

We replace $u(t, \cdot)$ by $\langle D \rangle^{-s} u(t, \cdot)$, where $\langle D \rangle = (1 - \Delta)^{\frac{1}{2}}$, and note that

$$(\partial_t^2 + P^2)\langle D \rangle^{-s} u = [P^2, \langle D \rangle^{-s}]u + \langle D \rangle^{-s}(\partial_t^2 + P^2)u.$$

The $S_{1, \frac{1}{2}}$ calculus shows that $[P^2, \langle D \rangle^{-s}] \in S_{1, \frac{1}{2}}^{1-s}$, where we also use that $\partial_x p(x, \xi) \in S_{1, \frac{1}{2}}^1$. Consequently, using Duhamel's principle as above we see that (2-13) is equivalent to showing that, for $u \in C^0 L^2 \cap C^1 H^{-1}$, we have

$$\|\langle D \rangle^{-s} u\|_{L_t^q L_x^r} \leq C (\|u\|_{L_t^\infty L_x^2} + \|\partial_t u\|_{L_t^\infty H_x^{-1}} + \|\langle D \rangle^{-s}(\partial_t^2 + P^2)u\|_{L_t^{\bar{q}'} L_x^{\bar{r}'}}).$$

By (2-9), with $\mu = 1 + C_1$ we have

$$\langle (P + \mu)f, f \rangle \geq \|f\|_{L^2}^2 \implies \|(P + \mu)f\|_{L^2} \geq \|f\|_{L^2} \text{ when } f \in H^1.$$

By elliptic estimates we have $\|(P + \mu)f\|_{L^2(\mathbb{R}^d)} \geq \|f\|_{H^1(\mathbb{R}^d)}$; consequently $(P + \mu)^{-1}$ exists as a map from $L^2(\mathbb{R}^d) \rightarrow H^1(\mathbb{R}^d)$. One can show that $(P + \mu)^{-1} \in \text{Op}(S_{1, \frac{1}{2}}^{-1})$, for example by [Bony 1997](#).

Note that since $(P + \mu)^2 - P^2 \in \text{Op}(S_{1, \frac{1}{2}}^1)$, the estimate remains unchanged if we replace P by $P + \mu$. We will therefore assume P is invertible, with $P^{-1} \in \text{Op}(S_{1, \frac{1}{2}}^{-1})$.

The remainder of this paper is devoted to constructing the exact evolution group $E(t) = \exp(-itP)$ for the self-adjoint operator P , and proving dispersive estimates for its kernel. The group $E(t)$ will satisfy following properties:

- $E(t)$ is a strongly continuous 1-parameter unitary group on $L^2(\mathbb{R}^d)$.
- $E(t)$ is strongly continuous with respect to t on $H^s(\mathbb{R}^d)$ for all $s \in \mathbb{R}$.
- $\partial_t E(t)$ is strongly continuous with respect to t from $H^s(\mathbb{R}^d)$ into $H^{s-1}(\mathbb{R}^d)$ for all $s \in \mathbb{R}$.
- $E(0)f = f$, and $\partial_t E(t)f = -iPE(t)f = -iE(t)Pf$ if $f \in H^s(\mathbb{R}^d)$ for some $s \in \mathbb{R}$.

The second and third conditions imply that $E(t)f \in C^0(H^s) \cap C^1(H^{s-1})$ if $f \in H^s(\mathbb{R}^d)$. For $s < 0$ we understand this to mean that $E(t)$ extends continuously to such an operator from $L^2(\mathbb{R}^d)$. It follows from the third and fourth conditions that $E(t)f \in C^j(H^{s-j})$ for all $s \in \mathbb{R}$ and all $j \in \mathbb{N}$. We now let

$$C(t) = \frac{1}{2}(E(t) + E(-t)), \quad S(t) = \frac{1}{2}(E(t) - E(-t))P^{-1}.$$

The solution u to the Cauchy problem with Sobolev data

$$(\partial_t^2 + P^2)u = F, \quad u(0) = f, \quad \partial_t u(0) = g,$$

is then given by

$$u(t) = C(t)f + S(t)g + \int_0^t S(t-s)F(s) ds.$$

The Strichartz estimates in [Theorem 2.3](#) are thus reduced to showing that, for $s, q, \tilde{q}, r, \tilde{r}$ as in the statement of [Theorem 1.1](#),

$$\begin{aligned} \|\langle D \rangle^{-s} E(t) f\|_{L_t^q L_x^r([0,1] \times \mathbb{R}^d)} &\leq C \|f\|_{L^2(\mathbb{R}^d)}, \\ \left\| \langle D \rangle^{-s} \int_0^t E(t-s) F(s, \cdot) \right\|_{L_t^q L_x^r([0,1] \times \mathbb{R}^d)} &\leq C \|\langle D \rangle^{1-s} F\|_{L_t^{\tilde{q}'} L_x^{\tilde{r}'([0,1] \times \mathbb{R}^d)}}. \end{aligned} \quad (2-14)$$

Here we have used that $\langle D \rangle^{1-s} P^{-1} \langle D \rangle^s$ is bounded on $L^{\tilde{r}'(\mathbb{R}^d)}$ since it is a Calderón–Zygmund operator.

Similar steps apply to the square function estimate. For that estimate it will be more convenient to work with smooth cutoffs of the solution. We fix $\phi \in C_c^\infty((-\frac{1}{2}, \frac{1}{2}))$ with $\phi(t) = 1$ if $|t| \leq \frac{1}{3}$. By energy conservation, the square function estimates of [Theorem 2.3](#) are then reduced to showing

$$\|\phi(t) \langle D \rangle^{-s_d} E(t) f\|_{L_x^{q_d} L_t^2(\mathbb{R}^d \times [0,1])} \leq C \|f\|_{L^2(\mathbb{R}^d)}. \quad (2-15)$$

3. Regularity of the geodesic and Hamiltonian flows

In this section we establish estimates for derivatives of all orders on the geodesic and Hamiltonian flows of the metrics g_k , as well as for spatial dilates $g_k(\varepsilon \cdot)$ for $\varepsilon \leq 1$. To operate in a general context we will consider a family of metrics g_M on \mathbb{R}^d that satisfy derivative estimates depending on the parameter $M \in [1, \infty)$.

For a sufficiently small constant c_d to be chosen depending only on the dimension d , we will assume a smallness condition,

$$\|\mathbf{R}_{ijkl}\|_{L^\infty} + \|(g_M)_{ij} - \delta_{ij}\|_{\text{Lip}} + \|\nabla_x^2 (g_M)_{ij}\|_{\text{BMO}} \leq c_d. \quad (3-1)$$

Here, \mathbf{R}_{ijkl} is the Riemann curvature tensor of g_M . This tensor, as well as the Christoffel symbols Γ_{ij}^n , depends on M , but to simplify notation we suppress the subscript M .

We additionally assume that, for constants C_β independent of M ,

$$\|\partial_x^\beta g_M^{ij}\|_{L^\infty} \leq C_\beta M^{|\beta|-1}, \quad |\beta| \geq 1, \quad (3-2)$$

$$\|\partial_x^\beta \mathbf{R}_{ijkl}\|_{L^\infty} \leq C_\beta M^{|\beta|}, \quad |\beta| \geq 0. \quad (3-3)$$

Let $\gamma(t, y, w)$ be the geodesic for g_M with initial conditions (y, w) :

$$\partial_t^2 \gamma^n = \sum_{ij} \Gamma_{ij}^n(\gamma) \dot{\gamma}^i \dot{\gamma}^j, \quad \gamma(0, y, w) = y, \quad \dot{\gamma}(0, y, w) = w,$$

where $\dot{\gamma} \equiv \partial_t \gamma$. Note that by (3-1)–(3-2) we have

$$\|\Gamma_{ij}^n\|_{L^\infty} \lesssim c_d, \quad \|\partial_x^\beta \Gamma_{ij}^n\|_{L^\infty} \leq C_\beta M^{|\beta|}, \quad |\beta| \geq 1, \quad (3-4)$$

where in this section $a \lesssim b$ means that $a \leq Cb$, where C depends only on the dimension d .

Theorem 3.1. *Suppose that \mathfrak{g}_M satisfies (3-1)–(3-3) for a suitably small constant c_d . Then there are constants $C_{\alpha,\beta}$, depending only on the constants C_β in (3-2)–(3-3), so that over the set $\frac{1}{2} \leq |w| \leq 2$ and $|t| \leq 1$,*

$$|\partial_y \gamma - \mathbf{I}| + |\partial_y \dot{\gamma}| + |\partial_w \dot{\gamma} - \mathbf{I}| \lesssim c_d, \quad |\partial_w \gamma - t \mathbf{I}| \lesssim c_d |t|, \tag{3-5}$$

and

$$|\partial_y^\beta \partial_w^\alpha \gamma(t, y, w)| + |\partial_y^\beta \partial_w^\alpha \dot{\gamma}(t, y, w)| \leq C_{\alpha,\beta} M^{|\alpha|+|\beta|-1}, \quad |\alpha| + |\beta| \geq 1.$$

Additionally,

$$|\partial_y^\beta \partial_w^\alpha \gamma(t, y, w)| \leq C_{\alpha,\beta} |t| M^{|\alpha|+|\beta|-1} \quad \text{if } |\alpha| \geq 1 \text{ or } |\beta| \geq 2.$$

Proof. We produce a (not necessarily orthonormal) frame $\{V_m\}_{m=1}^d$ along $\gamma(t, y, w)$ by parallel translation of the standard frame $\{\partial_m\}_{m=1}^d$. We label the resulting vector fields $V_m(t, y, w) = \sum_n v_m^n(t, y, w) \partial_n$. The dual frame $\{V^n\}_{n=1}^d$ under \mathfrak{g}_M is obtained by parallel translating $\sum_m \mathfrak{g}_M^{nm}(y) \partial_m$ along γ , so $v^{n,l}(t, y, w) = \sum_m \mathfrak{g}_M^{nm}(y) v_m^l(t, y, w)$, and derivative estimates for the functions $v^{n,l}$ will follow directly from those for v_m^l . We have

$$\partial_t v_m^n = -\Gamma_{ij}^n(\gamma) \dot{\gamma}^i v_m^j, \quad v_m^n(0, t, w) = \delta_m^n. \tag{3-6}$$

We expand the variation of the flow in the initial parameters using the frame

$$\begin{aligned} \partial_{y^k} \gamma &= \sum_m f_k^m(t, y, w) V_m = \sum_{mj} f_k^m(t, y, w) v_m^j(t, y, w) \partial_j, \\ \partial_{w^k} \gamma &= \sum_m h_k^m(t, y, w) V_m = \sum_{mj} h_k^m(t, y, w) v_m^j(t, y, w) \partial_j. \end{aligned} \tag{3-7}$$

By (3-6) we then have

$$\begin{aligned} \partial_{y^k} \dot{\gamma}^n &= \partial_t \partial_{y^k} \gamma^n = \sum_m (\partial_t f_k^m) v_m^n - \sum_{ij} \Gamma_{ij}^n(\gamma) \dot{\gamma}^i f_k^m v_m^j, \\ \partial_{w^k} \dot{\gamma}^n &= \partial_t \partial_{w^k} \gamma^n = \sum_m (\partial_t h_k^m) v_m^n - \sum_{ij} \Gamma_{ij}^n(\gamma) \dot{\gamma}^i h_k^m v_m^j. \end{aligned} \tag{3-8}$$

Since $D_t^2 \partial_{y^k} \gamma = \sum_m (\partial_t^2 f_k^m) V_m$, with D_t covariant differentiation in t , the Jacobi variation formula yields

$$\partial_t^2 f_k^m = \sum_n \left(\sum_{ijlp} R_{ijlp}(\gamma) \dot{\gamma}^i v_n^j \dot{\gamma}^l v^{m,p} \right) f_k^n, \tag{3-9}$$

with the following initial conditions, where the second holds by (3-8):

$$f_k^m(0, y, w) = \delta_k^m, \quad \partial_t f_k^m(0, y, w) = \sum_i \Gamma_{ik}^m(y) w^i.$$

Equation (3-9) holds with f replaced by h , with initial conditions

$$h_k^m(0, y, w) = 0, \quad \partial_t h_k^m(0, y, w) = \delta_k^m.$$

The bounds $|\dot{\gamma}| \lesssim 1$ and $|v| \lesssim 1$, together with (3-4), yield for $|t| \leq 1$,

$$|v_j^n - \delta_j^n| + |f_k^m - \delta_k^m| + |\partial_t f_k^m| + |\partial_t h_k^m - \delta_k^m| \lesssim c_d, \quad |h_k^m - t \delta_k^m| \lesssim c_d |t|.$$

Together with (3-4) and (3-7)–(3-8), these yield the bound (3-5).

Assume we have shown the following for $|\alpha| + |\beta| \leq N - 1$, where $N \geq 1$:

$$|\partial_y^\beta \partial_w^\alpha (v_j^n, f_k^m, h_k^m)| + |\partial_y^\beta \partial_w^\alpha (\partial_t f_k^m, \partial_t h_k^m)| \leq C_{\alpha, \beta} M^{|\alpha| + |\beta|}. \quad (3-10)$$

Using (3-7), (3-4), and (3-8), we conclude that if $1 \leq |\alpha| + |\beta| \leq N$,

$$|\partial_y^\beta \partial_w^\alpha \gamma^n| + |\partial_y^\beta \partial_w^\alpha \dot{\gamma}^n| \leq C_{\alpha, \beta} M^{|\alpha| + |\beta| - 1}.$$

By (3-6) and the Leibniz rule, for $|\alpha| + |\beta| = N$ we then can write

$$\partial_t \partial_y^\beta \partial_w^\alpha v_m^n = -\Gamma_{ij}^n(\gamma) \dot{\gamma}^i \partial_y^\beta \partial_w^\alpha v_m^j + \mathcal{O}(M^{|\alpha| + |\beta|}).$$

Similarly, by (3-9), for $|\alpha| + |\beta| = N$ we have

$$\partial_t^2 \partial_y^\beta \partial_w^\alpha f_k^m = \sum_n \left(\sum_{ijklp} \mathbf{R}_{ijklp}(\gamma) \dot{\gamma}^i v_n^j \dot{\gamma}^l v^{m,p} \right) \partial_y^\beta \partial_w^\alpha f_k^n + \mathcal{O}(M^{|\alpha| + |\beta|})$$

and the same for f replaced by h . By the initial conditions, we have

$$\partial_t^\beta \partial_w^\alpha (v_n^m, f_k^m, h_k^m, \partial_t h_k^m)|_{t=0} = 0, \quad |\partial_t \partial_y^\beta \partial_w^\alpha f_k^m(0, y, w)| \leq C_{\alpha, \beta} M^{|\alpha| + |\beta|}.$$

An application of Gronwall's lemma then yields, for $|\alpha| + |\beta| = N$,

$$|\partial_y^\beta \partial_w^\alpha (v_n^m, f_k^m, h_k^m)| + |\partial_y^\beta \partial_w^\alpha (\partial_t f_k^m, \partial_t h_k^m)| \leq C_{\alpha, \beta} M^{|\alpha| + |\beta|},$$

and (3-10) follows for $|\alpha| + |\beta| = N$ by (3-7) and (3-8), and hence all α, β by induction. As above, this implies the desired bounds for $\partial_y^\beta \partial_w^\alpha (\gamma, \dot{\gamma})$.

The last estimate of the theorem follows from the bound on $|\partial_t \partial_y^\beta \partial_w^\alpha \gamma|$, since $\partial_y^\beta \partial_w^\alpha \gamma(0, y, w) = 0$ if either $|\alpha| \geq 1$ or $|\beta| \geq 2$. \square

We now consider the related Hamiltonian flow. Let

$$p_M(x, \eta) = \left(\sum_{ij} g_M^{ij}(x) \eta_i \eta_j \right)^{\frac{1}{2}},$$

and consider the solution $(x(t, y, \eta), \xi(t, y, \eta))$ to Hamilton's equations

$$\dot{x} = (\nabla_\xi p_M)(x, \xi), \quad \dot{\xi} = -(\nabla_x p_M)(x, \xi), \quad x(0, y, \eta) = y, \quad \xi(0, y, \eta) = \eta.$$

These are related to the geodesic flow by

$$\begin{aligned} x^i(t, y, \eta) &= \gamma^i(t, y, w(y, \eta)), \\ \xi_j(t, y, \eta) &= p_M(y, \eta) \sum_j g_{M,ij}(\gamma) \dot{\gamma}^j(t, y, w(y, \eta)), \end{aligned}$$

where

$$w^i(y, \eta) = \frac{1}{p_M(y, \eta)} \sum_j g_M^{ij}(y) \eta_j.$$

It follows from (3-1) that

$$|\partial_y w| + |w - |\eta|^{-1} \eta| + |\partial_\eta w - (\mathbf{I} - |\eta|^{-2} \eta \otimes \eta)| \lesssim c_d,$$

and from (3-2) and homogeneity that

$$|\partial_y^\beta \partial_\eta^\alpha w(y, \eta)| \leq C_{\alpha, \beta} M^{|\beta|-1} |\eta|^{-|\alpha|}.$$

Observe that $\mathbf{I} - |\eta|^{-2} \eta \otimes \eta$ is equal to Π_η^\perp , the projection onto the plane perpendicular to η . We use this to deduce the following corollary of [Theorem 3.1](#).

Corollary 3.2. *Suppose that g_M satisfies (3-1)–(3-3) for a suitably small constant c_d . Then there are constants $C_{\alpha, \beta}$, depending only on the constants C_β in (3-2)–(3-3), so that for $|t| \leq 1$*

$$|\partial_y x - \mathbf{I}| + |\partial_\eta \xi - \mathbf{I}| \lesssim c_d, \quad |\partial_y \xi| + |\xi - \eta| \lesssim c_d |\eta|, \quad |\partial_\eta x - t \Pi_\eta^\perp| \lesssim c_d |t|,$$

and when $|\alpha| + |\beta| + m \geq 1$,

$$|\eta| |\partial_\eta^\alpha \partial_y^\beta \partial_t^m x(t, y, \eta)| + |\partial_\eta^\alpha \partial_y^\beta \partial_t^m \xi(t, y, \eta)| \leq C_{\alpha, \beta} M^{|\alpha|+|\beta|+m-1} |\eta|^{1-|\alpha|}.$$

Additionally,

$$|\partial_y^\beta \partial_\eta^\alpha x(t, y, \eta)| \leq C_{\alpha, \beta} |t| M^{|\alpha|+|\beta|-1} |\eta|^{-|\alpha|}, \quad |\alpha| \geq 1 \text{ or } |\beta| \geq 2.$$

Proof. The estimates other than those involving derivatives in t follow from [Theorem 3.1](#). Estimates on derivatives in t follow by induction using Hamilton's equations and the following consequence of (3-2):

$$|\partial_x^\beta \partial_\xi^\alpha (\nabla_\xi p_M)| + |\xi|^{-1} |\partial_x^\beta \partial_\xi^\alpha (\nabla_x p_M)| \leq C_{\alpha, \beta} M^{|\beta|} |\xi|^{-|\alpha|}. \quad \square$$

For the generating function $\varphi_k(t, x, \eta)$, we need to consider the function $y(t, x, \eta)$ that is the inverse of the map $y \rightarrow x(t, y, \eta)$.

Theorem 3.3. *Suppose that g_M satisfies (3-1)–(3-3) for a suitably small constant c_d . Then there are constants $C_{\alpha, \beta}$, depending only on the constants C_β in (3-2)–(3-3), so that if $|t| \leq 1$ and $\eta \neq 0$, the map $y \rightarrow x(t, y, \eta)$ is invertible. The inverse map $y(t, x, \eta)$ satisfies $|\partial_x y - \mathbf{I}| \lesssim c_d$, and*

$$|\partial_x^\beta \partial_\eta^\alpha y(t, x, \eta)| \leq C_{\alpha, \beta} M^{|\alpha|+|\beta|-1} |\eta|^{-|\alpha|}, \quad |\alpha| + |\beta| \geq 1.$$

Additionally,

$$|\partial_x^\beta \partial_\eta^\alpha y(t, x, \eta)| \leq C_{\alpha, \beta} |t| M^{|\alpha|+|\beta|-1} |\eta|^{-|\alpha|}, \quad |\alpha| \geq 1 \text{ or } |\beta| \geq 2.$$

Also, for the function $\xi(t, x, \eta) := \xi(t, y(t, x, \eta), \eta)$,

$$|\partial_x^\beta \partial_\eta^\alpha \xi(t, x, \eta)| \leq C_{\alpha, \beta} M^{|\alpha|+|\beta|-1} |\eta|^{1-|\alpha|}, \quad |\alpha| + |\beta| \geq 1.$$

Proof. We have $|x(t, y, \eta) - y| \lesssim |t|$, so for each $\eta \neq 0$ and $|t| \leq 1$ the map $y \rightarrow x$ is proper and hence a closed mapping. Since $|\partial_y x - I| \lesssim c_d$, it is an open mapping, and hence onto and one-to-one by connectivity and simple connectivity of \mathbb{R}^d . Thus $y \rightarrow x(t, y, \eta)$ is a diffeomorphism of \mathbb{R}^d , with inverse satisfying $|\partial_x y - I| \lesssim c_d$. The estimates of the theorem are then a consequence of the inverse function theorem and [Corollary 3.2](#). \square

4. Estimates for solutions of the eikonal equation

In this section we establish estimates on derivatives of the solution to the eikonal equation for g_k . For simplicity we consider $0 \leq t \leq 1$. Let g_k be the mollification of g at spatial scale $2^{-\frac{k}{2}}$ from [Section 2](#), and let φ_k be the solution to the eikonal equation

$$\partial_t \varphi_k(t, x, \eta) = -p_k(x, \nabla_x \varphi_k(t, x, \eta)), \quad \varphi_k(0, x, \eta) = \langle x, \eta \rangle.$$

Then $\varphi_k(t, x, \eta) = \sum_i \eta_i y_i(t, x, \eta)$, where $y(t, x, \eta)$ is as in [Theorem 3.3](#), and the estimates of that theorem hold with $M = 2^{\frac{k}{2}}$. Furthermore,

$$\partial_{\eta_j} \varphi_k(t, x, \eta) = y_j(t, x, \eta), \quad \partial_{x_j} \varphi_k(t, x, \eta) = \xi_j(t, x, \eta).$$

These identities follow from the fact that $\eta \cdot dy = \xi \cdot dx$ for the homogeneous symplectic transformation $(y, \eta) \rightarrow (x, \xi)$ at fixed t .

We then easily read off the following from [Theorem 3.3](#):

$$|\partial_x^\beta \varphi_k(t, x, \eta)| \leq C_\beta 2^{\frac{k}{2}(|\beta|-2)} |\eta|, \quad |\beta| \geq 2, \quad (4-1)$$

$$|\partial_x^\beta \partial_\eta \varphi_k(t, x, \eta)| \leq C_\beta t 2^{\frac{k}{2}(|\beta|-1)}, \quad |\beta| \geq 2, \quad (4-2)$$

$$|\partial_x^\beta \partial_\eta^\alpha \varphi_k(t, x, \eta)| \leq C_{\alpha, \beta} t 2^{\frac{k}{2}(|\alpha|+|\beta|-2)} |\eta|^{1-|\alpha|}, \quad |\alpha| \geq 2. \quad (4-3)$$

Additionally,

$$|\partial_x \partial_\eta \varphi_k(t, x, \eta)| \leq C. \quad (4-4)$$

The following shows that some estimates can be improved for derivatives in η , which is key to controlling the evolution operators for small t .

Theorem 4.1. *Assume that $|\alpha| \geq 2$ or $|\beta| \geq 2$. Then when $2^{-k} \leq t \leq 1$,*

$$|\langle \eta, \partial_\eta \rangle^j \partial_x^\beta \partial_\eta^\alpha \varphi_k(t, x, \eta)| \leq C_{j, \alpha, \beta} (t^{\frac{1}{2}} 2^{\frac{k}{2}})^{|\alpha|} 2^{\frac{k}{2}(|\beta|-2)} |\eta|^{1-|\alpha|},$$

and when $0 \leq t \leq 2^{-k}$,

$$|\langle \eta, \partial_\eta \rangle^j \partial_x^\beta \partial_\eta^\alpha \varphi_k(t, x, \eta)| \leq C_{j, \alpha, \beta} 2^{\frac{k}{2}(|\beta|-2)} |\eta|^{1-|\alpha|}.$$

Proof. By homogeneity it suffices to consider the case $j = 0$. If $|\alpha| \leq 1$, the estimates for all $0 \leq t \leq 1$ follow from (4-1)–(4-2). To handle $|\alpha| \geq 2$, we take a parameter ε with $2^{-\frac{k}{2}} \leq \varepsilon \leq 1$. Let $g_{\varepsilon, k}(x) = g_k(\varepsilon x)$, where g_k is the localization of g to frequency $2^{\frac{k}{2}}$. Similarly, let $p_{\varepsilon, k}(x, \xi) = p_k(\varepsilon x, \xi)$. Let $\varphi_{\varepsilon, k}$ be the solution to

$$\partial_t \varphi_{\varepsilon, k}(t, x, \eta) = -p_{\varepsilon, k}(x, \nabla_x \varphi_{\varepsilon, k}(t, x, \eta)), \quad \varphi_{\varepsilon, k}(0, x, \eta) = \langle x, \eta \rangle.$$

Then by homogeneity we have

$$\varphi_k(t, x, \eta) = \varepsilon \varphi_{\varepsilon, k}(\varepsilon^{-1}t, \varepsilon^{-1}x, \eta). \quad (4-5)$$

The metric $g_{\varepsilon, k}(x)$ is a mollification of $g(\varepsilon x)$ at scale $\varepsilon^{-1}2^{-\frac{k}{2}} \leq 1$. Since $g(\varepsilon x)$ is Lipschitz with bounded curvature, uniformly over $\varepsilon \in [0, 1]$, we can apply estimates (4-1)–(4-3) with $2^{\frac{k}{2}}$ replaced by $M = \varepsilon 2^{\frac{k}{2}}$.

For $2^{-\frac{k}{2}} \leq t \leq 1$ we take $\varepsilon = t$ in (4-5), and apply (4-3) with $M = t 2^{-\frac{k}{2}}$ to get

$$|\partial_x^\beta \partial_\eta^\alpha \varphi_k(t, x, \eta)| \leq C_{\alpha, \beta} t^{|\alpha|-1} 2^{\frac{k}{2}(|\alpha|+|\beta|-2)} |\eta|^{1-|\alpha|}.$$

For $|\alpha| \geq 2$ this implies the desired estimate.

For $0 \leq t \leq 2^{-\frac{k}{2}}$ we take $\varepsilon = 2^{-\frac{k}{2}}$ in (4-5), and apply (4-3) with $2^{\frac{k}{2}}$ replaced by 1 to get

$$|\partial_x^\beta \partial_\eta^\alpha \varphi_k(t, x, \eta)| \leq C_{\alpha, \beta} t 2^{\frac{k}{2}|\beta|} |\eta|^{1-|\alpha|}.$$

Since $t \leq t^{\frac{|\alpha|}{2}} 2^{\frac{k}{2}(|\alpha|-2)}$ for $t \geq 2^{-k}$ and $|\alpha| \geq 2$, and $t 2^{\frac{k}{2}|\beta|} \leq 2^{\frac{k}{2}(|\beta|-2)}$ for $0 \leq t \leq 2^{-k}$, this concludes the theorem for $0 \leq t \leq 2^{-\frac{k}{2}}$. \square

As a corollary we obtain the estimates we need for linearizing the phase function, and showing the symbols are slowly varying, for η in an appropriate conical region. Given a unit vector ν , and $2^{-k} \leq t \leq 1$, we define the dyadic/conic region

$$\Omega_{k,t}^\nu = \left\{ \eta : \frac{2}{3} 2^{k-1} \leq |\eta| \leq \frac{3}{2} 2^{k+2}, \quad |\nu - |\eta|^{-1} \eta| \leq \frac{1}{16} t^{-\frac{1}{2}} 2^{-\frac{k}{2}} \right\}. \quad (4-6)$$

Note that on this region, since $t^{-\frac{1}{2}} 2^{-\frac{k}{2}} \leq 1$,

$$|\eta| \geq \langle \nu, \eta \rangle \geq \frac{3}{4} |\eta|, \quad |\Pi_{\nu^\perp} \eta| \leq t^{-\frac{1}{2}} 2^{\frac{k}{2}},$$

where Π_{ν^\perp} is projection onto the hyperplane perpendicular to ν .

Corollary 4.2. *The following estimates hold if $\eta \in \Omega_{k,t}^\nu$ and $2^{-k} \leq t \leq 1$:*

$$|\langle \nu, \partial_\eta \rangle^j \partial_\eta^\alpha \partial_x^\beta (\partial_\eta^2 \varphi_k)(t, x, \eta)| \leq C_{j, \alpha, \beta} t 2^{-k} 2^{-kj} (t^{\frac{1}{2}} 2^{-\frac{k}{2}})^{|\alpha|} 2^{\frac{k}{2}|\beta|}, \quad (4-7)$$

$$\begin{aligned} & |\langle \nu, \partial_\eta \rangle^j \partial_\eta^\alpha \partial_x^\beta (\partial_\eta \partial_x \varphi_k)(t, x, \eta)| + 2^{-k} |\langle \nu, \partial_\eta \rangle^j \partial_\eta^\alpha \partial_x^\beta (\partial_x^2 \varphi_k)(t, x, \eta)| \\ & \leq C_{j, \alpha, \beta} 2^{-kj} (t^{\frac{1}{2}} 2^{-\frac{k}{2}})^{|\alpha|} 2^{\frac{k}{2}|\beta|}, \end{aligned} \quad (4-8)$$

and

$$|\langle \nu, \partial_\eta \rangle^j \partial_\eta^\alpha \partial_x^\beta (\varphi_k(t, x, \eta) - \eta \cdot \nabla_\eta \varphi_k(t, x, \nu))| \leq C_{j, \alpha, \beta} 2^{-kj} (t^{\frac{1}{2}} 2^{-\frac{k}{2}})^{|\alpha|} 2^{\frac{k}{2}|\beta|}. \quad (4-9)$$

For $0 \leq t \leq 2^{-k}$ these hold for η in the dyadic shell $\frac{2}{3} 2^{k-1} \leq |\eta| \leq \frac{3}{2} 2^{k+2}$ if t is replaced by 2^{-k} on the right-hand side.

Proof. We consider the estimate (4-7). Theorem 4.1 gives

$$|\langle \eta, \partial_\eta \rangle^j \partial_\eta^\alpha \partial_x^\beta (\partial_\eta^2 \varphi_k)(t, x, \eta)| \leq C_{j, \alpha, \beta} t 2^{-k} (t^{\frac{1}{2}} 2^{-\frac{k}{2}})^{|\alpha|} 2^{\frac{k}{2}|\beta|}.$$

After rotation we may assume that $\nu = (1, 0, \dots, 0)$. We proceed by induction in j , the case $j = 0$ being the same as above. Suppose then that (4-7) holds for $j < j_0$. We expand

$$\langle \eta, \partial_\eta \rangle^{j_0} = \eta_1^{j_0} \partial_{\eta_1}^{j_0} + \sum_{\substack{j+|\alpha| \leq j_0 \\ j < j_0}} c_{j_0, j, \alpha} \eta_1^j \eta'^{\alpha} \partial_{\eta_1}^j \partial_{\eta'}^{\alpha}$$

Since $\eta_1 \leq \frac{3}{2} 2^{k+2}$ and $|\eta'| \leq t^{-\frac{1}{2}} 2^{\frac{k}{2}}$ on $\Omega_{k,t}^\nu$, the induction hypothesis yields

$$|\eta_1^{j_0} \partial_{\eta_1}^{j_0} \partial_\eta^\alpha \partial_x^\beta (\partial_\eta^2 \varphi_k)(t, x, \eta)| \leq C_{j, \alpha, \beta} t 2^{-k} (t^{\frac{1}{2}} 2^{-\frac{k}{2}})^{|\alpha|} 2^{\frac{k}{2} |\beta|},$$

which establishes (4-7) for $j = j_0$, since $\eta_1 \geq 2^{k-2}$ on $\Omega_{k,t}^\nu$. Similar steps establish (4-8).

The estimate (4-9) follows from (4-7) if $|\alpha| \geq 2$, so it suffices to consider $|\alpha| \leq 1$. The proof for $|\beta| \neq 0$ will follow from the proof for $\beta = 0$ with φ_k replaced by $\partial_x^\beta \varphi_k$, so we assume $\beta = 0$. We then rotate to assume that $\nu = e_1$, in which case by homogeneity the estimate becomes

$$|\partial_{\eta_1}^j \partial_{\eta'}^\alpha (\varphi_k(t, x, \eta_1, \eta') - \varphi_k(t, x, \eta_1, 0) - \eta' \cdot \nabla_{\eta'} \varphi_k(t, x, \eta_1, 0))| \leq C_{j, \alpha} 2^{-kj} (t^{\frac{1}{2}} 2^{-\frac{k}{2}})^{|\alpha|}.$$

This estimate follows from a Taylor expansion argument together with (4-7), since $|\eta'| \leq t^{-\frac{1}{2}} 2^{\frac{k}{2}}$ on $\Omega_{k,t}^{e_1}$.

For $0 \leq t \leq 2^{-k}$ the desired estimates follow easily from [Theorem 4.1](#). \square

We also record estimates for time derivatives of φ_k , which will be used in establishing space-time energy estimates.

Corollary 4.3. *Assume that $2^{-k} \leq t \leq 1$. If $|\alpha| \geq 1$, then*

$$|\langle \eta, \partial_\eta \rangle^j \partial_\eta^\alpha \partial_t \varphi_k(t, x, \eta)| \leq C_{j, \alpha} (t^{\frac{1}{2}} 2^{\frac{k}{2}})^{|\alpha|-1} |\eta|^{1-|\alpha|},$$

and if $m + |\beta| \geq 2$,

$$|\langle \eta, \partial_\eta \rangle^j \partial_x^\beta \partial_\eta^\alpha \partial_t^m \varphi_k(t, x, \eta)| \leq C_{j, m, \alpha, \beta} (t^{\frac{1}{2}} 2^{\frac{k}{2}})^{|\alpha|} 2^{\frac{k}{2} (m+|\beta|-2)} |\eta|^{1-|\alpha|}.$$

If $0 \leq t \leq 2^{-k}$, both of these estimates hold with t replaced by 2^{-k} on the right-hand side.

Proof. By homogeneity we may assume $j = 0$. The estimates that involve no derivatives in t , that is, the second estimate with $m = 0$, hold by [Theorem 4.1](#). We assume both estimates hold for derivatives up to order $m \geq 0$ in t , and prove they hold for derivatives of order $m + 1$ in t . Write $\partial_t \varphi_k = p_k(x, \nabla_x \varphi_k)$, and observe that $\partial_x^\beta \partial_\eta^\alpha \partial_t^{m+1} \varphi_k$ can be written as a sum of terms of the form

$$(\partial_x^{\beta_0} \partial_\xi^\gamma p_k)(x, \nabla_x \varphi_k) (\partial_x^{\beta_1} \partial_\eta^{\alpha_1} \partial_t^{m_1} \nabla_x \varphi_k) \cdots (\partial_x^{\beta_{|\gamma|}} \partial_\eta^{\alpha_{|\gamma|}} \partial_t^{m_{|\gamma|}} \nabla_x \varphi_k),$$

where $\sum_{j=0}^{|\gamma|} \beta_j = \beta$, $\sum_{j=1}^{|\gamma|} \alpha_j = \alpha$, $\sum_{j=1}^{|\gamma|} m_j = m$. If $|\beta| = m = 0$, we must have $|\alpha_j| \geq 1$ for all j , and the first estimate of the corollary is a result of the following bounds from (2-7) and [Theorem 4.1](#):

$$\begin{aligned} |(\partial_\xi^\gamma p_k)(x, \nabla_x \varphi_k)| &\leq C_\gamma |\eta|^{1-|\gamma|}, \\ |\partial_\eta^{\alpha_j} \nabla_x \varphi_k| &\leq C_{\alpha_j} (t^{\frac{1}{2}} 2^{\frac{k}{2}})^{|\alpha_j|-1} |\eta|^{1-|\alpha_j|}. \end{aligned}$$

Assume that $|\beta| + m \geq 1$. If $|\beta_0| \geq 1$, then the second estimate of the corollary is a result of the following bounds from (2-7) and the induction assumption:

$$\begin{aligned} |(\partial_x^{\beta_0} \partial_\xi^\gamma p_k)(x, \nabla_x \varphi_k)| &\leq C_{\gamma, \beta_0} 2^{\frac{k}{2}(|\beta_0| - 1)} |\eta|^{1 - |\gamma|}, \\ |\partial_x^{\beta_j} \partial_\eta^{\alpha_j} \partial_t^{m_j} \nabla_x \varphi_k| &\leq C_{\alpha_j, \beta_j, m_j} 2^{\frac{k}{2}(|\beta_j| + m_j)} (t^{\frac{1}{2}} 2^{\frac{k}{2}})^{|\alpha_j|} |\eta|^{1 - |\alpha_j|}. \end{aligned}$$

Finally, if $|\beta_0| = 0$ then we may assume $|\beta_1| + m_1 \geq 1$, and use the bounds

$$\begin{aligned} |(\partial_\xi^\gamma p_k)(x, \nabla_x \varphi_k)| &\leq C_\gamma |\eta|^{1 - |\gamma|}, \\ |\partial_x^{\beta_1} \partial_\eta^{\alpha_1} \partial_t^{m_1} \nabla_x \varphi_k| &\leq C_{\alpha_1, \beta_1, m_1} 2^{\frac{k}{2}(|\beta_1| + m_1 - 1)} (t^{\frac{1}{2}} 2^{\frac{k}{2}})^{|\alpha_1|} |\eta|^{1 - |\alpha_1|}, \\ |\partial_x^{\beta_j} \partial_\eta^{\alpha_j} \partial_t^{m_j} \nabla_x \varphi_k| &\leq C_{\alpha_j, \beta_j, m_j} 2^{\frac{k}{2}(|\beta_j| + m_j)} (t^{\frac{1}{2}} 2^{\frac{k}{2}})^{|\alpha_j|} |\eta|^{1 - |\alpha_j|}. \end{aligned} \quad \square$$

5. Parametrix for the dyadically localized equation

In this section, we use the eikonal solution φ_k to produce an approximation to the wave group for P with data at frequency scale 2^k . In the next section we will use these approximations to produce the exact evolution group for P by iteration. For $k \geq 2$ we define

$$\tilde{P}_k = \frac{1}{2} \sum_{j=k-1}^{k+1} \beta_j(D) (p_j(x, D) + p_j(x, D)^*) \beta_j(D).$$

Let $\tilde{p}_k(x, \eta)$ denote the symbol of \tilde{P}_k . Recalling that $\beta_j^2 = \psi_j$,

$$\tilde{p}_k(x, \eta) = \sum_{j=k-1}^{k+1} p_j(x, \eta) \psi_j(\eta) + \sum_{j=k-1}^{k+1} q_j(x, \eta) \beta_j(\eta), \quad (5-1)$$

where $q_j \in S_{1, \frac{1}{2}}^0$, uniformly over j . For $|\eta| \in [\frac{3}{4} 2^k, \frac{4}{3} 2^{k+1}]$ we define

$$b_k(t, x, \eta) = e^{-i\varphi_k(t, x, \eta)} (\partial_t + i\tilde{P}_k) e^{i\varphi_k(t, x, \eta)},$$

where \tilde{P}_k acts on x .

We then define $W_k(t)$ for $k \geq 2$ by

$$(W_k(t)f)(x) = \frac{1}{(2\pi)^d} \int e^{i\varphi_k(t, x, \eta)} \psi_k(\eta) \hat{f}(\eta) d\eta. \quad (5-2)$$

It follows that $(\partial_t + i\tilde{P}_k)W_k(t) = B_k(t)$, where

$$(B_k(t)f)(x) = \frac{1}{(2\pi)^d} \int e^{i\varphi_k(t, x, \eta)} b_k(t, x, \eta) \psi_k(\eta) \hat{f}(\eta) d\eta. \quad (5-3)$$

Theorem 5.1. *For $|t| \leq 1$ the symbol $b_k(t, x, \eta)$ satisfies*

$$|\langle \eta, \partial_\eta \rangle^j \partial_\eta^\alpha \partial_x^\beta \partial_t^m b_k(t, x, \eta)| \leq C_{j, \alpha, \beta, m} \begin{cases} (t^{\frac{1}{2}} 2^{-\frac{k}{2}})^{|\alpha|} 2^{\frac{k}{2}(|\beta| + m)}, & |t| \geq 2^{-k}, \\ 2^{-k|\alpha|} 2^{\frac{k}{2}(|\beta| + m)}, & |t| \leq 2^{-k}. \end{cases}$$

Proof. The symbol $b_k(t, x, \eta)$ is given by the oscillatory integral

$$i \partial_t \varphi_k(t, x, \eta) + \frac{i}{(2\pi)^n} \int e^{i \langle x-y, \xi \rangle + i \varphi_k(t, y, \eta) - i \varphi_k(t, x, \eta)} \tilde{p}_k(x, \xi) dy d\xi, \quad (5-4)$$

where recall that we assume $|\eta| \in [\frac{3}{4}2^k, \frac{4}{3}2^{k+1}]$. We write

$$\varphi_k(t, y, \eta) - \varphi_k(t, x, \eta) = (y - x) \cdot V(t, x, y - x, \eta),$$

where

$$V(t, x, h, \eta) = \int_0^1 (\nabla_x \varphi_k)(t, x + sh, \eta) ds.$$

Then

$$V(t, x, 0, \eta) = \nabla_x \varphi_k(t, x, \eta), \quad \partial_{h_i} V_j(t, x, 0, \eta) = \frac{1}{2} \partial_{x_i} \partial_{x_j} \varphi_k(t, x, \eta).$$

We note $|V(t, x, h, \eta) - \eta| \leq \frac{1}{8}|\eta|$ by (3-1), and for $|\alpha| + |\beta| + m + |\gamma| \geq 1$ Corollary 4.3 yields

$$|\partial_\eta^\alpha \partial_x^\beta \partial_t^m \partial_h^\gamma V(t, x, h, \eta)| \leq C_{\alpha, \beta, m, \gamma} 2^{\frac{k}{2}(|\alpha| + |\beta| + m + |\gamma| - 1)} |\eta|^{1 - |\alpha|}. \quad (5-5)$$

We make the change of variables $y \rightarrow y + h$, followed by $\xi \rightarrow V(t, x, h, \eta) + \xi$, to write the integral term in (5-4) as

$$\int e^{-i \langle h, \xi \rangle} \tilde{p}_k(x, V(t, x, h, \eta) + \xi) dh d\xi. \quad (5-6)$$

We then decompose (5-6) using a smooth cutoff χ , supported in $|\xi| \leq 2$, with $\chi(\xi) = 1$ for $|\xi| \leq 1$. Specifically, we write

$$1 = \chi(2^{-k+4}\xi)(1 - \chi(h)) + (1 - \chi(2^{-k+4}\xi)) + \chi(h)\chi(2^{-k+4}\xi).$$

Since $\tilde{p}_k \in S_{1, \frac{1}{2}}^1$, the estimates (5-5) imply that if $|\eta| \in [\frac{3}{4}2^k, \frac{4}{3}2^{k+1}]$,

$$\begin{aligned} & |\partial_\xi^\theta \partial_\eta^\alpha \partial_x^\beta \partial_t^m \partial_h^\gamma \tilde{p}_k(x, V(t, x, h, \eta) + \xi) \chi(2^{-k+4}\xi)| \\ & \leq C_{\alpha, \beta, m, \gamma, \theta} \begin{cases} 2^{k(1 - |\alpha| - |\theta|)} 2^{\frac{k}{2}(|\alpha| + |\beta| + m + |\gamma| - 1)}, & |\alpha| + |\beta| + m + |\gamma| \geq 1, \\ 2^{k(1 - |\theta|)}, & |\alpha| + |\beta| + m + |\gamma| = 0. \end{cases} \end{aligned} \quad (5-7)$$

Consider first the term $r_1(t, x, \eta)$, defined by

$$\begin{aligned} & \int e^{-i \langle h, \xi \rangle} \tilde{p}_k(x, V(t, x, h, \eta) + \xi) \chi(2^{-k+4}\xi)(1 - \chi(h)) dh d\xi \\ & = \int e^{-i \langle h, \xi \rangle} \Delta_\xi^N (\tilde{p}_k(x, V(t, x, h, \eta) + \xi) \chi(2^{-k+4}\xi)) \times (1 - \chi(h)) |h|^{-2N} dh d\xi. \end{aligned}$$

The estimates (5-7) show that the integrand is bounded by $2^{k(1-2N)}|h|^{-2N}$, and it is supported where $|\xi| \leq 2^{k-3}$ and $|h| > 1$. Similar estimates on its derivatives in (x, η) yield that, for all N ,

$$|\partial_\eta^\alpha \partial_x^\beta \partial_t^m r_1(t, x, \eta)| \leq C_{N, \alpha, \beta, m} 2^{-kN}, \quad |\eta| \in [\frac{3}{4}2^k, \frac{4}{3}2^{k+1}]. \quad (5-8)$$

Next consider the term $r_2(t, x, \eta)$, defined by the integral

$$\begin{aligned} & \int e^{-i\langle h, \xi \rangle} \tilde{p}_k(x, V(t, x, h, \eta) + \zeta)(1 - \chi(2^{-k+4}\zeta)) dh d\zeta \\ &= \int e^{-i\langle h, \xi \rangle} (1 - \Delta_\xi)^n \Delta_h^N (\tilde{p}_k(x, V(t, x, h, \eta) + \zeta)(1 - \chi(2^{-k+4}\zeta)) |\zeta|^{-2N}) (1 + |h|^2)^{-n} dh d\zeta. \end{aligned}$$

The estimates (5-7) show that the integrand is bounded by a constant times $2^k(N + \frac{1}{2}) |\zeta|^{-2N} (1 + |h|^2)^{-n}$, and it is supported where $|\zeta| \geq 2^{k-4}$. It follows that $r_2(t, x, \eta)$ also satisfies the estimates (5-8).

Thus, up to rapidly decreasing terms, the symbol $b_k(t, x, \eta)$ is equal to

$$i \partial_t \varphi_k(t, x, \eta) + \frac{i}{(2\pi)^d} \int e^{-i\langle h, \xi \rangle} \tilde{p}_k(x, V(t, x, h, \eta) + \zeta) \chi(2^{-k+4}\zeta) \chi(h) dh d\zeta.$$

We take a Taylor expansion in ζ of \tilde{p}_k about $\zeta = 0$ to write the integral as

$$\sum_{|\gamma| < 2N} \frac{1}{\gamma!} \int e^{-i\langle h, \xi \rangle} D_h^\gamma ((\partial_\xi^\gamma \tilde{p}_k)(x, V(t, x, h, \eta)) \chi(h)) \chi(2^{-k+4}\zeta) dh d\zeta + r(t, x, \eta),$$

where $r(t, x, \eta)$ is given by

$$\sum_{|\gamma| = 2N} \int_0^1 (1-s)^{N-1} \int e^{-i\langle h, \xi \rangle} D_h^\gamma ((\partial_\xi^\gamma \tilde{p}_k)(x, V(t, x, h, \eta) + s\zeta) \chi(h)) \chi(2^{-k+4}\zeta) dh d\zeta ds.$$

The estimates (5-7) show that $|\partial_x^\beta \partial_\eta^\alpha r(t, x, \eta)| \leq C_{N, \alpha, \beta} 2^{k(d+1-\frac{1}{2}|\alpha|+\frac{1}{2}|\beta|-N)}$, provided that $|\eta| \in [\frac{3}{4}2^k, \frac{4}{3}2^{k+1}]$.

To handle the terms with $|\gamma| < 2N$, let $\phi(h) = 2^{-4d} \hat{\chi}(2^{-4}h)$, which has integral $(2\pi)^n$ and vanishing moments of all nonzero order, and write the γ term in the sum as

$$\int e^{-i\langle h, \xi \rangle} D_h^\gamma ((\partial_\xi^\gamma \tilde{p}_k)(x, V(t, x, h, \eta)) \chi(h)) 2^{nk} \phi(2^k h) dh d\zeta.$$

We Taylor expand $\tilde{p}_k(x, V(t, x, h, \eta)) \chi(h)$ to order N about $h = 0$. The N -th order remainder term will lead to a term bounded by $2^{k(1-\frac{1}{2}|\gamma|-N)}$, with similar estimates on derivatives in (x, η) . All terms with h^θ with $\theta \neq 0$ integrate to 0 by the moment condition. Therefore, since $\partial_t \varphi_k(t, x, \eta) = -p_k(x, \nabla_x \varphi_k(t, x, \eta))$, we can write $b_k(t, x, \eta)$ as $r(t, x, \eta)$ plus

$$i \left(-p_k(x, \nabla_x \varphi_k(t, x, \eta)) + \sum_{|\gamma| < 2N} \frac{1}{\gamma!} D_h^\gamma ((\partial_\xi^\gamma \tilde{p}_k)(x, V(t, x, h, \eta))|_{h=0}) \right). \quad (5-9)$$

If $|\eta| \in [\frac{3}{4}2^k, \frac{4}{3}2^{k+1}]$ then $(\psi_{k-1} + \psi_k + \psi_{k+1})(\nabla_x \varphi_k(t, x, \eta)) = 1$, so by (5-1) the $\gamma = 0$ term combines with $-p_k(t, x, \nabla_x \varphi_k(t, x, \eta))$ to give

$$\begin{aligned} & \sum_{j=k-1}^{k+1} (p_k - p_j)(x, \nabla_x \varphi_k(t, x, \eta)) \psi_j(\nabla_x \varphi_k(t, x, \eta)) \\ & \quad + \sum_{j=k-1}^{k+1} q_j(x, \nabla_x \varphi_k(t, x, \eta)) \beta_j(\nabla_x \varphi_k(t, x, \eta)). \end{aligned} \quad (5-10)$$

We will estimate this term similarly to the term $|\gamma| = 1$, using the following estimate, which is a consequence of (2-5):

$$|\partial_x^\beta \partial_\xi^\alpha (p_k - p_j)(x, \xi)| \leq C_{\alpha, \beta} 2^{k(\frac{1}{2}|\beta|-1)} |\xi|^{1-|\alpha|}. \quad (5-11)$$

The same estimates hold for the term $q_j(x, \xi) \in S_{1, \frac{1}{2}}^0$ when $|\xi| \approx 2^k$.

We now examine the terms in the sum when $|\gamma| \geq 1$. Observe that

$$\partial_h^\theta V(t, x, h, \eta)|_{h=0} = \frac{1}{1 + |\theta|} \partial_x^\theta \nabla_x \varphi_k(t, x, \eta).$$

The γ term in (5-9) is then a finite linear combination of terms of the form

$$(\partial_\xi^{\gamma+\sigma} \tilde{p}_k)(x, \nabla_x \varphi_k(t, x, \eta)) (\partial_x^{\theta_1} \nabla_x \varphi_k(t, x, \eta)) \cdots (\partial_x^{\theta_l} \nabla_x \varphi_k(t, x, \eta)),$$

where $\theta_1 + \cdots + \theta_l = \gamma$, each $\theta_i \neq 0$, and $l = |\sigma| \geq 1$.

By Corollary 4.3, when $\theta_i \neq 0$, $|\eta| \in [\frac{3}{4}2^k, \frac{4}{3}2^{k+1}]$, and $2^{-k} \leq t \leq 1$,

$$|\langle \eta, \partial_\eta \rangle^j \partial_\eta^\alpha \partial_x^\beta \partial_t^m (\partial_x^{\theta_i} \nabla_x \varphi_k(t, x, \eta))| \leq C_{j, \alpha, \beta, m, \theta} 2^{\frac{k}{2}(|\theta_i|+1)} (t^{\frac{1}{2}} 2^{-\frac{k}{2}})^{|\alpha|} 2^{\frac{k}{2}(|\beta|+m)}.$$

A recursion argument and (2-7) then show that, for $2^{-k} \leq t \leq 1$,

$$\begin{aligned} |\langle \eta, \partial_\eta \rangle^j \partial_\eta^\alpha \partial_x^\beta \partial_t^m ((\partial_\xi^{\gamma+\sigma} \tilde{p}_k)(x, \nabla_x \varphi_k) \partial_x^{\theta_1} \nabla_x \varphi_k \cdots \partial_x^{\theta_l} \nabla_x \varphi_k)| \\ \leq C_{j, \alpha, \beta, m, \gamma, \sigma} 2^{\frac{k}{2}(2-|\gamma|-|\sigma|)} (t^{\frac{1}{2}} 2^{-\frac{k}{2}})^{|\alpha|} 2^{\frac{k}{2}(|\beta|+m)}. \end{aligned}$$

The expression for $b_k(t, x, \eta)$ involves an asymptotic sum over $|\gamma| \geq 1$, where also $|\sigma| \geq 1$ in all terms, and the sum thus satisfies the statement of the theorem in the case $2^{-k} \leq t \leq 1$. The estimate for $0 \leq t \leq 2^{-k}$ follows similarly.

It remains to consider the term (5-10). Using (5-11) and a similar recursion argument, we obtain for the case $2^{-k} \leq t \leq 1$,

$$\begin{aligned} \left| \langle \eta, \partial_\eta \rangle^j \partial_\eta^\alpha \partial_x^\beta \partial_t^m \left(\sum_{i=k-1}^{k+1} (p_k - p_i)(x, \nabla_x \varphi_k(t, x, \eta)) \psi_i(\nabla_x \varphi_k(t, x, \eta)) \right) \right| \\ \leq C_{j, \alpha, \beta, m} (t^{\frac{1}{2}} 2^{-\frac{k}{2}})^{|\alpha|} 2^{\frac{k}{2}(|\beta|+m)}, \end{aligned}$$

and the proof for $0 \leq t \leq 2^{-k}$ is similar. \square

Repeating the proof of Corollary 4.2, we obtain the following.

Corollary 5.2. *The following estimates hold for $\eta \in \Omega_{k, t}^v$:*

$$|\langle v, \partial_\eta \rangle^j \partial_\eta^\alpha \partial_x^\beta \partial_t^m b_k(t, x, \eta)| \leq C_{j, \alpha, \beta, m} 2^{-kj} (t^{\frac{1}{2}} 2^{-\frac{k}{2}})^{|\alpha|} 2^{\frac{k}{2}(|\beta|+m)}.$$

For $0 \leq t \leq 2^{-k}$ these hold for η in the dyadic shell $\frac{2}{3}2^{k-1} \leq |\eta| \leq \frac{3}{2}2^{k+2}$ if t is replaced by 2^{-k} on the right-hand side.

6. Energy flow estimates

In this section we construct the exact wave group $\exp(-itP)$ via a convergent iteration based on the approximate wave group

$$W(t) = \sum_{k=2}^{\infty} W_k(t) + \psi_0(D) + \psi_1(D). \tag{6-1}$$

Recall from (5-2)–(5-3) that $(\partial_t + i\tilde{P}_k)W_k(t) = B_k(t)$ is of order 0. To show that $(\partial_t + iP)W(t)$ is of order 0 we will show that, for $|t| \leq 1$, $W_k(t)f$ remains localized in frequency to an appropriate dyadic shell at scale 2^k , modulo smoothing errors. This will yield

$$(\partial_t + iP)W(t) = \sum_{k=2}^{\infty} B_k(t) + R(t),$$

with $R(t)$ a smoothing error. Denoting the right-hand side by $B(t)$, since $W(0) = I$ the wave group can be obtained by convergent iteration of $W(t)$ and $B(t)$, using Sobolev mapping bounds for both. Dispersive estimates will then depend on showing that composite terms

$$W(t - s_1)B(s_1 - s_2) \cdots B(s_{n-1} - s_n)B(s_n), \quad t \geq s_1 \geq \cdots \geq s_n \geq 0,$$

have similar microlocal mapping properties to $W(t)$ and $B(t)$. For a fixed n we could show that this term has an oscillatory integral representation similar to that for $B(t)$, but at frequency scale 2^k we will need to consider n up to $n \sim 2^{k\sigma}$ for some $\sigma > 0$. To prove preservation of dyadic localization of the energy we then need to microlocalize the energy mapping of each term $B_k(s)$ to within $2^{k(1-\sigma)}$ of the Hamiltonian flow. For convenience we fix $\sigma = \frac{1}{4}$, though any $\sigma \in (0, \frac{1}{2})$ would work. We then consider frequency cutoffs with symbols $a(\eta) \in S_{\frac{3}{4}}^0$, that is,

$$|\partial_{\eta}^{\alpha} a(\eta)| \leq C_{\alpha} 2^{-\frac{3}{4}k|\alpha|}, \quad \text{supp}(a) \subset \{\eta : \frac{4}{5} 2^{k-1} < |\eta| < \frac{5}{4} 2^{k+2}\}. \tag{6-2}$$

Given any compact set $K \subset \{\eta : \frac{7}{8} 2^{k-1} < |\eta| < \frac{8}{7} 2^{k+2}\}$ and $\delta > 0$, there exists a cutoff a satisfying (6-2) such that $\text{supp}(a)$ is contained in the $\delta 2^{\frac{3k}{4}}$ neighborhood of K and $a = 1$ on the $\frac{1}{2}\delta 2^{\frac{3k}{4}}$ neighborhood of K , and such that the constants C_{α} depend on δ but are independent of K . Such an $a(\eta)$ can be obtained, for example, by convolving the support function of the $\frac{3}{4}\delta 2^{\frac{3k}{4}}$ neighborhood of K with an approximation to the identity supported in the $\frac{1}{8}\delta$ ball.

Lemma 6.1. *Suppose that a_1 and a_2 are cutoffs satisfying (6-2), and let K be the projection onto η of the image of $\mathbb{R}^d \times \text{supp}(a_1)$ under the Hamiltonian flow of p_k at time t . Assume that $a_2 = 1$ on the $\delta 2^{\frac{3k}{4}}$ neighborhood of K . Then for all N ,*

$$\|(1 - a_2(D))B_k(t)a_1(D)f\|_{H^N} \leq C_N 2^{-kN} \|f\|_{H^{-N}},$$

where the constant C_N depends only on N , the constants C_{α} in (6-2), and δ . The same holds with $B_k(t)$ replaced by $W_k(t)$.

Proof. We prove this using a modification of the Córdoba–Fefferman wave packet transform [1978]. We use the particular transform from [Smith 1998], which is based on a Schwartz function with Fourier

transform of compact support, instead of a Gaussian. Fix g a radial, real Schwartz function with $\|g\|_{L^2} = (2\pi)^{\frac{d}{2}}$ and $\text{supp}(\hat{g}) \subset \{|\zeta| < \frac{1}{4}\}$, and set

$$g_{x,\xi}(z) = 2^{\frac{kd}{4}} e^{i\langle \xi, z-x \rangle} g(2^{\frac{k}{2}}(z-x)).$$

For $f \in L^2(\mathbb{R}^d)$ define

$$(T_k f)(x, \xi) = \int f(z) \overline{g_{x,\xi}(z)} dz.$$

Then T_k is an isometry, with adjoint given by

$$(T_k^* F)(z) = \int F(x, \xi) g_{x,\xi}(z) dx d\xi.$$

Since $|\eta| \approx 2^k$ on the support of $a_1(\eta)$, it suffices to show that for all N ,

$$\|T_k \langle D \rangle^N (1 - a_2(D)) B_k(t) a_1(D) T_k^* F\|_{L^2(\mathbb{R}^{2d})} \leq C_N 2^{-2kN} \|F\|_{L^2(\mathbb{R}^{2d})}.$$

The operator on the left is given by the following integral kernel:

$$K_k(t, x', \xi'; x, \xi) = \int_{\mathbb{R}^d} (B_k(t) a_1(D) g_{x,\xi})(z) \overline{\langle D \rangle^N (1 - a_2(D)) g_{x',\xi'}(z)} dz.$$

Let $(x_t, \xi_t) = \chi_t(x, \xi)$, with χ_t the Hamiltonian flow for p_k . A simple integration by parts argument, using [Lemma 6.2](#) below, shows that for all N ,

$$\begin{aligned} |K_k(t, x', \xi'; x, \xi)| &\leq C_N 2^{kN} (1 + 2^{\frac{k}{2}} |x' - x_t| + 2^{-\frac{k}{2}} |\xi' - \xi_t|)^{-8N-2d-1} \\ &\leq C'_N 2^{-kN} (1 + 2^{\frac{k}{2}} |x' - x_t| + 2^{-\frac{k}{2}} |\xi' - \xi_t|)^{-2d-1}, \end{aligned} \quad (6-3)$$

where in deducing the second bound we used that the integrand vanishes unless $|\xi' - \xi_t| \geq \delta 2^{\frac{3k}{4}} - 2^{\frac{k}{2}}$. The desired L^2 bound then follows by the Schur test, using the fact that $(x, \xi) \rightarrow (x_t, \xi_t)$ is a volume-preserving diffeomorphism, which is homogeneous in ξ and bilipschitz on the cotangent bundle (uniformly over k). \square

Lemma 6.2. *Let $f_{x,\xi}(y) = 2^{\frac{kd}{4}} e^{i\langle \xi, y-x \rangle} f(2^{\frac{k}{2}}(y-x))$. Assume that f is a Schwartz function and $|\xi| \in [2^{k-1}, 2^{k+2}]$, and let $(x_t, \xi_t) = \chi_t(x, \xi)$. Then*

$$(B_k(t) f_{x,\xi})(z) = 2^{\frac{kd}{4}} e^{i\langle \xi_t, z-x_t \rangle} h(t, 2^{\frac{k}{2}}(z-x_t)),$$

where for all N, j, γ ,

$$|\partial_t^j \partial_z^\gamma h(t, z)| \leq C_{N,j,\gamma} 2^{\frac{k}{2}j} (1 + |z|)^{-N}.$$

For each N, j, γ , the constant $C_{N,j,\gamma}$ is bounded by a Schwartz seminorm of f , but is uniform over k, x, ξ .

Proof. Up to a factor of $(2\pi)^d$, the function $h(t, z)$ is given by the integral

$$\int e^{i\Phi(t,z,\eta)} b_k(t, x_t + 2^{-\frac{k}{2}}z, \xi + 2^{\frac{k}{2}}\eta) \psi_k(\xi + 2^{\frac{k}{2}}\eta) \hat{f}(\eta) d\eta,$$

where

$$\Phi(t, z, \eta) = \varphi_k(t, x_t + 2^{-\frac{k}{2}}z, \xi + 2^{\frac{k}{2}}\eta) - \langle x, \xi + 2^{\frac{k}{2}}\eta \rangle - \langle \xi_t, 2^{-\frac{k}{2}}z \rangle.$$

Since φ_k is the homogeneous generating function for χ_t , this equals

$$\varphi_k(t, x_t + 2^{-\frac{k}{2}}z, \xi + 2^{\frac{k}{2}}\eta) - \varphi_k(t, x_t, \xi) - 2^{\frac{k}{2}}\eta \cdot (\nabla_\eta \varphi_k)(t, x_t, \xi) - 2^{-\frac{k}{2}}z \cdot (\nabla_x \varphi_k)(t, x_t, \xi).$$

By [Corollary 3.2](#), [Theorem 4.1](#), [Corollary 4.3](#), and (4-4), the following estimates hold on the support of the integrand:

$$|\partial_t^j \partial_z^\beta \partial_\eta^\alpha \Phi(z, \eta)| \leq C_{\alpha, \beta, j} 2^{\frac{k}{2}j} \quad \text{if } |\alpha| + |\beta| \geq 2.$$

As Φ vanishes to second order at $z = \eta = 0$, on the region of integration we have

$$\begin{aligned} |\partial_t^j \Phi(z, \eta)| &\leq C_j 2^{\frac{k}{2}j} (1 + |z| + |\eta|)^2, \\ |\partial_t^j \nabla_{z, \eta} \Phi(z, \eta)| &\leq C_j 2^{\frac{k}{2}j} (1 + |z| + |\eta|). \end{aligned}$$

By [Theorem 3.3](#) we have $|\nabla_z \nabla_\eta \Phi(z, \eta) - \mathbf{I}| \lesssim c_d$, and since $|\nabla_\eta^2 \Phi(y, \eta)| \leq C$, we deduce that $|z| \leq C(|\nabla_\eta \Phi(z, \eta)| + |\eta|)$ and thus

$$\frac{1}{1 + |\nabla_\eta \Phi(y, \eta)|^2} \leq C \frac{1 + |\eta|^2}{1 + |z|^2}.$$

By [Corollary 3.2](#), [Theorem 5.1](#), and (6-2), since $|\xi + 2^{\frac{k}{2}}\eta| \approx 2^k$ we have

$$|\partial_t^j \partial_z^\beta \partial_\eta^\alpha (b_k(t, x_t + 2^{-\frac{k}{2}}z, \xi + 2^{\frac{k}{2}}\eta) \psi_k(\xi + 2^{\frac{k}{2}}\eta) \hat{f}(\eta))| \leq C_{N, \alpha, \beta, j} 2^{\frac{k}{2}j} (1 + |\eta|)^{-N}.$$

Integrating by parts with respect to the vector field

$$L = \frac{1 - i \nabla_\eta \Phi(z, \eta) \cdot \nabla_\eta}{1 + |\nabla_\eta \Phi(z, \eta)|^2}$$

then leads to the bounds on $\partial_t^j \partial_z^\beta h$ in the statement. □

The same argument also shows that the kernel of $T_k B_k(t) T_k^*$ satisfies (6-3) with $N = 0$, and in particular $B_k(t)$ is bounded on $L^2(\mathbb{R}^d)$, uniformly over k and $|t| \leq 1$. By applying [Lemma 6.1](#) with $a_1(\eta) = 1$ on the support of $\beta_k(\eta)$, and $a_2(\eta)$ supported in the annulus $|\eta| \in [2^{k-1}, 2^{k+2}]$, we then obtain the following by an orthogonality argument.

Lemma 6.3. *For all $s \in \mathbb{R}$ we have $\|\sum_{k=2}^\infty B_k(t) f\|_{H^s} \leq C_s \|f\|_{H^s}$, uniformly over $|t| \leq 1$.*

We can now show that $W(t)$ defined above is an approximate evolution operator for P .

Lemma 6.4. *Let $W(t)$ be defined by (6-1). Then*

$$(\partial_t + iP)W(t) = \sum_{k=2}^\infty B_k(t) + R(t),$$

where $R(t)$ is an integral kernel operator with kernel K satisfying

$$|\partial_x^\alpha \partial_y^\beta K(t, x, y)| \leq C_{N, \alpha, \beta} (1 + |x - y|)^{-N}.$$

In particular, $\|R(t) f\|_{H^N} \leq C_N \|f\|_{H^{-N}}$ for all N , uniformly over $|t| \leq 1$.

Proof. We take $a_k(\eta)$ supported in $\{\frac{4}{5}2^k < |\eta| < \frac{5}{4}2^{k+1}\}$, and equal to 1 where $\{\frac{7}{8}2^k < |\eta| < \frac{8}{7}2^{k+1}\}$, satisfying (6-2) with constants C_α independent of k . For c_d small enough, the condition of Lemma 6.1 with $a_1 = \psi_k$ and $a_2 = a_k$ is satisfied for all t with $|t| \leq 1$. We need show that the operator

$$\sum_{k=2}^{\infty} (P - \tilde{P}_k) W_k(t) = \sum_{k=2}^{\infty} (P - \tilde{P}_k)(1 - a_k(D)) W_k(t)$$

satisfies the conditions for $R(t)$, since $P \circ (\psi_0(D) + \psi_1(D))$ does. It suffices to show we can write

$$(P - \tilde{P}_k) W_k(t) = \text{Op}(R_k) \circ \psi_k(D),$$

with $R_k(t, x, y)$ an integral kernel satisfying, for all N ,

$$|\partial_x^\alpha \partial_y^\beta R_k(t, x, y)| \leq C_{N,\alpha,\beta} 2^{-kN} (1 + |x - y|)^{-N}.$$

Observe that $R_k = T_k^* \text{Op}(K_k) T_k$, where K_k satisfies (6-3), and vanishes unless $|\xi| \in [\frac{4}{5}2^k, \frac{5}{4}2^{k+1}]$ and $|\xi'| \notin [\frac{7}{8}2^k, \frac{8}{7}2^{k+1}]$. For c_d small this implies $|\xi_t| \in [\frac{5}{6}2^k, \frac{6}{5}2^{k+1}]$; hence $|\xi' - \xi_t| \geq 2^{k-4}$. Since $|x - x_t| \leq 2$ for $|t| \leq 1$, we have for all N

$$|K_k(t, x', \xi'; x, \xi)| \leq C_N 2^{-kN} (1 + 2^{-\frac{k}{2}} |\xi'|)^{-N} (1 + |x - x'|)^{-N}.$$

The operator T_k is given by a kernel satisfying for all N

$$|\partial_y^\alpha T_k(x, \xi; y)| \leq C_{N,\alpha} 2^{k(\frac{|\alpha|}{2} + \frac{d}{4})} (1 + 2^{\frac{k}{2}} |x - y|)^{-N}.$$

Since the volume of integration in ξ is less than $C_d 2^{kd}$, the estimate for $R_k(t, x, y)$ follows by composition. \square

We now write

$$\int R_k(t, x, y) (\psi_k(D) f)(y) dy = \frac{1}{(2\pi)^n} \int e^{i\varphi_k(t,x,\eta)} r_k(t, x, \eta) \psi_k(\eta) \hat{f}(\eta) d\eta,$$

with

$$r_k(t, x, \eta) = e^{-i\varphi_k(t,x,\eta)} \int R_k(t, x, y) e^{-i\langle y, \eta \rangle} dy.$$

Then for all N

$$|\partial_x^\beta \partial_\eta^\alpha r_k(t, x, \eta)| \leq C_{\alpha,\beta,N} 2^{-kN}, \quad 2^{k-1} \leq |\eta| \leq 2^{k+2},$$

and we can incorporate r_k into b_k , and hence $R_k(t)$ into $B_k(t)$. Thus we can write

$$(\partial_t + iP)W(t) = \sum_{k=2}^{\infty} B_k(t) + P \circ (\psi_0(D) + (D)) \equiv B(t).$$

We now can generate the exact wave group $E(t)$ for $\partial_t + iP$ by iteration,

$$E(t) = W(t) - \int_0^t W(t-s)B(s) ds + \int_0^t \int_0^s W(t-s)B(s-r)B(r) dr ds - \dots.$$

To write the iteration more concisely, let $\Lambda^m \subset \mathbb{R}_+^{m+1}$ be the m -simplex, consisting of $\mathbf{r} = (r_1, \dots, r_{m+1})$ with $r_j > 0$ for all j , and with $r_1 + \dots + r_{m+1} = 1$. Let $d\mathbf{r}$ be the measure on Λ^m induced by projection onto (r_1, \dots, r_m) . Then

$$E(t) = \sum_{m=0}^{\infty} (-t)^m \int_{\Lambda^m} W(tr_{m+1})B(tr_m) \cdots B(tr_1) d\mathbf{r}. \quad (6-4)$$

If C_s is an upper bound for the $H^s(\mathbb{R}^d)$ operator norm of both $W(t)$ and $B(t)$ for all $|t| \leq 1$, then the m -th term has $H^s(\mathbb{R}^d)$ operator norm at most $C_s^{m+1} t^m / m!$, and the following theorem holds.

Theorem 6.5. *The expansion (6-4) converges uniformly over $|t| \leq 1$, in the operator norm topology on $H^s(\mathbb{R}^d)$ for all $s \in \mathbb{R}$. The limit $E(t)$ is a one-parameter group of L^2 -unitary operators, and for $f \in H^s$, $F \in L^1([-1, 1], H^s)$, the solution to $(\partial_t + iP)u = F$, $u(0, \cdot) = f$ is given by*

$$u(t, \cdot) = E(t)f + \int_0^t E(t-s)F(s, \cdot) ds.$$

Our next two results show that if we localize $E(t)$ on the right to frequency scale 2^k , then modulo a smoothing operator error one can localize each of the terms $W(tr_j)$ and $B(tr_j)$ in (6-4) to frequencies of scale 2^k . We use the notation $\tilde{\psi}_k = \psi_{k-1} + \psi_k + \psi_{k+1}$, and define

$$\begin{aligned} \tilde{W}_k(t) &= \tilde{\psi}_k(D)(W_{k-1} + W_k + W_{k+1})(t), \\ \tilde{B}_k(t) &= \tilde{\psi}_k(D)(B_{k-1} + B_k + B_{k+1})(t). \end{aligned} \quad (6-5)$$

Lemma 6.6. *If $m+1 \leq 2^{\frac{k}{4}}$, then for all $N \geq 0$ the operator*

$$R_{k,\mathbf{r}}(t) = W(tr_{m+1})B(tr_m) \cdots B(tr_1)\psi_k(D) - \tilde{W}_k(tr_{m+1})\tilde{B}_k(tr_m) \cdots \tilde{B}_k(tr_1)\psi_k(D)$$

satisfies the following, with constant C_N independent of m, t, k , and \mathbf{r} :

$$\|R_{k,\mathbf{r}}(t)f\|_{H^N} \leq C_N 2^{-kN} \|f\|_{H^{-N}}.$$

Proof. Fix t and \mathbf{r} , and without loss of generality assume $t \geq 0$. We introduce a family of intermediate cutoffs $\psi_{k,j}(D)$ for $1 \leq j \leq m$, which depend on $t\mathbf{r}$. Define points $\frac{10}{9} \leq p'_{j-1} < p_j < p'_j \leq \frac{5}{4}$ as follows. Take c_0 and c_1 such that $p_0 = e^{c_0} = \frac{10}{9}$, and $e^{c_0+2c_1} = \frac{5}{4}$. For $j \geq 0$ we set

$$p_j = e^{c_0+c_1(r_1+\dots+r_j)t+c_1j} 2^{-\frac{k}{4}}, \quad p'_j = e^{c_0+c_1(r_1+\dots+r_j)t+c_1(j+\frac{1}{2})} 2^{-\frac{k}{4}}.$$

Thus ψ_k is supported where $|\eta| \in [p_0^{-1}2^k, p_0 2^{k+1}]$, and $\tilde{\psi}_k(\eta) = 1$ on the set $\{\eta: |\eta| \in [p'_m{}^{-1}2^k, p'_m 2^{k+1}]\}$. Also,

$$|p'_j - p_j| \geq \frac{1}{2} c_1 2^{-\frac{k}{4}}, \quad |p_{j+1} - p'_j| \geq c_1 r_{j+1} t + \frac{1}{2} c_1 2^{-\frac{k}{4}}.$$

Let $\psi_{k,0} = \psi_k$. By the comments following (6-2) we can construct functions $\psi_{k,j}(\xi)$ for $j \geq 1$ that satisfy (6-2), with constants C_α that depend only on the dimension d , such that

$$\begin{aligned} \text{supp}(\psi_{k,j}) &\subset \{\eta: |\eta| \in [p'_j{}^{-1}2^k, p'_j 2^{k+1}]\}, \quad j \geq 0, \\ \psi_{k,j}(\eta) &= 1 \quad \text{if } |\eta| \in [p_j{}^{-1}2^k, p_j 2^{k+1}], \quad j \geq 1. \end{aligned}$$

Let $c'_d = \sup_{x,\xi} (|\xi|^{-1} |\nabla_x p_k(x, \xi)|) \lesssim c_d$. Then for solutions to the Hamiltonian flow,

$$\exp(-c'_d tr_j) |\xi(s)| \leq |\xi(s + tr_j)| \leq \exp(c'_d tr_j) |\xi(s)|.$$

Then if $c'_d \leq c_1$, the condition of [Lemma 6.1](#) with $\delta = \frac{1}{4}c_1$ is satisfied for $a_2 = \psi_{k,j}$ and $a_1 = \psi_{k,j-1}$. Thus [Lemma 6.1](#) yields, for $j \geq 1$,

$$\|(1 - \psi_{k,j}(D))B(tr_j)\psi_{k,j-1}(D)\|_{H^s \rightarrow H^s} \leq C_{s,N} 2^{-kN} \quad \text{for all } s, N.$$

Since $B(t)\psi_{k,j}(D) = \tilde{B}_k(t)\psi_{k,j}(D)$, and the number of terms is at most $m \leq 2^{\frac{k}{4}}$, we can apply this repeatedly to write

$$\begin{aligned} W(tr_{m+1})B(tr_m) \cdots B(tr_1)\psi_k(D) \\ = \tilde{W}_k(tr_{m+1})\psi_{k,m}(D)\tilde{B}_k(tr_m) \cdots \psi_{k,1}(D)\tilde{B}_k(tr_1)\psi_k(D) + R_{k,r}(t), \end{aligned}$$

where $\|R_{k,r}(t)\|_{H^s \rightarrow H^s} \leq C_{s,N} 2^{-Nk}$ for all s, N . We then prove [Lemma 6.6](#) by observing that the same steps let us write

$$\begin{aligned} \tilde{W}_k(tr_{m+1})\psi_{k,m}(D)\tilde{B}_k(tr_m) \cdots \psi_{k,1}(D)\tilde{B}_k(tr_1)\psi_k(D) \\ = \tilde{W}_k(tr_{m+1})\tilde{B}_k(tr_m) \cdots \tilde{B}_k(tr_1)\psi_k(D) + R_{k,r}(t) \end{aligned}$$

for a similar $R_{k,r}(t)$. Since $R_{k,r}(t)$ is localized on the right at frequency 2^k , it follows that

$$\|R_{k,r}(t)\|_{H^{-N} \rightarrow H^N} \leq C_N 2^{-kN}$$

for all N . □

Corollary 6.7. *One can write*

$$E(t) = \sum_{k=0}^{\infty} \sum_{m=0}^{2^{k/4}} (-t)^m \int_{\Lambda^m} \tilde{W}_k(tr_{m+1})\tilde{B}_k(tr_m) \cdots \tilde{B}_k(tr_1)\psi_k(D) d\mathbf{r} + R(t),$$

where for all N we have $\|R(t)f\|_{H^N} \leq C_N \|f\|_{H^{-N}}$, uniformly over $|t| \leq 1$.

Proof. Consider

$$\sum_{m=2^{k/4}}^{\infty} (-t)^m \int_{\Lambda^m} W(tr_{m+1})B(tr_m) \cdots B(tr_1)\psi_k(D) d\mathbf{r}.$$

For $|t| \leq 1$ and all N , the $H^N \rightarrow H^N$ operator norm of this sum is bounded by the sum

$$\sum_{m \geq 2^{k/4}} C_N^{m+1}/m! \leq C_N 2^{-3kN}.$$

It is localized on the right at frequency 2^k , and thus maps $H^{-N} \rightarrow H^N$ with norm $\leq C_N 2^{-kN}$. □

The arguments leading to [Lemma 6.6](#) apply equally well to conic localization. We take a finite partition of unity on $\mathbb{R}^d \setminus \{0\}$,

$$1 = \sum_{\omega \in \mathbb{E}} a_\omega(D), \quad \text{supp}(a_\omega(\eta)) \subset \left\{ \eta : \left| \omega - \frac{\eta}{|\eta|} \right| \leq \frac{1}{32} \right\}.$$

Let $\tilde{a}_\omega(\eta)$ be a smooth, homogeneous cutoff such that

$$\tilde{a}_\omega(\eta) = 1 \quad \text{if} \quad \left| \omega - \frac{\eta}{|\eta|} \right| \leq \frac{1}{24}, \quad \text{supp}(\tilde{a}_\omega) \subset \left\{ \eta : \left| \omega - \frac{\eta}{|\eta|} \right| \leq \frac{1}{16} \right\}.$$

We define an angularly localized version of \tilde{W}_k , recalling (6-5),

$$\begin{aligned} \tilde{W}_k^\omega(t) &= \tilde{a}_\omega(D) \tilde{W}_k(t) \tilde{a}_\omega(D), \\ \tilde{B}_k^\omega(t) &= \tilde{a}_\omega(D) \tilde{B}_k(t) \tilde{a}_\omega(D). \end{aligned} \tag{6-6}$$

Then,

$$\begin{aligned} R_{k,r}^\omega(t) &= W(tr_{m+1})B(tr_m) \cdots B(tr_1)a_\omega(D)\psi_k(D) \\ &\quad - \tilde{W}_k^\omega(tr_{m+1})\tilde{B}_k^\omega(tr_m) \cdots \tilde{B}_k^\omega(tr_1)a_\omega(D)\psi_k(D) \end{aligned} \tag{6-7}$$

satisfies the conclusion of Lemma 6.6, and consequently, with $R(t)$ as in Corollary 6.7,

$$E(t) = \sum_{k=0}^{\infty} \sum_{\omega \in \Xi} \tilde{E}_k^\omega(t) + R(t),$$

where we define

$$\tilde{E}_k^\omega(t) = \sum_{m=0}^{2^{k/4}} (-t)^m \int_{\Lambda^m} \tilde{W}_k^\omega(tr_{m+1})\tilde{B}_k^\omega(tr_m) \cdots \tilde{B}_k^\omega(tr_1)a_\omega(D)\psi_k(D). \tag{6-8}$$

Lemma 6.8. *Let $f_{x,\xi}(y) = 2^{\frac{kd}{4}} e^{i\langle \xi, y-x \rangle} f(2^{\frac{k}{2}}(y-x))$. Assume that f is a Schwartz function, and let $(x_t, \xi_t) = \chi_t(x, \xi)$. Then one can write*

$$(\tilde{E}_k^\omega(t) f_{x,\xi})(z) = 2^{\frac{kd}{4}} e^{i\langle \xi_t, z-x_t \rangle} h(t, 2^{\frac{k}{2}}(z-x_t)) \equiv h(t, \cdot)_{x_t, \xi_t},$$

where for all N ,

$$|\partial_t^j \partial_z^\gamma h(t, z)| \leq C_{N,j,\gamma} 2^{\frac{k}{2}j} (1+|z|)^{-N}.$$

For each N, j, γ , the constant $C_{N,j,\gamma}$ is bounded by a Schwartz seminorm of f , but is uniform over k, x, ξ .

Proof. Let $K(s, y, \eta; x, \xi)$ denote the integral kernel of $T_k \tilde{B}_k^\omega(s) T_k^*$. Following the proof of (6-3) we can bound, with C_N uniform over k and s ,

$$|K(s, y, \eta; x, \xi)| \leq C_N (1 + 2^{\frac{k}{2}}|y-x_s| + 2^{-\frac{k}{2}}|\eta-\xi_s|)^{-N}.$$

Furthermore, this kernel vanishes unless $2^{k-2} \leq |\eta|, |\xi| \leq 2^{k+3}$. By the bilipschitz property of the Hamiltonian flow, for such η, ξ we have

$$|y_s - x_s| + 2^{-k}|\eta_s - \xi_s| \leq A|y-x| + 2^{-k}A|\eta-\xi|. \tag{6-9}$$

For $N \geq 2d+1$, we then bound the kernel of

$$T_k \tilde{B}_k^\omega(tr_1) \tilde{B}_k^\omega(tr_2) T_k^* = (T_k \tilde{B}_k^\omega(tr_1) T_k^*) (T_k \tilde{B}_k^\omega(tr_2) T_k^*)$$

by C_N^2 multiplied by the quantity

$$\begin{aligned} & \int (1 + 2^{\frac{k}{2}}|y - z_{tr_1}| + 2^{-\frac{k}{2}}|\eta - \xi_{tr_1}|)^{-N} (1 + 2^{\frac{k}{2}}|z - x_{tr_2}| + 2^{-\frac{k}{2}}|\zeta - \xi_{tr_2}|)^{-N} dz d\zeta \\ & \leq A_N (1 + 2^{\frac{k}{2}}|y - x_{t(r_1+r_2)}| + 2^{-\frac{k}{2}}|\eta - \xi_{t(r_1+r_2)}|)^{-N}. \end{aligned}$$

Similarly, for $\mathbf{r} \in \Lambda^m$ the operator $T_k \tilde{W}_k^\omega(tr_{m+1}) \tilde{B}_k^\omega(tr_m) \cdots \tilde{B}_k^\omega(tr_1) T_k^*$ has kernel bounded by

$$C_N (A_N C_N)^m (1 + 2^{\frac{k}{2}}|y - x_t| + 2^{-\frac{k}{2}}|\eta - \xi_t|)^{-N},$$

and summing over m gives the following bounds for the kernel of $T_k \tilde{E}_k^\omega(t) T_k^*$,

$$|\tilde{K}_k^\omega(t, y, \eta; x, \xi)| \leq C_N e^{tA_N C_N} (1 + 2^{\frac{k}{2}}|y - x_t| + 2^{-\frac{k}{2}}|\eta - \xi_t|)^{-N}.$$

Let $F = T_k(f_{x, \xi})$. Then

$$|F(\bar{x}, \bar{\xi})| \leq C_N (1 + 2^{\frac{k}{2}}|\bar{x} - x| + 2^{-\frac{k}{2}}|\bar{\xi} - \xi|)^{-N}.$$

Then $(\tilde{E}_k^\omega(t) f_{x, \xi})(z)$ is equal to

$$2^{\frac{k_d}{4}} \int \tilde{K}_k^\omega(t, y, \eta; \bar{x}, \bar{\xi}) F(\bar{x}, \bar{\xi}) e^{i\langle \eta, z-y \rangle} g(2^{\frac{k}{2}}(z-y)) d\bar{x} d\bar{\xi} dy d\eta.$$

The change of variables

$$\begin{aligned} (y, \eta) & \rightarrow (x_t + 2^{-\frac{k}{2}}y, \xi_t + 2^{\frac{k}{2}}\eta), \\ (\bar{x}, \bar{\xi}) & \rightarrow (x + 2^{-\frac{k}{2}}\bar{x}, \xi + 2^{\frac{k}{2}}\bar{\xi}) \end{aligned}$$

shows that $h(t, z) = 2^{-\frac{k_d}{4}} e^{-i2^{-k/2}\langle \xi_t, z \rangle} (\tilde{E}_k^\omega(t) f_{x, \xi})(x_t + 2^{-\frac{k}{2}}z)$ is equal to

$$\begin{aligned} & \int \tilde{K}_k^\omega(t, x_t + 2^{-\frac{k}{2}}y, \xi_t + 2^{\frac{k}{2}}\eta; x + 2^{-\frac{k}{2}}\bar{x}, \xi + 2^{\frac{k}{2}}\bar{\xi}) F(x + 2^{-\frac{k}{2}}\bar{x}, \xi + 2^{\frac{k}{2}}\bar{\xi}) \\ & \quad \times e^{-i2^{-\frac{k}{2}}\langle \xi_t, y \rangle} e^{i\langle \eta, z-y \rangle} g(z-y) d\bar{x} d\bar{\xi} dy d\eta. \end{aligned}$$

By the bilipschitz property (6-9) of χ_t we have

$$|y| + |\eta| \leq A|\bar{x}| + A|\bar{\xi}| + 2^{\frac{k}{2}}|(x_t + 2^{-\frac{k}{2}}y) - (x + 2^{-\frac{k}{2}}\bar{x})_t| + 2^{-\frac{k}{2}}|(\xi_t + 2^{\frac{k}{2}}\eta) - (\xi + 2^{\frac{k}{2}}\bar{\xi})_t|,$$

and conclude that

$$|\tilde{K}_k^\omega(t, x_t + 2^{-\frac{k}{2}}y, \xi_t + 2^{\frac{k}{2}}\eta; x + 2^{-\frac{k}{2}}\bar{x}, \xi + 2^{\frac{k}{2}}\bar{\xi})| \leq C_N (1 + |y| + |\eta|)^{-N} (1 + |\bar{x}| + |\bar{\xi}|)^N.$$

Together with the bound

$$|F(x + 2^{-\frac{k}{2}}\bar{x}, \xi + 2^{\frac{k}{2}}\bar{\xi})| \leq C_N (1 + |\bar{x}| + |\bar{\xi}|)^{-2N},$$

this leads to the following estimates on $\partial_z^\gamma h(t, z)$, which is the case $j = 0$:

$$|\partial_z^\gamma h(t, z)| \leq C_{N, \gamma} (1 + |z|)^{-N}. \quad (6-10)$$

The constant $C_{N, \gamma}$ is seen to be bounded by a Schwartz seminorm of f , but uniform over k, x, ξ .

To handle time derivatives we proceed by induction, and assume the estimates on $\partial_t^i \partial_z^\gamma h(t, z)$ hold for $0 \leq i \leq j$, and all γ . We write

$$\tilde{E}_k^\omega(t) f_{x,\xi} = \tilde{W}_k^\omega(t) f_{x,\xi} + \int_0^t \tilde{W}_k^\omega(t-s) \tilde{E}_k^\omega(s) f_{x,\xi} ds,$$

where on the right the term $\tilde{E}_k^\omega(t)$, defined in (6-8), has upper summation limit reduced by 1. This does not affect the validity of (6-10), since the proof of (6-10) is done separately for each value of m . By Lemma 6.2 the first term satisfies the conditions of the statement, since the proof of that lemma works equally well for $B_k(t)$ replaced by $\tilde{W}_k^\omega(t)$. The desired estimates on h are then a consequence of the following, for the given value of j and all γ :

$$\left| (\partial_z - i\xi_t)^\gamma (\partial_t + ip_k(x_t, \xi_t))^{j+1} \int_0^t \tilde{W}_k^\omega(t-s) \tilde{E}_k^\omega(s) f_{x,\xi} ds \right| \leq C_{N,j+1,\gamma} 2^{\frac{kd}{4}} 2^{\frac{k}{2}(|\gamma|+j+1)} (1 + 2^{\frac{k}{2}} |z - x_t|)^{-N}. \quad (6-11)$$

This is seen by noting that

$$\begin{aligned} e^{-i\langle \xi_t, z - x_t \rangle} (\partial_t + ip_k(x_t, \xi_t)) (e^{i\langle \xi_t, z - x_t \rangle} h(t, 2^{\frac{k}{2}}(z - x_t))) \\ = (\partial_t h)(t, 2^{\frac{k}{2}}(z - x_t)) - (i\nabla_x p_k(x_t, \xi_t) \cdot (z - x_t) h + 2^{\frac{k}{2}} (\nabla_\xi p_k)(x_t, \xi_t) \cdot \nabla_z h)(t, 2^{\frac{k}{2}}(z - x_t)). \end{aligned}$$

The latter terms are controlled by the spatial derivative bounds on h , and their time derivatives are controlled by the bounds

$$|\partial_t^i (\nabla_x p_k)(x_t, \xi_t)| \leq C_i 2^{k + \frac{k}{2}i}, \quad |\partial_t^i (\nabla_\xi p_k)(x_t, \xi_t)| \leq C_i 2^{\frac{k}{2}i},$$

which follow by Corollary 3.2 and (2-7).

To establish (6-11) we expand

$$\begin{aligned} (\partial_t + ip_k(x_t, \xi_t))^{j+1} \int_0^t \tilde{W}_k^\omega(t-s) \tilde{E}_k^\omega(s) f_{x,\xi} ds \\ = \sum_{i=0}^j (\partial_t + ip_k(x_t, \xi_t))^{j-i} [(\partial_r + ip_k(x_{t+r}, \xi_{t+r}))^i \tilde{W}_k^\omega(r) \tilde{E}_k^\omega(t) f_{x,\xi}]_{r=0} \\ + \int_0^t (\partial_t + ip_k(x_t, \xi_t))^{j+1} \tilde{W}_k^\omega(t-s) \tilde{E}_k^\omega(s) f_{x,\xi} ds. \end{aligned}$$

The latter term on the right is handled by Lemma 6.2, since we have already shown that $\tilde{E}_k^\omega(s) f_{x,\xi} = f(s, \cdot)_{x_s, \xi_s}$, where $f(s, \cdot)$ is a bounded family of Schwartz functions. The first term on the right expands into a sum of terms

$$[\partial_t^n (\partial_r + ip_k(x_{t+r}, \xi_{t+r}))^i \tilde{W}_k^\omega(r)]_{r=0} (\partial_t + ip_k(x_t, \xi_t))^{j-n-i} \tilde{E}_k^\omega(t) f_{x,\xi}. \quad (6-12)$$

We can write $[\partial_t^n (\partial_r + ip_k(x_{t+r}, \xi_{t+r}))^i \tilde{W}_k^\omega(r)]_{r=0}$ as a sum of terms

$$(\partial_t^{n_1} p_k(x_t, \xi_t)) \cdots (\partial_t^{n_m} p_k(x_t, \xi_t)) [(\partial_r + ip_k(x_{t+r}, \xi_{t+r}))^l \tilde{W}_k^\omega(r)]_{r=0},$$

where $n_1 + \dots + n_m + m + l = n + i$, and each $n_j \geq 1$. By [Lemma 6.2](#) and the induction assumption we can write

$$[(\partial_r + i p_k(x_{t+r}, \xi_{t+r}))^l \tilde{W}_k^\omega(r)]_{r=0} (\partial_t + i p_k(x_t, \xi_t))^{j-n-i} \tilde{E}_k^\omega(t) f_{x, \xi} = 2^{\frac{k}{2}(l+j-n-i)} f(t, \cdot)_{x_t, \xi_t}$$

for a bounded family of Schwartz functions $f(t, \cdot)$. The estimates

$$|\partial_t^{n_j} p_k(x_t, \xi_t)| \leq C_{n_j} 2^{\frac{k}{2}(n_j+1)}, \quad n_j \geq 1,$$

then show that the term in [\(6-12\)](#) is of the form $2^{\frac{k}{2}j} f(t, \cdot)_{x_t, \xi_t}$ for a bounded family of Schwartz functions $f(t, \cdot)$, which implies [\(6-11\)](#). \square

We use this to establish sideways energy estimates for $E(t)$, which state that if the initial data f is microlocalized to frequencies within a small angle of the codirection ω , then the L^2 norm of the restriction of $E(t)f$ to space-time hyperplanes perpendicular to ω is dominated by the L^2 norm of f . By rotation and translation invariance it suffices to consider $\omega = e_1$ and the plane $x_1 = 0$.

Theorem 6.9. *Suppose $\phi \in C_c^\infty((-\frac{1}{2}, \frac{1}{2}))$. Then*

$$\|\phi(t)(a_{e_1}(D)\psi_k(D)E(t)f)|_{x_1=0}\|_{L_{x'}^2, L_t^2} \leq C \|f\|_{L^2}$$

for a constant C that is independent of k .

Proof. By [Lemma 6.6](#) and the comments following [Corollary 6.7](#), it suffices to show that

$$\|\phi(t)(\tilde{E}_k^{e_1}(t)f)|_{x_1=0}\|_{L_{x'}^2, L_t^2} \leq C \|f\|_{L^2}.$$

For $\xi \in \mathbb{R}^d$ with $|\angle(\xi, e_1)| \leq \frac{1}{2}$ and $|s| \leq 2$, the null bicharacteristic curve $\gamma(s) \in (\mathbb{R}^{d+1})^*$ of $\tau + p_k(y, \eta)$ that passes over (x, ξ) at time $s = 0$ satisfies $\frac{4}{5} \leq |y'_1(s)| \leq \frac{5}{4}$. Consequently, if $|x_1| \leq \frac{3}{2}$ and $|\angle(\xi, e_1)| \leq \frac{1}{2}$ there is a unique value $s = s(x, \xi)$ in $\{s : |s| \leq 2\}$ such that $\gamma(s(x, \xi)) \in \{y_1 = 0\}$. We parametrize the cotangent bundle of $y_1 = 0$ by (t, y', τ, η') , and let T_k^0 be the wave packet transform acting on this plane. Observe that the integral kernel $\tilde{K}_k^{e_1}(t, y', \tau, \eta'; x, \xi)$ of $T_k^0(\phi(t)\tilde{E}_k^{e_1}(t)T_k^*)$ vanishes unless $|\angle(\xi, e_1)| \leq \frac{1}{2}$.

We show that if $|x_1| \leq \frac{3}{2}$, then

$$|\tilde{K}_k^{e_1}(t, y', \tau, \eta'; x, \xi)| \leq C_N (1 + 2^{\frac{k}{2}} |(t, y') - \Pi_{t, y'} \gamma(s(x, \xi))| + 2^{-\frac{k}{2}} |(\tau, \eta') - \Pi_{\tau, \eta'} \gamma(s(x, \xi))|)^{-N}. \quad (6-13)$$

The Schur test, and the fact that $(x, \xi) \rightarrow \gamma(s(x, \xi))|_{y_1=0}$ is a bilipschitz symplectic map, shows L^2 boundedness of $T_k^0 \phi(t) \tilde{E}_k^{e_1}(t) T_k^* \mathbb{1}_{|x_1| < \frac{3}{2}}$. We consider the case $|x_1| > \frac{3}{2}$ afterwards.

To prove [\(6-13\)](#), we use [Lemma 6.8](#) to express $\tilde{K}_k^{e_1}(t, y', \tau, \eta'; x, \xi)$ as

$$2^{\frac{kd}{2}} \int e^{i\langle \xi_s, (0, z') - x_s \rangle - i\tau(s-t) - i\langle \eta', z' - y' \rangle} h(s, 2^{\frac{k}{2}}((0, z') - x_s)) g(2^{\frac{k}{2}}(s-t, z' - y')) ds dz'.$$

Since γ is null, we have $\Pi_\tau \gamma(s) = -p(x_s, \xi_s) = -\langle \xi_s, \partial_s x_s \rangle$. We then note that

$$\begin{aligned} & (\partial_{z'} + i(\eta' - \xi'_s)) e^{i\langle \xi_s, (0, z') - x_s \rangle - i\tau(s-t) - i(\eta', z' - y')} \\ &= 0 \\ &= (\partial_s + i(\tau + p_k(x_s, \xi_s)) - i\langle \partial_s \xi_s, (0, z') - x_s \rangle) e^{i\langle \xi_s, (0, z') - x_s \rangle - i\tau(s-t) - i(\eta', z' - y')}. \end{aligned}$$

Applying each of $2^{-\frac{k}{2}} \partial_{z'}$, $2^{-\frac{k}{2}} \partial_s$, or $\langle \partial_s \xi_s, (0, z') - x_s \rangle$ to the amplitude term $h(\cdots)g(\cdots)$ preserves its form. An integration by parts argument, together with Schwartz bounds on h and g , then shows that the integral is dominated in absolute value by

$$C_N 2^{\frac{kd}{2}} \int (1 + 2^{-\frac{k}{2}} |\eta' - \xi'_s| + 2^{-\frac{k}{2}} |\tau + p_k(x_s, \xi_s)| + 2^{\frac{k}{2}} |s-t| + 2^{\frac{k}{2}} |z' - y'| + 2^{\frac{k}{2}} |(0, z') - x_s|)^{-N} ds dz'. \quad (6-14)$$

Note that $|(x_s)_1| \geq \frac{4}{5} |s - s(x, \xi)|$ as $|\partial_s(x_s)_1| \geq \frac{4}{5}$. Since $2^{-k} \xi_s$, $2^{-k} p_k(x_s, \xi_s)$, and x_s are all uniformly Lipschitz in s , the integral is in turn bounded by

$$\begin{aligned} C_{N+2n+1} 2^{\frac{kd}{2}} \int (1 + 2^{\frac{k}{2}} |s-t| + 2^{\frac{k}{2}} |z' - y'|)^{-2n-1} ds dz' \\ \times (1 + 2^{-\frac{k}{2}} |(\tau, \eta') - \Pi_{\tau, \eta'} \gamma(s(x, \xi))| + 2^{\frac{k}{2}} |(t, y') - \Pi_{t, y'} \gamma(s(x, \xi))|)^{-N}, \end{aligned}$$

which yields the estimate (6-13) for $|x_1| \leq \frac{3}{2}$.

If $|x_1| \geq \frac{3}{2}$, $|t| \leq 1$, we have $|(x_t)_1| \geq \frac{1}{6} |x_1| \geq \frac{1}{4} |t|$. By a proof similar to above, (6-14) then leads to the following bounds:

$$|\tilde{K}_k^{e_1}(t, y', \tau, \eta'; x, \xi) \mathbb{1}_{|x_1| \geq \frac{3}{2}}| \leq C_N (1 + 2^{-\frac{k}{2}} |\tau + p_k(x, \xi)| + 2^{-\frac{k}{2}} |\eta' - \xi'| + 2^{\frac{k}{2}} |x_1| + 2^{\frac{k}{2}} |y' - x'|)^{-N}.$$

Here we use, for example, that

$$|x_1| + |y' - x'| \lesssim |(x_1)_t| + |y' - x'| \lesssim |(x_1)_s| + |y' - x'_s| + |s - t|$$

by the above. The Schur test, and the fact that $(x, \xi_1, \xi') \rightarrow (x, p_k(x, \xi), \xi')$ is a diffeomorphism on $|\angle(\xi, e_1)| \leq \frac{1}{2}$, proves L^2 boundedness of the operator $T_k^0 \phi(t) \tilde{E}_k^{e_1}(t) T_k^* \mathbb{1}_{|x_1| \geq \frac{3}{2}}$. \square

We now turn to the proof of (2-14) for the operator $E(t)$, that is,

$$\begin{aligned} & \|\langle D \rangle^{-s} E(t) f\|_{L_t^q L_x^r([0,1] \times \mathbb{R}^d)} \leq C \|f\|_{L^2(\mathbb{R}^d)}, \\ & \left\| \langle D \rangle^{-s} \int_0^t E(t-s) F(s, \cdot) ds \right\|_{L_t^q L_x^r([0,1] \times \mathbb{R}^d)} \leq C \|\langle D \rangle^{1-s} F\|_{L_t^{\tilde{q}'} L_x^{\tilde{r}'}}([0,1] \times \mathbb{R}^d) \end{aligned}$$

for $s, q, \tilde{q}, r, \tilde{r}$ satisfying the conditions of Theorem 1.1. A consequence of Corollary 6.7 is that

$$a_\omega(D) \psi_k(D) E(t) = a_\omega(D) \psi_k(D) E(t) a'_\omega(D) \psi'_k(D) + a_\omega(D) \psi_k(D) R(t),$$

with $R(t)$ a smoothing operator, and $a'_\omega(\eta) \psi'_k(\eta)$ a $S_{1,0}^0$ cutoff to a $\delta 2^k$ neighborhood of the support of $a_\omega(\eta) \psi_k(\eta)$. Since $q, r \geq 2 \geq \tilde{q}', \tilde{r}'$, it suffices by Littlewood–Paley theory to prove that, for a constant C independent of k ,

$$\|a_\omega(D) \psi_k(D) E(t) f\|_{L_t^q L_x^r([0,1] \times \mathbb{R}^d)} \leq C 2^{ks} \|f\|_{L^2(\mathbb{R}^d)},$$

and that

$$\left\| \int_0^t a_\omega(D) \psi_k(D) E(t-s) a'_\omega(D) \psi'_k(D) F(s, \cdot) ds \right\|_{L_t^q L_x^r([0,1] \times \mathbb{R}^d)} \leq C 2^k \|F\|_{L_t^{\bar{q}'} L_x^{\bar{r}'}([0,1] \times \mathbb{R}^d)}.$$

Since $E(t)E^*(s) = E(t-s)$, we can apply [Keel and Tao 1998, Theorem 1.2] with a scaling of (t, x) by 2^k to conclude that these are implied by the estimate

$$\|a'_\omega(D) \psi'_k(D) E(t-s) a'_\omega(D) \psi'_k(D) f\|_{L^\infty(\mathbb{R}^d)} \leq C 2^{kd} (1 + 2^k |t-s|)^{-\frac{d-1}{2}} \|f\|_{L^1(\mathbb{R}^d)}.$$

By Corollary 6.7 and the comments following it, this estimate in turn is implied by proving the same estimate with $E(t-s)$ replaced by $\tilde{E}_k^\omega(t-s)$. Letting $\tilde{K}_k^\omega(t, x, y)$ be the integral kernel of $\tilde{E}_k^\omega(t)$, we need to show that

$$|\tilde{K}_k^\omega(t, x, y)| \leq C 2^{kd} (1 + 2^k |t|)^{-\frac{d-1}{2}}, \quad |t| \leq 1.$$

We in fact prove a stronger estimate, which captures the decay of the fundamental solution away from the light cone. We will show in Section 7 that, for all N , with $S_t(y)$ the geodesic sphere of radius $|t|$ centered at y , and $\text{dist}(x, S_t(y))$ the geodesic distance in \mathfrak{g}_k of x to the set $S_t(y)$,

$$|\tilde{K}_k^\omega(t, x, y)| \leq C_N 2^{kd} (1 + 2^k |t|)^{-\frac{d-1}{2}} (1 + 2^k |\text{dist}(x, S_t(y))|)^{-N}, \quad (6-15)$$

which will imply (2-14) by the above.

By similar steps and duality, estimate (2-15) reduces to proving that, for q_d and s_d as in Theorem 1.2, and $\phi \in C_c^\infty((-\frac{1}{2}, \frac{1}{2}))$,

$$\left\| \phi(t) \int \tilde{E}_k^\omega(t-s) \phi(s) F(s, \cdot) ds \right\|_{L_x^{q_d} L_t^2} \leq C 2^{2ks_d} \|F\|_{L_y^{q'_d} L_s^2}.$$

It suffices to prove this for $\omega = e_1$. We deduce from (6-15) that

$$|\tilde{K}_k^{e_1}(t, x, y)| \leq C_N 2^{kd} (1 + 2^k |x-y|)^{-\frac{d-1}{2}} (1 + 2^k |t - \text{dist}(x, y)|)^{-N},$$

which uses that $\text{dist}(x, S_t(y)) \geq |t - \text{dist}(x, y)|$ and $\text{dist}(x, y) \approx |x - y|$. As a consequence, letting $x = (x_1, x')$, we have

$$\begin{aligned} \left\| \phi(t) \int \tilde{K}_k^{e_1}(t-s, x_1, x', y_1, y') \phi(s) F(s, y_1, y') ds dy' \right\|_{L_{x'}^\infty L_t^2} \\ \leq C 2^{k(d-1)} (1 + 2^k |x_1 - y_1|)^{-\frac{d-1}{2}} \|F(\cdot, y_1, \cdot)\|_{L_{y'}^1 L_s^2}. \end{aligned}$$

On the other hand, writing $E(t-s) = E(t)E(s)^*$, Theorem 6.9 and the comments surrounding (6-7) show that

$$\left\| \phi(t) \int \tilde{K}_k^{e_1}(t-s, x_1, x', y_1, y') \phi(s) F(s, y_1, y') ds dy' \right\|_{L_{x'}^2 L_t^2} \leq C \|F(\cdot, y_1, \cdot)\|_{L_{y'}^2 L_s^2}.$$

Interpolation then yields

$$\begin{aligned} \left\| \phi(t) \int \tilde{K}_k^{e_1}(t-s, x_1, x', y_1, y') \phi(s) F(s, y_1, y') ds dy' \right\|_{L_{x'}^{q_d} L_t^2} \\ \leq C 2^{2ks_d} |x_1 - y_1|^{-1 + \frac{1}{q_d} - \frac{1}{q_d}} \|F(\cdot, y_1, \cdot)\|_{L_{y'}^{q'_d} L^2_3}, \end{aligned}$$

and an application of the Hardy–Littlewood inequality yields the desired bound.

7. Wave packets and dispersive estimates

This section is devoted to the proof of (6-15) for $|t| \leq 1$. Without loss of generality we assume $0 \leq t \leq 1$ throughout to simplify notation.

To motivate the proof we recall Fefferman’s analysis [1973] of $\exp(-i|D|)$, the wave group for the Euclidean laplacian at $t = 1$. Consider

$$K_k(x) = (2\pi)^{-n} \int e^{i\langle x, \eta \rangle - i|\eta|} \psi_k(\eta) d\eta.$$

Following [Fefferman 1973], we have the decomposition $\psi_k(\eta) = \sum_\nu \psi_k^\nu(\eta)$, where ψ_k^ν equals ψ_k multiplied by a homogeneous cutoff to a conic neighborhood of angle $2^{-\frac{k}{2}}$ about the direction $\nu \in \mathbb{S}^{d-1}$, and ν varies over a discrete set of directions separated by distance $2^{-\frac{k}{2}}$. The function ψ_k^ν behaves like a scaled cutoff to a rectangle of dimension $2^k \times (2^{\frac{k}{2}})^{d-1}$ in that

$$|\langle \nu, \partial_\eta \rangle^m \partial_\eta^\alpha \psi_k^\nu(\eta)| \leq C_{m,\alpha} 2^{-k(m + \frac{|\alpha|}{2})},$$

with constants independent of k . The angular width is selected since one can write

$$e^{-i|\eta|} \psi_k^\nu(\eta) = e^{-i\langle \nu, \eta \rangle} a_k^\nu(\eta),$$

where a_k^ν satisfies the same derivative estimates as ψ_k^ν . Then we have the decomposition

$$K_k(x) = \sum_\nu f_k^\nu(x - \nu), \quad \text{where } \hat{f}_k^\nu(\eta) = a_k^\nu(\eta).$$

The function $f_k^\nu(x - \nu)$ is concentrated in a rectangle centered at ν , of dimension 2^{-k} along the ν -direction and $2^{-\frac{k}{2}}$ in perpendicular directions. By the spacing of the indices ν these rectangles are essentially disjoint, and simple geometry shows that, for all N ,

$$|K_k(x)| \leq C_N 2^{k(\frac{d+1}{2})} (1 + 2^k ||x| - 1|)^{-N}.$$

If $2^{-k} \leq t \leq 1$, the above argument can be scaled by t to decompose the kernel of $\exp(-it|D|)$. This gives a t -dependent splitting $\psi_k = \sum_\nu \psi_{k,t}^\nu$, where now $\psi_{k,t}^\nu$ is localized to a cone of angle $t^{-\frac{1}{2}} 2^{-\frac{k}{2}}$, and the $f_{k,t}^\nu(x - t\nu)$ are concentrated in a rectangle of dimensions 2^{-k} and $t^{\frac{1}{2}} 2^{-\frac{k}{2}}$, centered at $t\nu$. These rectangles are again mutually disjoint, leading to bounds

$$|K_k(t, x)| \leq C_N 2^{k(\frac{d+1}{2})} t^{-\binom{d-1}{2}} (1 + 2^k ||x| - t|)^{-N}.$$

For $0 \leq t \leq 2^{-k}$, the symbol $e^{-it|\eta|}$ is a classical symbol, and the kernel has the same size as $\hat{\psi}_k(-x)$, or as $K_k(t, x)$ at $t = 2^{-k}$.

The decomposition of [Fefferman 1973] was used in [Seeger, Sogge, and Stein 1991] to estimate the kernel of oscillatory integral operators with nondegenerate phase functions, for example $\exp(-iP)$ for a smooth metric. The key ingredient is that the phase function $\varphi(x, \eta)$ can be linearized in η over the support of each ψ_k^ν , up to an error that behaves like an appropriate amplitude function.

Getting the correct kernel estimates for $t \ll 1$ requires better estimates on the phase function for it to linearize over the support of $\psi_{k,t}^\nu$. The needed estimates are precisely those of (4-7), and the corresponding estimates for amplitudes are those of (4-8).

The proof of the estimates in (6-15) for a single term $\tilde{W}_k^\omega(t)$ or $\tilde{B}_k^\omega(t)$ would follow along the lines of [Seeger, Sogge, and Stein 1991], using the decomposition $\psi_{k,t}^\nu$, together with (4-7)–(4-8). We need, however, to prove these estimates for a product of arbitrarily many terms $\prod_j \tilde{B}_k^\omega(tr_j)$, where $\sum r_j = 1$. It is still appropriate to use the partition $\psi_{k,t}^\nu$ for each term; however, we need a function space argument in order to handle a product of terms since there is no hope for controlling the operator product using a symbol calculus. We therefore work with a wave packet frame and function spaces using weighted norms in that frame that grow with the distance to a given point (x_0, ν_0) on the cosphere bundle. We prove that the operator $\tilde{B}_k^\omega(s)$ is bounded from the space weighted at (x_0, ν_0) to the space weighted at its time- s flowout (x_s, ν_s) . These function space estimates iterate and yield a convergent sum, which is sufficient to prove the bounds in (6-15).

The wave packet frame. We will establish (6-15) for $2^{-k} \leq t \leq 1$; the proof for $0 \leq t \leq 2^{-k}$ follows by using the same proof as for $t = 2^{-k}$. We consider t to be fixed for this section and suppress the dependence of the frame on t ; however, we note that all constants are uniform over $t \in [0, 1]$.

We prove the estimate by studying the behavior of $\tilde{E}_k^\omega(t)$ in a family of wave packets that form a frame for functions that are frequency localized at scale 2^k . The wave packet frame that we use at scale 2^k is essentially a spatial dilation by t^{-1} of the scale $t2^k$ parabolic wave packets of [Smith 1998]. The only difference is that our frame covers more than one dyadic region, but we provide the details here for completeness.

We will be expanding functions with Fourier transform supported in the annulus

$$A_k = \left\{ \eta : \frac{4}{3} 2^{k-1} \leq |\eta| \leq \frac{5}{4} 2^{k+2} \right\}.$$

Let $A'_k = \left\{ \eta : \frac{2}{3} 2^{k-1} \leq |\eta| \leq \frac{3}{2} 2^{k+2} \right\}$. We construct a partition of unity on A_k , supported in A'_k , of the form

$$1 = \sum_{\nu \in \Upsilon_{k,t}} \beta_{k,t}^\nu(\eta)^2 \quad \text{when } \eta \in A_k, \quad \text{supp}(\beta_{k,t}^\nu) \subset \Omega_{k,t}^\nu,$$

where $\Upsilon_{k,t}$ is a collection of unit vectors separated by $t^{-\frac{1}{2}} 2^{-\frac{k}{2}}$, and $\beta_{k,t}^\nu(\eta)$ satisfies the following estimates:

$$|\langle \nu, \partial_\eta \rangle^j \partial_\eta^\alpha \beta_{k,t}^\nu(\eta)| \leq C_{j,\alpha} 2^{-kj} (t^{-\frac{1}{2}} 2^{\frac{k}{2}})^{-|\alpha|}. \quad (7-1)$$

Observe that $\Omega_{k,t}^\nu$, defined in (4-6), is contained in a rectangle of dimension 2^{k+3} along the direction ν , and $t^{-\frac{1}{2}} 2^{\frac{k}{2}}$ along the directions orthogonal to ν . For each ν , let $\Xi_{k,t}^\nu$ be a rectangular lattice in \mathbb{R}^n

with spacing $2\pi \cdot 2^{-k-3}$ along the ν -direction and spacing $2\pi \cdot t^{\frac{1}{2}} 2^{-\frac{k}{2}}$ in directions orthogonal to ν . Let $\Gamma_{k,t} = \{(x, \nu) : x \in \Xi_{k,t}^\nu, \nu \in \Upsilon_{k,t}\}$, which is a discrete subset of the cosphere bundle $S^*(\mathbb{R}^d)$. We use $\gamma = (x, \nu)$ to denote a variable in $S^*(\mathbb{R}^d)$, and for $\gamma \in \Gamma_{k,t}$ we set

$$\hat{\phi}_\gamma(\eta) = 2^{-\frac{3}{2}} 2^{-k(\frac{d+1}{4})} t^{\frac{d-1}{4}} e^{-i\langle x, \eta \rangle} \beta_{k,t}^\nu(\eta).$$

Then, with $\langle \nu^\perp, \partial_y \rangle$ denoting derivatives in directions perpendicular to ν ,

$$\begin{aligned} & |\langle \nu^\perp, \partial_y \rangle^\alpha \partial_y^\beta \phi_\gamma(y)| \\ & \leq C_{N,\alpha,\beta} 2^{k(\frac{d+1}{4})} t^{-\frac{d-1}{4}} 2^{k|\beta|} (t^{-\frac{1}{2}} 2^{\frac{k}{2}})^{|\alpha|} (1 + 2^k |\langle \nu, y - x \rangle| + t^{-1} 2^k |y - x|^2)^{-N}. \end{aligned} \quad (7-2)$$

Functions $f \in L^2(\mathbb{R}^n)$ with $\text{supp}(\hat{f}) \subset A_k$ admit an expansion in $\{\phi_\gamma\}_{\gamma \in \Gamma_{k,t}}$,

$$f = \sum_{\gamma \in \Gamma_{k,t}} c_\gamma \phi_\gamma, \quad c_\gamma = \int \overline{\phi_\gamma(y)} f(y) dy.$$

We define a pseudodistance function on the cosphere bundle $S^*(\mathbb{R}^d)$ by

$$d_t(x, \nu; x', \nu') = |\langle \nu, x - x' \rangle| + |\langle \nu', x - x' \rangle| + t|\nu - \nu'|^2 + t^{-1}|x - x'|^2.$$

This is the parabolic pseudodistance of [Smith 1998] scaled like the wave packet frame, and it satisfies, for all $t > 0$,

$$d_t(\gamma; \gamma'') \leq 4d_t(\gamma; \gamma') + 4d_t(\gamma'; \gamma''). \quad (7-3)$$

It is also approximately invariant under the Hamiltonian flow χ_s for $s \leq t$. This was proven for $C^{1,1}$ metrics in [Smith 1998]; we provide the proof here for metrics of bounded curvature.

Lemma 7.1. *For some C and all $0 \leq s \leq t \leq 1$, and χ_s the projected Hamiltonian flow map for any metric g_M satisfying (3-1)–(3-3), we have*

$$C^{-1} d_t(\gamma; \gamma') \leq d_t(\chi_s(\gamma); \chi_s(\gamma')) \leq C d_t(\gamma; \gamma').$$

Proof. Let $\eta = \nu$ and $\eta' = \nu'$. If (x_s, ξ_s) is the (nonprojected) Hamiltonian flow of (x, η) , then $|\xi_s - \eta| \lesssim c_d$, so we can replace ν_s by ξ_s in the distance function. From Corollary 3.2, when $|\eta| = 1$ we have the bounds $|\partial_\eta x_s| \lesssim s$, $|\partial_x x_s| + |\partial_x \xi_s| + |\partial_\eta \xi_s| \lesssim 1$, and we deduce

$$|x'_s - x_s| + t|\xi'_s - \xi_s| \lesssim |x' - x| + t|\eta' - \eta|.$$

Applying this also to χ_{-s} we obtain

$$t^{-1}|x'_s - x_s|^2 + t|\xi'_s - \xi_s|^2 \approx t^{-1}|x' - x|^2 + t|\eta' - \eta|^2.$$

By symmetry it thus suffices to show that

$$|\langle \eta, x' - x \rangle - \langle \xi_s, x'_s - x_s \rangle| \lesssim t^{-1}|x'_s - x_s|^2 + t|\eta' - \eta|^2. \quad (7-4)$$

Let φ be the phase function for \mathfrak{g}_M , and write $x = \nabla_\eta \varphi(s, x_s, \eta)$ and $\xi_s = \nabla_x \varphi(s, x_s, \eta)$. By homogeneity,

$$\begin{aligned} \langle \eta, x' - x \rangle - \langle \xi_s, x'_s - x_s \rangle &= \langle \eta, \nabla_\eta \varphi(x, x'_s, \eta') - \nabla_\eta \varphi(s, x_s, \eta) \rangle - \langle \nabla_x \varphi(s, x_s, \eta), x'_s - x_s \rangle \\ &= \varphi(s, x'_s, \eta') - \varphi(s, x_s, \eta) - \langle x'_s - x_s, \nabla_x \varphi(s, x_s, \eta) \rangle - \langle \eta' - \eta, \nabla_\eta \varphi(s, x'_s, \eta') \rangle. \end{aligned}$$

Observe that, by [Theorem 3.3](#),

$$\begin{aligned} |\langle \eta' - \eta, \nabla_\eta \varphi(s, x'_s, \eta') - \nabla_\eta \varphi(s, x_s, \eta) \rangle| &\lesssim |\eta' - \eta| (|x'_s - x_s| + t|\eta' - \eta|) \\ &\lesssim t^{-1} |x'_s - x_s|^2 + t|\eta' - \eta|^2. \end{aligned}$$

Consequently, it suffices to show that the error bound for the first-order Taylor expansion of $\varphi(s, x'_s, \eta') - \varphi(s, x_s, \eta)$ is bounded by the right-hand side of (7-4). The estimates (4-1)–(4-4) give $|\partial_x^2 \varphi_k| \lesssim 1$, $|\partial_x \partial_\eta \varphi_k| \lesssim 1$, $|\partial_\eta^2 \varphi_k| \lesssim |s|$, and hence the remainder is dominated by

$$|x'_s - x_s|^2 + |x'_s - x_s| |\eta' - \eta| + t|\eta' - \eta|^2 \leq \frac{3}{2} t^{-1} |x'_s - x_s|^2 + \frac{3}{2} t|\eta' - \eta|^2,$$

giving the desired bound. \square

For any given integer $M \geq 0$ and point $\gamma_0 \in S^*(\mathbb{R}^d)$, we define a weighted norm space

$$\|f\|_{M, \gamma_0}^2 = \sum_\gamma (1 + 2^k d_t(\gamma; \gamma_0))^{2M} |c_\gamma(f)|^2, \quad c_\gamma(f) = \int \overline{\phi_\gamma(y)} f(y) dy.$$

For dyadically localized f , this norm roughly measures how far f is from being a wave packet centered at γ_0 . In the next subsection we will prove the following theorem.

Theorem 7.2. *Suppose that $0 \leq s \leq t \leq 1$, $\gamma_0 \in \Gamma_{k,t}$, and $\chi_s(\gamma_0) = (x_s, \nu_s)$, where χ_s is the projected Hamiltonian flow for \mathfrak{g}_k . Then for all l, β, N , there are constants $C_{l, \beta, N}$ so that*

$$\begin{aligned} |\langle \nu_s^\perp, \partial_x \rangle^\alpha \partial_x^\beta (\tilde{B}_k^\omega(s) \phi_{\gamma_0})(x)| \\ \leq C_{N, \alpha, \beta} 2^{k(\frac{d+1}{4})} t^{-\frac{d-1}{4}} 2^{k|\beta|} (t^{-\frac{1}{2}} 2^{\frac{k}{2}})^{|\alpha|} (1 + 2^k |\langle \nu_s, x - x_s \rangle| + t^{-1} 2^k |x - x_s|^2)^{-N}. \end{aligned} \quad (7-5)$$

In the remainder of this subsection we deduce (6-15) from [Theorem 7.2](#). First we deduce $\|\cdot\|_{M, \chi_s(\gamma)}$ mapping properties for $\tilde{B}_k^\omega(s)$ from (7-5). The left-hand side of (7-5) vanishes unless $\angle(\omega, \gamma) \leq \frac{1}{4}$, so we may assume $\angle(\omega, \gamma_s) \leq \frac{1}{2}$.

Lemma 7.3. *Suppose that \hat{f} is supported in the set $\{n : \angle(\eta, \nu_0) \leq \frac{1}{2}\}$, and for all N, α, β we have*

$$|\langle \nu_0^\perp, \partial_y \rangle^\alpha \partial_y^\beta f(y)| \leq C_{N, \alpha, \beta} 2^{k(\frac{d+1}{4})} t^{-\frac{d-1}{4}} 2^{k|\beta|} (t^{-\frac{1}{2}} 2^{\frac{k}{2}})^{|\alpha|} (1 + 2^k |\langle \nu_0, y - x_0 \rangle| + t^{-1} 2^k |y - x_0|^2)^{-N}.$$

Let $\gamma_0 = (x_0, \nu_0)$. Then for all $M \geq 0$ we have $\|f\|_{M, \gamma_0} \leq C_M$, where C_M depends on only a finite number of the $C_{N, \alpha, \beta}$.

Proof. Without loss of generality we assume that $\nu_0 = e_1$. By the derivative estimates we have

$$|\hat{f}(\eta)| \leq C_N 2^{-k(\frac{d+1}{4})} t^{\frac{d-1}{4}} (1 + 2^{-k} |\eta_1| + 2^{-k} t |\eta'|^2)^{-N},$$

where for each N the value of C_N depends on only a finite number of $C_{N,\alpha,\beta}$. Since $\hat{\phi}_\gamma$ is supported where $|\eta'| \geq 2^{k-4}|v - e_1|$, by Plancherel's theorem we obtain for all N , and similar C_N ,

$$|c_\gamma(f)| \leq C_N (1 + 2^k t |v - e_1|^2)^{-2N}, \quad c_\gamma(f) = \int \overline{\phi_\gamma(y)} f(y) dx. \quad (7-6)$$

By the pointwise estimates on $f(y)$ and $\phi_\gamma(y)$, we have

$$|c_\gamma(f)| \leq C_N 2^{k(\frac{d+1}{2})} t^{-\frac{d-1}{2}} \int (1 + 2^k d_t(y, e_1; \gamma_0))^{-2N-d} (1 + 2^k d_t(y, v; \gamma))^{-2N-d} dy.$$

By (7-3), noting that $d_t(y, e_1; y, v) = t|v - e_1|^2$, we have

$$\frac{1}{16} d_t(v, \gamma_0) \leq d_t(y, e_1; \gamma_0) + d_t(y, v; \gamma) + t|v - e_1|^2.$$

Together with (7-6), this implies $|c_\gamma(f)| \leq C_N (1 + 2^k d_t(\gamma; \gamma_0))^{-N}$. The lemma then follows from the bound

$$\sup_{\gamma'} \sum_{\gamma \in \Gamma_{k,t}} (1 + 2^k d_t(\gamma; \gamma'))^{-d-1} \leq C_d, \quad (7-7)$$

which follows from estimate (2.3) in [Smith 1998] after rescaling. \square

The converse to Lemma 7.3 also holds; we need it only for $\alpha = \beta = 0$, and prove that version in the proof of Corollary 7.5 below.

An immediate consequence of Theorem 7.2 and Lemma 7.3 is decay estimates on the matrix coefficients of $\tilde{B}_k^\omega(s)$. Precisely, for all N we have

$$\left| \int \overline{\phi_\gamma(y)} (\tilde{B}_k^\omega(s) \phi_{\gamma'})(y) dy \right| \leq C_N (1 + 2^k d_t(\gamma; \chi_s(\gamma')))^{-N}. \quad (7-8)$$

We then use this to prove boundedness of $\tilde{B}_k^\omega(s)$ in the weighted norm spaces via the following lemma.

Lemma 7.4. *Suppose that $M \geq 0$, $0 \leq s \leq t \leq 1$, and $T : \mathcal{S}(\mathbb{R}^d) \rightarrow \mathcal{S}'(\mathbb{R}^d)$ is a linear map such that the matrix coefficients*

$$a(\gamma, \gamma') = \int \overline{\phi_\gamma(y)} (T \phi_{\gamma'})(y) dy$$

satisfy the bound

$$|a(\gamma, \gamma')| \leq (1 + 2^k d_t(\gamma; \chi_s(\gamma')))^{-(M+d+1)}.$$

Then, uniformly over $\gamma_0 \in \mathcal{S}^(\mathbb{R}^d)$, we have $\|Tf\|_{M, \chi_s(\gamma_0)} \leq C_M \|f\|_{M, \gamma_0}$.*

Proof. It follows from (7-7) that

$$\sup_{\gamma'} \sum_{\gamma} |a(\gamma, \gamma')| \leq C,$$

and, since $d_t(\gamma; \chi_s(\gamma')) \approx d_t(\chi_{-s}(\gamma); \gamma')$ by Lemma 7.1, we also have

$$\sup_{\gamma} \sum_{\gamma'} |a(\gamma, \gamma')| \leq C,$$

where C is independent of s, t and k . By Schur's lemma we conclude

$$\|\tilde{B}_k^\omega(s)f\|_{0,\chi_s(\gamma_0)} \leq C\|f\|_{0,\gamma_0}.$$

The weighted case $M \geq 1$ follows by noting that

$$(1 + 2^k d_t(\gamma; \chi_s(\gamma_0))) \lesssim (1 + 2^k d_t(\gamma; \chi_s(\gamma'))) (1 + 2^k d_t(\gamma'; \gamma_0)),$$

which follows from

$$d_t(\gamma; \chi_s(\gamma_0)) \leq 4d_t(\gamma; \chi_s(\gamma')) + 4d_t(\chi_s(\gamma'); \chi_s(\gamma_0)),$$

and the fact that $d_t(\chi_s(\gamma'); \chi_s(\gamma_0)) \approx d_t(\gamma', \gamma_0)$. \square

Corollary 7.5. *Let χ_s denote the time- s projected Hamiltonian flow for g_k . Then for $0 \leq s \leq t \leq 1$, and all $M \geq 0$,*

$$\|\tilde{E}_k^\omega(s)f\|_{M,\chi_s(\gamma_0)} \leq C_M \|f\|_{M,\gamma_0},$$

with constant C_M independent of s, t, γ_0, ω , and k .

Proof. By Lemma 7.4 and the estimate (7-8), which holds also for $\tilde{W}_k^\omega(s)$ by the same proof, we have

$$\|\tilde{B}_k^\omega(s)f\|_{M,\chi_s(\gamma_0)} \leq C_M \|f\|_{M,\gamma_0}.$$

The formula (6-8) for $\tilde{E}_k^\omega(t)$ and the group property of χ_s then show that

$$\|\tilde{E}_k^\omega(s)f\|_{M,\chi_s(\gamma)} \leq \sum_{m=0}^{\infty} \frac{s^m C_M^{m+1}}{m!} \|f\|_{M,\gamma} = C_M e^{sC_M} \|f\|_{M,\gamma}. \quad \square$$

We conclude this section by deriving the bound (6-15) from Corollary 7.5. Write $\tilde{K}_k^\omega(t, x, y) = (\tilde{E}_k^\omega(t)\delta_y)(x)$. Since $\tilde{E}_k^\omega(t)$ has the factor $\psi_k(D)$ on the right, we may write

$$(\tilde{E}_k^\omega(t)\delta_y)(x) = \sum_{\nu \in \Upsilon_{k,t}} (\tilde{E}_k^\omega(t)\beta_{k,t}^\nu(D)^2\delta_y)(x).$$

The function $\beta_{k,t}^\nu(D)^2\delta_y$ has Fourier transform $e^{-i\langle y, \eta \rangle} \beta_{k,t}^\nu(\eta)^2$. Up to a normalization factor, this behaves like the frame element ϕ_γ at $\gamma = (y, \nu)$, and it is easy to verify that for all M

$$\|\beta_{k,t}^\nu(D)^2\delta_y\|_{M,\gamma} \leq C_M 2^{k(\frac{d+1}{4})} t^{-\frac{d-1}{4}}.$$

By Corollary 7.5, letting $\gamma_t \equiv (x_t, \nu_t) = \chi_t(y, \nu)$ we have

$$\|\tilde{E}_k^\omega(t)\beta_{k,t}^\nu(D)^2\delta_y\|_{M,\gamma_t} \leq C_M 2^{k(\frac{d+1}{4})} t^{-\frac{d-1}{4}}. \quad (7-9)$$

This implies that, for all N ,

$$|(\tilde{E}_k^\omega(t)\beta_{k,t}^\nu(D)^2\delta_y)(x)| \leq C_N 2^{k(\frac{d+1}{2})} t^{-\frac{d-1}{2}} (1 + 2^k |\langle \nu_t, x - x_t \rangle| + 2^k t^{-1} |x - x_t|^2)^{-N}.$$

We see this using (7-9), that the frame coefficients $\{c_{\gamma'}\}$ of $\tilde{E}_k^\omega(t)\beta_{k,t}^\nu(D)^2\delta_y$ satisfy for all M

$$|c_{\gamma'}| \leq C_M 2^{k(\frac{d+1}{4})} t^{-\frac{d-1}{4}} (1 + 2^k d_t(\gamma'; \gamma_t))^{-M}.$$

From estimates (7-2) on $|\phi_\gamma(x)|$, we follow the proof of [Smith 1998, Lemma 2.5] with $\gamma' = (x', v')$ to bound $|(\tilde{E}_k^\omega(t)\beta_{k,t}^v(D)^2\delta_y)(x)|$ by

$$\begin{aligned} C_M 2^k \left(\frac{d+1}{2}\right) t^{-\frac{d-1}{2}} \sum_{\gamma' \in \Gamma_{k,t}} (1 + 2^k d_t(\gamma'; \gamma_t))^{-M} (1 + 2^k d_t((x, v'); \gamma'))^{-M} \\ \leq C_M 2^k \left(\frac{d+1}{2}\right) t^{-\frac{d-1}{2}} \sum_{v' \in \Upsilon_{k,t}} (1 + 2^k d_t((x, v'); \gamma_t))^{-M} \\ \leq C_M 2^k \left(\frac{d+1}{2}\right) t^{-\frac{d-1}{2}} (1 + 2^k |\langle v_t, x - x_t \rangle| + 2^k t^{-1} |x - x_t|^2)^{-M + \frac{d}{2}}. \end{aligned}$$

Deriving (6-15) from Corollary 7.5 then reduces to showing that

$$\sum_{v \in \Upsilon_{k,t}} (1 + 2^k |\langle v_t, x - x_t \rangle| + 2^k t^{-1} |x - x_t|^2)^{-N-d} \leq C_N (1 + 2^k \text{dist}(x, S_t(y)))^{-N}.$$

If $(\tilde{x}_t, \tilde{v}_t) = \chi_t(y, \tilde{v})$ and $(x_t, v_t) = \chi_t(y, v)$, then by Corollary 3.2

$$\frac{4}{5} t \leq \frac{|\tilde{x}_t - x_t|}{|\tilde{v} - v|} \leq \frac{5}{4} t.$$

Consequently, the points x_t are separated by $t^{\frac{1}{2}} 2^{-\frac{k}{2}}$ for $v \in \Upsilon_{k,t}$, and thus

$$\sum_{v \in \Upsilon_{k,t}} (1 + 2^k t^{-1} |x - x_t|^2)^{-d} \leq C.$$

It therefore suffices to show that, for c_d small enough, and for each $v \in \mathbb{S}^{d-1}$,

$$|\langle v_t, x - x_t \rangle| + t^{-1} |x - x_t|^2 \geq \frac{1}{4} \text{dist}(x, S_t(y)). \quad (7-10)$$

Here, $x_t \in S_t(y)$ for each v , and v_t is the unit normal to $S_t(y)$ at the point x_t , in that $\langle v_t, \partial_{\eta_j} x_t |_{\eta=v} \rangle = 0$, which follows by homogeneity.

We observe that, by scaling, it suffices to prove (7-10) in the case $t = 1$. Precisely, $(t^{-1}x_t, v_t)$ is the image at time 1 of $(t^{-1}y, v)$ under the projected Hamiltonian flow for the metric $g_k(t \cdot)$, and $t^{-1}S_t(y)$ is the corresponding unit geodesic sphere centered at $t^{-1}y$; hence the two sides of (7-10) dilate by the same factor t . Furthermore, the metric $g_k(t \cdot)$ satisfies conditions (3-1)–(3-3) with $M = t 2^{\frac{k}{2}} \leq 2^{\frac{k}{2}}$.

Without loss of generality we assume $v = e_1$ and $y = 0$. We introduce the notation $(x(\omega), n(\omega)) = \chi_1(0, \omega)$ to denote the mapping of the unit sphere \mathbb{S}^{d-1} onto the unit conormal bundle of $S_1(0)$. By Corollary 3.2, this map is C^1 -close to the map $\omega \rightarrow (\omega, \omega)$; precisely

$$|x(\omega) - \omega| + |\nabla_\omega x(\omega) - \Pi_\omega^\perp| + |n(\omega) - \omega| + |\nabla_\omega n(\omega) - \Pi_\omega^\perp| \lesssim c_d.$$

As a consequence we may parametrize $S_1(0) \cap \{x_1 > 0, |x'| \leq \frac{1}{2}\}$ as a graph $x_1 = F(x')$, where

$$|\partial_x^\alpha (F(x') - \sqrt{1 - |x'|^2})| \lesssim c_d, \quad |\alpha| \leq 2, \quad |x'| \leq \frac{1}{2}. \quad (7-11)$$

This holds for $|\alpha| \leq 1$ by C^1 closeness of $x(\omega)$ to ω , and for $|\alpha| = 2$ since

$$\nabla_{x'} F(x') = -n'(\omega(x'))/n_1(\omega(x'))$$

is C^1 close to $-x'/\sqrt{1-|x'|^2}$.

The bound (7-10) is equivalent to proving, for $x = (x_1, x') \in \mathbb{R}^d$,

$$\min_{\omega} |x - x(\omega)| \leq 4(|\langle n(e_1), x - x(e_1) \rangle| + |x - x(e_1)|^2).$$

We assume that $|x - x(e_1)| \leq \frac{1}{4}$; hence $|x'| \leq \frac{1}{2}$, as the bound is immediate otherwise. The left-hand side is bounded above by $|x_1 - F(x')|$, and the bound then follows by the Taylor expansion of $F(x')$ about $x'(e_1)$,

$$\begin{aligned} |x_1 - F(x')| &\leq |x_1 - F(x'(e_1)) - \langle x' - x'(e_1), \nabla_{x'} F(x'(e_1)) \rangle| + |x' - x'(e_1)|^2 \\ &= n_1(e_1)^{-1} |\langle n(e_1), x - x(e_1) \rangle| + |x' - x'(e_1)|^2, \end{aligned}$$

where we use that $\|\nabla_{x'}^2 F\| \leq 2$ for $|x'| \leq \frac{1}{2}$ by (7-11) if c_d is small, and $F(x'(e_1)) = x_1(e_1)$.

Proof of Theorem 7.2. We follow the key idea of [Seeger, Sogge, and Stein 1991], that the action of a Fourier integral operator on a function f whose Fourier transform is suitably localized can be decomposed as a pseudodifferential operator acting on f , followed by a change of coordinates. Suitably localized means that the phase function can be written as a phase that is linear in η plus a term that satisfies the estimates of a zero-order symbol on the support of $\hat{f}(\eta)$. Here we take $f = \phi_\gamma$, with \hat{f} supported in the set $\Omega_{k,t}^\nu$ defined by (4-6), and the zero-order symbol estimates are those of Corollary 5.2. The estimates of Corollary 4.2 will be used to establish the linearization of φ_k on $\Omega_{k,t}^\nu$.

We prove Theorem 7.2 with $\tilde{B}_k^\omega(s)\varphi_\gamma$ replaced by $B_k(s)$; recall the definition (6-6) and (6-5). The operator $\tilde{a}_\omega(D)\tilde{\psi}_k(D)$ is a mollifier on spatial scale 2^{-k} and commutes with differentiation, and hence preserves the estimates of Theorem 7.2, and $\tilde{a}_\omega(D)\phi_\gamma$ satisfies the same conditions as ϕ_γ . The terms $B_{k\pm 1}(s)$ will follow the same proof as for $B_k(s)$.

Without loss of generality we assume $\gamma_0 = (0, e_1)$. We need to establish the bounds of Theorem 7.2 for the function

$$(B_k(s)\phi_{\gamma_0})(x) = 2^{-\frac{3}{2}} 2^{-k(\frac{d+1}{4})} t^{\frac{d-1}{4}} \int e^{i\varphi_k(s,x,\eta)} b_k(s, x, \eta) \beta_{k,t}^{\epsilon_1}(\eta) d\eta.$$

We can express this in the form

$$(B_k(s)\phi_{\gamma_0})(x) = 2^{-\frac{3}{2}} 2^{-k(\frac{d+1}{4})} t^{\frac{d-1}{4}} \int e^{i(y(s,x),\eta)} e^{ih(s,x,\eta)} b_k(s, x, \eta) \beta_{k,t}^{\epsilon_1}(\eta) d\eta,$$

where $y(s, x) = \nabla_\eta \varphi_k(s, x, e_1)$, and where by (4-9) on the support of $\beta_{k,t}^{\epsilon_1}$ the function $h(s, x, \eta) = \varphi_k(s, x, \eta) - \eta \cdot \partial_\eta \varphi_k(s, x, e_1)$ satisfies

$$|\partial_{\eta_1}^j \partial_{\eta'}^\alpha \partial_x^\beta h(s, x, \eta)| \leq C_{j,\alpha,\beta} 2^{-kj} (t^{\frac{1}{2}} 2^{-\frac{k}{2}})^{|\alpha|} 2^{\frac{k}{2}|\beta|}.$$

This, together with Corollary 5.2 and (7-1), leads to the estimates

$$|\partial_{\eta_1}^j \partial_{\eta'}^\alpha \partial_x^\beta (e^{ih(s,x,\eta)} b_k(s, x, \eta) \beta_{k,t}^{\epsilon_1}(\eta))| \leq C_{j,\alpha,\beta} 2^{-kj} (t^{\frac{1}{2}} 2^{-\frac{k}{2}})^{|\alpha|} 2^{\frac{k}{2}|\beta|}. \quad (7-12)$$

We now express

$$(B_k(s)\phi_{\gamma_0})(x) = 2^k \left(\frac{d+1}{4}\right) t^{-\frac{d-1}{4}} F(x, y(s, x)),$$

where

$$F(x, y) = 2^{-\frac{3}{2}} 2^{-k \left(\frac{d+1}{2}\right)} t^{\frac{d-1}{2}} \int e^{i\langle y, \eta \rangle} e^{ih(s, x, \eta)} b_k(s, x, \eta) \beta_{k,t}^{e_1}(\eta) d\eta.$$

The estimates (7-12) and integration by parts leads to the bounds

$$|\partial_{y_1}^j \partial_{y'}^\alpha \partial_x^\beta F(x, y)| \leq C_{N,j,\alpha,\beta} 2^{\frac{k}{2}|\gamma|} 2^{k|\beta|} (t^{-\frac{1}{2}} 2^{\frac{k}{2}})^{|\alpha|} (1 + 2^k |y_1| + t^{-1} 2^k |y|^2)^{-N}.$$

We now use the chain rule to express x -derivatives of the composition of $F(x, y)$ with $y = y(s, x)$ as a sum of terms,

$$\partial_{x_i} F(x, y(s, x)) = (\partial_{x_i} F)(x, y(s, x)) + (\nabla_y F)(x, y(s, x)) \cdot \partial_{x_i} y(s, x).$$

The ∂_{x_i} in the first term on the right counts as a factor of $2^{\frac{k}{2}}$ in the derivative estimates, which is better than the conclusion of [Theorem 7.2](#). Similar considerations apply to terms in the expansion of higher-order derivatives. Since we will estimate individually each term arising in such an expansion, we therefore can consider functions F that are functions of only y . That is, we assume for all N that

$$|\partial_{y_1}^j \partial_{y'}^\alpha F(y)| \leq C_{N,\alpha,\beta} 2^{kj} (t^{-\frac{1}{2}} 2^{\frac{k}{2}})^{|\alpha|} (1 + 2^k |y_1| + t^{-1} 2^k |y|^2)^{-N}, \tag{7-13}$$

and prove that the composition with $y(s, x)$ satisfies for all N

$$|\partial_x^\alpha \partial_x^\beta F(y(s, x))| \leq C_{N,\alpha,\beta} 2^{k|\beta|} (t^{-\frac{1}{2}} 2^{\frac{k}{2}})^{|\alpha|} (1 + 2^k |\langle v_s, x - x_s \rangle| + t^{-1} 2^k |x - x_s|^2)^{-N}, \tag{7-14}$$

where $\partial_x \equiv \langle v_s^\perp, \partial_x \rangle$ denotes derivatives in directions perpendicular to v_s .

Since $y(s, x_s) = 0$, and the map $x \rightarrow y(s, x)$ is a globally bilipschitz map of \mathbb{R}^d , with uniform bounds on the map and its inverse, we have

$$|y(s, x)|^2 \approx |x - x_s|^2,$$

with the ratio of the two sides close to 1 for c_d small. For a constant c close to 1, we also have

$$c v_s = (\nabla_x \varphi_k)(s, x_s, e_1) = (\nabla_x \partial_{\eta_1} \varphi_k)(s, x_s, e_1) = (\nabla_x y_1)(s, x_s).$$

We also have the equality $y_1(s, x) = \varphi_k(s, x, e_1)$ by homogeneity, which by (4-1) implies

$$|\partial_x^\beta y_1(s, x)| \leq \begin{cases} C, & |\beta| = 1, \\ C 2^{\frac{k}{2}(|\beta|-2)}, & |\beta| \geq 2. \end{cases} \tag{7-15}$$

Together with a first-order Taylor expansion these imply that, for $0 < t \leq 1$,

$$|y_1(s, x)| + t^{-1} |y(s, x)|^2 \approx |\langle v_s, x - x_s \rangle| + t^{-1} |x - x_s|^2, \tag{7-16}$$

with uniform bounds on the ratios. Together with (7-13) this gives (7-14) for $j = \alpha = 0$.

To bound derivatives, we use the chain rule to express $\partial_x^\alpha \partial_x^\beta F(y(s, x))$ as a sum of terms of the form

$$(\partial_{y_1}^m \partial_{y'}^\theta F)(\partial_x^{\alpha_1} \partial_x^{\beta_1} y_1) \cdots (\partial_x^{\alpha_m} \partial_x^{\beta_m} y_1) (\partial_x^{\alpha_{m+1}} \partial_x^{\beta_{m+1}} y') \cdots (\partial_x^{\alpha_{m+|\theta|}} \partial_x^{\beta_{m+|\theta|}} y'),$$

where

$$\alpha = \sum_{j=1}^{m+|\theta|} \alpha_j, \quad \beta = \sum_{j=1}^{m+|\theta|} \beta_j, \quad m + |\theta| \leq |\alpha| + |\beta|.$$

The estimate (7-14) then follows from (7-13) and (7-16), together with the following bounds for the derivatives of $y(s, x)$ for $|\alpha| + |\beta| \geq 1$, and where $2^{-k} \leq t \leq 1$:

$$\begin{aligned} |\partial_x^\alpha \partial_x^\beta y_1(s, x)| &\leq C_{\alpha, \beta} 2^{k(|\beta|-1)} (t^{-\frac{1}{2}} 2^{\frac{k}{2}})^{|\alpha|} (1 + t^{-\frac{1}{2}} 2^{\frac{k}{2}} |x - x_s|), \\ |\partial_x^\alpha \partial_x^\beta y'(s, x)| &\leq C_{\alpha, \beta} 2^{k|\beta|} (t^{-\frac{1}{2}} 2^{\frac{k}{2}})^{|\alpha|-1}. \end{aligned}$$

The second of these holds by the stronger bound of $C_{\alpha, \beta} 2^{\frac{k}{2}(|\alpha|+|\beta|-1)}$ from Theorem 3.3, where if $|\alpha| = 0$ we use that $2^{-k} \leq (t^{-\frac{1}{2}} 2^{\frac{k}{2}})^{-1}$ and $|\beta| \geq 1$. For the first, if $|\alpha| = 1$ and $|\beta| = 0$, we use (7-15) and that $(\partial_x y_1)(s, x_s) = 0$ to see that $|\partial_x y_1(s, x)| \leq C |x - x_s|$. If $|\alpha| \geq 2$ or $|\beta| \geq 1$ then the estimate follows directly from (7-15). \square

References

- [Bony 1997] J.-M. Bony, “Caractérisations des opérateurs pseudo-différentiels”, exposé XXIII in *Séminaire sur les Équations aux Dérivées Partielles, 1996-1997*, École Polytech., Palaiseau, France, 1997. [MR](#) [Zbl](#)
- [Carleson 1962] L. Carleson, “Interpolations by bounded analytic functions and the corona problem”, *Ann. of Math. (2)* **76** (1962), 547–559. [MR](#) [Zbl](#)
- [Coifman and Meyer 1978] R. R. Coifman and Y. Meyer, *Au-delà des opérateurs pseudo-différentiels*, Astérisque **57**, Soc. Math. France, Paris, 1978. [MR](#) [Zbl](#)
- [Córdoba and Fefferman 1978] A. Córdoba and C. Fefferman, “Wave packets and Fourier integral operators”, *Comm. Partial Differential Equations* **3**:11 (1978), 979–1005. [MR](#) [Zbl](#)
- [DeTurck and Kazdan 1981] D. M. DeTurck and J. L. Kazdan, “Some regularity theorems in Riemannian geometry”, *Ann. Sci. École Norm. Sup. (4)* **14**:3 (1981), 249–260. [MR](#) [Zbl](#)
- [Fefferman 1973] C. Fefferman, “A note on spherical summation multipliers”, *Israel J. Math.* **15** (1973), 44–52. [MR](#) [Zbl](#)
- [Gilbarg and Trudinger 1983] D. Gilbarg and N. S. Trudinger, *Elliptic partial differential equations of second order*, 2nd ed., Grundlehren der Math. Wissenschaften **224**, Springer, 1983. [MR](#) [Zbl](#)
- [Ginibre and Velo 1997] J. Ginibre and G. Velo, “Generalized Strichartz inequalities for the wave equation in homogeneous Besov spaces”, pp. 135–144 in *Nonlinear waves* (Sapporo, 1995), edited by R. Agemi et al., GAKUTO Int. Ser. Math. Sci. Appl. **10**, Gakkōtoshō, Tokyo, 1997. [MR](#) [Zbl](#)
- [Keel and Tao 1998] M. Keel and T. Tao, “Endpoint Strichartz estimates”, *Amer. J. Math.* **120**:5 (1998), 955–980. [MR](#) [Zbl](#)
- [Lindblad and Sogge 1995] H. Lindblad and C. D. Sogge, “On existence and scattering with minimal regularity for semilinear wave equations”, *J. Funct. Anal.* **130**:2 (1995), 357–426. [MR](#) [Zbl](#)
- [Mockenhaupt, Seeger and Sogge 1993] G. Mockenhaupt, A. Seeger, and C. D. Sogge, “Local smoothing of Fourier integral operators and Carleson–Sjölin estimates”, *J. Amer. Math. Soc.* **6**:1 (1993), 65–130. [MR](#) [Zbl](#)
- [Seeger, Sogge, and Stein 1991] A. Seeger, C. D. Sogge, and E. M. Stein, “Regularity properties of Fourier integral operators”, *Ann. of Math. (2)* **134**:2 (1991), 231–251. [MR](#) [Zbl](#)
- [Smith 1998] H. F. Smith, “A parametrix construction for wave equations with $C^{1,1}$ coefficients”, *Ann. Inst. Fourier (Grenoble)* **48**:3 (1998), 797–835. [MR](#) [Zbl](#)
- [Smith 2006a] H. F. Smith, “Sharp $L^2 \rightarrow L^q$ bounds on spectral projectors for low regularity metrics”, *Math. Res. Lett.* **13**:5-6 (2006), 967–974. [MR](#) [Zbl](#)
- [Smith 2006b] H. F. Smith, “Spectral cluster estimates for $C^{1,1}$ metrics”, *Amer. J. Math.* **128**:5 (2006), 1069–1103. [MR](#) [Zbl](#)

- [Smith and Sogge 1994] H. F. Smith and C. D. Sogge, “On Strichartz and eigenfunction estimates for low regularity metrics”, *Math. Res. Lett.* **1**:6 (1994), 729–737. [MR](#) [Zbl](#)
- [Sogge 1988] C. D. Sogge, “Concerning the L^p norm of spectral clusters for second-order elliptic operators on compact manifolds”, *J. Funct. Anal.* **77**:1 (1988), 123–138. [MR](#) [Zbl](#)
- [Sogge 1993] C. D. Sogge, *Fourier integrals in classical analysis*, Cambridge Tracts in Math. **105**, Cambridge Univ. Press, 1993. [MR](#) [Zbl](#)
- [Stein 1970] E. M. Stein, *Singular integrals and differentiability properties of functions*, Princeton Math. Series **30**, Princeton Univ. Press, 1970. [MR](#) [Zbl](#)
- [Stein 1993] E. M. Stein, *Harmonic analysis: real-variable methods, orthogonality, and oscillatory integrals*, Princeton Math. Series **43**, Princeton Univ. Press, 1993. [MR](#) [Zbl](#)
- [Strichartz 1970] R. S. Strichartz, “A priori estimates for the wave equation and some applications”, *J. Funct. Anal.* **5**:2 (1970), 218–235. [MR](#) [Zbl](#)
- [Strichartz 1977] R. S. Strichartz, “Restrictions of Fourier transforms to quadratic surfaces and decay of solutions of wave equations”, *Duke Math. J.* **44**:3 (1977), 705–714. [MR](#) [Zbl](#)
- [Tataru 2000] D. Tataru, “Strichartz estimates for operators with nonsmooth coefficients and the nonlinear wave equation”, *Amer. J. Math.* **122**:2 (2000), 349–376. [MR](#) [Zbl](#)
- [Tataru 2001] D. Tataru, “Strichartz estimates for second order hyperbolic operators with nonsmooth coefficients, II”, *Amer. J. Math.* **123**:3 (2001), 385–423. [MR](#) [Zbl](#)
- [Tataru 2002] D. Tataru, “Strichartz estimates for second order hyperbolic operators with nonsmooth coefficients, III”, *J. Amer. Math. Soc.* **15**:2 (2002), 419–442. [MR](#) [Zbl](#)
- [Taylor 2000] M. E. Taylor, *Tools for PDE*, Math. Surveys and Monographs **81**, Amer. Math. Soc., Providence, RI, 2000. [MR](#) [Zbl](#)

Received 8 Aug 2018. Revised 12 Sep 2018. Accepted 22 Oct 2018.

YUANLONG CHEN: ylchen88@uw.edu

Department of Mathematics, University of Washington, Seattle, WA, United States

HART F. SMITH: hfsmith@uw.edu

Department of Mathematics, University of Washington, Seattle, WA, United States

EDITORS-IN-CHIEF

- Charles L. Epstein University of Pennsylvania
cle@math.upenn.edu
- Maciej Zworski University of California at Berkeley
zworski@math.berkeley.edu

EDITORIAL BOARD

- Sir John M. Ball University of Oxford
ball@maths.ox.ac.uk
- Michael P. Brenner Harvard University
brenner@seas.harvard.edu
- Charles Fefferman Princeton University
cf@math.princeton.edu
- Susan Friedlander University of Southern California
susanfri@usc.edu
- Anna Gilbert University of Michigan
annacg@umich.edu
- Leslie F. Greengard Courant Institute, New York University, and
Flatiron Institute, Simons Foundation
greengard@cims.nyu.edu
- Yan Guo Brown University
yan_guo@brown.edu
- Claude Le Bris CERMICS - ENPC
lebris@cermics.enpc.fr
- Robert J. McCann University of Toronto
mccann@math.toronto.edu
- Michael O'Neil Courant Institute, New York University
oneil@cims.nyu.edu
- Jill Pipher Brown University
jill_pipher@brown.edu
- Johannes Sjöstrand Université de Dijon
johannes.sjostrand@u-bourgogne.fr
- Vladimir Šverák University of Minnesota
sverak@math.umn.edu
- Daniel Tataru University of California at Berkeley
tataru@berkeley.edu
- Michael I. Weinstein Columbia University
miw2103@columbia.edu
- Jon Wilkening University of California at Berkeley
wilken@math.berkeley.edu
- Enrique Zuazua DeustoTech-Bilbao, and
Universidad Autónoma de Madrid
enrique.zuazua@deusto.es

PRODUCTION

- Silvio Levy (Scientific Editor)
production@msp.org

Cover image: The figure shows the outgoing scattered field produced by scattering a plane wave, coming from the northwest, off of the (stylized) letters P A A. The total field satisfies the homogeneous Dirichlet condition on the boundary of the letters. It is based on a numerical computation by Mike O'Neil of the Courant Institute.

See inside back cover or msp.org/paa for submission instructions.

The subscription price for 2019 is US \$495/year for the electronic version, and \$555/year (+\$25, if shipping outside the US) for print and electronic. Subscriptions, requests for back issues and changes of subscriber address should be sent to MSP.

Pure and Applied Analysis (ISSN 2578-5885 electronic, 2578-5893 printed) at Mathematical Sciences Publishers, 798 Evans Hall #3840, c/o University of California, Berkeley, CA 94720-3840 is published continuously online. Periodical rate postage paid at Berkeley, CA 94704, and additional mailing offices.

PAA peer review and production are managed by EditFlow[®] from MSP.

PUBLISHED BY

 **mathematical sciences publishers**
nonprofit scientific publishing
<http://msp.org/>

© 2019 Mathematical Sciences Publishers

PURE and APPLIED ANALYSIS

vol. 1 no. 1 2019

- Semiclassical resolvent estimates for bounded potentials 1
FRÉDÉRIC KLOPP and MARTIN VOGEL
- The quantum Sabine law for resonances in transmission problems 27
JEFFREY GALKOWSKI
- Dispersive estimates for the wave equation on Riemannian manifolds of bounded curvature 101
YUANLONG CHEN and HART F. SMITH