


PURE and APPLIED ANALYSIS

PAM

JORY GRIFFIN AND JENS MARKLOF

**QUANTUM TRANSPORT IN A LOW-DENSITY PERIODIC
POTENTIAL: HOMOGENISATION VIA HOMOGENEOUS FLOWS**



vol. 1 no. 4 2019

QUANTUM TRANSPORT IN A LOW-DENSITY PERIODIC POTENTIAL: HOMOGENISATION VIA HOMOGENEOUS FLOWS

JORY GRIFFIN AND JENS MARKLOF

We show that the time evolution of a quantum wavepacket in a periodic potential converges in a combined high-frequency/Boltzmann–Grad limit, up to second order in the coupling constant, to terms that are compatible with the linear Boltzmann equation. This complements results of Eng and Erdős for low-density random potentials, where convergence to the linear Boltzmann equation is proved in all orders. We conjecture, however, that the linear Boltzmann equation fails in the periodic setting for terms of order 4 and higher. Our proof uses Floquet–Bloch theory, multivariable theta series and equidistribution theorems for homogeneous flows. Compared with other scaling limits traditionally considered in homogenisation theory, the Boltzmann–Grad limit requires control of the quantum dynamics for longer times, which are inversely proportional to the total scattering cross-section of the single-site potential.

1. Introduction	571
2. Momentum representation	577
3. Bloch projections	578
4. Duhamel’s principle	580
5. The Boltzmann–Grad limit	584
6. Orders 0 and 1	586
7. Equidistribution of horocycles	587
8. Mean value theorems for theta functions	591
9. Order 2	596
10. Higher-order theta functions	600
11. Error terms	603
12. Averages over α	606
Appendix	608
Acknowledgements	612
References	613

1. Introduction

The analysis of wave transport in periodic media plays an important role in explaining numerous physical phenomena, most notably in solid state physics, continuum mechanics and optics. A key challenge is the derivation of macroscopic transport equations from the underlying microscopic laws, and to thus describe effects on scales which are several orders of magnitude above the length scale given by the period of the medium. Semiclassical analysis and homogenisation theory have produced a remarkable

MSC2010: 37A17, 82C10.

Keywords: theta functions, homogeneous dynamics, quantum transport.




Figure 1. Illustration of a wavepacket at time $t = 0$ with wavelength h in a \mathbb{Z}^d -periodic potential with interaction regions of diameter $2r$. For small r , the classical mean-free path length in this setting is of the order r^{1-d} .

collection of results in scaling limits where the characteristic wavelength is either much larger than the period (low-frequency homogenisation) or of the same or smaller order (high-frequency homogenisation); see for example [Allaire and Piatnitski 2005; Benoit and Gloria 2017; Bensoussan et al. 1978; Birman and Suslina 2003; Craster et al. 2010; Gérard 1991; Gérard et al. 1997; Harutyunyan et al. 2016; Markowich et al. 1994; Panati et al. 2003].

Here we study the limit when the diameter $2r$ of the interaction region in each fundamental cell is significantly smaller than the period, and the wavelength h is comparable to the interaction region; see Figure 1.

Such a scaling, which is not traditionally discussed in high-frequency homogenisation, is motivated by the desire to understand the Boltzmann–Grad limit of particle transport in crystals. This problem is currently only understood (a) in the case of zero quasimomentum [Castella 1999; 2001; Castella and Plagne 2002], (b) in the classical limit [Caglioti and Golse 2010; Marklof and Strömbergsson 2008; 2010; 2011a; 2011b], and (c) when the medium is random rather than periodic, in both the classical [Gallavotti 1969; Spohn 1978; Boldrighini et al. 1983] and quantum settings [Eng and Erdős 2005] (see also [Erdős and Yau 2000; Spohn 1977] for the weak-coupling limit and [Bal et al. 1999; 2002; 2011] for related models). In the random setting — classical and quantum — the limit transport equation is proved to be the *linear Boltzmann equation*, as predicted in [Lorentz 1905].

The linear Boltzmann equation for a particle density $f(t, \mathbf{x}, \mathbf{y})$ at time t , where \mathbf{x} denotes position and \mathbf{y} momentum, is given by

$$\partial_t f(t, \mathbf{x}, \mathbf{y}) + \mathbf{y} \cdot \nabla_{\mathbf{x}} f(t, \mathbf{x}, \mathbf{y}) = \rho(\mathbf{x}) \int_{\mathbb{R}^d} \Sigma(\mathbf{y}, \mathbf{y}') [f(t, \mathbf{x}, \mathbf{y}') - f(t, \mathbf{x}, \mathbf{y})] d\mathbf{y}', \quad (1-1)$$

subject to initial data $f(0, \mathbf{x}, \mathbf{y}) = a(\mathbf{x}, \mathbf{y})$. The collision kernel $\Sigma(\mathbf{y}, \mathbf{y}')$ is determined by the single-site scattering potential, and can be interpreted as the rate of particles with velocity \mathbf{y} being scattered to

velocity \mathbf{y}' (or vice versa). The quantity $\rho(\mathbf{x})$ denotes the macroscopic scatterer density at \mathbf{x} , which for a homogeneous medium means $\rho(\mathbf{x})$ is constant. In the absence of scatterers $\rho(\mathbf{x}) = 0$, and the solution of (1-1) is $f(t, \mathbf{x}, \mathbf{y}) = a(\mathbf{x} - t\mathbf{y}, \mathbf{y})$, which is consistent with free transport. In the case of a single scatterer, classical and semiclassical scattering theory yields a linear Boltzmann equation with $\rho(\mathbf{x}) = \delta(\mathbf{x})$ [Nier 1995]. See also [Nier 1996], in particular Section 7.2, for the case when $\rho(\mathbf{x})$ is an infinite superposition of point masses in dimension $d = 1$.

The principal result of the present work establishes convergence in the Boltzmann–Grad limit for the quantum periodic setting, at least up to second order in the coupling constant. Perhaps surprisingly, and unlike the classical case [Golse 2008], this limit is compatible with the linear Boltzmann equation. We nevertheless conjecture that higher-order terms in the coupling constant are incompatible, and that in particular the limit process does not satisfy the linear Boltzmann equation. A heuristic description of the full limit process will be provided elsewhere [Griffin and Marklof \geq 2019].

A technical step in this paper is to generalise the limit theorems for multivariable theta series, which were employed in the proof of the Berry–Tabor conjecture for the Laplacian on tori with quasiperiodic boundary conditions [Marklof 2002; 2003]. Crucial ingredients in the proofs of these statements are equidistribution results for homogeneous flows against unbounded test functions, which require estimates on the escape of mass into the cusp of the relevant homogeneous space. The results in [Marklof 2002; 2003] are based on Ratner’s measure classification theorem and are therefore ineffective. The recent paper [Strömbergsson and Vishe 2018] provides effective rate-of-convergence estimates in this context (we will not need these for the present study).

Given initial data f_0 in the Schwartz class $\mathcal{S}(\mathbb{R}^d)$ and scaling parameter $h > 0$, the quantum amplitude $f(t, \mathbf{x})$ at time t is given by the Schrödinger equation

$$i \frac{h}{2\pi} \partial_t f(t, \mathbf{x}) = H_{h,\lambda} f(t, \mathbf{x}), \quad f(0, \mathbf{x}) = f_0(\mathbf{x}), \tag{1-2}$$

with quantum Hamiltonian

$$H_{h,\lambda} = H_{h,0} + \lambda \text{Op}(V), \quad H_{h,0} = -\frac{h^2}{8\pi^2} \Delta. \tag{1-3}$$

Here Δ is the standard Laplacian in \mathbb{R}^d , and $\text{Op}(V)$ denotes multiplication by the \mathbb{Z}^d -periodic potential

$$V(\mathbf{x}) = V_r(\mathbf{x}) = \sum_{\mathbf{m} \in \mathbb{Z}^d} W(r^{-1}(\mathbf{x} + \mathbf{m})), \tag{1-4}$$

with a fixed single-site potential W . We will assume from here onwards that $d \geq 2$ and that $W \in \mathcal{S}(\mathbb{R}^d)$ is real-valued.

We expect that our analysis can be extended to scatterer configurations where \mathbb{Z}^d is replaced by an arbitrary Euclidean lattice \mathcal{L} of full rank in \mathbb{R}^d , and more generally to locally finite \mathcal{L} -periodic point sets. This requires, however, a substantial generalisation of the asymptotics discussed in Section 7, which are based on limit theorems for the pair correlation of general positive definite quadratic forms. The latter are currently understood, in the necessary scaling regime, only in dimension $d = 2$ [Eskin et al. 2005; Margulis and Mohammadi 2011].

The quantities $r, \lambda > 0$ are scaling parameters which we will refer to as the scattering radius and coupling constant respectively. The operator $H_{h,\lambda}$ can be realised as the Weyl quantisation of the classical Hamiltonian $H_\lambda^{\text{cl}}(\mathbf{x}, \mathbf{y}) = \frac{1}{2}\|\mathbf{y}\|^2 + \lambda V(\mathbf{x})$. The solution of (1-2) can be represented as $f(t, \mathbf{x}) = U_{h,\lambda}(t)f_0(\mathbf{x})$ with

$$U_{h,\lambda}(t) = e(-H_{h,\lambda}t/h), \quad e(z) := e^{2\pi iz}. \quad (1-5)$$

To characterise the asymptotic behaviour of the quantum dynamics, it will be convenient to use the time evolution of linear operators $A(t)$ (“quantum observables”) given by the Heisenberg evolution

$$A(t) = U_{h,\lambda}(t) A U_{h,\lambda}(t)^{-1}. \quad (1-6)$$

We will use the L^2 inner product

$$\langle a, b \rangle = \int_{\mathbb{R}^d \times \mathbb{R}^d} a(\mathbf{x}, \mathbf{y}) \overline{b(\mathbf{x}, \mathbf{y})} \, d\mathbf{x} \, d\mathbf{y}, \quad (1-7)$$

and the Hilbert–Schmidt inner product

$$\langle A, B \rangle_{\text{HS}} = \text{Tr} AB^\dagger. \quad (1-8)$$

As is standard in semiclassics, we will measure momentum in units of h , and use the rescaling $a(\mathbf{x}, \mathbf{y}) \mapsto h^{d/2}a(\mathbf{x}, h\mathbf{y})$; the normalisation is chosen so that the L^2 -norm is preserved. In the classical picture of a point particle moving through an infinite field of scatterers, the *Boltzmann–Grad* scaling limit is one in which the radius of the scatterers is taken to zero, while space and time are simultaneously rescaled in order to ensure the mean-free path length and mean-free flight time remain finite. The classical mean-free path length scales like r^{1-d} , and so we define the *semiclassical Boltzmann–Grad scaling* of $a \in \mathcal{S}(\mathbb{R}^d \times \mathbb{R}^d)$ by

$$D_{r,h}a(\mathbf{x}, \mathbf{y}) = r^{d(d-1)/2} h^{d/2} a(r^{d-1}\mathbf{x}, h\mathbf{y}), \quad (1-9)$$

where again the normalisation is chosen so that $D_{r,h}$ preserves the inner product (1-7). To ensure that the mean-free flight time remains of constant order as $r \rightarrow 0$ we similarly rescale time by a factor of r^{1-d} .

We denote by $\text{Op}(a)$ the standard Weyl quantisation of $a \in \mathcal{S}(\mathbb{R}^d \times \mathbb{R}^d)$:

$$\text{Op}(a)f(\mathbf{x}) = \int_{\mathbb{R}^d \times \mathbb{R}^d} a\left(\frac{1}{2}(\mathbf{x} + \mathbf{x}'), \mathbf{y}\right) e(i(\mathbf{x} - \mathbf{x}') \cdot \mathbf{y}) f(\mathbf{x}') \, d\mathbf{x}' \, d\mathbf{y}, \quad (1-10)$$

where $f \in \mathcal{S}(\mathbb{R}^d)$. We furthermore define the corresponding scaled quantisation by $\text{Op}_{r,h} = \text{Op} \circ D_{r,h}$, and set $\text{Op}_h = \text{Op}_{1,h}$.

Throughout this paper we will consider the scaling limit where the quantum wavelength is of the same order as the scattering radius r ; i.e., $h = h_0 r$ where h_0 is a fixed constant. By a simple scaling argument, we may assume without loss of generality that $h_0 = 1$.

Conjecture 1.1. *There exists a family of linear operators $L(t) : L^1(\mathbb{R}^d \times \mathbb{R}^d) \rightarrow L^1(\mathbb{R}^d \times \mathbb{R}^d)$ such that*

(i) *for all $a, b \in \mathcal{S}(\mathbb{R}^d \times \mathbb{R}^d)$, $A = \text{Op}_{r,h}(a)$, $B = \text{Op}_{r,h}(b)$, $\lambda > 0$ and $t > 0$,*

$$\lim_{h=r \rightarrow 0} \langle A(tr^{1-d}), B \rangle_{\text{HS}} = \langle L(t)a, b \rangle, \quad (1-11)$$

(ii) *$L(t)a(\mathbf{x}, \mathbf{y})$ is in general **not** a solution of the linear Boltzmann equation.*

Section A in the appendix provides an interpretation of $\langle A(tr^{1-d}), B \rangle_{\text{HS}}$ in terms of the phase-space distribution of a solution $f(t, \mathbf{x})$ of the Schrödinger equation (1-2) with initial condition

$$f_0(\mathbf{x}) = r^{d(d-1)/2} \phi(r^{d-1} \mathbf{x}) e(\mathbf{p} \cdot \mathbf{x} / h) \tag{1-12}$$

for $\phi \in \mathcal{S}(\mathbb{R}^d)$ and $\mathbf{p} \in \mathbb{R}^d$. A schematic drawing of the initial wavepacket f_0 is given in Figure 1 (shown is the positive real part of f_0).

In the case of random (rather than periodic) scatterer configurations, [Eng and Erdős 2005] proves convergence to a limit $L(t)a(\mathbf{x}, \mathbf{y})$, which in fact is a solution to the linear Boltzmann equation with the standard quantum mechanical collision kernel

$$\Sigma(\mathbf{y}, \mathbf{y}') = 8\pi^2 \delta(\|\mathbf{y}\|^2 - \|\mathbf{y}'\|^2) |T(\mathbf{y}, \mathbf{y}')|^2. \tag{1-13}$$

Here $T(\mathbf{y}, \mathbf{y}')$ is the kernel of the T -matrix in momentum representation. It is related to the quantum scattering cross-section by the formula (see [Nier 1995, Appendix A])

$$\sigma(\mathbf{y}, \mathbf{y}') = 4\pi^2 \|\mathbf{y}\|^{d-3} |T(\mathbf{y}, \mathbf{y}')|^2. \tag{1-14}$$

The Born approximation for the T -matrix yields Fermi's golden rule,

$$\Sigma_2(\mathbf{y}, \mathbf{y}') = 8\pi^2 \delta(\|\mathbf{y}\|^2 - \|\mathbf{y}'\|^2) |\widehat{W}(\mathbf{y} - \mathbf{y}')|^2, \tag{1-15}$$

where \widehat{W} is the Fourier transform of the single-site potential W .

We will use a perturbative approach to provide evidence for Conjecture 1.1: The present paper establishes convergence up to second order in the coupling constant λ , where all terms are consistent with the linear Boltzmann equation. Based on this analysis, we develop in [Griffin and Marklof ≥ 2019] a heuristic model for higher-order terms, some of which do not match the linear Boltzmann equation; this provides support for the second assertion of Conjecture 1.1. To formulate the main theorem of the present paper, consider the formal expansion

$$L(t) \sim \sum_{n=0}^{\infty} L_n(t) \lambda^n, \tag{1-16}$$

and define the linear operators L_0, L_1 and L_2 acting on functions in $\mathcal{S}(\mathbb{R}^d \times \mathbb{R}^d)$ by

$$L_0(t)a(\mathbf{x}, \mathbf{y}) = a(\mathbf{x} - t\mathbf{y}, \mathbf{y}), \quad L_1(t)a(\mathbf{x}, \mathbf{y}) = 0, \tag{1-17}$$

$$L_2(t)a(\mathbf{x}, \mathbf{y}) = \int_0^t \int_{\mathbb{R}^d} \Sigma_2(\mathbf{y}, \mathbf{y}') [a(\mathbf{x} - s\mathbf{y} - (t-s)\mathbf{y}', \mathbf{y}') - a(\mathbf{x} - t\mathbf{y}, \mathbf{y})] d\mathbf{y}' ds. \tag{1-18}$$

Relations (1-16)–(1-18) are consistent with $L(t)$ generating solutions of the linear Boltzmann equation with $\rho(\mathbf{x}) = 1$.

Our main result is as follows.

Theorem 1.2. *Let $t > 0$ and $a, b \in \mathcal{S}(\mathbb{R}^d \times \mathbb{R}^d)$, $A = \text{Op}_{r,h}(a)$, $B = \text{Op}_{r,h}(b)$. Then there exist linear operators $A_0^{(r)}(t), A_1^{(r)}(t), A_2^{(r)}(t)$ such that for $h = r \in (0, 1]$*

$$\langle A(tr^{1-d}), B \rangle_{\text{HS}} = \sum_{n=0}^2 \langle A_n^{(r)}(tr^{1-d}), B \rangle_{\text{HS}} \lambda^n + \sum_{n=3}^6 O(r^{-nd/2} \lambda^n) \tag{1-19}$$

and

$$\lim_{h=r \rightarrow 0} \langle A_n^{(r)}(tr^{1-d}), B \rangle_{\text{HS}} = \langle L_n(t)a, b \rangle \quad (n = 0, 1, 2). \tag{1-20}$$

The notation $f(x) = O(g(x))$ means “there is a positive constant C such that $|f(x)| \leq C |g(x)|$.” We will also use $f(x) \ll g(x)$ synonymously, and subscript O_ϵ or \ll_ϵ to highlight the dependence of the implied constant $C = C_\epsilon$ on a parameter ϵ .

The key point here is to view the first sum on the right-hand side of (1-19) as the first three terms of a formal power series expansion in λ , which according to (1-20) each converge to the corresponding terms of the conjectured limit (1-16). The second sum in (1-19) provides an error estimate that allows an interpretation beyond a formal power series, but this is only of secondary interest.

We will actually prove a stronger result than Theorem 1.2. For a given quasimomentum $\alpha \in [0, 1)^d$, consider the Bloch functions $\varphi_m^\alpha(\mathbf{x}) = e(i(\mathbf{m} + \alpha) \cdot \mathbf{x})$, $\mathbf{m} \in \mathbb{Z}^d$, and define the projection Π_α acting on $f \in \mathcal{S}(\mathbb{R}^d)$ by

$$\Pi_\alpha f(\mathbf{x}) = \sum_{\mathbf{m} \in \mathbb{Z}^d} \langle f, \varphi_m^\alpha \rangle \varphi_m^\alpha(\mathbf{x}), \tag{1-21}$$

with inner product

$$\langle f, g \rangle = \int_{\mathbb{R}^d} f(\mathbf{x}) \overline{g(\mathbf{x})} \, d\mathbf{x}. \tag{1-22}$$

Note that, by Poisson summation,

$$\Pi_\alpha f(\mathbf{x}) = \sum_{\mathbf{m} \in \mathbb{Z}^d} e(-\mathbf{m} \cdot \alpha) f(\mathbf{x} + \mathbf{m}), \tag{1-23}$$

and hence by integrating over $\alpha \in [0, 1)^d$ one regains $f(\mathbf{x})$. We will refer to Π_α as a *Bloch projection* and α as a *Bloch vector* or *quasimomentum*. Instead of (1-19) we consider now

$$\langle \Pi_\alpha A(tr^{1-d}), B \rangle_{\text{HS}}. \tag{1-24}$$

As we will see, the behaviour of (1-24) in the limit $h = r \rightarrow 0$ depends on the number-theoretic properties of α . We call a vector $\alpha = (\alpha_1, \dots, \alpha_d) \in \mathbb{R}^d$ *Diophantine of type κ* if there exists a constant $C > 0$ such that

$$\max_j \left| \alpha_j - \frac{m_j}{q} \right| > \frac{C}{q^\kappa} \tag{1-25}$$

for all $m_1, \dots, m_d, q \in \mathbb{Z}$, $q > 0$. The smallest possible value for κ is $\kappa = 1 + \frac{1}{d}$. In this case α is called *badly approximable*.

Theorem 1.3. *Suppose α is Diophantine of type $\kappa < (d - 1)/(d - 2)$ and the components of $(1, i\alpha)$ are linearly independent over \mathbb{Q} . Let $t > 0$ and $a, b \in \mathcal{S}(\mathbb{R}^d \times \mathbb{R}^d)$, $A = \text{Op}_{r,h}(a)$, $B = \text{Op}_{r,h}(b)$. Then there exist linear operators $A_0^{(r,\alpha)}(t)$, $A_1^{(r,\alpha)}(t)$, $A_2^{(r,\alpha)}(t)$ such that for $h = r \in (0, 1]$*

$$\langle \Pi_\alpha A(tr^{1-d}), B \rangle_{\text{HS}} = \sum_{n=0}^2 \langle A_n^{(r,\alpha)}(tr^{1-d}), B \rangle_{\text{HS}} \lambda^n + \sum_{n=3}^6 O(r^{-nd/2} \lambda^n) \tag{1-26}$$

and

$$\lim_{h=r \rightarrow 0} \langle A_n^{(r,\alpha)}(tr^{1-d}), B \rangle_{\text{HS}} = \langle L_n(t)a, b \rangle \quad (n = 0, 1, 2). \tag{1-27}$$

Since the set of Diophantine $\alpha \in [0, 1)^d$ has full Lebesgue measure, Theorem 1.2 may be viewed as an averaged (and thus weaker) version of Theorem 1.3. The convergence in (1-27) is however highly nonuniform in α , and the derivation of Theorem 1.2 from Theorem 1.3 requires nontrivial dominated convergence estimates.

In his PhD thesis [Griffin 2017], the first author established a version of Theorem 1.3 for the small-scatter problem on the torus $\mathbb{T}^d = \mathbb{R}^d / \mathbb{Z}^d$ with quasiperiodic boundary conditions $f(\mathbf{x} + \mathbf{m}) = e(\mathbf{m} \cdot \alpha) f(\mathbf{x})$ ($\mathbf{m} \in \mathbb{Z}^d$) for observables that do not depend on position \mathbf{x} . This in particular complements results in [Castella 1999; Castella and Plagne 2002] where $\alpha = \mathbf{0}$, and furthermore provides a discussion of the expansion terms leading to a failure of the linear Boltzmann equation. The key observation in those papers is that due to the large mean degeneracy of the spectrum of the Laplacian on the torus \mathbb{T}^d , the semiclassical Boltzmann–Grad limit diverges; a different normalisation then yields a nonuniversal limit, which in particular is not consistent with the linear Boltzmann equation. It is interesting to note that adding a suitably chosen damping term allows one to recover the linear Boltzmann equation even in this singular case [Castella 2001; 2002]. The small-scatterer problem in rectangular domains (Sinai billiards) has also been investigated in the context of quantum chaos; here the smooth potential is replaced by a disc with Dirichlet boundary conditions [Berry 1981; Dahlqvist and Vattay 1998].

This paper is organised as follows. Sections 2 and 3 provide basic background and notation on Weyl calculus in momentum representation and Floquet–Bloch theory. Section 4 uses the Duhamel principle to obtain a perturbation series in λ . We then apply the Boltzmann–Grad scaling in Section 5. The zeroth- and first-order terms are elementary, and are calculated in Section 6. Terms of second order require equidistribution results for horocycles (Section 7) and mean value theorems for theta functions (Section 8), which build on the papers [Marklof 2002; 2003]. The second-order terms are computed in Section 9. The estimates of the error term in Theorem 1.3 require analogous results for higher-dimensional theta functions (Section 10) and are presented in Section 11. The proof of Theorem 1.3 is given at the end of Section 11. Section 12 concludes with the proof of Theorem 1.2.

2. Momentum representation

We have so far represented quantum wave amplitudes f in the position representation. It will in fact be more convenient to work with its Fourier transform \hat{f} , which represents the wave amplitude as a function of the quantum particle’s momentum. Set $e(x) = \exp(2\pi i x)$, and define the Fourier transform $\hat{f} = \mathcal{F} f$ of f by

$$\hat{f}(\mathbf{y}) = \mathcal{F} f(\mathbf{y}) = \int_{\mathbb{R}^d} e(-\mathbf{y} \cdot \mathbf{x}) f(\mathbf{x}) \, d\mathbf{x}. \tag{2-1}$$

The Fourier transform of a linear operator A on $L^2(\mathbb{R}^d)$ is then naturally defined by

$$\hat{A} = \mathcal{F} A \mathcal{F}^{-1}. \tag{2-2}$$

Explicitly, the corresponding Schwartz kernel satisfies

$$\hat{A}(\mathbf{y}, \mathbf{y}') = \int_{\mathbb{R}^{2d}} A(\mathbf{x}, \mathbf{x}') e(-\mathbf{x} \cdot \mathbf{y} + \mathbf{x}' \cdot \mathbf{y}') \, d\mathbf{x} \, d\mathbf{x}'. \tag{2-3}$$

The Schwartz kernel of the Fourier transform of $\text{Op}(a)$ is given by

$$\begin{aligned}\widehat{\text{Op}}(a)(\mathbf{y}, \mathbf{y}') &= \int_{\mathbb{R}^d} a(\mathbf{x}, \tfrac{1}{2}(\mathbf{y} + \mathbf{y}')) e(-\mathbf{x} \cdot (\mathbf{y} - \mathbf{y}')) d\mathbf{x} \\ &= \tilde{a}(\mathbf{y} - \mathbf{y}', \tfrac{1}{2}(\mathbf{y} + \mathbf{y}')), \end{aligned} \quad (2-4)$$

where \tilde{a} denotes the Fourier transform of a in the first variable only; i.e.,

$$\tilde{a}(\boldsymbol{\eta}, \mathbf{y}) = \int_{\mathbb{R}^d} a(\mathbf{x}, \mathbf{y}) e(-\mathbf{x} \cdot \boldsymbol{\eta}) d\mathbf{x}. \quad (2-5)$$

The definition given above extends to larger function spaces by standard arguments [Folland 1989]. Two notable special cases occur when a is a function exclusively of either \mathbf{x} or \mathbf{y} . In the first case when $a = a(\mathbf{x})$ we have $\widehat{\text{Op}}(a)(\mathbf{y}, \mathbf{y}') = \hat{a}(\mathbf{y} - \mathbf{y}')$, and in the second case when $a = a(\mathbf{y})$ we obtain $\widehat{\text{Op}}(a)(\mathbf{y}, \mathbf{y}') = a(\mathbf{y}) \delta_{\mathbf{0}}(\mathbf{y} - \mathbf{y}')$. The choice $a = L_0(t)V$ in (2-4) yields for instance

$$\widehat{\text{Op}}(L_0(t)V)(\mathbf{y}, \mathbf{y}') = r^d \sum_{\mathbf{m} \in \mathbb{Z}^d} \widehat{W}(r\mathbf{m}) e(-\tfrac{1}{2}t\mathbf{m} \cdot (\mathbf{y} + \mathbf{y}')) \delta_{\mathbf{m}}(\mathbf{y} - \mathbf{y}'), \quad (2-6)$$

where $\delta_{\mathbf{m}}$ denotes the Dirac delta mass at the point \mathbf{m} .

The quantisations of the Hamilton functions H_0^{cl} and H_{λ}^{cl} are denoted by $H_0 = \text{Op } H_0^{\text{cl}} = -\frac{1}{8\pi^2} \Delta$ and $H_{\lambda} = \text{Op } H_{\lambda}^{\text{cl}} = H_0 + \lambda \text{Op } V$ respectively. The Schrödinger equation for the time evolution of the wave amplitude $f(t, \mathbf{x})$ can then be written (in units where Planck's constant is 1)

$$\frac{i}{2\pi} \partial_t f(t, \mathbf{x}) = H_{\lambda} f(t, \mathbf{x}), \quad f(0, \mathbf{x}) = f_0(\mathbf{x}), \quad (2-7)$$

which has the solution

$$f(t, \mathbf{x}) = U_{\lambda}(t) f_0(\mathbf{x}), \quad U_{\lambda}(t) := e(-H_{\lambda}t). \quad (2-8)$$

The relation to the corresponding operators in the Introduction is

$$H_{h,\lambda} = h^2 H_{\lambda/h^2}, \quad U_{h,\lambda}(t) = U_{\lambda/h^2}(ht). \quad (2-9)$$

It will be more convenient to work with $U_{\lambda}(t)$ in what follows, and then later appeal to (2-9).

Since H_0^{cl} is a quadratic polynomial, we have the exact Egorov property,

$$U_0(t) \text{Op}(a) U_0(-t) = \text{Op}(L_0(t)a). \quad (2-10)$$

In momentum representation the kernel of the operator \widehat{H}_0 takes the form

$$\widehat{H}_0(\mathbf{y}, \mathbf{y}') = \tfrac{1}{2} \|\mathbf{y}\|^2 \delta_{\mathbf{0}}(\mathbf{y} - \mathbf{y}') \quad (2-11)$$

and thus also

$$\widehat{U}_0(t)(\mathbf{y}, \mathbf{y}') = e(-\tfrac{1}{2}t \|\mathbf{y}\|^2) \delta_{\mathbf{0}}(\mathbf{y} - \mathbf{y}'). \quad (2-12)$$

3. Bloch projections

As is standard in the study of periodic potentials, we use the fact that any solution to our Schrödinger equation can be decomposed into quasiperiodic functions parametrised by quasimomentum $\boldsymbol{\alpha} \in \mathbb{T}^d = \mathbb{R}^d / \mathbb{Z}^d$

(Floquet–Bloch decomposition). For $f \in \mathcal{S}(\mathbb{R}^d)$ the function $\psi(\mathbf{x}) = \Pi_\alpha f(\mathbf{x})$ satisfies, for every $\mathbf{k} \in \mathbb{Z}^d$,

$$\psi(\mathbf{x} + \mathbf{k}) = e(\mathbf{k} \cdot \boldsymbol{\alpha})\psi(\mathbf{x}). \tag{3-1}$$

We denote by \mathcal{H}_α the Hilbert space of functions that satisfy the quasiperiodicity condition (3-1) and have finite L^2 -norm with respect to the inner product

$$\langle \psi, \varphi \rangle_\alpha = \int_{\mathbb{T}^d} \psi(\mathbf{x}) \overline{\varphi(\mathbf{x})} \, d\mathbf{x}. \tag{3-2}$$

We define the corresponding Hilbert–Schmidt product for linear operators from $L^2(\mathbb{R}^d)$ to \mathcal{H}_α by

$$\langle A, B \rangle_{\text{HS}, \alpha} = \text{Tr} AB^\dagger = \int_{\mathbb{T}^d} \left(\int_{\mathbb{R}^d} A(\mathbf{x}, \mathbf{x}') \overline{B(\mathbf{x}, \mathbf{x}')} \, d\mathbf{x}' \right) d\mathbf{x}. \tag{3-3}$$

Lemma 3.1. *If $f, g \in \mathcal{S}(\mathbb{R}^d)$, then $\Pi_\alpha f, \Pi_\alpha g \in \mathcal{H}_\alpha \cap C^\infty(\mathbb{R}^d)$ and*

$$\langle \Pi_\alpha f, g \rangle = \langle f, \Pi_\alpha g \rangle = \langle \Pi_\alpha f, \Pi_\alpha g \rangle_\alpha = \sum_{\mathbf{m} \in \mathbb{Z}^d} \hat{f}(\mathbf{m} + \boldsymbol{\alpha}) \overline{\hat{g}(\mathbf{m} + \boldsymbol{\alpha})}. \tag{3-4}$$

Proof. We have by (1-23)

$$\langle \Pi_\alpha f, \Pi_\alpha g \rangle_\alpha = \sum_{\mathbf{m} \in \mathbb{Z}^d} e(\mathbf{m} \cdot \boldsymbol{\alpha}) \int_{\mathbb{T}^d} (\Pi_\alpha f)(\mathbf{x}) \overline{g(\mathbf{x} + \mathbf{m})} \, d\mathbf{x}. \tag{3-5}$$

Using the invariance (3-1) of $\Pi_\alpha f$, we see that the summation and integration can be combined to an integral over \mathbb{R}^d which equals $\langle \Pi_\alpha f, g \rangle$. The final identity follows directly from the definition (1-21), which yields

$$\langle \Pi_\alpha f, g \rangle = \sum_{\mathbf{m} \in \mathbb{Z}^d} \langle f, \varphi_{\mathbf{m}}^\alpha \rangle \langle \varphi_{\mathbf{m}}^\alpha, g \rangle = \sum_{\mathbf{m} \in \mathbb{Z}^d} \hat{f}(\mathbf{m} + \boldsymbol{\alpha}) \overline{\hat{g}(\mathbf{m} + \boldsymbol{\alpha})}, \tag{3-6}$$

concluding the proof. □

Note that for the Fourier transform,

$$\widehat{\Pi_\alpha f}(\mathbf{y}) = \sum_{\mathbf{m} \in \mathbb{Z}^d} f(\mathbf{m} + \boldsymbol{\alpha}) \delta_{\mathbf{m} + \boldsymbol{\alpha}}(\mathbf{y}). \tag{3-7}$$

Lemma 3.2. *If A, B have Schwartz kernel in $\mathcal{S}(\mathbb{R}^d \times \mathbb{R}^d)$, then $\Pi_\alpha A, \Pi_\alpha B$ are linear operators $L^2(\mathbb{R}^d) \rightarrow \mathcal{H}_\alpha$, and*

$$\begin{aligned} \langle \Pi_\alpha A, B \rangle_{\text{HS}} &= \langle A, \Pi_\alpha B \rangle_{\text{HS}} = \langle \Pi_\alpha A, \Pi_\alpha B \rangle_{\text{HS}, \alpha} \\ &= \sum_{\mathbf{m} \in \mathbb{Z}^d} \int_{\mathbb{R}^d} \widehat{A}(\mathbf{m} + \boldsymbol{\alpha}, \mathbf{y}) \overline{\widehat{B}(\mathbf{m} + \boldsymbol{\alpha}, \mathbf{y})} \, d\mathbf{y}. \end{aligned} \tag{3-8}$$

Proof. This is analogous to the proof of Lemma 3.1. By (1-21), we have

$$[\Pi_\alpha B](\mathbf{x}, \mathbf{x}') = \sum_{\mathbf{m} \in \mathbb{Z}^d} e(-\mathbf{m} \cdot \boldsymbol{\alpha}) B(\mathbf{x} + \mathbf{m}, \mathbf{x}'), \tag{3-9}$$

and so

$$\begin{aligned} \langle \Pi_\alpha A, \Pi_\alpha B \rangle_{\text{HS}, \alpha} &= \sum_{\mathbf{m} \in \mathbb{Z}^d} e(\mathbf{m} \cdot \boldsymbol{\alpha}) \int_{\mathbb{T}^d} \left(\int_{\mathbb{R}^d} [\Pi_\alpha A](x, x') \overline{B(x + \mathbf{m}, x')} dx' \right) dx \\ &= \int_{\mathbb{R}^d} \left(\int_{\mathbb{R}^d} [\Pi_\alpha A](x, x') \overline{B(x, x')} dx' \right) dx = \langle \Pi_\alpha A, B \rangle_{\text{HS}}, \end{aligned} \quad (3-10)$$

where we have used the identity $[\Pi_\alpha A](x + \mathbf{m}, x') = e(\mathbf{m} \cdot \boldsymbol{\alpha})[\Pi_\alpha A](x, x')$; see (3-1). The proof of $\langle A, \Pi_\alpha B \rangle_{\text{HS}} = \langle \Pi_\alpha A, \Pi_\alpha B \rangle_{\text{HS}, \alpha}$ is analogous. Finally, in view of (2-3) and (3-7) we have

$$[\widehat{\Pi_\alpha A}](y, y') = \sum_{\mathbf{m} \in \mathbb{Z}^d} \delta_{\mathbf{m} + \boldsymbol{\alpha}}(y) \widehat{A}(\mathbf{m} + \boldsymbol{\alpha}, y'), \quad (3-11)$$

which yields

$$\begin{aligned} \langle \Pi_\alpha A, B \rangle_{\text{HS}} &= \int_{\mathbb{R}^{2d}} \sum_{\mathbf{m} \in \mathbb{Z}^d} \delta_{\mathbf{m} + \boldsymbol{\alpha}}(y) \widehat{A}(\mathbf{m} + \boldsymbol{\alpha}, y') \overline{\widehat{B}(y, y')} dy dy' \\ &= \sum_{\mathbf{m} \in \mathbb{Z}^d} \int_{\mathbb{R}^d} \widehat{A}(\mathbf{m} + \boldsymbol{\alpha}, y) \overline{\widehat{B}(\mathbf{m} + \boldsymbol{\alpha}, y)} dy, \end{aligned} \quad (3-12)$$

completing the proof. \square

We denote by Δ^α the standard Laplacian acting on \mathcal{H}_α , and set

$$H_\lambda^\alpha = H_0^\alpha + \lambda \text{Op}(V), \quad U_\lambda^\alpha(t) = e(-H_\lambda^\alpha t). \quad (3-13)$$

Lemma 3.3. For $f \in \mathcal{S}(\mathbb{R}^d)$,

$$\Pi_\alpha U_\lambda(t) f = U_\lambda^\alpha(t) \Pi_\alpha f. \quad (3-14)$$

Proof. We have the commutation relations

$$\Pi_\alpha H_0 = H_0^\alpha \Pi_\alpha, \quad \Pi_\alpha \text{Op}(V) = \text{Op}(V) \Pi_\alpha. \quad (3-15)$$

Consider the time derivative of the left-hand side of (3-14),

$$\begin{aligned} \partial_t \Pi_\alpha U_\lambda(t) f &= -2\pi i \Pi_\alpha (H_0 + \lambda \text{Op}(V)) U_\lambda(t) f \\ &= -2\pi i (H_0^\alpha + \lambda \text{Op}(V)) \Pi_\alpha U_\lambda(t) f. \end{aligned} \quad (3-16)$$

Thus the left-hand side of (3-14) is the unique solution to

$$\partial_t g(t, y) = -2\pi i H_\lambda^\alpha g(t, y) \quad (3-17)$$

with initial condition $g(0, y) := \Pi_\alpha f(y)$. The right-hand side of (3-14) solves the same PDE, and the proof is complete. \square

4. Duhamel's principle

Duhamel's principle provides an explicit expansion of the solution in terms of the coupling constant λ . By truncating the expansion at order 2, we will be left with theta functions that, in a certain scaling limit, can

be treated with the tools of homogeneous dynamics. The explicit error terms can be handled separately. Our first aim is to work out the time evolution of unscaled observables,

$$U_\lambda(t) \text{Op}(a) U_\lambda(-t), \tag{4-1}$$

perturbatively in λ . We first study the problem in the interaction picture; i.e., consider

$$U_\lambda(t) U_0(-t) \text{Op}(a) U_0(t) U_\lambda(-t). \tag{4-2}$$

Note that in view of the Egorov property (2-10) this is equivalent to the original problem upon replacing a by $L_0(t)a$. We define the operators $K(t)$ and $R(t)$ for $t \in \mathbb{R}$ by

$$K(t) = U_0(t) \text{Op}(V) U_0(-t) \quad \text{and} \quad R(t) = U_\lambda(t) U_0(-t). \tag{4-3}$$

Furthermore, for $s = (s_1, \dots, s_n)$ and $\ell \leq n$ we denote by $K_{\ell,n}(s)$ the product

$$K_{\ell,n}(s) = K(s_\ell) \cdots K(s_n). \tag{4-4}$$

Then

$$\langle \Pi_\alpha U_\lambda(t) U_0(-t) \text{Op}(a) U_0(t) U_\lambda(-t), \text{Op}(b) \rangle_{\text{HS}} = \langle \Pi_\alpha R(t) \text{Op}(a) R(t)^{-1}, \text{Op}(b) \rangle_{\text{HS}}. \tag{4-5}$$

Duhamel's principle asserts that

$$R(t) = I - 2\pi i \lambda \int_0^t R(s) K(s) ds, \tag{4-6}$$

and iterating this expression N times yields

$$R(t) = \sum_{n=0}^N \lambda^n R_n(t) + \lambda^{N+1} R_{N+1,\varepsilon}(t), \tag{4-7}$$

where $R_0(t) = I$ and

$$R_n(t) = (-2\pi i)^n \int_{0 < s_1 < \dots < s_n < t} K_{1,n}(s) ds \quad (n \geq 1). \tag{4-8}$$

The error term is similarly given by

$$R_{N+1,\varepsilon}(t) = (-2\pi i)^{N+1} \int_{0 < s_1 < \dots < s_{N+1} < t} R(s_1) K_{1,N+1}(s) ds. \tag{4-9}$$

The inverse of $R(t)$ can be calculated by taking the Hermitian conjugate. It is given by

$$R(t)^{-1} = \sum_{n=0}^N \lambda^n R_n^-(t) + \lambda^{N+1} R_{N+1,\varepsilon}^-(t), \tag{4-10}$$

where $R_0^-(t) = I$,

$$R_n^-(t) = (2\pi i)^n \int_{0 < s_n < \dots < s_1 < t} K_{1,n}(s) ds \quad (n \geq 1), \tag{4-11}$$

and the error term is

$$R_{N+1,\varepsilon}^-(t) = (2\pi i)^{N+1} \int_{0 < s_{N+1} < \dots < s_1 < t} K_{1,N+1}(s) R(s_{N+1})^{-1} ds. \quad (4-12)$$

We have also used the fact that $\text{Op}(V) = \text{Op}(V)^\dagger$ (since V is real-valued) and thus $K(t) = K(t)^\dagger$. Our methods will permit explicit calculation of the terms in this expansion up to order 2, and so specializing to the case $N = 2$ the expansion takes the following form

$$\langle \Pi_\alpha U_\lambda(t) U_0(-t) \text{Op}(a) U_0(t) U_\lambda(-t), \text{Op}(b) \rangle_{\text{HS}} = \sum_{n=0}^6 \lambda^n Q_n(t, a, b), \quad (4-13)$$

with the main terms Q_0 to Q_2 given by

$$\begin{aligned} Q_0(t, a, b) &= \langle \Pi_\alpha \text{Op}(a), \text{Op}(b) \rangle_{\text{HS}}, \\ Q_1(t, a, b) &= \langle \Pi_\alpha R_1(t) \text{Op}(a), \text{Op}(b) \rangle_{\text{HS}} + \langle \Pi_\alpha \text{Op}(a) R_1^-(t), \text{Op}(b) \rangle_{\text{HS}}, \\ Q_2(t, a, b) &= \langle \Pi_\alpha R_2(t) \text{Op}(a), \text{Op}(b) \rangle_{\text{HS}} + \langle \Pi_\alpha R_1(t) \text{Op}(a) R_1^-(t), \text{Op}(b) \rangle_{\text{HS}} \\ &\quad + \langle \Pi_\alpha \text{Op}(a) R_2^-(t), \text{Op}(b) \rangle_{\text{HS}}. \end{aligned} \quad (4-14)$$

The error terms Q_3 through Q_6 are given by

$$\begin{aligned} Q_3(t, a, b) &= \langle \Pi_\alpha R_{3,\varepsilon}(t) \text{Op}(a), \text{Op}(b) \rangle_{\text{HS}} + \langle \Pi_\alpha R_2(t) \text{Op}(a) R_1^-(t), \text{Op}(b) \rangle_{\text{HS}} \\ &\quad + \langle \Pi_\alpha R_1(t) \text{Op}(a) R_2^-(t), \text{Op}(b) \rangle_{\text{HS}} + \langle \Pi_\alpha \text{Op}(a) R_{3,\varepsilon}^-(t), \text{Op}(b) \rangle_{\text{HS}}, \\ Q_4(t, a, b) &= \langle \Pi_\alpha R_{3,\varepsilon}(t) \text{Op}(a) R_1^-(t), \text{Op}(b) \rangle_{\text{HS}} \\ &\quad + \langle \Pi_\alpha R_2(t) \text{Op}(a) R_2^-(t), \text{Op}(b) \rangle_{\text{HS}} + \langle \Pi_\alpha R_1(t) \text{Op}(a) R_{3,\varepsilon}^-(t), \text{Op}(b) \rangle_{\text{HS}}, \\ Q_5(t, a, b) &= \langle \Pi_\alpha R_{3,\varepsilon}(t) \text{Op}(a) R_2^-(t), \text{Op}(b) \rangle_{\text{HS}} + \langle \Pi_\alpha R_2(t) \text{Op}(a) R_{3,\varepsilon}^-(t), \text{Op}(b) \rangle_{\text{HS}}, \\ Q_6(t, a, b) &= \langle \Pi_\alpha R_{3,\varepsilon}(t) \text{Op}(a) R_{3,\varepsilon}^-(t), \text{Op}(b) \rangle_{\text{HS}}. \end{aligned} \quad (4-15)$$

We will treat these error terms in the following way. First of all, Lemma 4.1 shows that all of the Q_j can be bounded above by quantities which are independent of $U_\lambda(t)$, and depend only on the free evolution $U_0(t)$. Then after rescaling, the resulting quantities, which we denote by $\mathcal{J}_{\ell,n}$, can be treated with similar techniques to those used in the computation of the limit of the second-order terms.

Define

$$\mathcal{J}_{\ell,n}(t, a) = (2\pi)^n \int_{\substack{0 < s_1 < \dots < s_\ell < t \\ 0 < s_n < \dots < s_{\ell+1} < t}} \|\Pi_\alpha K_{1,\ell}(s) \text{Op}(a) K_{\ell+1,n}(s)\|_{\text{HS},\alpha} ds. \quad (4-16)$$

Lemma 4.1. For $a, b \in \mathcal{S}(\mathbb{R}^d)$,

$$\begin{aligned} |\langle \Pi_\alpha R_\ell(t) \text{Op}(a) R_{n-\ell}^-(t), \text{Op}(b) \rangle_{\text{HS}}| &\leq \mathcal{J}_{\ell,n}(t, a) \|\Pi_\alpha \text{Op}(b)\|_{\text{HS},\alpha}, \\ |\langle \Pi_\alpha R_\ell(t) \text{Op}(a) R_{n-\ell,\varepsilon}^-(t), \text{Op}(b) \rangle_{\text{HS}}| &\leq \mathcal{J}_{\ell,n}(t, a) \|\Pi_\alpha \text{Op}(b)\|_{\text{HS},\alpha}, \\ |\langle \Pi_\alpha R_{\ell,\varepsilon}(t) \text{Op}(a) R_{n-\ell}^-(t), \text{Op}(b) \rangle_{\text{HS}}| &\leq \mathcal{J}_{\ell,n}(t, a) \|\Pi_\alpha \text{Op}(b)\|_{\text{HS},\alpha}, \\ |\langle \Pi_\alpha R_{\ell,\varepsilon}(t) \text{Op}(a) R_{n-\ell,\varepsilon}^-(t), \text{Op}(b) \rangle_{\text{HS}}| &\leq \mathcal{J}_{\ell,n}(t, a) \|\Pi_\alpha \text{Op}(b)\|_{\text{HS},\alpha}. \end{aligned} \quad (4-17)$$

Proof. For the first bound, note that by Lemma 3.2 and direct computation we have

$$\begin{aligned}
 & |\langle \Pi_\alpha R_\ell(t) \text{Op}(a) R_{n-\ell}^-(t), \text{Op}(b) \rangle_{\text{HS}}| \\
 &= |\langle \Pi_\alpha R_\ell(t) \text{Op}(a) R_{n-\ell}^-(t), \Pi_\alpha \text{Op}(b) \rangle_{\text{HS}, \alpha}| \\
 &= (2\pi)^n \left| \int_{\substack{0 < s_1 < \dots < s_\ell < t \\ 0 < s_n < \dots < s_{\ell+1} < t}} \langle \Pi_\alpha K_{1,\ell}(s) \text{Op}(a) K_{\ell+1,n}(s), \Pi_\alpha \text{Op}(b) \rangle_{\text{HS}, \alpha} ds \right| \\
 &\leq (2\pi)^n \int_{\substack{0 < s_1 < \dots < s_\ell < t \\ 0 < s_n < \dots < s_{\ell+1} < t}} |\langle \Pi_\alpha K_{1,\ell}(s) \text{Op}(a) K_{\ell+1,n}(s), \Pi_\alpha \text{Op}(b) \rangle_{\text{HS}, \alpha}| ds. \quad (4-18)
 \end{aligned}$$

The bound then follows by an application of the Cauchy–Schwarz inequality. For the second bound we similarly have

$$\begin{aligned}
 & \langle \Pi_\alpha R_\ell(t) \text{Op}(a) R_{n-\ell, \mathcal{E}}^-(t), \text{Op}(b) \rangle_{\text{HS}} \\
 &\leq (2\pi)^n \int_{\substack{0 < s_1 < \dots < s_\ell < t \\ 0 < s_n < \dots < s_{\ell+1} < t}} |\langle \Pi_\alpha K_{1,\ell}(s) \text{Op}(a) K_{\ell+1,n}(s) R(s_n)^{-1}, \Pi_\alpha \text{Op}(b) \rangle_{\text{HS}, \alpha}| ds. \quad (4-19)
 \end{aligned}$$

The result then follows by applying Cauchy–Schwarz and using that $R(s_n)$ is unitary. For the third bound we have

$$\begin{aligned}
 & \langle \Pi_\alpha R_{\ell, \mathcal{E}}(t) \text{Op}(a) R_{n-\ell}^-(t), \text{Op}(b) \rangle_{\text{HS}} \\
 &\leq (2\pi)^n \int_{\substack{0 < s_1 < \dots < s_\ell < t \\ 0 < s_n < \dots < s_{\ell+1} < t}} |\langle \Pi_\alpha R(s_1) K_{1,\ell}(s) \text{Op}(a) K_{\ell+1,n}(s), \Pi_\alpha \text{Op}(b) \rangle_{\text{HS}, \alpha}| ds. \quad (4-20)
 \end{aligned}$$

This time the bound follows by first applying Lemma 3.3, then the Cauchy–Schwarz inequality and finally using the unitarity of $R(s)$. The last bound follows by combining the arguments for the second and third bounds. \square

Let us introduce the shorthand

$$\mathcal{T}_{\ell,n}(\mathbf{y}) = \begin{cases} \prod_{j=\ell}^n e^{-\frac{1}{2}(s_{j+1} - s_j) \|\mathbf{y} - \mathbf{m}_j\|^2} \widehat{W}(r(\mathbf{m}_{j+1} - \mathbf{m}_j)) & (l \leq n), \\ 1 & (l > n). \end{cases} \quad (4-21)$$

Lemma 4.2. *The kernel of $\widehat{K}_{\ell,n}(s) = \mathcal{F}K_{\ell,n}(s)\mathcal{F}^{-1}$ is explicitly given by*

$$\begin{aligned}
 & [\widehat{K}_{\ell,n}(s)](\mathbf{y}, \mathbf{y}') \\
 &= r^{(n-\ell+1)d} \sum_{\mathbf{m}_\ell, \dots, \mathbf{m}_n \in \mathbb{Z}^d} e^{-\frac{1}{2} s_\ell \|\mathbf{y}\|^2} \widehat{W}(r\mathbf{m}_\ell) \mathcal{T}_{\ell,n-1}(\mathbf{y}) e^{\frac{1}{2} s_n \|\mathbf{y} - \mathbf{m}_n\|^2} \delta_{\mathbf{m}_n}(\mathbf{y} - \mathbf{y}'). \quad (4-22)
 \end{aligned}$$

Proof. We have

$$\begin{aligned}
 & \widehat{K}_{\ell,n}(s) f(\mathbf{y}) = \widehat{K}(s_\ell) \cdots \widehat{K}(s_n) f(\mathbf{y}) \\
 &= \mathcal{F}U_0(s_\ell) \text{Op}(V)U_0(s_{\ell+1} - s_\ell) \cdots U_0(s_n - s_{n-1}) \text{Op}(V)U_0(-s_n)\mathcal{F}^{-1} f(\mathbf{y}), \quad (4-23)
 \end{aligned}$$

and

$$\mathcal{F}U_0(s) \text{Op}(V)\mathcal{F}^{-1} f(\mathbf{y}) = r^d e^{-\frac{1}{2}s\|\mathbf{y}\|^2} \sum_{\mathbf{m} \in \mathbb{Z}^d} \widehat{W}(r\mathbf{m}) f(\mathbf{y} - \mathbf{m}). \quad (4-24)$$

By iterating we thus see

$$\begin{aligned}
\widehat{K}_{\ell,n}(s) f(\mathbf{y}) &= \widehat{K}(s_\ell) \cdots \widehat{K}(s_n) f(\mathbf{y}) \\
&= r^{(n-\ell+1)d} e\left(-\frac{1}{2}s_\ell \|\mathbf{y}\|^2\right) \\
&\quad \times \sum_{\mathbf{m}_\ell, \dots, \mathbf{m}_n \in \mathbb{Z}^d} \widehat{W}(r\mathbf{m}_\ell) e\left(-\frac{1}{2}(s_{\ell+1}-s_\ell) \|\mathbf{y}-\mathbf{m}_\ell\|^2\right) \widehat{W}(r\mathbf{m}_{\ell+1}) \\
&\quad \quad \times e\left(-\frac{1}{2}(s_{\ell+2}-s_{\ell+1}) \|\mathbf{y}-\mathbf{m}_\ell-\mathbf{m}_{\ell+1}\|^2\right) \widehat{W}(r\mathbf{m}_{\ell+2}) \\
&\quad \quad \quad \times \cdots \times e\left(-\frac{1}{2}(s_n-s_{n-1}) \|\mathbf{y}-\mathbf{m}_\ell-\cdots-\mathbf{m}_{n-1}\|^2\right) \widehat{W}(r\mathbf{m}_n) \\
&\quad \quad \quad \times e\left(\frac{1}{2}s_n \|\mathbf{y}-\mathbf{m}_\ell-\cdots-\mathbf{m}_n\|^2\right) f(\mathbf{y}-\mathbf{m}_\ell-\cdots-\mathbf{m}_n). \tag{4-25}
\end{aligned}$$

We then make the variable substitutions $\mathbf{m}_j = \tilde{\mathbf{m}}_j - \sum_{i=\ell}^{j-1} \mathbf{m}_i$ for $j = \ell + 1, \dots, n$. Note that this gives $\mathbf{y} - \mathbf{m}_\ell - \cdots - \mathbf{m}_j = \mathbf{y} - \tilde{\mathbf{m}}_j$ and also $\mathbf{m}_j = \tilde{\mathbf{m}}_j - \tilde{\mathbf{m}}_{j-1}$. Inserting these new variables, dropping the tildes, and using the definition of $\mathcal{T}_{\ell,n}$ yields the result. \square

5. The Boltzmann–Grad limit

Recall the semiclassical Boltzmann–Grad scaling (1-9) given by

$$D_{r,h}a(\mathbf{x}, \mathbf{y}) = r^{d(d-1)/2} h^{d/2} a(r^{d-1}\mathbf{x}, h\mathbf{y}). \tag{5-1}$$

Performing the Fourier transform in the \mathbf{x} -variable yields the expression

$$\widetilde{D}_{r,h}\tilde{a}(\boldsymbol{\eta}, \mathbf{y}) = \widetilde{(D_{r,h}a)}(\boldsymbol{\eta}, \mathbf{y}) = r^{-d(d-1)/2} h^{d/2} \tilde{a}(r^{1-d}\boldsymbol{\eta}, h\mathbf{y}), \tag{5-2}$$

and thus after quantizing the rescaled observables we see

$$\widehat{\text{Op}}(D_{r,h}a)(\mathbf{y}, \mathbf{y}') = r^{-d(d-1)/2} h^{d/2} \tilde{a}(r^{1-d}(\mathbf{y} - \mathbf{y}'), \frac{h}{2}(\mathbf{y} + \mathbf{y}')). \tag{5-3}$$

Note that after this rescaling we have the relation

$$\begin{aligned}
D_{r,h}L_0(t)a(\mathbf{x}, \mathbf{y}) &= r^{d(d-1)/2} h^{d/2} L_0(t)a(r^{d-1}\mathbf{x}, h\mathbf{y}) \\
&= r^{d(d-1)/2} h^{d/2} a(r^{d-1}\mathbf{x} - th\mathbf{y}, h\mathbf{y}) \\
&= D_{r,h}a(\mathbf{x} - thr^{1-d}\mathbf{y}, \mathbf{y}) \\
&= L_0(thr^{1-d})D_{r,h}a(\mathbf{x}, \mathbf{y}) \tag{5-4}
\end{aligned}$$

and so the Egorov property (2-10) becomes

$$U_0(thr^{1-d}) \text{Op}(D_{r,h}a) U_0(-thr^{1-d}) = \text{Op}(D_{r,h}L_0(t)a). \tag{5-5}$$

Given a linear operator A on $L^2(\mathbb{R}^d)$ with Schwartz kernel in $\mathcal{S}(\mathbb{R}^d \times \mathbb{R}^d)$, we define the partial trace

$$\text{Tr}_\alpha A = \sum_{\mathbf{m} \in \mathbb{Z}^d} \widehat{A}(\mathbf{m} + \boldsymbol{\alpha}, \mathbf{m} + \boldsymbol{\alpha}), \tag{5-6}$$

and note that in view of Lemma 3.2, $\langle \Pi_\alpha A, B \rangle_{\text{HS}} = \text{Tr}_\alpha AB^\dagger$. Let us furthermore define $\mathcal{I}_{\ell,n}$, implicitly dependent on r and h , by

$$\mathcal{I}_{\ell,n}(\mathbf{s}) = \begin{cases} \text{Tr}_\alpha[\text{Op}(D_{r,h}a) \text{Op}(D_{r,h}b)] & (\ell = n = 0), \\ \text{Tr}_\alpha[K_{1,\ell}(\mathbf{s}) \text{Op}(D_{r,h}a) K_{\ell+1,n}(\mathbf{s}) \text{Op}(D_{r,h}b)] & (1 \leq \ell < n), \\ \text{Tr}_\alpha[K_{1,n}(\mathbf{s}) \text{Op}(D_{r,h}a) \text{Op}(D_{r,h}b)] & (0 < \ell = n), \\ \text{Tr}_\alpha[\text{Op}(D_{r,h}a) K_{1,n}(\mathbf{s}) \text{Op}(D_{r,h}b)] & (\ell = 0 < n). \end{cases} \quad (5-7)$$

In view of (4-14), we have for $n = 0, 1, 2$

$$Q_n(t, D_{r,h}a, D_{r,h}\bar{b}) = (2\pi i)^n \sum_{\ell=0}^n (-1)^\ell \int_{\substack{0 < s_1 < \dots < s_\ell < t \\ 0 < s_n < \dots < s_{\ell+1} < t}} \mathcal{I}_{\ell,n}(\mathbf{s}) \, d\mathbf{s}. \quad (5-8)$$

(We work with \bar{b} rather than b to simplify the notation in the calculations that follow.) In other words, the $\mathcal{I}_{\ell,n}$ are precisely the expressions that appear in the expansion of

$$\langle \Pi_\alpha U_\lambda(t) U_0(-t) \text{Op}(D_{r,h}a) U_0(t) U_\lambda(-t), \text{Op}(D_{r,h}\bar{b}) \rangle_{\text{HS}}; \quad (5-9)$$

see (4-13).

Let us write down the $\mathcal{I}_{\ell,n}$ explicitly. For $1 \leq \ell < n$, we show in Section B in the appendix that one has

$$\begin{aligned} \mathcal{I}_{\ell,n}(\mathbf{s}) = r^{nd} h^d \int_{\mathbb{R}^d} \sum_{\mathbf{m}_1, \dots, \mathbf{m}_n} e(-\frac{1}{2} s_1 \|\mathbf{m}_n + \boldsymbol{\alpha}\|^2) \widehat{W}(r(\mathbf{m}_n - \mathbf{m}_1)) \mathcal{T}_{1,\ell-1}^-(\boldsymbol{\alpha}) e(\frac{1}{2} s_\ell \|\mathbf{m}_\ell + \boldsymbol{\alpha}\|^2) \\ \times \tilde{a}(-\boldsymbol{\eta}, h(\mathbf{m}_\ell + \boldsymbol{\alpha} + \frac{1}{2} r^{d-1} \boldsymbol{\eta})) e(-\frac{1}{2} s_{\ell+1} \|\mathbf{m}_\ell + \boldsymbol{\alpha} + r^{d-1} \boldsymbol{\eta}\|^2) \\ \times \widehat{W}(r(\mathbf{m}_\ell - \mathbf{m}_{\ell+1})) \mathcal{T}_{\ell+1,n-1}^-(\boldsymbol{\alpha} + r^{d-1} \boldsymbol{\eta}) e(\frac{1}{2} s_n \|\mathbf{m}_n + \boldsymbol{\alpha} + r^{d-1} \boldsymbol{\eta}\|^2) \\ \times \tilde{b}(\boldsymbol{\eta}, h(\mathbf{m}_n + \boldsymbol{\alpha} + \frac{1}{2} r^{d-1} \boldsymbol{\eta})) \, d\boldsymbol{\eta} + O(r^\infty), \end{aligned} \quad (5-10)$$

with the definition

$$\mathcal{T}_{\ell,n}^-(\mathbf{y}) = \begin{cases} \prod_{j=\ell}^n e(\frac{1}{2} (s_j - s_{j+1}) \|\mathbf{y} + \mathbf{m}_j\|^2) \widehat{W}(r(\mathbf{m}_j - \mathbf{m}_{j+1})) & (\ell \leq n), \\ 1 & (\ell > n). \end{cases} \quad (5-11)$$

The symbol $O(r^\infty)$ is a shorthand for “ $O_\beta(r^\beta)$ for any $\beta \geq 1$.” It follows more immediately from the definition of $K_{\ell,n}$ that for $\ell = n$

$$\begin{aligned} \mathcal{I}_{n,n}(\mathbf{s}) = r^{nd} h^d \int_{\mathbb{R}^d} \sum_{\mathbf{m}_1, \dots, \mathbf{m}_n \in \mathbb{Z}^d} e(-\frac{1}{2} s_1 \|\mathbf{m}_n + \boldsymbol{\alpha}\|^2) \widehat{W}(r(\mathbf{m}_n - \mathbf{m}_1)) \\ \times \mathcal{T}_{1,n-1}^-(\boldsymbol{\alpha}) e(\frac{1}{2} s_n \|\mathbf{m}_n + \boldsymbol{\alpha}\|^2) \tilde{a}(-\boldsymbol{\eta}, h(\mathbf{m}_n + \boldsymbol{\alpha} + \frac{1}{2} r^{d-1} \boldsymbol{\eta})) \\ \times \tilde{b}(\boldsymbol{\eta}, h(\mathbf{m}_n + \boldsymbol{\alpha} + \frac{1}{2} r^{d-1} \boldsymbol{\eta})) \, d\boldsymbol{\eta} + O(r^\infty), \end{aligned} \quad (5-12)$$

and for $\ell = 0$

$$\begin{aligned} \mathcal{I}_{0,n}(\mathbf{s}) \\ = r^{nd} h^d \int_{\mathbb{R}^d} \sum_{\mathbf{m}_1, \dots, \mathbf{m}_n \in \mathbb{Z}^d} \tilde{a}(-\boldsymbol{\eta}, h(\mathbf{m}_n + \boldsymbol{\alpha} + \frac{1}{2} r^{d-1} \boldsymbol{\eta})) \\ \times e(-\frac{1}{2} s_1 \|\mathbf{m}_n + \boldsymbol{\alpha} + r^{d-1} \boldsymbol{\eta}\|^2) \widehat{W}(r(\mathbf{m}_n - \mathbf{m}_1)) \mathcal{T}_{1,n-1}^-(\boldsymbol{\alpha} + r^{d-1} \boldsymbol{\eta}) \\ \times e(\frac{1}{2} s_n \|\mathbf{m}_n + \boldsymbol{\alpha} + r^{d-1} \boldsymbol{\eta}\|^2) \tilde{b}(\boldsymbol{\eta}, h(\mathbf{m}_n + \boldsymbol{\alpha} + \frac{1}{2} r^{d-1} \boldsymbol{\eta})) \, d\boldsymbol{\eta} + O(r^\infty). \end{aligned} \quad (5-13)$$

6. Orders 0 and 1

The asymptotics for zeroth- and first-order terms follows from the Poisson summation formula.

Lemma 6.1.
$$\mathcal{I}_{0,0} = \int_{\mathbb{R}^d \times \mathbb{R}^d} a(x, y)b(x, y) \, dx dy + O(h^\infty). \quad (6-1)$$

Proof. We have (by Lemma 3.2)

$$\mathcal{I}_{0,0} = h^d \sum_{\mathbf{m} \in \mathbb{Z}^d} \int_{\mathbb{R}^d} \tilde{a}(-\boldsymbol{\eta}, h(\mathbf{m} + \boldsymbol{\alpha} + \frac{1}{2}r^{d-1}\boldsymbol{\eta})) \tilde{b}(\boldsymbol{\eta}, h(\mathbf{m} + \boldsymbol{\alpha} + \frac{1}{2}r^{d-1}\boldsymbol{\eta})) \, d\boldsymbol{\eta}. \quad (6-2)$$

Since \tilde{a} and \tilde{b} are in the Schwartz class, applying Poisson summation in \mathbf{m} gives

$$\begin{aligned} \mathcal{I}_{0,0} &= \int_{\mathbb{R}^d \times \mathbb{R}^d} \tilde{a}(-\boldsymbol{\eta}, \mathbf{y}) \tilde{b}(\boldsymbol{\eta}, \mathbf{y}) \, d\boldsymbol{\eta} \, d\mathbf{y} + O(h^\infty) \\ &= \int_{\mathbb{R}^d \times \mathbb{R}^d} a(\mathbf{x}, \mathbf{y})b(\mathbf{x}, \mathbf{y}) \, dx \, dy + O(h^\infty), \end{aligned} \quad (6-3)$$

completing the proof. □

Recall that the mean-free flight time is of the order of r^{1-d} , and that according to (2-9) we should consider time in units of h . This suggests the rescaling $t \rightarrow hr^{1-d}t$, and thus, by the Egorov property (5-5), we obtain for the propagated symbol

$$\begin{aligned} \text{Tr}_\alpha[U_0(thr^{1-d}) \text{Op}(D_{r,h}a)U_0(-thr^{1-d}) \text{Op}(D_{r,h}b)] \\ &= \text{Tr}_\alpha[\text{Op}(D_{r,h}L_0(t)a) \text{Op}(D_{r,h}b)] \\ &= \int_{\mathbb{R}^d \times \mathbb{R}^d} (L_0(t)a)(\mathbf{x}, \mathbf{y})b(\mathbf{x}, \mathbf{y}) \, dx \, dy + O(h^\infty) \\ &= \int_{\mathbb{R}^d \times \mathbb{R}^d} a(\mathbf{x} - t\mathbf{y}, \mathbf{y})b(\mathbf{x}, \mathbf{y}) \, dx \, dy + O(h^\infty) \end{aligned} \quad (6-4)$$

uniformly for all t in a fixed compact interval. It is worth noting that this is precisely the answer one would expect: at order 0 the potential does not appear, which means the solution simply displays free evolution. We see this is true by virtue of the fact that the initial density has simply been translated in position space for time t with momentum \mathbf{y} .

Lemma 6.2.
$$\mathcal{I}_{0,1}(s_1) - \mathcal{I}_{1,1}(s_1) = O(r^d h^\infty + r^\infty). \quad (6-5)$$

Proof. By (5-12),

$$\begin{aligned} \mathcal{I}_{1,1}(s_1) &= r^d h^d \widehat{W}(\mathbf{0}) \sum_{\mathbf{m} \in \mathbb{Z}^d} \int_{\mathbb{R}^d} \tilde{a}(-\boldsymbol{\eta}, h(\mathbf{m} + \boldsymbol{\alpha} + \frac{1}{2}r^{d-1}\boldsymbol{\eta})) \tilde{b}(\boldsymbol{\eta}, h(\mathbf{m} + \boldsymbol{\alpha} + \frac{1}{2}r^{d-1}\boldsymbol{\eta})) \, d\boldsymbol{\eta} + O(r^\infty) \\ &= r^d \widehat{W}(\mathbf{0}) \int_{\mathbb{R}^d \times \mathbb{R}^d} \tilde{a}(-\boldsymbol{\eta}, \mathbf{y}) \tilde{b}(\boldsymbol{\eta}, \mathbf{y}) \, d\boldsymbol{\eta} \, d\mathbf{y} + O(r^d h^\infty + r^\infty), \end{aligned} \quad (6-6)$$

again by Poisson summation. Similarly, using (5-13),

$$\begin{aligned} \mathcal{I}_{0,1}(s_1) &= r^{nd} h^d \widehat{W}(\mathbf{0}) \sum_{\mathbf{m}_1 \in \mathbb{Z}^d} \int_{\mathbb{R}^d} \tilde{a}(-\boldsymbol{\eta}, h(\mathbf{m}_1 + \boldsymbol{\alpha} + \frac{1}{2}r^{d-1}\boldsymbol{\eta})) \tilde{b}(\boldsymbol{\eta}, h(\mathbf{m}_1 + \boldsymbol{\alpha} + \frac{1}{2}r^{d-1}\boldsymbol{\eta})) d\boldsymbol{\eta} + O(r^\infty) \\ &= r^d \widehat{W}(\mathbf{0}) \int_{\mathbb{R}^d \times \mathbb{R}^d} \tilde{a}(-\boldsymbol{\eta}, \mathbf{y}) \tilde{b}(\boldsymbol{\eta}, \mathbf{y}) d\boldsymbol{\eta} d\mathbf{y} + O(r^d h^\infty + r^\infty), \end{aligned} \tag{6-7}$$

completing the proof. □

Indeed, in the expansion (5-8) the terms $\mathcal{I}_{1,1}(s_1)$ and $\mathcal{I}_{0,1}(s_1)$ appear with opposite sign and therefore cancel up to an error $O(r^d h^\infty + r^\infty)$. The total error term after integrating over s_1 is thus obtained by multiplying this by the integration range of size $hr^{1-d}t$.

7. Equidistribution of horocycles

At second order we will use the fact that the $\mathcal{I}_{\ell,n}$ can be written as functions on some noncompact, finite-volume manifold. Specifically, consider the semidirect product group $G = \text{SL}(2, \mathbb{R}) \ltimes \mathbb{R}^{2d}$ with multiplication law

$$(M, \boldsymbol{\xi})(M', \boldsymbol{\xi}') = (MM', \boldsymbol{\xi} + M\boldsymbol{\xi}'), \tag{7-1}$$

where $M, M' \in \text{SL}(2, \mathbb{R})$ and $\boldsymbol{\xi}, \boldsymbol{\xi}' \in \mathbb{R}^d \times \mathbb{R}^d$; the action of $\text{SL}(2, \mathbb{R})$ on $\mathbb{R}^d \times \mathbb{R}^d$ is defined canonically as

$$M\boldsymbol{\xi} = \begin{pmatrix} ax + by \\ cx + dy \end{pmatrix}, \quad M = \begin{pmatrix} a & b \\ c & d \end{pmatrix}, \quad \boldsymbol{\xi} = \begin{pmatrix} x \\ y \end{pmatrix}, \tag{7-2}$$

where $\mathbf{x}, \mathbf{y} \in \mathbb{R}^d$. A convenient parametrisation of $\text{SL}(2, \mathbb{R})$ can be obtained by means of the Iwasawa decomposition

$$M = n_-(u) \Phi^{-\log v} \mathcal{R}(\phi), \tag{7-3}$$

with

$$n_-(u) = \begin{pmatrix} 1 & u \\ 0 & 1 \end{pmatrix}, \quad \Phi^t = \begin{pmatrix} e^{-t/2} & 0 \\ 0 & e^{t/2} \end{pmatrix}, \quad \mathcal{R}(\phi) = \begin{pmatrix} \cos \phi & -\sin \phi \\ \sin \phi & \cos \phi \end{pmatrix}. \tag{7-4}$$

This decomposition is unique for $\tau = u + iv \in \mathfrak{H}$, $\phi \in [0, 2\pi)$, where \mathfrak{H} denotes the upper half-plane $\mathfrak{H} = \{\tau \in \mathbb{C} : \text{Im } \tau > 0\}$. We will use the notation $M = (\tau, \phi)$ and $(M, \boldsymbol{\xi}) = (\tau, \phi, \boldsymbol{\xi})$ interchangeably. With this, we have for instance $n_-(u)\Phi^{-2\log r} = (u + ir^2, 0)$ and

$$\left(1, \begin{pmatrix} \mathbf{0} \\ \mathbf{y} \end{pmatrix}\right) n_-(u)\Phi^{-2\log r} = \left(u + ir^2, 0, \begin{pmatrix} \mathbf{0} \\ \mathbf{y} \end{pmatrix}\right). \tag{7-5}$$

Throughout this section, let Γ be a subgroup of $\text{SL}(2, \mathbb{Z}) \ltimes (\frac{1}{2}\mathbb{Z})^{2d}$ of finite index. The Haar measure on G induces a G -invariant measure on $\Gamma \backslash G$, which will be denoted by μ . Since Γ is a lattice in G , we have (by definition) $0 < \mu(\Gamma \backslash G) < \infty$.

Proposition 7.1. *Fix $\mathbf{y} \in \mathbb{R}^d \setminus \mathbb{Q}^d$ so that the components of $(1, \mathbf{y})$ are linearly independent over \mathbb{Q} . Let $w : \mathbb{R} \rightarrow \mathbb{R}$ be piecewise continuous with compact support. Let $F : \Gamma \backslash G \times \mathbb{R} \rightarrow \mathbb{R}$ be bounded continuous,*

and F_r be a sequence of continuous, uniformly bounded functions $\Gamma \backslash G \times \mathbb{R} \rightarrow \mathbb{R}$ such that $F_r \rightarrow F_0$ uniformly on compacta as $r \rightarrow 0$. Then, for $\sigma \geq 0$, we have

$$\lim_{r \rightarrow 0} r^\sigma \int_{\mathbb{R}} F_r \left(\left(u + ir^2, 0, \begin{pmatrix} \mathbf{0} \\ \mathbf{y} \end{pmatrix} \right), r^\sigma u \right) w(r^\sigma u) du = \frac{1}{\mu(\Gamma \backslash G)} \int_{\Gamma \backslash G} \int_{\mathbb{R}} F_0(g, u) w(u) du d\mu(g). \quad (7-6)$$

Proof. The proof of Theorem 5.1 in [Marklof 2002] tells us that for $F : \Gamma \backslash G \rightarrow \mathbb{R}$ bounded continuous, we have

$$\lim_{r \rightarrow 0} r^\sigma \int_{\mathbb{R}} F \left(\left(u + ir^2, 0, \begin{pmatrix} \mathbf{0} \\ \mathbf{y} \end{pmatrix} \right) \right) w(r^\sigma u) du = \frac{1}{\mu(\Gamma \backslash G)} \int_{\Gamma \backslash G} F d\mu \int_{\mathbb{R}} w(u) du. \quad (7-7)$$

The claim now follows from the same argument as [Marklof and Strömbergsson 2010, Theorem 5.3]. \square

We define the subgroup Γ_∞ by

$$\Gamma_\infty = \left\{ \begin{pmatrix} 1 & m \\ 0 & 1 \end{pmatrix} : m \in \mathbb{Z} \right\} \subset \mathrm{SL}(2, \mathbb{Z}) \quad (7-8)$$

and for $\gamma = \begin{pmatrix} a & b \\ c & d \end{pmatrix}$ use the notation

$$v_\gamma := \mathrm{Im}(\gamma\tau) = \frac{v}{|c\tau + d|^2}, \quad \mathbf{y}_\gamma := c\mathbf{x} + d\mathbf{y}. \quad (7-9)$$

Then, with χ_R the characteristic function of $[R, \infty)$ we define the characteristic function $X_R : \mathfrak{H} \rightarrow \mathbb{R}_{\geq 0}$ by

$$X_R(\tau) = \sum_{\gamma \in (\Gamma_\infty \cup -\Gamma_\infty) \backslash \mathrm{SL}(2, \mathbb{Z})} \chi_R(v_\gamma). \quad (7-10)$$

Note that by construction X_R is $\mathrm{SL}(2, \mathbb{Z})$ -invariant. For $f : \mathbb{R} \rightarrow \mathbb{R}_{\geq 0}$ of rapid decay at $\pm\infty$ and $\beta \in \mathbb{R}$, the function $\Psi_{R,f}^\beta : G \rightarrow \mathbb{R}_{\geq 0}$ is defined by

$$\Psi_{R,f}^\beta(\tau, \xi) = \sum_{\gamma \in \Gamma_\infty \backslash \mathrm{SL}(2, \mathbb{Z})} \sum_{\mathbf{m} \in \mathbb{Z}^d} f((\mathbf{y}_\gamma + \mathbf{m})v_\gamma^{1/2}) v_\gamma^{\beta d/2} \chi_R(v_\gamma), \quad (7-11)$$

and for convenience when $\beta = 1$ we write $\Psi_{R,f} := \Psi_{R,f}^1$. The function $\Psi_{R,f}^\beta$ is left-invariant under $\mathrm{SL}(2, \mathbb{Z}) \times \left(\frac{1}{2}\mathbb{Z}\right)^{2d}$. Both X_R and $\Psi_{R,f}^\beta$ can thus be viewed as functions on G and, since Γ is a finite-index subgroup of $\mathrm{SL}(2, \mathbb{Z}) \times \left(\frac{1}{2}\mathbb{Z}\right)^{2d}$, are also left Γ -invariant.

Proposition 7.2 [Marklof 2002, Proposition 6.4]. *Let \mathbf{y} be Diophantine of type κ , $w : \mathbb{R} \rightarrow \mathbb{R}$ piecewise continuous with compact support, and $0 < \epsilon < 1$ and $0 < \epsilon' < 1/(\kappa - 1)$. Then, for every $R \geq 1$,*

$$\limsup_{r \rightarrow 0} r^{d-2} \int_{|u| > r^{2-\epsilon}} \Psi_{R,f} \left(u + ir^2, \begin{pmatrix} \mathbf{0} \\ \mathbf{y} \end{pmatrix} \right) w(r^{d-2}u) du \ll_{\epsilon, \epsilon'} R^{-(1/(\kappa-1)-d+2)/2} + R^{-\epsilon'/2}. \quad (7-12)$$

Note that the term $R^{-\epsilon'/2}$ is only relevant for $d = 2$. The expression vanishes as $R \rightarrow \infty$ if $\kappa < (d - 1)/(d - 2)$. The following generalisation to $\beta < 1$ holds. Note the range of integration is now over all $u \in \mathbb{R}$.

Proposition 7.3. *Let $0 \leq \beta < 1$, \mathbf{y} be Diophantine of type κ , $w : \mathbb{R} \rightarrow \mathbb{R}$ piecewise continuous with compact support. Then, for every $R \geq 1$,*

$$\limsup_{r \rightarrow 0} r^{d-2} \int_{\mathbb{R}} \Psi_{R,f}^\beta \left(u + ir^2, \begin{pmatrix} \mathbf{0} \\ \mathbf{y} \end{pmatrix} \right) w(r^{d-2}u) \, du \ll R^{-(1/(\kappa-1)-\beta d+2)/2} + R^{(\beta-1)d/2}. \quad (7-13)$$

The right-hand side vanishes as $R \rightarrow \infty$ if and only if

$$\kappa < \begin{cases} \infty & (\beta \leq 2/d), \\ (\beta d - 1)/(\beta d - 2) & (\beta > 2/d). \end{cases} \quad (7-14)$$

In practice, we want both Propositions 7.2 and 7.3 to hold simultaneously. We do this by taking $\kappa < (d - 1)/(d - 2)$ and use the fact that for $2/d \leq \beta < 1$ we have $(\beta d - 1)/(\beta d - 2) > (d - 1)/(d - 2)$.

Proof. Writing $\tau = u + iv$ and $v = r^2$ we have the explicit representation

$$\begin{aligned} \Psi_{R,f}^\beta \left(\tau, \begin{pmatrix} \mathbf{0} \\ \mathbf{y} \end{pmatrix} \right) &= 2 \sum_{\mathbf{m} \in \mathbb{Z}^d} f \left(\mathbf{m} \frac{v^{1/2}}{|\tau|} \right) \frac{v^{\beta d/2}}{|\tau|^{\beta d}} \chi_R \left(\frac{v}{|\tau|^2} \right) \\ &\quad + 2 \sum_{\substack{(c,d) \in \mathbb{Z}^2 \\ \gcd(c,d)=1 \\ c>0,d \neq 0}} \sum_{\mathbf{m} \in \mathbb{Z}^d} f \left((d\mathbf{y} + \mathbf{m}) \frac{v^{1/2}}{|c\tau + d|} \right) \frac{v^{\beta d/2}}{|c\tau + d|^{\beta d}} \chi_R \left(\frac{v}{|c\tau + d|^2} \right). \end{aligned} \quad (7-15)$$

For the first term we make the substitution $u = vt$ in the integral, which yields

$$\begin{aligned} 2v^{d/2-1} \int_{\mathbb{R}} w(v^{d/2-1}u) \sum_{\mathbf{m} \in \mathbb{Z}^d} f \left(\mathbf{m} \frac{v^{1/2}}{|\tau|} \right) \frac{v^{\beta d/2}}{|\tau|^{\beta d}} \chi_R \left(\frac{v}{|\tau|^2} \right) \, du \\ = 2v^{(1-\beta)d/2} \int_{\mathbb{R}} \frac{w(v^{d/2}t)}{(1+t^2)^{\beta d/2}} \sum_{\mathbf{m} \in \mathbb{Z}^d} f \left(\frac{\mathbf{m}}{v^{1/2}(1+t^2)^{1/2}} \right) \chi_R \left(\frac{1}{v(1+t^2)} \right) \, dt. \end{aligned} \quad (7-16)$$

Under the assumption that $0 < \beta < 1$ we have

$$\frac{v^{(1-\beta)d/2}}{(1+t^2)^{\beta d/2}} \chi_R \left(\frac{1}{v(1+t^2)} \right) \leq \frac{R^{(\beta-1)d/2}}{(1+t^2)^{d/2}} \chi_R \left(\frac{1}{v(1+t^2)} \right) \quad (7-17)$$

and thus obtain the bound

$$\begin{aligned} \limsup_{v \rightarrow 0} 2v^{d/2-1} \int_{\mathbb{R}} w(v^{d/2-1}u) \sum_{\mathbf{m} \in \mathbb{Z}^d} f \left(\mathbf{m} \frac{v^{1/2}}{|\tau|} \right) \frac{v^{\beta d/2}}{|\tau|^{\beta d}} \chi_R \left(\frac{v}{|\tau|^2} \right) \, du \\ \leq 2R^{(\beta-1)d/2} w(0) f(\mathbf{0}) \int_{\mathbb{R}} \frac{dt}{(1+t^2)^{d/2}} + O(R^{-\infty}). \end{aligned} \quad (7-18)$$

For the second term, using

$$\frac{v^{\beta d/2}}{|c\tau + d|^{\beta d}} \chi_R \left(\frac{v}{|c\tau + d|^2} \right) \leq \frac{v^{d/2}}{|c\tau + d|^d} R^{(\beta-1)d/2} \chi_R \left(\frac{v}{|c\tau + d|^2} \right), \quad (7-19)$$

we see that

$$\sum_{\substack{(c,d) \in \mathbb{Z}^2 \\ \gcd(c,d)=1 \\ c>0, d \neq 0}} \sum_{\mathbf{m} \in \mathbb{Z}^d} v^{d/2-1} \int_{\mathbb{R}} f\left((d\mathbf{y} + \mathbf{m}) \frac{v^{1/2}}{|c\tau + d|}\right) \frac{v^{\beta d/2}}{|c\tau + d|^{\beta d}} \chi_R\left(\frac{v}{|c\tau + d|^2}\right) w(v^{d/2-1}u) \, du \quad (7-20)$$

is bounded above by

$$R^{(\beta-1)d/2} \sum_{\substack{(c,d) \in \mathbb{Z}^2 \\ \gcd(c,d)=1 \\ c>0, d \neq 0}} \sum_{\mathbf{m} \in \mathbb{Z}^d} v^{d/2-1} \int_{\mathbb{R}} f\left((d\mathbf{y} + \mathbf{m}) \frac{v^{1/2}}{|c\tau + d|}\right) \frac{v^{d/2}}{|c\tau + d|^d} \chi_R\left(\frac{v}{|c\tau + d|^2}\right) w(v^{d/2-1}u) \, du. \quad (7-21)$$

This reduces the problem to the same calculation as in the proof of Proposition 7.2, which yields that (7-21) is bounded above by

$$R^{(\beta-1)d/2} (R^{-(1/(\kappa-1)-d+2)/2} + 1) = R^{-(1/(\kappa-1)-\beta d+2)/2} + R^{((\beta-1)d)/2}, \quad (7-22)$$

completing the proof. □

Fix a compact interval $A \subset \mathbb{R}$. We say $F : \Gamma \backslash G \times \mathbb{R} \rightarrow \mathbb{C}$ is dominated by $\Psi_{R,f}$ on $\Gamma \backslash G \times A$ if there are positive constants L, R_0 such that $|F((\tau, \phi, \xi), u')| X_R(\tau) \leq L(1 + \Psi_{R,f}(\tau, \phi, \xi))$ for all $(\tau, \phi, \xi) \in G, u' \in A$ and $R \geq R_0$. A sequence of functions $F_r : \Gamma \backslash G \times \mathbb{R} \rightarrow \mathbb{C}$ is uniformly dominated if L, R_0 are independent of r .

Proposition 7.4. *Assume \mathbf{y} is Diophantine of type $\kappa < (d - 1)/(d - 2)$ with the components of $(1, {}^t\mathbf{y})$ linearly independent over \mathbb{Q} . Let $w : \mathbb{R} \rightarrow \mathbb{R}$ be piecewise continuous with compact support. Let $F_0 : \Gamma \backslash G \times \mathbb{R} \rightarrow \mathbb{R}$ be continuous and dominated by $\Psi_{R,f}$ on $\Gamma \backslash G \times \text{supp } w$. Let F_r be a sequence of continuous functions $\Gamma \backslash G \times \mathbb{R} \rightarrow \mathbb{R}$ uniformly dominated by $\Psi_{R,f}$ on $\Gamma \backslash G \times \text{supp } w$ such that $F_r \rightarrow F_0$ uniformly on compacta as $r \rightarrow 0$. Then for any $0 < \epsilon < 2$ we have*

$$\lim_{r \rightarrow 0} r^{d-2} \int_{|u| > r^{2-\epsilon}} F_r\left(\left(u + ir^2, 0, \begin{pmatrix} \mathbf{0} \\ \mathbf{y} \end{pmatrix}, r^{d-2}u\right), w(r^{d-2}u) \, du = \frac{1}{\mu(\Gamma \backslash G)} \int_{\mathbb{R}} \int_{\Gamma \backslash G} F_0(g, u) w(u) \, d\mu(g) \, du. \quad (7-23)$$

Proof. (This follows the proof of [Marklof 2002, Theorem 6.8/Corollary 6.10].) We may assume without loss of generality that F_r and w are real-valued and nonnegative. Set

$$J_{r,R}((\tau, \phi, \xi), u') = F_r((\tau, \phi, \xi), u')(1 - X_R(\tau)). \quad (7-24)$$

Then $J_{r,R}$ is bounded and thus

$$\int_{|u| > r^{2-\epsilon}} J_{r,R}\left(\left(u + ir^2, 0, \begin{pmatrix} \mathbf{0} \\ \mathbf{y} \end{pmatrix}, r^{d-2}u\right), w(r^{d-2}u) \, du = \int_{\mathbb{R}} J_{r,R}\left(\left(u + ir^2, 0, \begin{pmatrix} \mathbf{0} \\ \mathbf{y} \end{pmatrix}, r^{d-2}u\right), w(r^{d-2}u) \, du + O(r^{2-\epsilon}). \quad (7-25)$$

By Proposition 7.1, which (by a standard probabilistic argument) extends to functions such as $J_{r,R}$ whose points of discontinuity are contained in a set of μ -measure zero (alternatively simply smooth the characteristic function χ_R to make $J_{r,R}$ continuous),

$$\begin{aligned} \lim_{r \rightarrow 0} r^{d-2} \int_{\mathbb{R}} J_{r,R} \left(\left(u + ir^2, 0, \begin{pmatrix} \mathbf{0} \\ \mathbf{y} \end{pmatrix} \right), r^{d-2}u \right) w(r^{d-2}u) du \\ = \frac{1}{\mu(\Gamma \setminus G)} \int_{\mathbb{R}} \int_{\Gamma \setminus G} J_{0,R}(g, u') w(u') d\mu(g) du'. \end{aligned} \quad (7-26)$$

Furthermore, $F_0((\tau, \phi, \xi), u') X_R(\tau) \leq L X_R(\tau) + L \Psi_{R,f}(\tau, \xi)$ for large R , and hence

$$\int_{\mathbb{R}} \int_{\Gamma \setminus G} F_0((\tau, \phi, \xi), u') X_R(\tau) w(u') d\mu du' \leq \int_{\mathbb{R}} w(u') du' \int_{\Gamma \setminus G} (L X_R + L \Psi_{R,f}) d\mu \ll R^{-1}; \quad (7-27)$$

see [Marklof 2002, §6.2]. Combining this with the result for $J_{0,R}$ yields

$$\int_{\mathbb{R}} \int_{\Gamma \setminus G} J_{0,R}(g, u') w(u') d\mu(g) du' = \int_{\mathbb{R}} \int_{\Gamma \setminus G} F_0(g, u') w(u') d\mu(g) du' + O(R^{-1}). \quad (7-28)$$

In summary, we have shown thus far that

$$\begin{aligned} \liminf_{r \rightarrow 0} r^{d-2} \int_{|u| > r^{2-\epsilon}} F_r \left(\left(u + ir^2, 0, \begin{pmatrix} \mathbf{0} \\ \mathbf{y} \end{pmatrix} \right), r^{d-2}u \right) w(r^{d-2}u) du \\ \geq \lim_{r \rightarrow 0} r^{d-2} \int_{|u| > r^{2-\epsilon}} J_{r,R} \left(\left(u + ir^2, 0, \begin{pmatrix} \mathbf{0} \\ \mathbf{y} \end{pmatrix} \right), r^{d-2}u \right) w(r^{d-2}u) du \\ = \frac{1}{\mu(\Gamma \setminus G)} \int_{\mathbb{R}} \int_{\Gamma \setminus G} F_0(g, u') w(u') d\mu(g) du' + O(R^{-1}) \end{aligned} \quad (7-29)$$

for every $R \geq R_0$. For the upper bound we use that

$$F_r((\tau, \phi, \xi), u') \leq F_r((\tau, \phi, \xi), u')(1 - X_R(\tau)) + L X_R(\tau) + L \Psi_{R,f}(\tau, \xi). \quad (7-30)$$

We proceed as above for the first two terms, and apply Proposition 7.2 to the third to obtain

$$\begin{aligned} \limsup_{r \rightarrow 0} r^{d-2} \int_{|u| > r^{2-\epsilon}} F_r \left(\left(u + ir^2, 0, \begin{pmatrix} \mathbf{0} \\ \mathbf{y} \end{pmatrix} \right), r^{d-2}u \right) w(r^{d-2}u) du \\ \leq \frac{1}{\mu(\Gamma \setminus G)} \int_{\mathbb{R}} \int_{\Gamma \setminus G} F_0(g, u') d\mu(g) du' + O(R^{-(1/(\kappa-1)-d+2)/2} + R^{-\epsilon'/2}) \end{aligned} \quad (7-31)$$

for every $R \geq R_0$. □

8. Mean value theorems for theta functions

For $f \in \mathcal{S}(\mathbb{R}^d \times \mathbb{R}^d)$ and $\phi \in \mathbb{R}$, define f_ϕ by

$$f_\phi(\mathbf{y}_1, \mathbf{y}_2) = \begin{cases} f(\mathbf{y}_1, \mathbf{y}_2) & (\phi = 0 \pmod{2\pi}), \\ f(-\mathbf{y}_1, -\mathbf{y}_2) & (\phi = \pi \pmod{2\pi}), \\ \int_{\mathbb{R}^{2d}} G_\phi(\mathbf{y}_1, \mathbf{y}_2, \mathbf{x}_1, \mathbf{x}_2) f(\mathbf{x}_1, \mathbf{x}_2) d\mathbf{x}_1 d\mathbf{x}_2 & (\phi \neq 0 \pmod{\pi}), \end{cases} \quad (8-1)$$

where

$$G_\phi(\mathbf{y}_1, \mathbf{y}_2, \mathbf{x}_1, \mathbf{x}_2) = |\sin \phi|^{-d} e^{\left(\frac{\frac{1}{2}(\|\mathbf{y}_1\|^2 + \|\mathbf{x}_1\|^2 - \|\mathbf{y}_2\|^2 - \|\mathbf{x}_2\|^2) \cos \phi - \mathbf{y}_1 \cdot \mathbf{x}_1 + \mathbf{y}_2 \cdot \mathbf{x}_2}{\sin \phi} \right)}. \quad (8-2)$$

Lemma 8.1. *If $f \in \mathcal{S}(\mathbb{R}^d \times \mathbb{R}^d)$ then $f_\phi \in \mathcal{S}(\mathbb{R}^d \times \mathbb{R}^d)$.*

Proof. If $\phi = 0 \pmod{\pi}$ then the result is immediate. For fixed $\phi \neq 0 \pmod{\pi}$, define

$$g(\mathbf{x}_1, \mathbf{x}_2) = e^{\left(\frac{\frac{1}{2}(\|\mathbf{x}_1\|^2 - \|\mathbf{x}_2\|^2) \cos \phi}{\sin \phi} \right)} f(\mathbf{x}_1, \mathbf{x}_2) \quad (8-3)$$

and its Fourier transform

$$I(\mathbf{y}_1, \mathbf{y}_2) = |\sin \phi|^{-d} \int_{\mathbb{R}^{2d}} g(\mathbf{x}_1, \mathbf{x}_2) e^{\left(\frac{-\mathbf{y}_1 \cdot \mathbf{x}_1 + \mathbf{y}_2 \cdot \mathbf{x}_2}{\sin \phi} \right)} d\mathbf{x}_1 d\mathbf{x}_2. \quad (8-4)$$

Note that

$$f_\phi(\mathbf{y}_1, \mathbf{y}_2) = e^{\left(\frac{\frac{1}{2}(\|\mathbf{y}_1\|^2 - \|\mathbf{y}_2\|^2) \cos \phi}{\sin \phi} \right)} I(\mathbf{y}_1, \mathbf{y}_2). \quad (8-5)$$

Now $f \in \mathcal{S}(\mathbb{R}^d \times \mathbb{R}^d)$ implies $g \in \mathcal{S}(\mathbb{R}^d \times \mathbb{R}^d)$ (since all derivatives of the exponential factor in (8-3) grow at most polynomially), which implies $I \in \mathcal{S}(\mathbb{R}^d \times \mathbb{R}^d)$ (since the Fourier transform preserves Schwartz class; use integration by parts), which in turn implies $f_\phi \in \mathcal{S}(\mathbb{R}^d \times \mathbb{R}^d)$ (by the first argument). \square

The following lemma provides rapid decay that is uniform in ϕ .

Lemma 8.2. *If $f \in \mathcal{S}(\mathbb{R}^d \times \mathbb{R}^d)$, then for all multi-indices $\beta_1, \beta_2 \in \mathbb{Z}_{\geq 0}^d$ and for every $T > 1$*

$$\sup_{\mathbf{y}_1, \mathbf{y}_2, \phi} (1 + \|\mathbf{y}_1\|)^T (1 + \|\mathbf{y}_2\|)^T |\partial_{\mathbf{y}_1}^{\beta_1} \partial_{\mathbf{y}_2}^{\beta_2} f_\phi(\mathbf{y}_1, \mathbf{y}_2)| < \infty. \quad (8-6)$$

Proof. The proof of Lemma 8.1 shows that

$$\sup_{\mathbf{y}_1, \mathbf{y}_2, \phi \in I} (1 + \|\mathbf{y}_1\|)^T (1 + \|\mathbf{y}_2\|)^T |\partial_{\mathbf{y}_1}^{\beta_1} \partial_{\mathbf{y}_2}^{\beta_2} f_\phi(\mathbf{y}_1, \mathbf{y}_2)| < \infty \quad (8-7)$$

for any closed interval I not containing $\phi = 0 \pmod{\pi}$. As in the proof of [Marklof 2003, Lemma 4.3], we represent $f_{\phi+\pi/2} = \int_{\mathbb{R}^{2d}} G_\phi(\mathbf{y}_1, \mathbf{y}_2, \mathbf{x}_1, \mathbf{x}_2) f_{\pi/2}(\mathbf{x}_1, \mathbf{x}_2) d\mathbf{x}_1 d\mathbf{x}_2$ using the Fourier transform $f_{\pi/2}$ of f . Since $f_{\pi/2} \in \mathcal{S}(\mathbb{R}^d \times \mathbb{R}^d)$, we see that (8-7) holds for any closed interval not containing $\phi = \frac{\pi}{2} \pmod{\pi}$. Both cases taken together, this shows that (8-7) holds in fact for every closed interval I , and so in particular for $I = [0, 2\pi]$. This proves the claim in view of the 2π -periodicity of f_ϕ . \square

We define the theta function $\Theta_f : G \mapsto \mathbb{C}$ by

$$\Theta_f\left(u + iv, \phi, \begin{pmatrix} \mathbf{x} \\ \mathbf{y} \end{pmatrix}\right) = v^{d/2} \sum_{\mathbf{m}_1, \mathbf{m}_2 \in \mathbb{Z}^d} f_\phi(v^{1/2}(\mathbf{m}_1 - \mathbf{y}), v^{1/2}(\mathbf{m}_2 - \mathbf{y})) \times e^{\left(\frac{1}{2} u (\|\mathbf{m}_1 - \mathbf{y}\|^2 - \|\mathbf{m}_2 - \mathbf{y}\|^2) + \mathbf{x} \cdot (\mathbf{m}_1 - \mathbf{m}_2) \right)}. \quad (8-8)$$

Since $f_\phi \in \mathcal{S}(\mathbb{R}^d \times \mathbb{R}^d)$ we have that $\Theta_f \in C^\infty(G)$. Let

$$\Gamma = \left\{ \left(\begin{pmatrix} a & b \\ c & d \end{pmatrix}, \begin{pmatrix} abs \\ cds \end{pmatrix} + \mathbf{m} \right) : \begin{pmatrix} a & b \\ c & d \end{pmatrix} \in \mathrm{SL}(2, \mathbb{Z}), \mathbf{m} \in \mathbb{Z}^{2d} \right\} \subset G, \quad (8-9)$$

with $s = (\frac{1}{2}, \frac{1}{2}, \dots, \frac{1}{2}) \in \mathbb{R}^d$. Then Γ is of finite index in $SL(2, \mathbb{Z}) \times (\frac{1}{2}\mathbb{Z})^{2d}$, and Θ_f is left Γ invariant; see [Marklof 2003, Proposition 4.9]. That is, $\Theta_f \in C^\infty(\Gamma \backslash G)$.

Proposition 8.3. *Let $f \in \mathcal{S}(\mathbb{R}^d \times \mathbb{R}^d)$. Then*

$$\Theta_f(u + iv, \phi, \xi) = v^{d/2} \sum_{\mathbf{m} \in \mathbb{Z}^d} f_\phi((\mathbf{m} - \mathbf{y}) v^{1/2}, (\mathbf{m} - \mathbf{y}) v^{1/2}) + O(v^{-\infty}) \tag{8-10}$$

uniformly for all $(u + iv, \phi, \xi) \in G$ with $v > \frac{1}{2}$.

Proof. See [Marklof 2003, Proposition 4.10]. □

Corollary 8.4. *Let $f \in \mathcal{S}(\mathbb{R}^d \times \mathbb{R}^d)$; then for all $T > 1$ we have that Θ_f is dominated by $\Psi_{R, \bar{f}}$ for*

$$\bar{f}(\mathbf{x}) = (1 + \|\mathbf{x}\|)^{-2T}. \tag{8-11}$$

Proof. This follows from Proposition 8.3 and Lemma 8.2 (with $\beta_1 = \beta_2 = \mathbf{0}$). □

Proposition 8.5. *Assume \mathbf{y} is Diophantine of type $\kappa < (d - 1)/(d - 2)$ with the components of $(1, {}^t\mathbf{y})$ linearly independent over \mathbb{Q} . Let $w : \mathbb{R} \rightarrow \mathbb{R}$ be piecewise continuous, continuous at 0, with compact support. Then*

$$\begin{aligned} \lim_{r \rightarrow 0} r^{d-2} \int_{\mathbb{R}} \Theta_f\left(u + ir^2, 0, \begin{pmatrix} \mathbf{0} \\ \mathbf{y} \end{pmatrix}\right) w(r^{d-2}u) du \\ = 2 w(0) \int_{\mathbb{R}^d \times \mathbb{R}^d} f(\mathbf{y}_1, \mathbf{y}_2) \delta(\|\mathbf{y}_1\|^2 - \|\mathbf{y}_2\|^2) d\mathbf{y}_1 d\mathbf{y}_2 + \int_{\mathbb{R}^d} f(\mathbf{y}_1, \mathbf{y}_1) d\mathbf{y}_1 \int_{\mathbb{R}} w(u) du. \end{aligned} \tag{8-12}$$

Proof. Fix $0 < \epsilon < 1$, and split the integration over u into the regions $|u| < r^{2-\epsilon}$ and $|u| > r^{2-\epsilon}$. In the first region, the proof of [Marklof 2002, Lemma 7.3] shows that

$$\begin{aligned} r^{d-2} \int_{|u| < r^{2-\epsilon}} \Theta_f\left(u + ir^2, 0, \begin{pmatrix} \mathbf{0} \\ \mathbf{y} \end{pmatrix}\right) w(r^{d-2}u) du \\ = r^{-2} \int_{|u| < r^{2-\epsilon}} \left(\int_{\mathbb{R}^d \times \mathbb{R}^d} f(\mathbf{y}_1, \mathbf{y}_2) e^{i\frac{1}{2}(\|\mathbf{y}_1\|^2 - \|\mathbf{y}_2\|^2)r^{-2}u} d\mathbf{y}_1 d\mathbf{y}_2 \right) w(r^{d-2}u) du + o(1) \\ = 2 w(0) \int_{\mathbb{R}^d \times \mathbb{R}^d} f(\mathbf{y}_1, \mathbf{y}_2) \delta(\|\mathbf{y}_1\|^2 - \|\mathbf{y}_2\|^2) d\mathbf{y}_1 d\mathbf{y}_2 + o(1). \end{aligned} \tag{8-13}$$

Since Θ_f is dominated by $\Psi_{R, f}$, for the region $|u| > r^{2-\epsilon}$ we can apply Proposition 7.4 and note that the limit can be written as

$$\frac{1}{\mu(\Gamma \backslash G)} \int_{\Gamma \backslash G} \Theta_f d\mu \int_{\mathbb{R}} w(u) du = \int_{\mathbb{R}^d} f(\mathbf{y}_1, \mathbf{y}_1) d\mathbf{y}_1 \int_{\mathbb{R}} w(u) du; \tag{8-14}$$

see [Marklof 2002, Lemma 7.2]. □

We will now deal with f that depend continuously on additional parameters $u \in \mathbb{R}, \boldsymbol{\eta} \in \mathbb{R}^d$. We denote by $\tilde{\mathcal{S}}$ the class of functions $f \in C(\mathbb{R}^d \times \mathbb{R}^d \times \mathbb{R} \times \mathbb{R}^d)$ with the property that for every multi-index

$\beta_1, \beta_2 \in \mathbb{Z}_{\geq 0}^d$ the derivative $\partial_{y_1}^{\beta_1} \partial_{y_2}^{\beta_2} f(\mathbf{y}_1, \mathbf{y}_2, u, \boldsymbol{\eta})$ (a) exists, (b) is continuous (in all variables), and (c) is rapidly decaying, i.e.,

$$\sup_{y_1, y_2, u, \boldsymbol{\eta}} (1 + \|\mathbf{y}_1\|)^T (1 + \|\mathbf{y}_2\|)^T (1 + |u|)^T (1 + \|\boldsymbol{\eta}\|)^T |\partial_{y_1}^{\beta_1} \partial_{y_2}^{\beta_2} f(\mathbf{y}_1, \mathbf{y}_2, u, \boldsymbol{\eta})| < \infty \tag{8-15}$$

for every $T > 1$. For $f \in \tilde{\mathcal{S}}$ we define $f_\phi \in \tilde{\mathcal{S}}$ in analogy with (8-1) by

$$f_\phi(\mathbf{y}_1, \mathbf{y}_2, u, \boldsymbol{\eta}) = \begin{cases} f(\mathbf{y}_1, \mathbf{y}_2, u, \boldsymbol{\eta}) & (\phi = 0 \pmod{2\pi}), \\ f(-\mathbf{y}_1, -\mathbf{y}_2, u, \boldsymbol{\eta}) & (\phi = \pi \pmod{2\pi}), \\ \int_{\mathbb{R}^{2d}} G_\phi(\mathbf{y}_1, \mathbf{y}_2, \mathbf{x}_1, \mathbf{x}_2) f(\mathbf{x}_1, \mathbf{x}_2, u, \boldsymbol{\eta}) \, d\mathbf{x}_1 \, d\mathbf{x}_2 & (\phi \neq 0 \pmod{\pi}). \end{cases} \tag{8-16}$$

The fact that $f_\phi \in \tilde{\mathcal{S}}$ follows from the same argument as in Lemma 8.1. We also have the following.

Lemma 8.6. *If $f \in \tilde{\mathcal{S}}$, then for all multi-indices $\beta_1, \beta_2 \in \mathbb{Z}_{\geq 0}^d$ and every $T > 1$*

$$\sup_{y_1, y_2, u, \boldsymbol{\eta}, \phi} (1 + \|\mathbf{y}_1\|)^T (1 + \|\mathbf{y}_2\|)^T (1 + |u|)^T (1 + \|\boldsymbol{\eta}\|)^T |\partial_{y_1}^{\beta_1} \partial_{y_2}^{\beta_2} f_\phi(\mathbf{y}_1, \mathbf{y}_2, u, \boldsymbol{\eta})| < \infty. \tag{8-17}$$

Proof. This is analogous to the proof of Lemma 8.2. □

Given $f \in \tilde{\mathcal{S}}$, we define the theta function

$$\Theta_f(g, u, \boldsymbol{\eta}) = \Theta_{f(\cdot, u, \boldsymbol{\eta})}(g), \tag{8-18}$$

with $\Theta_{f(\cdot, u, \boldsymbol{\eta})}$ as defined in (8-8) (with $u, \boldsymbol{\eta}$ fixed). In view of Lemma 8.6, we have $\Theta_f \in C(\Gamma \backslash G \times \mathbb{R} \times \mathbb{R}^d)$. We further define

$$F_r(g, u) = \int_{\mathbb{R}^d} \Theta_f\left(g\left(1, \begin{pmatrix} \mathbf{0} \\ \frac{1}{2}r^d \boldsymbol{\eta} \end{pmatrix}\right), u, \boldsymbol{\eta}\right) \, d\boldsymbol{\eta}. \tag{8-19}$$

Proposition 8.7. *Let $f \in \tilde{\mathcal{S}}$. Then*

$$F_r(u + iv, \phi, \boldsymbol{\xi}, u') = v^{d/2} \sum_{\mathbf{m} \in \mathbb{Z}^d} \int_{\mathbb{R}^d} f_\phi(v^{1/2}(\mathbf{m} - \mathbf{y}), v^{1/2}(\mathbf{m} - \mathbf{y}), u', \boldsymbol{\eta}) \, d\boldsymbol{\eta} + O(r^d) + O(v^{-\infty}) \tag{8-20}$$

uniformly for all $(u + iv, \phi, \boldsymbol{\xi}) \in G, u' \in \mathbb{R}$, with $v > \frac{1}{2}$ and $r < 1$.

Proof. Note that

$$(u + iv, \phi, \boldsymbol{\xi})\left(1, \begin{pmatrix} \mathbf{0} \\ \frac{1}{2}r^d \boldsymbol{\eta} \end{pmatrix}\right) = \left(u + iv, \phi, \begin{pmatrix} \mathbf{x} + \mathbf{x}_{\tau, \phi, \boldsymbol{\eta}} \\ \mathbf{y} + \mathbf{y}_{\tau, \phi, \boldsymbol{\eta}} \end{pmatrix}\right), \tag{8-21}$$

where

$$\begin{aligned} \mathbf{x}_{\tau, \phi, \boldsymbol{\eta}} &= -\frac{1}{2}v^{1/2}r^d \boldsymbol{\eta} \sin \phi + \frac{1}{2}uv^{-1/2}r^d \boldsymbol{\eta} \cos \phi, \\ \mathbf{y}_{\tau, \phi, \boldsymbol{\eta}} &= \frac{1}{2}v^{-1/2}r^d \boldsymbol{\eta} \cos \phi. \end{aligned} \tag{8-22}$$

We thus have

$$\begin{aligned} F_r(u + iv, \phi, \boldsymbol{\xi}, u') &= \int_{\mathbb{R}^d} v^{d/2} \sum_{\mathbf{m}_1, \mathbf{m}_2 \in \mathbb{Z}^d} f_\phi(v^{1/2}(\mathbf{m}_1 - \mathbf{y} - \mathbf{y}_{\tau, \phi, \boldsymbol{\eta}}), v^{1/2}(\mathbf{m}_2 - \mathbf{y} - \mathbf{y}_{\tau, \phi, \boldsymbol{\eta}}), u', \boldsymbol{\eta}) \\ &\quad \times e\left(\frac{1}{2}u(\|\mathbf{m}_1 - \mathbf{y} - \mathbf{y}_{\tau, \phi, \boldsymbol{\eta}}\|^2 - \|\mathbf{m}_2 - \mathbf{y} - \mathbf{y}_{\tau, \phi, \boldsymbol{\eta}}\|^2)\right) \\ &\quad \times e((\mathbf{x} + \mathbf{x}_{\tau, \phi, \boldsymbol{\eta}}) \cdot (\mathbf{m}_1 - \mathbf{m}_2)) \, d\boldsymbol{\eta}. \end{aligned} \tag{8-23}$$

Choose $\mathbf{m} \in \mathbb{Z}^d$ such that $\mathbf{m} \in [-\frac{1}{2}, \frac{1}{2}]^d + \mathbf{y} + \mathbf{y}_{\tau, \phi, \eta}$. Then, for any $T \geq 1$ and for all $\mathbf{m}_1 \neq \mathbf{m}$,

$$\begin{aligned} f_\phi(v^{1/2}(\mathbf{m}_1 - \mathbf{y} - \mathbf{y}_{\tau, \phi, \eta}), v^{1/2}(\mathbf{m}_2 - \mathbf{y} - \mathbf{y}_{\tau, \phi, \eta}), u', \eta) \\ = O_T(v^{-T}(1 + \|\mathbf{m}_1\|^{-2T})(1 + \|\mathbf{m}_2\|^{-2T})(1 + \|\eta\|^{-2T})). \end{aligned} \quad (8-24)$$

The same is true for $\mathbf{m}_2 \neq \mathbf{m}$. Therefore

$$\begin{aligned} F_r(u + iv, \phi, \xi, u') \\ = v^{d/2} \sum_{\mathbf{m} \in \mathbb{Z}^d} \int_{\mathbb{R}^d} f_\phi(v^{1/2}(\mathbf{m} - \mathbf{y} - \mathbf{y}_{\tau, \phi, \eta}), v^{1/2}(\mathbf{m} - \mathbf{y} - \mathbf{y}_{\tau, \phi, \eta}), u', \eta) d\eta + O(v^{-\infty}). \end{aligned} \quad (8-25)$$

The result follows from applying Taylor’s theorem and using Lemma 8.6 to conclude that the error term is small uniformly in u' and ϕ . □

Lemma 8.8. *Fix $T > d$. Then:*

- (1) *The sequence $(F_r)_r$ of continuous functions $\Gamma \backslash G \times \mathbb{R} \rightarrow \mathbb{C}$ is uniformly dominated by $\Psi_{R, \bar{f}}$, where $\bar{f}(\mathbf{y}) = (1 + \|\mathbf{y}\|)^{-2T}$.*
- (2) *$F_r \rightarrow F_0$ uniformly on compacta.*

Proof. The set of $(u + iv, \phi, \xi) \in G$ with $v > \frac{1}{2}$ contains a fundamental domain of Γ in G . Therefore, by Proposition 8.7 we have for $r < 1$ that

$$\begin{aligned} F_r(u + iv, \phi, \xi, u') &\ll 1 + v^{d/2} \sum_{\mathbf{m} \in \mathbb{Z}^d} \int_{\mathbb{R}^d} f_\phi((\mathbf{m} - \mathbf{y})v^{1/2}, (\mathbf{m} - \mathbf{y})v^{1/2}, u', \eta) d\eta \\ &\ll 1 + v^{d/2} \sum_{\mathbf{m} \in \mathbb{Z}^d} \bar{f}((\mathbf{m} - \mathbf{y})v^{1/2}) \int_{\mathbb{R}^d} (1 + \|\eta\|)^{-T} d\eta \\ &\ll 1 + \Psi_{R, \bar{f}}(\tau, \xi). \end{aligned} \quad (8-26)$$

The first result is thus proved. The second result follows from the continuity of Θ_f and Lemma 8.6. □

Proposition 8.9. *Let $f \in \tilde{\mathcal{S}}$, and assume \mathbf{y} is Diophantine of type $\kappa < (d - 1)/(d - 2)$ with the components of $(1, {}^t\mathbf{y})$ linearly independent over \mathbb{Q} . Let $w : \mathbb{R} \rightarrow \mathbb{R}$ be piecewise continuous, continuous at 0, with compact support. Then*

$$\begin{aligned} \lim_{r \rightarrow 0} r^{d-2} \int_{\mathbb{R}} F_r\left(\left(u + ir^2, 0, \begin{pmatrix} \mathbf{0} \\ \mathbf{y} \end{pmatrix}\right), r^{d-2}u\right) w(r^{d-2}u) du \\ = 2 w(0) \int_{(\mathbb{R}^d)^3} f(\mathbf{y}_1, \mathbf{y}_2, 0, \eta) \delta(\|\mathbf{y}_1\|^2 - \|\mathbf{y}_2\|^2) d\mathbf{y}_1 d\mathbf{y}_2 d\eta \\ + \int_{\mathbb{R}^d \times \mathbb{R} \times \mathbb{R}^d} f(\mathbf{y}_1, \mathbf{y}_1, u, \eta) w(u) d\mathbf{y}_1 du d\eta. \end{aligned} \quad (8-27)$$

Proof. This is analogous to the proof of Proposition 8.5. □

9. Order 2

In this section we show how the terms at order λ^2 can be written as averages over theta functions of the form (8-19). We assume throughout this section that α is Diophantine of type $\kappa < (d-1)/(d-2)$ with the components of $(1, \alpha)$ linearly independent over \mathbb{Q} .

The cases $\ell = 2$ and $\ell = 0$. The cases $\ell = 0$ and 2 are similar and we treat them together. First, from (5-12) we have that $\mathcal{I}_{2,2}$ can be written

$$\mathcal{I}_{2,2}(s_1, s_2) = r^{2d} h^d \sum_{\mathbf{m}_1, \mathbf{m}_2 \in \mathbb{Z}^d} \int_{\mathbb{R}^d} |\widehat{W}(r(\mathbf{m}_2 - \mathbf{m}_1))|^2 e^{\frac{1}{2}(s_2 - s_1)(\|\mathbf{m}_2 + \alpha\|^2 - \|\mathbf{m}_1 + \alpha\|^2)} \times \tilde{a}(-\eta, h(\mathbf{m}_2 + \alpha + \frac{1}{2}r^{d-1}\eta)) \tilde{b}(\eta, h(\mathbf{m}_2 + \alpha + \frac{1}{2}r^{d-1}\eta)) d\eta + O(r^\infty), \quad (9-1)$$

which we express as

$$\begin{aligned} \mathcal{I}_{2,2}(s_1, s_2) &= r^{2d} h^d \sum_{\mathbf{m}_1, \mathbf{m}_2 \in \mathbb{Z}^d} \int_{\mathbb{R}^d} |\widehat{W}(r(\mathbf{m}_2 - \mathbf{m}_1))|^2 e^{-\frac{1}{2}(s_2 - s_1)r^{d-1}(\mathbf{m}_2 - \mathbf{m}_1) \cdot \eta} \\ &\quad \times e^{\frac{1}{2}(s_2 - s_1)(\|\mathbf{m}_2 + \alpha + \frac{1}{2}r^{d-1}\eta\|^2 - \|\mathbf{m}_1 + \alpha + \frac{1}{2}r^{d-1}\eta\|^2)} \\ &\quad \times \tilde{a}(-\eta, h(\mathbf{m}_2 + \alpha + \frac{1}{2}r^{d-1}\eta)) \tilde{b}(\eta, h(\mathbf{m}_2 + \alpha + \frac{1}{2}r^{d-1}\eta)) d\eta + O(r^\infty). \end{aligned} \quad (9-2)$$

In the same way we can see from (5-13) that $\mathcal{I}_{0,2}$ can be written

$$\begin{aligned} \mathcal{I}_{0,2}(s_1, s_2) &= r^{2d} h^d \sum_{\mathbf{m}_1, \mathbf{m}_2 \in \mathbb{Z}^d} \int_{\mathbb{R}^d} \tilde{a}(-\eta, h(\mathbf{m}_2 + \alpha + \frac{1}{2}r^{d-1}\eta)) e^{-\frac{1}{2}s_1\|\mathbf{m}_2 + \alpha + r^{d-1}\eta\|^2} \widehat{W}(r(\mathbf{m}_2 - \mathbf{m}_1)) \\ &\quad \times e^{-\frac{1}{2}(s_2 - s_1)\|\mathbf{m}_1 + \alpha + r^{d-1}\eta\|^2} \widehat{W}(r(\mathbf{m}_1 - \mathbf{m}_2)) \\ &\quad \times e^{\frac{1}{2}s_2\|\mathbf{m}_2 + \alpha + r^{d-1}\eta\|^2} \tilde{b}(\eta, h(\mathbf{m}_2 + \alpha + \frac{1}{2}r^{d-1}\eta)) d\eta + O(r^\infty), \end{aligned} \quad (9-3)$$

which we express as

$$\begin{aligned} \mathcal{I}_{0,2}(s_1, s_2) &= r^{2d} h^d \sum_{\mathbf{m}_1, \mathbf{m}_2 \in \mathbb{Z}^d} \int_{\mathbb{R}^d} |\widehat{W}(r(\mathbf{m}_2 - \mathbf{m}_1))|^2 e^{\frac{1}{2}(s_2 - s_1)r^{d-1}(\mathbf{m}_2 - \mathbf{m}_1) \cdot \eta} \\ &\quad \times e^{\frac{1}{2}(s_2 - s_1)(\|\mathbf{m}_2 + \alpha + \frac{1}{2}r^{d-1}\eta\|^2 - \|\mathbf{m}_1 + \alpha + \frac{1}{2}r^{d-1}\eta\|^2)} \\ &\quad \times \tilde{a}(-\eta, h(\mathbf{m}_2 + \alpha + \frac{1}{2}r^{d-1}\eta)) \tilde{b}(\eta, h(\mathbf{m}_2 + \alpha + \frac{1}{2}r^{d-1}\eta)) d\eta + O(r^\infty). \end{aligned} \quad (9-4)$$

We can then combine these two terms in the following way: First define $\mathcal{I}_{+,2}$ as

$$\begin{aligned} \mathcal{I}_{+,2}(s_1, s_2) &= r^{2d} h^d \sum_{\mathbf{m}_1, \mathbf{m}_2 \in \mathbb{Z}^d} \int_{\mathbb{R}^d} |\widehat{W}(r(\mathbf{m}_2 - \mathbf{m}_1))|^2 e^{-\frac{1}{2}|s_2 - s_1|r^{d-1}(\mathbf{m}_2 - \mathbf{m}_1) \cdot \eta} \\ &\quad \times e^{\frac{1}{2}(s_2 - s_1)(\|\mathbf{m}_2 + \alpha + \frac{1}{2}r^{d-1}\eta\|^2 - \|\mathbf{m}_1 + \alpha + \frac{1}{2}r^{d-1}\eta\|^2)} \\ &\quad \times \tilde{a}(-\eta, h(\mathbf{m}_2 + \alpha + \frac{1}{2}r^{d-1}\eta)) \tilde{b}(\eta, h(\mathbf{m}_2 + \alpha + \frac{1}{2}r^{d-1}\eta)) d\eta \end{aligned} \quad (9-5)$$

and note that

$$\mathcal{I}_{+,2}(s_1, s_2) = \begin{cases} \mathcal{I}_{2,2}(s_1, s_2) + O(r^\infty) & \text{if } s_1 \leq s_2, \\ \mathcal{I}_{0,2}(s_1, s_2) + O(r^\infty) & \text{if } s_1 \geq s_2. \end{cases} \quad (9-6)$$

Therefore, after inserting the integration over s_1 and s_2 we obtain

$$\begin{aligned} \int_{0 < s_1 < s_2 < hr^{1-d}t} \mathcal{I}_{2,2}(s_1, s_2) ds_1 ds_2 + \int_{0 < s_2 < s_1 < hr^{1-d}t} \mathcal{I}_{0,2}(s_1, s_2) ds_1 ds_2 \\ = \int_0^{hr^{1-d}t} \int_0^{hr^{1-d}t} \mathcal{I}_{+,2}(s_1, s_2) ds_1 ds_2 + O(r^\infty). \end{aligned} \quad (9-7)$$

Note that we measure time in units of hr^{1-d} as in the treatment of the zeroth-order term.

Lemma 9.1. *Let $\mathcal{I}_{+,2}$ be defined as above and set $h = r$. Then,*

$$\int_0^{hr^{1-d}t} \int_0^{hr^{1-d}t} \mathcal{I}_{+,2}(s_1, s_2) ds_1 ds_2 = r^{d+2} \int_{-r^{2-d}t}^{r^{2-d}t} F_r \left(\left(u + ir^2, 0, \begin{pmatrix} \mathbf{0} \\ -\boldsymbol{\alpha} \end{pmatrix} \right), r^{d-2}u \right) du, \quad (9-8)$$

with F_r as defined in (8-19), with the choice

$$f(\mathbf{y}_1, \mathbf{y}_2, u, \boldsymbol{\eta}) = e\left(\frac{1}{2}(u + |u|)(\mathbf{y}_2 - \mathbf{y}_1) \cdot \boldsymbol{\eta}\right) (t - |u|) \chi_{[-t, t]}(u) |\widehat{W}(\mathbf{y}_2 - \mathbf{y}_1)|^2 \tilde{a}(\boldsymbol{\eta}, \mathbf{y}_2) \tilde{b}(-\boldsymbol{\eta}, \mathbf{y}_2). \quad (9-9)$$

Proof. In the case $h = r$ the left-hand side of (9-8) reads (after the variable substitution $\boldsymbol{\eta} \mapsto -\boldsymbol{\eta}$)

$$\begin{aligned} r^{3d} \int_0^{r^{2-d}t} \int_0^{r^{2-d}t} \int_{\mathbb{R}^d} \sum_{\mathbf{m}_1, \mathbf{m}_2 \in \mathbb{Z}^d} |\widehat{W}(r(\mathbf{m}_2 - \mathbf{m}_1))|^2 e\left(\frac{1}{2}|s_2 - s_1|r^{d-1}(\mathbf{m}_2 - \mathbf{m}_1) \cdot \boldsymbol{\eta}\right) \\ \times e\left(\frac{1}{2}(s_2 - s_1)(\|\mathbf{m}_2 + \boldsymbol{\alpha} - \frac{1}{2}r^{d-1}\boldsymbol{\eta}\|^2 - \|\mathbf{m}_1 + \boldsymbol{\alpha} - \frac{1}{2}r^{d-1}\boldsymbol{\eta}\|^2)\right) \\ \times \tilde{a}(\boldsymbol{\eta}, r(\mathbf{m}_2 + \boldsymbol{\alpha} - \frac{1}{2}r^{d-1}\boldsymbol{\eta})) \tilde{b}(-\boldsymbol{\eta}, r(\mathbf{m}_2 + \boldsymbol{\alpha} - \frac{1}{2}r^{d-1}\boldsymbol{\eta})) d\boldsymbol{\eta} ds_1 ds_2. \end{aligned} \quad (9-10)$$

We then use the relation

$$\int_0^t \int_0^t f(s_2 - s_1) ds_1 ds_2 = \int_{-t}^t (t - |u|) f(u) du \quad (9-11)$$

to rewrite the above as

$$\begin{aligned} r^{2d+2} \int_{-r^{2-d}t}^{r^{2-d}t} \int_{\mathbb{R}^d} \sum_{\mathbf{m}_1, \mathbf{m}_2 \in \mathbb{Z}^d} |\widehat{W}(r(\mathbf{m}_2 - \mathbf{m}_1))|^2 e\left(\frac{1}{2}|u|r^{d-1}(\mathbf{m}_2 - \mathbf{m}_1) \cdot \boldsymbol{\eta}\right) \\ \times (t - r^{d-2}|u|) e\left(\frac{1}{2}u(\|\mathbf{m}_1 + \boldsymbol{\alpha} - \frac{1}{2}r^{d-1}\boldsymbol{\eta}\|^2 - \|\mathbf{m}_2 + \boldsymbol{\alpha} - \frac{1}{2}r^{d-1}\boldsymbol{\eta}\|^2)\right) \\ \times \tilde{a}(\boldsymbol{\eta}, r(\mathbf{m}_2 + \boldsymbol{\alpha} - \frac{1}{2}r^{d-1}\boldsymbol{\eta})) \tilde{b}(-\boldsymbol{\eta}, r(\mathbf{m}_2 + \boldsymbol{\alpha} - \frac{1}{2}r^{d-1}\boldsymbol{\eta})) d\boldsymbol{\eta} du \\ = r^{d+2} \int_{-r^{2-d}t}^{r^{2-d}t} \int_{\mathbb{R}^d} \Theta_f \left(\left(u + ir^2, 0, \begin{pmatrix} \frac{1}{2}ur^{d-1}\boldsymbol{\eta} \\ -\boldsymbol{\alpha} + \frac{1}{2}r^{d-1}\boldsymbol{\eta} \end{pmatrix} \right), r^{d-2}u, \boldsymbol{\eta} \right) d\boldsymbol{\eta} du, \end{aligned} \quad (9-12)$$

with f as in (9-9). Noting that

$$\left(u + ir^2, 0, \begin{pmatrix} \frac{1}{2}ur^{d-1}\boldsymbol{\eta} \\ -\boldsymbol{\alpha} + \frac{1}{2}r^{d-1}\boldsymbol{\eta} \end{pmatrix} \right) = \left(u + ir^2, 0, \begin{pmatrix} \mathbf{0} \\ -\boldsymbol{\alpha} \end{pmatrix} \right) \left(i, 0, \begin{pmatrix} \mathbf{0} \\ \frac{1}{2}r^d\boldsymbol{\eta} \end{pmatrix} \right), \quad (9-13)$$

the result follows. \square

Note that in view of (2-9) we should consider the rescaling of the coupling constant $\lambda \rightarrow \lambda h^{-2}$, or equivalently of the potential itself $W \rightarrow h^{-2}W$. At second order the potential appears as $|\widehat{W}|^2$, and so we must rescale our terms by a factor of h^{-4} .

Proposition 9.2. *Let $\mathcal{I}_{+,2}$ be defined as above. Then*

$$\begin{aligned} \lim_{h=r \rightarrow 0} h^{-4} \int_0^{hr^{1-d}t} \int_0^{hr^{1-d}t} \mathcal{I}_{+,2}(s_1, s_2) ds_1 ds_2 \\ = 2t \int_{(\mathbb{R}^d)^3} |\widehat{W}(\mathbf{y}_2 - \mathbf{y}_1)|^2 a(\mathbf{x}, \mathbf{y}_2) b(\mathbf{x}, \mathbf{y}_2) \delta(\|\mathbf{y}_1\|^2 - \|\mathbf{y}_2\|^2) d\mathbf{x} d\mathbf{y}_1 d\mathbf{y}_2 \\ + t^2 |\widehat{W}(\mathbf{0})|^2 \int_{\mathbb{R}^d \times \mathbb{R}^d} a(\mathbf{x}, \mathbf{y}) b(\mathbf{x}, \mathbf{y}) d\mathbf{x} d\mathbf{y}. \end{aligned} \quad (9-14)$$

Proof. By Proposition 8.9 and Lemma 9.1 we have that the limit in (9-14) is given by

$$2 \int_{(\mathbb{R}^d)^3} f(\mathbf{y}_1, \mathbf{y}_2, 0, \boldsymbol{\eta}) \delta(\|\mathbf{y}_1\|^2 - \|\mathbf{y}_2\|^2) d\mathbf{y}_1 d\mathbf{y}_2 d\boldsymbol{\eta} + \int_{-t}^t \int_{\mathbb{R}^d \times \mathbb{R}^d} f(\mathbf{y}, \mathbf{y}, u, \boldsymbol{\eta}) d\mathbf{y} d\boldsymbol{\eta} du. \quad (9-15)$$

We have for the first term

$$\begin{aligned} 2 \int_{(\mathbb{R}^d)^3} f(\mathbf{y}_1, \mathbf{y}_2, 0, \boldsymbol{\eta}) \delta(\|\mathbf{y}_1\|^2 - \|\mathbf{y}_2\|^2) d\mathbf{y}_1 d\mathbf{y}_2 d\boldsymbol{\eta} \\ = 2t \int_{(\mathbb{R}^d)^3} |\widehat{W}(\mathbf{y}_2 - \mathbf{y}_1)|^2 \tilde{a}(\boldsymbol{\eta}, \mathbf{y}_2) \tilde{b}(-\boldsymbol{\eta}, \mathbf{y}_2) \delta(\|\mathbf{y}_1\|^2 - \|\mathbf{y}_2\|^2) d\mathbf{y}_1 d\mathbf{y}_2 d\boldsymbol{\eta} \\ = 2t \int_{(\mathbb{R}^d)^3} |\widehat{W}(\mathbf{y}_2 - \mathbf{y}_1)|^2 a(\mathbf{x}, \mathbf{y}_2) b(\mathbf{x}, \mathbf{y}_2) \delta(\|\mathbf{y}_1\|^2 - \|\mathbf{y}_2\|^2) d\mathbf{x} d\mathbf{y}_1 d\mathbf{y}_2. \end{aligned} \quad (9-16)$$

Similarly for the second term we obtain

$$\begin{aligned} \int_{-t}^t \int_{\mathbb{R}^d \times \mathbb{R}^d} f(\mathbf{y}, \mathbf{y}, u, \boldsymbol{\eta}) d\mathbf{y} d\boldsymbol{\eta} du = \int_{-t}^t (t - |u|) \int_{\mathbb{R}^d \times \mathbb{R}^d} |\widehat{W}(\mathbf{0})|^2 \tilde{a}(\boldsymbol{\eta}, \mathbf{y}) \tilde{b}(-\boldsymbol{\eta}, \mathbf{y}) d\mathbf{y} d\boldsymbol{\eta} du \\ = t^2 |\widehat{W}(\mathbf{0})|^2 \int_{\mathbb{R}^d \times \mathbb{R}^d} a(\mathbf{x}, \mathbf{y}) b(\mathbf{x}, \mathbf{y}) d\mathbf{x} d\mathbf{y}, \end{aligned} \quad (9-17)$$

completing the proof. \square

The case $\ell = 1$.

Lemma 9.3. *For $h = r$,*

$$\int_0^{hr^{1-d}t} \int_0^{hr^{1-d}t} \mathcal{I}_{1,2}(s_1, s_2) ds_1 ds_2 = r^{d+2} \int_{-r^{2-d}t}^{r^{2-d}t} F_r \left(\left((u + ir^2, 0, (\mathbf{0} - \boldsymbol{\alpha})) \right), r^{d-2}u \right) du + O(r^\infty), \quad (9-18)$$

with F_r as defined in (8-19), where

$$\begin{aligned} f(\mathbf{y}_1, \mathbf{y}_2, u, \boldsymbol{\eta}) = \frac{1}{2} \left(\int_{|u|}^{2t-|u|} e\left(\frac{1}{2}(u-u')\boldsymbol{\eta} \cdot (\mathbf{y}_2 - \mathbf{y}_1)\right) du' \right) \chi_{[-t,t]}(u) \\ \times |\widehat{W}(\mathbf{y}_1 - \mathbf{y}_2)|^2 \tilde{a}(\boldsymbol{\eta}, \mathbf{y}_1) \tilde{b}(-\boldsymbol{\eta}, \mathbf{y}_2). \end{aligned} \quad (9-19)$$

Proof. As before, we start from (5-10). For $\mathcal{I}_{1,2}$ this yields the explicit formula

$$\begin{aligned} \mathcal{I}_{1,2}(s_1, s_2) \\ = r^{2d} h^d \sum_{\mathbf{m}_1, \mathbf{m}_2 \in \mathbb{Z}^d} \int_{\mathbb{R}^d} e\left(-\frac{1}{2} s_1 \|\mathbf{m}_2 + \boldsymbol{\alpha}\|^2\right) \widehat{W}(r(\mathbf{m}_2 - \mathbf{m}_1)) e\left(\frac{1}{2} s_1 \|\mathbf{m}_1 + \boldsymbol{\alpha}\|^2\right) \tilde{a}(-\boldsymbol{\eta}, h(\mathbf{m}_1 + \boldsymbol{\alpha} + \frac{1}{2} r^{d-1} \boldsymbol{\eta})) \\ \times e\left(-\frac{1}{2} s_2 \|\mathbf{m}_1 + \boldsymbol{\alpha} + r^{d-1} \boldsymbol{\eta}\|^2\right) \widehat{W}(r(\mathbf{m}_1 - \mathbf{m}_2)) e\left(\frac{1}{2} s_2 \|\mathbf{m}_2 + \boldsymbol{\alpha} + r^{d-1} \boldsymbol{\eta}\|^2\right) \\ \times \tilde{b}(\boldsymbol{\eta}, h(\mathbf{m}_2 + \boldsymbol{\alpha} + \frac{1}{2} r^{d-1} \boldsymbol{\eta})) d\boldsymbol{\eta} + O(r^\infty). \end{aligned} \quad (9-20)$$

We then note that we can write

$$\begin{aligned} & s_1 \|\mathbf{m}_1 + \boldsymbol{\alpha}\|^2 - s_2 \|\mathbf{m}_1 + \boldsymbol{\alpha} + r^{d-1} \boldsymbol{\eta}\|^2 \\ &= (s_1 - s_2) \|\mathbf{m}_1 + \boldsymbol{\alpha} + \frac{1}{2} r^{d-1} \boldsymbol{\eta}\|^2 - (s_1 + s_2) r^{d-1} \boldsymbol{\eta} \cdot (\mathbf{m}_1 + \boldsymbol{\alpha}) - \frac{1}{4} s_1 r^{2d-2} \|\boldsymbol{\eta}\|^2 - \frac{3}{4} s_2 r^{2d-2} \|\boldsymbol{\eta}\|^2 \end{aligned} \quad (9-21)$$

and similarly

$$\begin{aligned} & -s_1 \|\mathbf{m}_2 + \boldsymbol{\alpha}\|^2 + s_2 \|\mathbf{m}_2 + \boldsymbol{\alpha} + r^{d-1} \boldsymbol{\eta}\|^2 \\ &= (s_2 - s_1) \|\mathbf{m}_2 + \boldsymbol{\alpha} + \frac{1}{2} r^{d-1} \boldsymbol{\eta}\|^2 + (s_1 + s_2) r^{d-1} \boldsymbol{\eta} \cdot (\mathbf{m}_2 + \boldsymbol{\alpha}) + \frac{1}{4} s_1 r^{2d-2} \|\boldsymbol{\eta}\|^2 + \frac{3}{4} s_2 r^{2d-2} \|\boldsymbol{\eta}\|^2. \end{aligned} \quad (9-22)$$

We then insert these expressions into the exponential and make the variable substitutions $s_1 - s_2 = u_1$, $s_1 + s_2 = u_2$, and $\boldsymbol{\eta} \mapsto -\boldsymbol{\eta}$ to obtain

$$\begin{aligned} & \frac{1}{2} r^{d+2} h^d \int_{-hr^{1-d}t}^{hr^{1-d}t} \left(\int_{r^{d-2}|u_1|}^{2hr^{-1}t-r^{d-2}|u_1|} \sum_{\mathbf{m}_1, \mathbf{m}_2 \in \mathbb{Z}^d} \int_{\mathbb{R}^d} |\widehat{W}(r(\mathbf{m}_2 - \mathbf{m}_1))|^2 e(-\frac{1}{2} u_2 r \boldsymbol{\eta} \cdot (\mathbf{m}_2 - \mathbf{m}_1)) \right. \\ & \quad \times e(\frac{1}{2} u_1 (\|\mathbf{m}_1 + \boldsymbol{\alpha} - \frac{1}{2} r^{d-1} \boldsymbol{\eta}\|^2 - \|\mathbf{m}_2 + \boldsymbol{\alpha} - \frac{1}{2} r^{d-1} \boldsymbol{\eta}\|^2)) \\ & \quad \times \tilde{a}(\boldsymbol{\eta}, h(\mathbf{m}_1 + \boldsymbol{\alpha} - \frac{1}{2} r^{d-1} \boldsymbol{\eta})) \\ & \quad \left. \times \tilde{b}(-\boldsymbol{\eta}, h(\mathbf{m}_2 + \boldsymbol{\alpha} - \frac{1}{2} r^{d-1} \boldsymbol{\eta})) \right) du_2 \Big) du_1 \\ &= r^{d+2} \int_{-r^{2-d}t}^{r^{2-d}t} \int_{\mathbb{R}^d} \Theta_f \left(\left(u_1 + ir^2, 0, \left(\frac{1}{2} u_1 r^{d-1} \boldsymbol{\eta}, -\boldsymbol{\alpha} + \frac{1}{2} r^{d-1} \boldsymbol{\eta} \right) \right), r^{d-2} u_1, \boldsymbol{\eta} \right) d\boldsymbol{\eta} du_1, \end{aligned} \quad (9-23)$$

with f as in (9-19). The statement follows from (9-13). □

Proposition 9.4.

$$\begin{aligned} & \lim_{h \rightarrow 0} h^{-4} \int_0^{hr^{1-d}t} \int_0^{hr^{1-d}t} \mathcal{I}_{1,2}(s_1, s_2) ds_1 ds_2 \\ &= 2 \int_0^t \int_{(\mathbb{R}^d)^3} |\widehat{W}(\mathbf{y}_2 - \mathbf{y}_1)|^2 \delta(\|\mathbf{y}_1\|^2 - \|\mathbf{y}_2\|^2) a(\mathbf{x} - s(\mathbf{y}_2 - \mathbf{y}_1), \mathbf{y}_1) b(\mathbf{x}, \mathbf{y}_2) d\mathbf{y}_1 d\mathbf{y}_2 d\mathbf{x} ds \\ & \quad + t^2 |\widehat{W}(\mathbf{0})|^2 \int_{\mathbb{R}^d \times \mathbb{R}^d} a(\mathbf{x}, \mathbf{y}) b(\mathbf{x}, \mathbf{y}) d\mathbf{x} d\mathbf{y}. \end{aligned} \quad (9-24)$$

Proof. By Proposition 8.9 and Lemma 9.3 we have that the limit in (9-24) is the sum of two terms. The first one can be written

$$\begin{aligned} & 2 \int_{(\mathbb{R}^d)^3} f(\mathbf{y}_1, \mathbf{y}_2, 0, \boldsymbol{\eta}) \delta(\|\mathbf{y}_1\|^2 - \|\mathbf{y}_2\|^2) d\mathbf{y}_1 d\mathbf{y}_2 d\boldsymbol{\eta} \\ &= 2 \int_{(\mathbb{R}^d)^3} |\widehat{W}(\mathbf{y}_2 - \mathbf{y}_1)|^2 \tilde{a}(\boldsymbol{\eta}, \mathbf{y}_1) \tilde{b}(-\boldsymbol{\eta}, \mathbf{y}_2) \delta(\|\mathbf{y}_1\|^2 - \|\mathbf{y}_2\|^2) \left(\int_0^t e(-u' \boldsymbol{\eta} \cdot (\mathbf{y}_2 - \mathbf{y}_1)) du' \right) d\mathbf{y}_1 d\mathbf{y}_2 d\boldsymbol{\eta} \\ &= 2 \int_0^t \int_{(\mathbb{R}^d)^3} |\widehat{W}(\mathbf{y}_2 - \mathbf{y}_1)|^2 a(\mathbf{x} - s(\mathbf{y}_2 - \mathbf{y}_1), \mathbf{y}_1) b(\mathbf{x}, \mathbf{y}_2) \delta(\|\mathbf{y}_1\|^2 - \|\mathbf{y}_2\|^2) d\mathbf{y}_1 d\mathbf{y}_2 d\mathbf{x} ds. \end{aligned} \quad (9-25)$$

The second term takes the form

$$\begin{aligned} \frac{1}{2} \int_{-t}^t \int_{\mathbb{R}^d \times \mathbb{R}^d} f(\mathbf{y}, \mathbf{y}, u, \boldsymbol{\eta}) \, d\mathbf{y} \, d\boldsymbol{\eta} \, du' \, du &= \int_{-t}^t (t - |u|) \int_{\mathbb{R}^d \times \mathbb{R}^d} |\widehat{W}(\mathbf{0})|^2 \tilde{a}(\boldsymbol{\eta}, \mathbf{y}) \tilde{b}(-\boldsymbol{\eta}, \mathbf{y}) \, d\mathbf{y} \, d\boldsymbol{\eta} \, du \\ &= t^2 |\widehat{W}(\mathbf{0})|^2 \int_{\mathbb{R}^d \times \mathbb{R}^d} a(\mathbf{x}, \mathbf{y}) b(\mathbf{x}, \mathbf{y}) \, d\mathbf{x} \, d\mathbf{y}. \end{aligned} \quad (9-26)$$

This completes the proof. \square

Thus, combining $\mathcal{I}_{j,2}$ for $j = 0, 1, 2$ yields the following limiting expression for the second-order terms.

Corollary 9.5.

$$\begin{aligned} \lim_{h=r \rightarrow 0} h^{-4} &\left[- \int_0^{hr^{1-d}t} \int_0^{s_2} \mathcal{I}_{2,2}(s_1, s_2) \, ds_1 \, ds_2 \right. \\ &\quad \left. + \int_0^{hr^{1-d}t} \int_0^{hr^{1-d}t} \mathcal{I}_{1,2}(s_1, s_2) \, ds_1 \, ds_2 - \int_0^{hr^{1-d}t} \int_{s_2}^{hr^{1-d}t} \mathcal{I}_{0,2}(s_1, s_2) \, ds_1 \, ds_2 \right] \\ &= 2 \int_0^t \int_{(\mathbb{R}^d)^3} |\widehat{W}(\mathbf{y}_2 - \mathbf{y}_1)|^2 \delta(\|\mathbf{y}_1\|^2 - \|\mathbf{y}_2\|^2) \\ &\quad \times [a(\mathbf{x} - s(\mathbf{y}_2 - \mathbf{y}_1), \mathbf{y}_1) - a(\mathbf{x}, \mathbf{y}_2)] b(\mathbf{x}, \mathbf{y}_2) \, d\mathbf{y}_1 d\mathbf{y}_2 \, d\mathbf{x} \, ds. \end{aligned} \quad (9-27)$$

Now replacing a by the time-evolved symbol $L_0(t)a$ yields, in place of (9-27),

$$\begin{aligned} 2 \int_0^t \int_{(\mathbb{R}^d)^3} |\widehat{W}(\mathbf{y}_2 - \mathbf{y}_1)|^2 \delta(\|\mathbf{y}_1\|^2 - \|\mathbf{y}_2\|^2) \\ \times [a(\mathbf{x} - (t-s)\mathbf{y}_1 - s\mathbf{y}_2, \mathbf{y}_1) - a(\mathbf{x} - t\mathbf{y}_2, \mathbf{y}_2)] b(\mathbf{x}, \mathbf{y}_2) \, d\mathbf{x} \, d\mathbf{y}_1 \, d\mathbf{y}_2 \, ds. \end{aligned} \quad (9-28)$$

10. Higher-order theta functions

In order to prove bounds on the error terms (4-15) in the Duhamel expansion we will need to define higher-order theta functions, that is, generalisations of the theta function given in (8-8) that live on the product space $(\Gamma \backslash G)^k$. Specifically, for $f \in \mathcal{S}(\mathbb{R}^{d \times k} \times \mathbb{R}^{d \times k})$, we denote by $\Theta_f^{(k)} : (\Gamma \backslash G)^k \rightarrow \mathbb{C}$ the theta function

$$\begin{aligned} \Theta_f^{(k)}(\boldsymbol{\tau}, \boldsymbol{\phi}, \boldsymbol{\Xi}) &= \det(\mathbf{v})^{d/2} \sum_{\mathbf{M}, \mathbf{M}' \in \mathbb{Z}^{d \times k}} f_{\boldsymbol{\phi}}((\mathbf{M} - \mathbf{Y})\mathbf{v}^{1/2}, (\mathbf{M}' - \mathbf{Y})\mathbf{v}^{1/2}) \\ &\quad \times e(\text{Tr}[\frac{1}{2} {}^t(\mathbf{M} - \mathbf{Y})(\mathbf{M} - \mathbf{Y})\mathbf{u} - \frac{1}{2} {}^t(\mathbf{M}' - \mathbf{Y})(\mathbf{M}' - \mathbf{Y})\mathbf{u} + {}^t(\mathbf{M} - \mathbf{M}')\mathbf{X}]), \end{aligned} \quad (10-1)$$

or more explicitly,

$$\begin{aligned} \Theta_f^{(k)}(\boldsymbol{\tau}, \boldsymbol{\phi}, \boldsymbol{\Xi}) &= \sum_{\substack{\mathbf{m}_1, \dots, \mathbf{m}_k \in \mathbb{Z}^d \\ \mathbf{m}'_1, \dots, \mathbf{m}'_k \in \mathbb{Z}^d}} f_{\boldsymbol{\phi}}(v_1^{1/2}(\mathbf{m}_1 - \mathbf{y}_1), \dots, v_k^{1/2}(\mathbf{m}_k - \mathbf{y}_k), v_1^{1/2}(\mathbf{m}'_1 - \mathbf{y}_1), \dots, v_k^{1/2}(\mathbf{m}'_k - \mathbf{y}_k)) \\ &\quad \times \prod_{j=1}^k v_j^{d/2} e(\frac{1}{2} u_j (\|\mathbf{m}_j - \mathbf{y}_j\|^2 - \|\mathbf{m}'_j - \mathbf{y}_j\|^2) + \mathbf{x}_j \cdot (\mathbf{m}_j - \mathbf{m}'_j)), \end{aligned} \quad (10-2)$$

where we use the natural notation

$$\begin{aligned} \tau &= \mathbf{u} + i\mathbf{v}, \quad \mathbf{u} = \text{diag}(u_1, \dots, u_k), \quad u_j \in \mathbb{R}, \\ \mathbf{v} &= \text{diag}(v_1, \dots, v_k), \quad v_j \in \mathbb{R}_{>0}, \quad \boldsymbol{\phi} = (\phi_1, \dots, \phi_k) \in [0, 2\pi)^k, \\ \boldsymbol{\Xi} &= (\boldsymbol{\xi}_1, \dots, \boldsymbol{\xi}_k) = \left(\begin{pmatrix} \mathbf{x}_1 \\ \mathbf{y}_1 \end{pmatrix}, \dots, \begin{pmatrix} \mathbf{x}_k \\ \mathbf{y}_k \end{pmatrix} \right) \in \mathbb{R}^{2d \times k}, \\ \mathbf{X} &= (\mathbf{x}_1, \dots, \mathbf{x}_k) \in \mathbb{R}^{d \times k}, \quad \mathbf{Y} = (\mathbf{y}_1, \dots, \mathbf{y}_k) \in \mathbb{R}^{d \times k}, \\ \mathbf{M} &= (\mathbf{m}_1, \dots, \mathbf{m}_k) \in \mathbb{Z}^{d \times k} \end{aligned} \tag{10-3}$$

and

$$f_\phi(\mathbf{Y}, \mathbf{Y}') = \int_{\mathbb{R}^{d \times k} \times \mathbb{R}^{d \times k}} G_\phi(\mathbf{Y}, \mathbf{Y}', \mathbf{Z}, \mathbf{Z}') f(\mathbf{Z}, \mathbf{Z}') d\mathbf{Z} d\mathbf{Z}', \tag{10-4}$$

with

$$G_\phi(\mathbf{Y}, \mathbf{Y}', \mathbf{Z}, \mathbf{Z}') = \prod_{j=1}^k |\sin \phi_j|^{-d} e^{\left(\frac{\frac{1}{2}(\|\mathbf{y}_j\|^2 + \|\mathbf{z}_j\|^2 - \|\mathbf{y}'_j\|^2 - \|\mathbf{z}'_j\|^2) \cos \phi_j - \mathbf{y}_j \cdot \mathbf{z}_j + \mathbf{y}'_j \cdot \mathbf{z}'_j}{\sin \phi_j} \right)}. \tag{10-5}$$

For $\phi = 0 \pmod{\pi}$ we define f_ϕ by generalising (8-16) in the analogous way. In the special case where $f = \prod_{j=1}^k f_j$, with $f_j \in \mathcal{S}(\mathbb{R}^d \times \mathbb{R}^d)$, the function $\Theta_f^{(k)}$ becomes the product of k independent theta functions of the form (8-8). In a similar vein as earlier, we wish to consider a generalisation of this theta function in which the function f is allowed to depend directly on $\mathbf{u} \in \mathbb{R}^k$ and some new parameters $\boldsymbol{\eta} \in \mathbb{R}^d$ and $\omega \in \mathbb{R}$.

We denote by $\tilde{\mathcal{S}}_k$ the class of functions $f \in C(\mathbb{R}^{d \times k} \times \mathbb{R}^{d \times k} \times \mathbb{R}^k \times \mathbb{R}^d \times \mathbb{R})$ with the property that for every multi-index $\boldsymbol{\beta}_1, \boldsymbol{\beta}_2 \in \mathbb{Z}_{\geq 0}^{d \times k}$, the derivative

$$\partial_{\mathbf{Y}_1}^{\boldsymbol{\beta}_1} \partial_{\mathbf{Y}_2}^{\boldsymbol{\beta}_2} f(\mathbf{Y}_1, \mathbf{Y}_2, \mathbf{u}, \boldsymbol{\eta}, \omega)$$

(a) exists, (b) is continuous (in all variables), and (c) is rapidly decaying, i.e.,

$$\sup_{\mathbf{Y}_1, \mathbf{Y}_2, \mathbf{u}, \boldsymbol{\eta}, \omega} (1 + \|\mathbf{Y}_1\|)^T (1 + \|\mathbf{Y}_2\|)^T (1 + |\mathbf{u}|)^T (1 + \|\boldsymbol{\eta}\|)^T (1 + |\omega|)^T |\partial_{\mathbf{Y}_1}^{\boldsymbol{\beta}_1} \partial_{\mathbf{Y}_2}^{\boldsymbol{\beta}_2} f(\mathbf{Y}_1, \mathbf{Y}_2, \mathbf{u}, \boldsymbol{\eta}, \omega)| < \infty \tag{10-6}$$

for every $T > 1$.

We then consider the test function $f = f(\mathbf{Y}, \mathbf{Y}', \mathbf{u}, \boldsymbol{\eta}, \omega)$ in $\tilde{\mathcal{S}}_k$ and set

$$\Theta_f^{(k)}(g, \mathbf{u}, \boldsymbol{\eta}, \omega) := \Theta_{f(\cdot, \cdot, \mathbf{u}, \boldsymbol{\eta}, \omega)}^{(k)}(g). \tag{10-7}$$

We now proceed to state some results in direct analogy with Section 8.

Lemma 10.1 (cf. Lemma 8.6). *If $f \in \tilde{\mathcal{S}}_k$, then for all multi-indices $\boldsymbol{\beta}_1, \boldsymbol{\beta}_2 \in \mathbb{Z}_{\geq 0}^{d \times k}$ and every $T > 1$*

$$\begin{aligned} \sup_{\mathbf{Y}_1, \mathbf{Y}_2, \mathbf{u}, \boldsymbol{\eta}, \omega, \phi} (1 + \|\mathbf{Y}_1\|)^T (1 + \|\mathbf{Y}_2\|)^T (1 + |\mathbf{u}|)^T \\ \times (1 + \|\boldsymbol{\eta}\|)^T (1 + |\omega|)^T |\partial_{\mathbf{Y}_1}^{\boldsymbol{\beta}_1} \partial_{\mathbf{Y}_2}^{\boldsymbol{\beta}_2} f_\phi(\mathbf{Y}_1, \mathbf{Y}_2, \mathbf{u}, \boldsymbol{\eta}, \omega)| < \infty. \end{aligned} \tag{10-8}$$

Proof. The proof is analogous to those of Lemmas 8.2 and 8.6. □

Now, let us use the shorthand

$$z_k(\boldsymbol{\eta}) := \left(\left(1, \begin{pmatrix} \mathbf{0} \\ \frac{1}{2}\boldsymbol{\eta} \end{pmatrix} \right), \dots, \left(1, \begin{pmatrix} \mathbf{0} \\ \frac{1}{2}\boldsymbol{\eta} \end{pmatrix} \right) \right) \in G^k,$$

and further define

$$F_r^{k,\beta}(g, \mathbf{u}) := \int_{\mathbb{R}} \left| \int_{\mathbb{R}^d} \Theta_f^{(k)}(g z_k(r^d \boldsymbol{\eta}), \mathbf{u}, \boldsymbol{\eta}, \omega) d\boldsymbol{\eta} \right|^\beta d\omega. \quad (10-9)$$

Proposition 10.2 (cf. Proposition 8.7). *Let $0 < \beta < 1$ and $f \in \tilde{\mathcal{S}}_k$. Then,*

$$\begin{aligned} & F_r^{k,\beta}(\mathbf{u} + i\mathbf{v}, \boldsymbol{\phi}, \boldsymbol{\Xi}, \mathbf{u}') \\ &= \det(\mathbf{v})^{\beta d/2} \sum_{\mathbf{M} \in \mathbb{Z}^{d \times k}} \int_{\mathbb{R}} \left| \int_{\mathbb{R}^d} f_{\boldsymbol{\phi}}((\mathbf{M} - \mathbf{Y})\mathbf{v}^{1/2}, (\mathbf{M} - \mathbf{Y})\mathbf{v}^{1/2}, \mathbf{u}', \boldsymbol{\eta}, \omega) d\boldsymbol{\eta} \right|^\beta d\omega + O(r^d) + \sum_{j=1}^k O(v_j^{-\infty}) \end{aligned} \quad (10-10)$$

uniformly for all $(\mathbf{u} + i\mathbf{v}, \boldsymbol{\phi}, \boldsymbol{\Xi}) \in (\Gamma \backslash G)^k$, $\mathbf{u}' \in \mathbb{R}^k$ with $v_j > \frac{1}{2}$ for all j and $r < 1$.

Proof. The proof is analogous to that of Proposition 8.7. \square

Recall the definitions of $\Psi_{R,f}^\beta$ and \bar{f} in (7-11) and (8-11).

Lemma 10.3. *Fix $T > d$. Then:*

(1) *There is a constant C such that for all $r < 1$*

$$|F_r^{k,\beta}(\boldsymbol{\tau}, \boldsymbol{\phi}, \boldsymbol{\Xi}, \mathbf{u}')| < C \prod_{j=1}^k (1 + \Psi_{1/2, \bar{f}}^\beta(\boldsymbol{\tau}_j, \boldsymbol{\xi}_j)). \quad (10-11)$$

(2) $F_r^{k,\beta} \rightarrow F_0^{k,\beta}$ *uniformly on compacta.*

Proof. The proof is analogous to that of Lemma 8.8 with Lemma 10.1 in place of Lemma 8.6. \square

In the following, we denote by I_k the $k \times k$ identity matrix.

Proposition 10.4. *Let $0 < \beta < 1$ and $f \in \tilde{\mathcal{S}}_k$. Assume for $j = 1, \dots, k$ that \mathbf{y}_j is Diophantine of type $\kappa < (d-1)/(d-2)$ with the components of $(1, {}^t\mathbf{y}_j)$ linearly independent over \mathbb{Q} . Let $w : \mathbb{R}^k \rightarrow \mathbb{R}$ be bounded with compact support. Then,*

$$\limsup_{r \rightarrow 0} r^{k(d-2)} \int_{\mathbb{R}^k} F_r^{k,\beta} \left(\mathbf{u} + ir^2 I_k, \mathbf{0}, \begin{pmatrix} \mathbf{0} \\ \mathbf{Y} \end{pmatrix}, r^{d-2} \mathbf{u} \right) w(r^{d-2} \mathbf{u}) d\mathbf{u} < \infty. \quad (10-12)$$

Proof. Applying Lemma 10.3 yields

$$\begin{aligned} & \limsup_{r \rightarrow 0} r^{k(d-2)} \int_{\mathbb{R}^k} F_r^{k,\beta} \left(\mathbf{u} + ir^2 I_k, \mathbf{0}, \begin{pmatrix} \mathbf{0} \\ \mathbf{Y} \end{pmatrix}, r^{d-2} \mathbf{u} \right) w(r^{d-2} \mathbf{u}) d\mathbf{u} \\ & < C \limsup_{r \rightarrow 0} r^{k(d-2)} \int_{\mathbb{R}^k} \left[\prod_{j=1}^k \left(1 + \Psi_{1/2, \bar{f}}^\beta \left(u_j + ir^2 I_k, \begin{pmatrix} \mathbf{0} \\ \mathbf{y}_j \end{pmatrix} \right) \right) \right] w(r^{d-2} \mathbf{u}) d\mathbf{u}. \end{aligned} \quad (10-13)$$

The function w has compact support, so fix L such that the cube $(-L, L)^k$ contains the support of w , and denote by χ_L the characteristic function of the interval $(-L, L)$. We can then bound the above expression by

$$C \sup |w| \limsup_{r \rightarrow 0} \prod_{j=1}^k \left(r^{d-2} \int_{\mathbb{R}} \left(1 + \Psi_{1/2, \bar{f}}^\beta \left(u_j + ir^2 I_k, \begin{pmatrix} \mathbf{0} \\ \mathbf{y}_j \end{pmatrix} \right) \right) \chi_L(r^{d-2} u_j) du_j \right). \quad (10-14)$$

The result then follows by applying Proposition 7.3. □

11. Error terms

In this section we prove upper bounds on the error terms (4-15) in the semiclassical Boltzmann–Grad scaling, i.e., for $Q_n(hr^{1-d}t, D_{r,ha}, D_{r,hb})$, where relevant cases are $n = 3, 4, 5, 6$. Lemma 4.1 tells us that

$$|Q_n(hr^{1-d}t, D_{r,ha}, D_{r,hb})| \leq \sum_{\ell=n-3}^3 \mathcal{J}_{\ell,n}(hr^{1-d}t, D_{r,ha}) \|\Pi_\alpha \text{Op}_{r,h}(b)\|_{\text{HS},\alpha}. \quad (11-1)$$

The term $\|\Pi_\alpha \text{Op}_{r,h}(b)\|_{\text{HS},\alpha}$ has a uniform upper bound; see Lemma 6.1. Hence the key is to estimate (recall (4-16) and Lemma 3.2)

$$\begin{aligned} & \mathcal{J}_{\ell,n}(hr^{1-d}t, D_{r,ha}) \\ &= (2\pi)^n \int_{\substack{0 < s_1 < \dots < s_\ell < thr^{1-d} \\ 0 < s_n < \dots < s_{\ell+1} < thr^{1-d}}} (\text{Tr}_\alpha [K_{1,\ell}(s)^\dagger K_{1,\ell}(s) \text{Op}_{r,h}(a) K_{\ell+1,n}(s) K_{\ell+1,n}(s)^\dagger \text{Op}_{r,h}(\bar{a})])^{1/2} ds. \end{aligned} \quad (11-2)$$

A straightforward computation (see Section C in the appendix) yields the expression

$$\begin{aligned} & \text{Tr}_\alpha [K_{1,\ell}(s)^\dagger K_{1,\ell}(s) \text{Op}_{r,h}(a) K_{\ell+1,n}(s) K_{\ell+1,n}(s)^\dagger \text{Op}_{r,h}(\bar{a})] \\ &= r^{2nd} h^d \sum_{\substack{\mathbf{m}_0, \mathbf{m}_1, \dots, \mathbf{m}_n \in \mathbb{Z}^d \\ \mathbf{m}'_1, \dots, \mathbf{m}'_n \in \mathbb{Z}^d}} \mathbb{1}[\mathbf{m}'_n - \mathbf{m}_n + \mathbf{m}_\ell - \mathbf{m}'_\ell = 0] \\ & \quad \times \int_{\mathbb{R}^d} \widehat{W}(r(\mathbf{m}_0 - \mathbf{m}_1)) \mathcal{T}_{1,\ell-1}^-(\boldsymbol{\alpha}) e^{(\frac{1}{2} s_\ell (\|\mathbf{m}_\ell + \boldsymbol{\alpha}\|^2 - \|\mathbf{m}'_\ell + \boldsymbol{\alpha}\|^2))} \\ & \quad \times \widehat{W}(r(\mathbf{m}'_1 - \mathbf{m}_0)) \overline{\mathcal{T}}_{1,\ell-1}^-(\boldsymbol{\alpha}) \tilde{a}(\boldsymbol{\eta}, h(\mathbf{m}_\ell + \boldsymbol{\alpha} - \frac{1}{2} r^{d-1} \boldsymbol{\eta})) \\ & \quad \times \widehat{W}(r(\mathbf{m}_\ell - \mathbf{m}_{\ell+1})) \mathcal{T}_{\ell+1,n-1}^-(\boldsymbol{\alpha} - r^{d-1} \boldsymbol{\eta}) \\ & \quad \times e^{(\frac{1}{2} s_{\ell+1} (\|\mathbf{m}'_\ell + \boldsymbol{\alpha} - r^{d-1} \boldsymbol{\eta}\|^2 - \|\mathbf{m}_\ell + \boldsymbol{\alpha} - r^{d-1} \boldsymbol{\eta}\|^2))} \\ & \quad \times \widehat{W}(r(\mathbf{m}'_{\ell+1} - \mathbf{m}'_\ell)) \overline{\mathcal{T}}_{\ell+1,n-1}^-(\boldsymbol{\alpha} - r^{d-1} \boldsymbol{\eta}) \\ & \quad \times \tilde{a}(-\boldsymbol{\eta}, h(\mathbf{m}'_\ell + \boldsymbol{\alpha} - \frac{1}{2} r^{d-1} \boldsymbol{\eta})) d\boldsymbol{\eta} + O(r^\infty), \end{aligned} \quad (11-3)$$

where

$$\begin{aligned} \mathcal{T}_{\ell,n}^-(\mathbf{y}) &= \begin{cases} \prod_{j=\ell}^n e^{(\frac{1}{2} (s_j - s_{j+1}) \|\mathbf{y} + \mathbf{m}_j\|^2)} \widehat{W}(r(\mathbf{m}_j - \mathbf{m}_{j+1})) & (l \leq n), \\ 1 & (l > n), \end{cases} \\ \overline{\mathcal{T}}_{\ell,n}^-(\mathbf{y}) &= \begin{cases} \prod_{j=\ell}^n e^{(\frac{1}{2} (s_{j+1} - s_j) \|\mathbf{y} + \mathbf{m}'_j\|^2)} \widehat{W}(r(\mathbf{m}'_{j+1} - \mathbf{m}'_j)) & (l \leq n), \\ 1 & (l > n). \end{cases} \end{aligned} \quad (11-4)$$

Let us focus on the exponential factors in (11-3); they are

$$\begin{aligned} & \left(\prod_{j=1}^{\ell-1} e\left(\frac{1}{2}(s_j - s_{j+1})(\|\mathbf{m}_j + \boldsymbol{\alpha}\|^2 - \|\mathbf{m}'_j + \boldsymbol{\alpha}\|^2)\right) \right) \\ & \times e\left(\frac{1}{2}s_\ell(\|\mathbf{m}_\ell + \boldsymbol{\alpha}\|^2 - \|\mathbf{m}'_\ell + \boldsymbol{\alpha}\|^2)\right) e\left(\frac{1}{2}s_{\ell+1}(\|\mathbf{m}'_\ell + \boldsymbol{\alpha} - r^{d-1}\boldsymbol{\eta}\|^2 - \|\mathbf{m}_\ell + \boldsymbol{\alpha} - r^{d-1}\boldsymbol{\eta}\|^2)\right) \\ & \times \left(\prod_{j=\ell+1}^{n-1} e\left(\frac{1}{2}(s_j - s_{j+1})(\|\mathbf{m}_j + \boldsymbol{\alpha} - r^{d-1}\boldsymbol{\eta}\|^2 - \|\mathbf{m}'_j + \boldsymbol{\alpha} - r^{d-1}\boldsymbol{\eta}\|^2)\right) \right). \end{aligned} \quad (11-5)$$

We write the above as

$$\begin{aligned} & \left(\prod_{j=1}^{\ell-1} e\left(\frac{1}{2}(s_j - s_{j+1})(\|\mathbf{m}_j + \boldsymbol{\alpha} - \frac{1}{2}r^{d-1}\boldsymbol{\eta}\|^2 - \|\mathbf{m}'_j + \boldsymbol{\alpha} - \frac{1}{2}r^{d-1}\boldsymbol{\eta}\|^2 + r^{d-1}(\mathbf{m}_j - \mathbf{m}'_j) \cdot \boldsymbol{\eta})\right) \right) \\ & \times e\left(\frac{1}{2}(s_\ell - s_{\ell+1})(\|\mathbf{m}_\ell + \boldsymbol{\alpha} - \frac{1}{2}r^{d-1}\boldsymbol{\eta}\|^2 - \|\mathbf{m}'_\ell + \boldsymbol{\alpha} - \frac{1}{2}r^{d-1}\boldsymbol{\eta}\|^2)\right) e\left(\frac{1}{2}(s_\ell + s_{\ell+1})r^{d-1}\boldsymbol{\eta} \cdot (\mathbf{m}_\ell - \mathbf{m}'_\ell)\right) \\ & \times \left(\prod_{j=\ell+1}^{n-1} e\left(\frac{1}{2}(s_j - s_{j+1})(\|\mathbf{m}_j + \boldsymbol{\alpha} - \frac{1}{2}r^{d-1}\boldsymbol{\eta}\|^2 - \|\mathbf{m}'_j + \boldsymbol{\alpha} - \frac{1}{2}r^{d-1}\boldsymbol{\eta}\|^2 - r^{d-1}(\mathbf{m}_j - \mathbf{m}'_j) \cdot \boldsymbol{\eta})\right) \right). \end{aligned} \quad (11-6)$$

Note that this product of exponentials is independent of the variables \mathbf{m}_0 , \mathbf{m}_n and \mathbf{m}'_n , and so the entire dependence on these variables is in the product of \widehat{W} -terms. In (11-3) we can therefore separately evaluate the threefold sum

$$\sum_{\substack{\mathbf{m}_0, \mathbf{m}_n, \mathbf{m}'_n \\ \mathbf{m}'_n - \mathbf{m}_n + \mathbf{m}_\ell - \mathbf{m}'_\ell = 0}} \widehat{W}(r(\mathbf{m}_0 - \mathbf{m}_1)) \widehat{W}(r(\mathbf{m}'_1 - \mathbf{m}_0)) \widehat{W}(r(\mathbf{m}_{n-1} - \mathbf{m}_n)) \widehat{W}(r(\mathbf{m}'_n - \mathbf{m}'_{n-1})), \quad (11-7)$$

which is equal to

$$\sum_{\mathbf{m}_0, \mathbf{m}_n} \widehat{W}(r(\mathbf{m}_0 - \mathbf{m}_1)) \widehat{W}(r(\mathbf{m}'_1 - \mathbf{m}_0)) \widehat{W}(r(\mathbf{m}_{n-1} - \mathbf{m}_n)) \widehat{W}(r(\mathbf{m}_n + \mathbf{m}'_\ell - \mathbf{m}_\ell - \mathbf{m}'_{n-1})). \quad (11-8)$$

Applying the Poisson summation formula to the sums over \mathbf{m}_0 and \mathbf{m}_n yields

$$r^{-2d} \sum_{\mathbf{k}_0, \mathbf{k}_n} \iint_{\mathbb{R}^{2d}} \widehat{W}(\mathbf{y}_0 - r\mathbf{m}_1) \widehat{W}(r\mathbf{m}'_1 - \mathbf{y}_0) \widehat{W}(r\mathbf{m}_{n-1} - \mathbf{y}_n) \\ \times \widehat{W}(\mathbf{y}_n + r(\mathbf{m}'_\ell - \mathbf{m}_\ell - \mathbf{m}'_{n-1})) e(r^{-1}\mathbf{k}_0 \cdot \mathbf{y}_0 + r^{-1}\mathbf{k}_n \cdot \mathbf{y}_n) d\mathbf{y}_0 d\mathbf{y}_n. \quad (11-9)$$

Since $W \in \mathcal{S}(\mathbb{R}^d)$, we have for any $T_1, T_2 \geq 1$ that (11-9) equals

$$r^{-2d} \mathcal{W}(r(\mathbf{m}'_1 - \mathbf{m}_1)) \mathcal{W}(r(\mathbf{m}'_\ell - \mathbf{m}_\ell + \mathbf{m}_{n-1} - \mathbf{m}'_{n-1})) \\ + O_T(r^{T_1}(1 + r\|\mathbf{m}_1 - \mathbf{m}'_1\|)^{-T_2}(1 + r\|\mathbf{m}'_\ell - \mathbf{m}_\ell + \mathbf{m}_{n-1} - \mathbf{m}'_{n-1}\|)^{-T_2}), \quad (11-10)$$

with

$$\mathcal{W}(\mathbf{t}) = \int_{\mathbb{R}^d} \widehat{W}(\mathbf{t} - \mathbf{y}) \widehat{W}(\mathbf{y}) d\mathbf{y}. \quad (11-11)$$

The error term in (11-10), after applying the remaining \mathbf{m}_j -sums, yields therefore a total contribution of order $O(r^\infty)$ for $h = r \in (0, 1]$. In order to write (11-3) as a higher-order theta function, we change

variable by the linear map $A : \mathbb{R}^n \rightarrow \mathbb{R}^n$, $s \mapsto \omega = As$, given by

$$\omega_j = s_j - s_{j+1} \quad (j = 1, \dots, n-1), \quad \omega_n = s_\ell + s_{\ell+1}. \quad (11-12)$$

The corresponding determinant equals 2, and hence A is invertible. Let

$$\mathcal{Q} = \{s \in \mathbb{R}^n : 0 < s_1 < \dots < s_\ell < 1, 0 < s_n < \dots < s_{\ell+1} < 1\}. \quad (11-13)$$

Then, for $h = r$ and $\omega = (\mathbf{u}, \omega) \in \mathbb{R}^{n-1} \times \mathbb{R}$,

$$\begin{aligned} & \mathcal{J}_{\ell,n}(hr^{1-d}t, D_{r,ha}) \\ &= (2\pi)^n r^{nd/2} \int_{\mathbb{R}^{n-1}} \int_{\mathbb{R}} \mathbb{1}(r^{d-2}(\mathbf{u}, \omega) \in A\mathcal{Q}) \\ & \quad \times \left| \int_{\mathbb{R}^d} \Theta_{f_*}^{(n-1)}(g_r(\mathbf{u}, \boldsymbol{\alpha}) z_{n-1}(r^d \boldsymbol{\eta}), r^{d-2} \mathbf{u}, \boldsymbol{\eta}, r^{d-2} \omega) d\boldsymbol{\eta} \right|^{1/2} d\omega d\mathbf{u} + O(r^\infty), \end{aligned} \quad (11-14)$$

with

$$g_r(\mathbf{u}, \boldsymbol{\alpha}) = \left(\mathbf{u} + ir^2 I_k, \mathbf{0}, \left(\begin{pmatrix} 0 \\ -\boldsymbol{\alpha} \end{pmatrix}, \dots, \begin{pmatrix} 0 \\ -\boldsymbol{\alpha} \end{pmatrix} \right) \right) \in G^{n-1},$$

and $\Theta_{f_*}^{(k)}$ as in (10-7) with $k = n - 1$ and test function

$$\begin{aligned} f_*(\mathbf{Y}, \mathbf{Y}', \mathbf{u}, \boldsymbol{\eta}, \omega) &= \mathcal{W}(y'_1 - y_1) \mathcal{W}(y'_\ell - y_\ell + y_{n-1} - y'_{n-1}) \\ & \quad \times \left(\prod_{j=1}^{n-2} \widehat{W}(y_j - y_{j+1}) \widehat{W}(y'_{j+1} - y'_j) \right) \tilde{a}(\boldsymbol{\eta}, y_\ell) \tilde{a}(-\boldsymbol{\eta}, y'_\ell) \\ & \quad \times \left(\prod_{j=\ell+1}^{n-1} e(-u_j(y_j - y'_j) \cdot \boldsymbol{\eta}) \right) e\left(\frac{1}{2}(\omega - u_\ell) \boldsymbol{\eta} \cdot (y_\ell - y'_\ell)\right), \end{aligned} \quad (11-15)$$

where $\mathbf{Y}, \mathbf{Y}' \in \mathbb{R}^{d \times (n-1)}$ are given by

$$\mathbf{Y} = (y_1, \dots, y_{n-1}), \quad \mathbf{Y}' = (y'_1, \dots, y'_{n-1}). \quad (11-16)$$

In order to apply the results in Section 10, we however require f_* to be continuous and compactly supported in \mathbf{u} , and rapidly decaying in ω . To achieve this, note that we can find f with precisely these properties by setting

$$f(\mathbf{Y}, \mathbf{Y}', \mathbf{u}, \boldsymbol{\eta}, \omega) = (\iota(\mathbf{u}, \omega))^2 f_*(\mathbf{Y}, \mathbf{Y}', \mathbf{u}, \boldsymbol{\eta}, \omega), \quad (11-17)$$

with $\iota : \mathbb{R}^n \rightarrow \mathbb{R}_{\geq 0}$ smooth and compactly supported such that $\iota(\mathbf{u}, \omega) > (2\pi)^n$ on the domain of integration. We then have, instead of (11-14),

$$\begin{aligned} & \mathcal{J}_{\ell,n}(hr^{1-d}t, D_{r,ha}) \\ & \leq r^{nd/2} \int_{\mathbb{R}^{n-1}} \int_{\mathbb{R}} \mathbb{1}(r^{d-2}(\mathbf{u}, \omega) \in A\mathcal{Q}) \\ & \quad \times \left| \int_{\mathbb{R}^d} \Theta_f^{(n-1)}(g_r(\mathbf{u}, \boldsymbol{\alpha}) z_{n-1}(r^d \boldsymbol{\eta}), r^{d-2} \mathbf{u}, \boldsymbol{\eta}, r^{d-2} \omega) d\boldsymbol{\eta} \right|^{1/2} d\omega d\mathbf{u} + O(r^\infty), \end{aligned} \quad (11-18)$$

and thus after the variable substitution $\omega \mapsto r^{2-d}\omega$,

$$\begin{aligned} & \mathcal{J}_{\ell,n}(hr^{1-d}t, D_{r,ha}) \\ & \leq r^{nd/2}r^{2-d} \int_{\mathbb{R}^{n-1}} w(r^{d-2}\mathbf{u}) \\ & \quad \times \int_{\mathbb{R}} \left| \int_{\mathbb{R}^d} \Theta_f^{(n-1)}(g_r(\mathbf{u}, \boldsymbol{\alpha}) z_{n-1}(r^d \boldsymbol{\eta}), r^{d-2}\mathbf{u}, \boldsymbol{\eta}, \omega) d\boldsymbol{\eta} \right|^{1/2} d\omega d\mathbf{u} + O(r^\infty), \end{aligned} \quad (11-19)$$

with

$$w(\mathbf{u}) = \sup_{\omega \in \mathbb{R}} \mathbb{1}((\mathbf{u}, \omega) \in A\mathcal{Q}), \quad (11-20)$$

which is bounded and has compact support.

Lemma 11.1. *Under the assumptions of Theorem 1.3, for $h = r < 1$,*

$$\mathcal{J}_{\ell,n}(hr^{1-d}t, D_{r,ha}) = O(r^{-nd/2+2n}). \quad (11-21)$$

Proof. For $F_r^{k,\beta}$ as in (10-9), we have

$$\int_{\mathbb{R}} \left| \int_{\mathbb{R}^d} \Theta_f^{(n-1)}(g z_{n-1}(r^d \boldsymbol{\eta}), r^{d-2}\mathbf{u}, \boldsymbol{\eta}, \omega) d\boldsymbol{\eta} \right|^{1/2} d\omega = F_r^{n-1,1/2}(g, r^{d-2}\mathbf{u}). \quad (11-22)$$

Thus, applying Proposition 10.4 we see that the right-hand side of (11-19) is bounded above by a constant times

$$r^{nd/2} \times r^{2-d} \times r^{-(n-1)(d-2)} = r^{-nd/2+2n}, \quad (11-23)$$

completing the proof. \square

Proof of Theorem 1.3. We recall the rescaling of t and λ in (2-9). The existence of the operators $A_n^{(r,\boldsymbol{\alpha})}(tr^{1-d})$ follows from the Duhamel expansion in (4-13). The error term follows from Lemmas 4.1 and 11.1, remembering that λ should be rescaled $\lambda \mapsto \lambda/h^2$ as in (2-9). Finally, the convergence of the operators $A_n^{(r,\boldsymbol{\alpha})}(tr^{1-d})$ in the limit $r \rightarrow 0$ is proved by combining Lemma 6.1, Lemma 6.2 and Corollary 9.5. \square

12. Averages over $\boldsymbol{\alpha}$

In this section we give the analogous results required to prove Theorem 1.2. First recall that Proposition 7.1 tells us that for $\mathbf{y} \in \mathbb{R}^d \setminus \mathbb{Q}^d$ with the components of $(1, {}^t\mathbf{y})$ linearly independent, and $(F_r)_{r \geq 0}$ a sequence of uniformly bounded, continuous functions we have

$$\lim_{r \rightarrow 0} r^\sigma \int_{\mathbb{R}} F_r \left(\left(u + ir^2, 0, \begin{pmatrix} \mathbf{0} \\ \mathbf{y} \end{pmatrix} \right), r^\sigma u \right) w(r^\sigma u) du = \frac{1}{\mu(\Gamma \setminus G)} \int_{\Gamma \setminus G} \int_{\mathbb{R}} F(g, u) w(u) du d\mu(g). \quad (12-1)$$

Note that since the F_r are uniformly bounded and continuous, and $w \in L^1(\mathbb{R})$, the integral over u is bounded uniformly in r and \mathbf{y} . Since the statement (12-1) holds for a full measure set of $\mathbf{y} \in [0, 1)^d$, one

can apply dominated convergence to conclude

$$\lim_{r \rightarrow 0} r^\sigma \int_{[0,1]^d} \int_{\mathbb{R}} F_r \left(\left(u + ir^2, 0, \begin{pmatrix} \mathbf{0} \\ \mathbf{y} \end{pmatrix} \right) \right), r^\sigma u) w(r^\sigma u) \, du \, d\mathbf{y} = \frac{1}{\mu(\Gamma \backslash G)} \int_{\Gamma \backslash G} \int_{\mathbb{R}} F(g, u) w(u) \, du \, d\mu(g). \quad (12-2)$$

Thus we now just need to consider the case of unbounded test functions. It follows from (7-15) that

$$\int_{\mathbf{y} \in [0,1]^d} \Psi_{R,f}^\beta \left(\tau, \begin{pmatrix} \mathbf{0} \\ \mathbf{y} \end{pmatrix} \right) \, d\mathbf{y} = 2 \sum_{\mathbf{m} \in \mathbb{Z}^d} f \left(\mathbf{m} \frac{v^{1/2}}{|\tau|} \right) \frac{v^{\beta d/2}}{|\tau|^{\beta d}} \chi_R \left(\frac{v}{|\tau|^2} \right) + 2 \int_{\mathbb{R}^d} f(\mathbf{y}) \, d\mathbf{y} \sum_{\substack{(c,d) \in \mathbb{Z}^2 \\ \gcd(c,d)=1 \\ c>0, d \neq 0}} \frac{v^{(\beta-1)d/2}}{|c\tau + d|^{(\beta-1)d}} \chi_R \left(\frac{v}{|c\tau + d|^2} \right). \quad (12-3)$$

Since for $0 \leq \beta \leq 1$

$$\frac{v^{(\beta-1)d/2}}{|c\tau + d|^{(\beta-1)d}} \leq R^{(\beta-1)d/2}, \quad (12-4)$$

we have

$$\int_{\mathbf{y} \in [0,1]^d} \Psi_{R,f}^\beta \left(\tau, \begin{pmatrix} \mathbf{0} \\ \mathbf{y} \end{pmatrix} \right) \, d\mathbf{y} \leq 2 \sum_{\mathbf{m} \in \mathbb{Z}^d} f \left(\mathbf{m} \frac{v^{1/2}}{|\tau|} \right) \frac{v^{\beta d/2}}{|\tau|^{\beta d}} \chi_R \left(\frac{v}{|\tau|^2} \right) + R^{(\beta-1)d/2} \chi_R(\tau) \int_{\mathbb{R}^d} f(\mathbf{y}) \, d\mathbf{y}. \quad (12-5)$$

This allows us to prove the following \mathbf{y} -averaged versions of Propositions 7.2 and 7.3.

Proposition 12.1. *Let $w : \mathbb{R} \rightarrow \mathbb{R}$ be piecewise continuous with compact support, and $0 < \epsilon < 1$. Then, for every $R \geq 1$,*

$$\limsup_{r \rightarrow 0} r^{d-2} \int_{|u| > r^{2-\epsilon}} \int_{[0,1]^d} \Psi_{R,f} \left(u + ir^2, \begin{pmatrix} \mathbf{0} \\ \mathbf{y} \end{pmatrix} \right) w(r^{d-2}u) \, d\mathbf{y} \, du \ll R^{-1}. \quad (12-6)$$

Proof. When $\beta = 1$, the first term in the right-hand side of (12-5) vanishes as $v \rightarrow 0$; see [Marklof 2002, §6.6.1]. By the equidistribution of closed horocycles and the fact that X_R is bounded and piecewise constant, we have for $R \geq 1$ that

$$\begin{aligned} \lim_{r \rightarrow 0} r^{d-2} \int_{\mathbb{R}} X_R(u + ir^2) w(r^{d-2}u) \, du &= \frac{3}{\pi} \int_{\mathbb{R}} w(x) \, dx \int_{\mathrm{SL}(2, \mathbb{R}) \backslash \mathfrak{H}} X_R(u + iv) \frac{du \, dv}{v^2} \\ &= \frac{3}{\pi} \int_{\mathbb{R}} w(x) \, dx \int_R^\infty \frac{dv}{v^2} = \frac{3}{\pi R} \int_{\mathbb{R}} w(x) \, dx, \end{aligned} \quad (12-7)$$

completing the proof. □

Proposition 12.2. *Let $w : \mathbb{R} \rightarrow \mathbb{R}$ be piecewise continuous with compact support, and $0 \leq \beta < 1$. Then, for every $R \geq 1$,*

$$\limsup_{r \rightarrow 0} r^{d-2} \int_{\mathbb{R}} \int_{[0,1]^d} \Psi_{R,f}^\beta \left(u + ir^2, \begin{pmatrix} \mathbf{0} \\ \mathbf{y} \end{pmatrix} \right) w(r^{d-2}u) \, d\mathbf{y} \, du \ll R^{(\beta-1)d/2}. \quad (12-8)$$

Proof. The first term in the right-hand side of (12-5) has already been estimated in the proof of Proposition 7.3. For the remaining terms the statement now follows from the observation that X_R is a bounded function. \square

Proof of Theorem 1.2. The convergence of the operators $A_n^{(r)}(r^{1-d}t)$ follows in the cases $n = 0, 1$ directly from the calculations in Section 6 for fixed α . Using Proposition 12.1 one can prove an α -averaged version of Proposition 8.9, and hence prove the convergence of $A_2^{(r)}(r^{1-d}t)$ as in Corollary 9.5, with $\mathbf{y} = -\alpha$. All that remains is the bound on the error terms. One first proves the α -averaged version of Proposition 10.4, with $\mathbf{y}_j = -\alpha$, by using Proposition 12.2. The remaining analysis proceeds identically to Section 11. \square

Appendix

A. The following proposition explains how Conjecture 1.1 and Theorem 1.2 yield information on the phase-space distribution of the wavepacket $f^{(\mathbf{p})}(t) = U_{h,\lambda}(t)f_0^{(\mathbf{p})}$ with an initial wavepacket $f_0^{(\mathbf{p})}$ of the form (see Figure 1)

$$f_0^{(\mathbf{p})}(\mathbf{x}) = r^{d(d-1)/2} \phi(r^{d-1}\mathbf{x}) e(\mathbf{p} \cdot \mathbf{x}/h), \quad (\text{A-1})$$

where $\phi \in \mathcal{S}(\mathbb{R}^d)$ is assumed to have unit L^2 -norm, and $\mathbf{p} \in \mathbb{R}^d$.

We use the shorthand

$$A(t) = U_{h,\lambda}(t) \text{Op}_{r,h}(a) U_{h,\lambda}(t)^{-1}, \quad B = \text{Op}_{r,h}(b). \quad (\text{A-2})$$

Proposition A.1. *Let $f_0^{(\mathbf{p})}$, $f^{(\mathbf{p})}(t)$ be as above, $w \in \mathcal{S}(\mathbb{R}^d)$ and $b \in \mathcal{S}(\mathbb{R}^d \times \mathbb{R}^d)$. Set*

$$a(\mathbf{x}, \mathbf{y}) = |\phi(\mathbf{x})|^2 w(\mathbf{y}). \quad (\text{A-3})$$

Then

$$r^{-d(d-1)/2} h^{-d/2} \int_{\mathbb{R}^d} \langle f^{(\mathbf{p})}(t), B f^{(\mathbf{p})}(t) \rangle w(\mathbf{p}) \, d\mathbf{p} = \langle A(t), B \rangle_{\text{HS}} + O(r^{d-1}h) \quad (\text{A-4})$$

uniformly in $r, h, t > 0$.

(The prefactor $r^{-d(d-1)/2} h^{-d/2}$ in (A-4) compensates for the L^2 -normalisation of $B = \text{Op}_{r,h}(b)$ in (1-9), which is not suitable in the present setting.)

Proof. Consider the linear operator $F_{r,h}^{(\mathbf{p})} : L^2(\mathbb{R}^d) \rightarrow L^2(\mathbb{R}^d)$ with Schwartz kernel

$$F_{r,h}^{(\mathbf{p})}(\mathbf{x}, \mathbf{x}') = f_0^{(\mathbf{p})}(\mathbf{x}) \overline{f_0^{(\mathbf{p})}(\mathbf{x}')} = r^{d(d-1)} \phi(r^{d-1}\mathbf{x}) \overline{\phi(r^{d-1}\mathbf{x}')} e(\mathbf{p} \cdot (\mathbf{x} - \mathbf{x}')/h). \quad (\text{A-5})$$

Using the Fourier transform \hat{w} of w yields

$$F_{r,h}(\mathbf{x}, \mathbf{x}') = \int F_{r,h}^{(\mathbf{p})}(\mathbf{x}, \mathbf{x}') w(\mathbf{p}) \, d\mathbf{p} = r^{d(d-1)} \phi(r^{d-1}\mathbf{x}) \overline{\phi(r^{d-1}\mathbf{x}')} \hat{w}((\mathbf{x}' - \mathbf{x})/h) \quad (\text{A-6})$$

and by Taylor's theorem we have

$$\phi(r^{d-1}\mathbf{x}) = \phi\left(\frac{1}{2}r^{d-1}(\mathbf{x} + \mathbf{x}')\right) + R_{r,h}(\mathbf{x}, \mathbf{x}'), \quad (\text{A-7})$$

with remainder

$$R_{r,h}(\mathbf{x}, \mathbf{x}') = \frac{1}{2} r^{d-1} \int_0^1 (\mathbf{x} - \mathbf{x}') \cdot \nabla \phi\left(\frac{1}{2} r^{d-1} ((\mathbf{x} + \mathbf{x}') + s(\mathbf{x} - \mathbf{x}'))\right) \, ds. \quad (\text{A-8})$$

We can express this term in the form

$$R_{r,h}(\mathbf{x}, \mathbf{x}') = \frac{1}{2}r^{d-1}h S_b\left(\frac{1}{2}r^{d-1}(\mathbf{x} + \mathbf{x}'), (\mathbf{x} - \mathbf{x}')/h\right), \quad b = \frac{1}{2}r^{d-1}h, \quad (\text{A-9})$$

with

$$S_b(\mathbf{x}, \mathbf{y}) = \int_0^1 \mathbf{y} \cdot \nabla \phi(\mathbf{x} + sb\mathbf{y}) \, ds. \quad (\text{A-10})$$

Now

$$F_{r,h}(\mathbf{x}, \mathbf{x}') = r^{d(d-1)} \left| \phi\left(\frac{1}{2}r^{d-1}(\mathbf{x} + \mathbf{x}')\right) \right|^2 \hat{w}((\mathbf{x}' - \mathbf{x})/h) + E_{r,h}(\mathbf{x}, \mathbf{x}'), \quad (\text{A-11})$$

with

$$\begin{aligned} E_{r,h}(\mathbf{x}, \mathbf{x}') = r^{d(d-1)} \hat{w}((\mathbf{x}' - \mathbf{x})/h) \{ & \phi\left(\frac{1}{2}r^{d-1}(\mathbf{x} + \mathbf{x}')\right) \overline{R_{r,h}(\mathbf{x}', \mathbf{x})} \\ & + R_{r,h}(\mathbf{x}, \mathbf{x}') \overline{\phi\left(\frac{1}{2}r^{d-1}(\mathbf{x} + \mathbf{x}')\right)} + R_{r,h}(\mathbf{x}, \mathbf{x}') \overline{R_{r,h}(\mathbf{x}', \mathbf{x})} \}. \end{aligned} \quad (\text{A-12})$$

On account of (A-9),

$$E_{r,h}(\mathbf{x}, \mathbf{x}') = r^{(d+1)(d-1)}h W_b\left(\frac{1}{2}r^{d-1}(\mathbf{x} + \mathbf{x}'), (\mathbf{x} - \mathbf{x}')/h\right), \quad b = \frac{1}{2}r^{d-1}h, \quad (\text{A-13})$$

with

$$W_b(\mathbf{x}, \mathbf{y}) = \frac{1}{2} \hat{w}(\mathbf{y}) \{ \phi(\mathbf{x}) \overline{S_b(\mathbf{x}, -\mathbf{y})} + S_b(\mathbf{x}, \mathbf{y}) \overline{\phi(\mathbf{x})} + b S_b(\mathbf{x}, \mathbf{y}) \overline{S_b(\mathbf{x}, -\mathbf{y})} \}. \quad (\text{A-14})$$

We rewrite (A-11) as

$$F_{r,h}(\mathbf{x}, \mathbf{x}') = r^{d(d-1)}h^d \int_{\mathbb{R}^d} \left| \phi\left(\frac{1}{2}r^{d-1}(\mathbf{x} + \mathbf{x}')\right) \right|^2 w(h\mathbf{y}) e(i(\mathbf{x} - \mathbf{x}') \cdot \mathbf{y}) \, d\mathbf{y} + E_{r,h}(\mathbf{x}, \mathbf{x}'), \quad (\text{A-15})$$

and so, for a as in (A-3),

$$F_{r,h} = r^{d(d-1)/2}h^{d/2} \text{Op}_{r,h}(a) + E_{r,h}. \quad (\text{A-16})$$

We conclude

$$\begin{aligned} r^{-d(d-1)/2}h^{-d/2} \int_{\mathbb{R}^d} \langle f^{(\mathbf{p})}(t), B f^{(\mathbf{p})}(t) \rangle w(\mathbf{p}) \, d\mathbf{p} \\ = r^{-d(d-1)/2}h^{-d/2} \langle U_{h,\lambda}(t) F_{r,h} U_{h,\lambda}(t)^{-1}, B \rangle_{\text{HS}} \\ = \langle U_{h,\lambda}(t) \text{Op}_{r,h}(a) U_{h,\lambda}(t)^{-1}, B \rangle_{\text{HS}} + O(r^{d-1}h), \end{aligned} \quad (\text{A-17})$$

where the error term follows from the upper bounds

$$|\langle U_{h,\lambda}(t) E_{r,h} U_{h,\lambda}(t)^{-1}, B \rangle_{\text{HS}}| \leq \|E_{r,h}\|_{\text{HS}} \|B\|_{\text{HS}}, \quad (\text{A-18})$$

and

$$\begin{aligned} \|E_{r,h}\|_{\text{HS}} &= r^{(d+1)(d-1)}h \left(\int_{\mathbb{R}^d \times \mathbb{R}^d} \left| W_b\left(\frac{1}{2}r^{d-1}(\mathbf{x} + \mathbf{x}'), (\mathbf{x} - \mathbf{x}')/h\right) \right|^2 \, d\mathbf{x} \, d\mathbf{x}' \right)^{1/2} \\ &= r^{(1+d/2)(d-1)}h^{d/2+1} \left(\int_{\mathbb{R}^d \times \mathbb{R}^d} |W_b(\mathbf{x}, \mathbf{y})|^2 \, d\mathbf{x} \, d\mathbf{y} \right)^{1/2}, \end{aligned} \quad (\text{A-19})$$

with

$$\lim_{b \rightarrow 0} \int_{\mathbb{R}^d \times \mathbb{R}^d} |W_b(\mathbf{x}, \mathbf{y})|^2 \, d\mathbf{x} \, d\mathbf{y} = \int_{\mathbb{R}^d \times \mathbb{R}^d} |W_0(\mathbf{x}, \mathbf{y})|^2 \, d\mathbf{x} \, d\mathbf{y} < \infty. \quad (\text{A-20})$$

This completes the proof. \square

B. In this section we compute the expression (5-10) for $\mathcal{I}_{\ell,n}$. Recall that

$$\begin{aligned} & [\widehat{K}_{\ell,n}(\mathbf{s})](\mathbf{y}, \mathbf{y}') \\ &= r^{(n-\ell+1)d} \sum_{\mathbf{m}_\ell, \dots, \mathbf{m}_n \in \mathbb{Z}^d} e\left(-\frac{1}{2} s_\ell \|\mathbf{y}\|^2\right) \widehat{W}(r\mathbf{m}_\ell) \mathcal{T}_{\ell,n-1}(\mathbf{y}) e\left(\frac{1}{2} s_n \|\mathbf{y} - \mathbf{m}_n\|^2\right) \delta_{\mathbf{m}_n}(\mathbf{y} - \mathbf{y}'). \end{aligned} \quad (\text{B-1})$$

Hence we have for $1 \leq \ell \leq n-1$ that

$$\begin{aligned} \mathcal{I}_{\ell,n}(\mathbf{s}) &= \text{Tr}_\alpha[K_{1,\ell}(\mathbf{s}) \text{Op}(D_{r,h}a) K_{\ell+1,n}(\mathbf{s}) \text{Op}(D_{r,h}b)] \\ &= r^{nd} r^{-d(d-1)} h^d \int_{\mathbb{R}^d} \sum_{\mathbf{m}_0, \dots, \mathbf{m}_n} e\left(-\frac{1}{2} s_1 \|\mathbf{m}_0 + \boldsymbol{\alpha}\|^2\right) \widehat{W}(r\mathbf{m}_1) \mathcal{T}_{1,\ell-1}(\mathbf{m}_0 + \boldsymbol{\alpha}) e\left(\frac{1}{2} s_\ell \|\mathbf{m}_0 + \boldsymbol{\alpha} - \mathbf{m}_\ell\|^2\right) \\ &\quad \times \tilde{a}\left(r^{1-d}(\mathbf{m}_0 + \boldsymbol{\alpha} - \mathbf{m}_\ell - \boldsymbol{\eta}), \frac{h}{2}(\mathbf{m}_0 + \boldsymbol{\alpha} - \mathbf{m}_\ell + \boldsymbol{\eta})\right) \\ &\quad \times e\left(-\frac{1}{2} s_{\ell+1} \|\boldsymbol{\eta}\|^2\right) \widehat{W}(r\mathbf{m}_{\ell+1}) \mathcal{T}_{\ell+1,n-1}(\boldsymbol{\eta}) e\left(\frac{1}{2} s_n \|\boldsymbol{\eta} - \mathbf{m}_n\|^2\right) \\ &\quad \times \tilde{b}\left(r^{1-d}(\boldsymbol{\eta} - \mathbf{m}_n - \mathbf{m}_0 - \boldsymbol{\alpha}), \frac{h}{2}(\boldsymbol{\eta} - \mathbf{m}_n + \mathbf{m}_0 + \boldsymbol{\alpha})\right) \, d\boldsymbol{\eta}. \end{aligned} \quad (\text{B-2})$$

We then make the variable substitution $\boldsymbol{\eta} \rightarrow r^{d-1}\boldsymbol{\eta} + \mathbf{m}_0 + \boldsymbol{\alpha} - \mathbf{m}_\ell$ so that \tilde{a} has first argument $-\boldsymbol{\eta}$. This leaves \tilde{b} with first argument $\boldsymbol{\eta} - r^{1-d}(\mathbf{m}_n + \mathbf{m}_\ell)$, and, by the rapid decay of \tilde{a} and \tilde{b} , the leading-order terms come from when $\mathbf{m}_n + \mathbf{m}_\ell = 0$, and we incur an error of order r^∞ . We thus have

$$\begin{aligned} \mathcal{I}_{\ell,n}(\mathbf{s}) &= r^{nd} h^d \int_{\mathbb{R}^d} \sum_{\mathbf{m}_0, \dots, \mathbf{m}_n} \mathbb{1}[\mathbf{m}_n + \mathbf{m}_\ell = 0] \\ &\quad \times e\left(-\frac{1}{2} s_1 \|\mathbf{m}_0 + \boldsymbol{\alpha}\|^2\right) \widehat{W}(r\mathbf{m}_1) \mathcal{T}_{1,\ell-1}(\mathbf{m}_0 + \boldsymbol{\alpha}) e\left(\frac{1}{2} s_\ell \|\mathbf{m}_0 + \boldsymbol{\alpha} - \mathbf{m}_\ell\|^2\right) \\ &\quad \times \tilde{a}\left(-\boldsymbol{\eta}, h(\mathbf{m}_0 + \boldsymbol{\alpha} - \mathbf{m}_\ell + \frac{1}{2} r^{d-1}\boldsymbol{\eta})\right) e\left(-\frac{1}{2} s_{\ell+1} \|\mathbf{m}_0 + \boldsymbol{\alpha} - \mathbf{m}_\ell + r^{d-1}\boldsymbol{\eta}\|^2\right) \\ &\quad \times \widehat{W}(r\mathbf{m}_{\ell+1}) \mathcal{T}_{\ell+1,n-1}(\mathbf{m}_0 + \boldsymbol{\alpha} + r^{d-1}\boldsymbol{\eta} - \mathbf{m}_\ell) e\left(\frac{1}{2} s_n \|\mathbf{m}_0 + \boldsymbol{\alpha} + r^{d-1}\boldsymbol{\eta}\|^2\right) \\ &\quad \times \tilde{b}\left(\boldsymbol{\eta}, h(\mathbf{m}_0 + \boldsymbol{\alpha} + \frac{1}{2} r^{d-1}\boldsymbol{\eta})\right) \, d\boldsymbol{\eta} + O(r^\infty). \end{aligned} \quad (\text{B-3})$$

Finally, we make the substitutions $\mathbf{m}_j \rightarrow \mathbf{m}_0 - \mathbf{m}_j$ for $j = 1, \dots, \ell$ followed by $\mathbf{m}_j \rightarrow \mathbf{m}_\ell - \mathbf{m}_j$ for $j = \ell + 1, \dots, n$ to obtain

$$\begin{aligned} \mathcal{I}_{\ell,n}(\mathbf{s}) &= r^{nd} h^d \int_{\mathbb{R}^d} \sum_{\mathbf{m}_0, \dots, \mathbf{m}_n} \mathbb{1}[\mathbf{m}_n = \mathbf{m}_0] \\ &\quad \times e\left(-\frac{1}{2} s_1 \|\mathbf{m}_0 + \boldsymbol{\alpha}\|^2\right) \widehat{W}(r(\mathbf{m}_0 - \mathbf{m}_1)) \mathcal{T}_{1,\ell-1}^-(\boldsymbol{\alpha}) e\left(\frac{1}{2} s_\ell \|\mathbf{m}_\ell + \boldsymbol{\alpha}\|^2\right) \\ &\quad \times \tilde{a}\left(-\boldsymbol{\eta}, h(\mathbf{m}_\ell + \boldsymbol{\alpha} + \frac{1}{2} r^{d-1}\boldsymbol{\eta})\right) e\left(-\frac{1}{2} s_{\ell+1} \|\mathbf{m}_\ell + \boldsymbol{\alpha} + r^{d-1}\boldsymbol{\eta}\|^2\right) \\ &\quad \times \widehat{W}(r(\mathbf{m}_\ell - \mathbf{m}_{\ell+1})) \mathcal{T}_{\ell+1,n-1}^-(\boldsymbol{\alpha} + r^{d-1}\boldsymbol{\eta}) e\left(\frac{1}{2} s_n \|\mathbf{m}_0 + \boldsymbol{\alpha} + r^{d-1}\boldsymbol{\eta}\|^2\right) \\ &\quad \times \tilde{b}\left(\boldsymbol{\eta}, h(\mathbf{m}_0 + \boldsymbol{\alpha} + \frac{1}{2} r^{d-1}\boldsymbol{\eta})\right) \, d\boldsymbol{\eta} + O(r^\infty). \end{aligned} \quad (\text{B-4})$$

This proves (5-10).

C. This section establishes relation (11-3), which is needed in the analysis of $\mathcal{J}_{\ell,n}(t, a)$. First we compute the kernel of $\widehat{K}_{\ell,n}^\dagger = \mathcal{F}K_{\ell,n}^\dagger\mathcal{F}^{-1}$. By taking the complex conjugate and switching \mathbf{y} and \mathbf{y}' in (4-22), we obtain

$$[\widehat{K}_{\ell,n}(\mathbf{s})^\dagger](\mathbf{y}, \mathbf{y}') = r^{(n-\ell+1)d} \sum_{\mathbf{m}'_\ell, \dots, \mathbf{m}'_n \in \mathbb{Z}^d} e\left(\frac{1}{2} s_\ell \|\mathbf{y} + \mathbf{m}'_n\|^2\right) \widehat{W}(-r\mathbf{m}'_\ell) \times \overline{\mathcal{T}}_{\ell,n-1}(\mathbf{y} + \mathbf{m}'_n) e\left(-\frac{1}{2} s_n \|\mathbf{y}\|^2\right) \delta_{\mathbf{m}'_n}(\mathbf{y}' - \mathbf{y}), \quad (\text{C-1})$$

where

$$\overline{\mathcal{T}}_{\ell,n}(\mathbf{y}) = \prod_{j=\ell}^n e\left(\frac{1}{2} (s_{j+1} - s_j) \|\mathbf{y} - \mathbf{m}'_j\|^2\right) \widehat{W}(r(\mathbf{m}'_j - \mathbf{m}'_{j+1})). \quad (\text{C-2})$$

Thus, using the formulae for the kernels of $\widehat{K}_{\ell,n}$, $\widehat{K}_{\ell,n}^\dagger$ and $\widehat{\text{Op}}_{r,h}$ we have

$$\begin{aligned} & [\widehat{K}_{\ell,n}(\mathbf{s})^\dagger \widehat{K}_{\ell,n}(\mathbf{s}) \widehat{\text{Op}}_{r,h}(a)](\mathbf{y}, \mathbf{y}') \\ &= r^{2(n-\ell+1)d} \sum_{\mathbf{m}_\ell, \dots, \mathbf{m}_n \in \mathbb{Z}^d} \sum_{\mathbf{m}'_\ell, \dots, \mathbf{m}'_n \in \mathbb{Z}^d} e\left(\frac{1}{2} s_\ell \|\mathbf{y} + \mathbf{m}'_n\|^2\right) \widehat{W}(-r\mathbf{m}'_\ell) \overline{\mathcal{T}}_{\ell,n-1}(\mathbf{y} + \mathbf{m}'_n) e\left(-\frac{1}{2} s_n \|\mathbf{y}\|^2\right) \\ & \quad \times e\left(-\frac{1}{2} s_\ell \|\mathbf{y} + \mathbf{m}'_n\|^2\right) \widehat{W}(r\mathbf{m}_\ell) \mathcal{T}_{\ell,n-1}(\mathbf{y} + \mathbf{m}'_n) e\left(\frac{1}{2} s_n \|\mathbf{y} + \mathbf{m}'_n - \mathbf{m}_n\|^2\right) \\ & \quad \times \tilde{a}\left(r^{1-d}(\mathbf{y} - \mathbf{m}_n + \mathbf{m}'_n - \mathbf{y}'), \frac{\hbar}{2}(\mathbf{y} - \mathbf{m}_n + \mathbf{m}'_n + \mathbf{y}')\right), \end{aligned} \quad (\text{C-3})$$

and similarly

$$\begin{aligned} & [\widehat{K}_{\ell,n}(\mathbf{s}) \widehat{K}_{\ell,n}(\mathbf{s})^\dagger \widehat{\text{Op}}_{r,h}(a)](\mathbf{y}, \mathbf{y}') \\ &= r^{2(n-\ell+1)d} \sum_{\mathbf{m}_\ell, \dots, \mathbf{m}_n \in \mathbb{Z}^d} \sum_{\mathbf{m}'_\ell, \dots, \mathbf{m}'_n \in \mathbb{Z}^d} e\left(-\frac{1}{2} s_\ell \|\mathbf{y}\|^2\right) \widehat{W}(r\mathbf{m}_\ell) \mathcal{T}_{\ell,n-1}(\mathbf{y}) e\left(\frac{1}{2} s_n \|\mathbf{y} - \mathbf{m}_n\|^2\right) \\ & \quad \times e\left(\frac{1}{2} s_\ell \|\mathbf{y} - \mathbf{m}_n + \mathbf{m}'_n\|^2\right) \widehat{W}(-r\mathbf{m}'_\ell) \\ & \quad \times \overline{\mathcal{T}}_{\ell,n-1}(\mathbf{y} - \mathbf{m}_n + \mathbf{m}'_n) e\left(-\frac{1}{2} s_n \|\mathbf{y} - \mathbf{m}_n\|^2\right) \\ & \quad \times \tilde{a}\left(r^{1-d}(\mathbf{y} + \mathbf{m}'_n - \mathbf{m}_n - \mathbf{y}'), \frac{\hbar}{2}(\mathbf{y} + \mathbf{m}'_n - \mathbf{m}_n + \mathbf{y}')\right). \end{aligned} \quad (\text{C-4})$$

Combining these yields explicitly

$$\begin{aligned} & \text{Tr}_\alpha [K_{1,\ell}(\mathbf{s})^\dagger K_{1,\ell}(\mathbf{s}) \text{Op}_{r,h}(a) K_{\ell+1,n}(\mathbf{s}) K_{\ell+1,n}(\mathbf{s})^\dagger \text{Op}_{r,h}(\bar{a})] \\ &= r^{2nd-d(d-1)} h^d \sum_{\substack{\mathbf{m}_0, \mathbf{m}_1, \dots, \mathbf{m}_n \in \mathbb{Z}^d \\ \mathbf{m}'_1, \dots, \mathbf{m}'_n \in \mathbb{Z}^d}} \int_{\mathbb{R}^d} \widehat{W}(r\mathbf{m}_1) \mathcal{T}_{1,\ell-1}(\mathbf{m}_0 + \mathbf{m}'_\ell + \boldsymbol{\alpha}) e\left(\frac{1}{2} s_\ell \|\mathbf{m}_0 + \mathbf{m}'_\ell - \mathbf{m}_\ell + \boldsymbol{\alpha}\|^2\right) \\ & \quad \times \widehat{W}(-r\mathbf{m}'_1) \overline{\mathcal{T}}_{1,\ell-1}(\mathbf{m}_0 + \mathbf{m}'_\ell + \boldsymbol{\alpha}) e\left(-\frac{1}{2} s_\ell \|\mathbf{m}_0 + \boldsymbol{\alpha}\|^2\right) \\ & \quad \times \tilde{a}\left(r^{1-d}(\mathbf{m}_0 + \mathbf{m}'_\ell - \mathbf{m}_\ell + \boldsymbol{\alpha} - \mathbf{y}), \frac{\hbar}{2}(\mathbf{m}_0 + \mathbf{m}'_\ell - \mathbf{m}_\ell + \boldsymbol{\alpha} + \mathbf{y})\right) \\ & \quad \times e\left(-\frac{1}{2} s_{\ell+1} \|\mathbf{y}\|^2\right) \widehat{W}(r\mathbf{m}_{\ell+1}) \mathcal{T}_{\ell+1,n-1}(\mathbf{y}) \\ & \quad \times e\left(\frac{1}{2} s_{\ell+1} \|\mathbf{y} + \mathbf{m}'_n - \mathbf{m}_n\|^2\right) \widehat{W}(-r\mathbf{m}'_{\ell+1}) \overline{\mathcal{T}}_{\ell+1,n-1}(\mathbf{y} + \mathbf{m}'_n - \mathbf{m}_n) \\ & \quad \times \tilde{a}\left(r^{1-d}(\mathbf{y} + \mathbf{m}'_n - \mathbf{m}_n - \mathbf{m}_0 - \boldsymbol{\alpha}), \frac{\hbar}{2}(\mathbf{y} + \mathbf{m}'_n - \mathbf{m}_n + \mathbf{m}_0 + \boldsymbol{\alpha})\right) d\mathbf{y}. \end{aligned} \quad (\text{C-5})$$

Now we make the substitution $\mathbf{y} = r^{d-1}\boldsymbol{\eta} + \mathbf{m}_0 + \boldsymbol{\alpha} + \mathbf{m}'_\ell - \mathbf{m}_\ell$ so that the first argument of \tilde{a} becomes $-\boldsymbol{\eta}$. Now \tilde{a} has first argument $\boldsymbol{\eta} + r^{1-d}(\mathbf{m}'_n - \mathbf{m}_n + \mathbf{m}'_\ell - \mathbf{m}_\ell)$, and hence (using the rapid decay of \tilde{a}) we have

that $\mathbf{m}'_n - \mathbf{m}_n + \mathbf{m}'_\ell - \mathbf{m}_\ell = 0$. This yields the expression

$$\begin{aligned}
& \text{Tr}_\alpha[K_{1,\ell}(\mathbf{s})^\dagger K_{1,\ell}(\mathbf{s}) \text{Op}_{r,h}(a) K_{\ell+1,n}(\mathbf{s}) K_{\ell+1,n}(\mathbf{s})^\dagger \text{Op}_{r,h}(\bar{a})] \\
&= r^{2nd} h^d \sum_{\substack{\mathbf{m}_0, \mathbf{m}_1, \dots, \mathbf{m}_n \in \mathbb{Z}^d \\ \mathbf{m}'_1, \dots, \mathbf{m}'_n \in \mathbb{Z}^d}} \mathbb{1}[\mathbf{m}'_n - \mathbf{m}_n + \mathbf{m}'_\ell - \mathbf{m}_\ell = 0] \\
&\quad \times \int_{\mathbb{R}^d} \widehat{W}(r\mathbf{m}_1) \mathcal{T}_{1,\ell-1}(\mathbf{m}_0 + \mathbf{m}'_\ell + \boldsymbol{\alpha}) e^{(\frac{1}{2} s_\ell \|\mathbf{m}_0 + \mathbf{m}'_\ell - \mathbf{m}_\ell + \boldsymbol{\alpha}\|^2)} \\
&\quad \times \widehat{W}(-r\mathbf{m}'_1) \bar{\mathcal{T}}_{1,\ell-1}(\mathbf{m}_0 + \mathbf{m}'_\ell + \boldsymbol{\alpha}) e^{(-\frac{1}{2} s_\ell \|\mathbf{m}_0 + \boldsymbol{\alpha}\|^2)} \\
&\quad \times \tilde{a}(-\boldsymbol{\eta}, h(\mathbf{m}_0 + \mathbf{m}'_\ell - \mathbf{m}_\ell + \boldsymbol{\alpha} + \frac{1}{2} r^{d-1} \boldsymbol{\eta})) \\
&\quad \times e^{(-\frac{1}{2} s_{\ell+1} \|r^{d-1} \boldsymbol{\eta} + \mathbf{m}_0 + \boldsymbol{\alpha} + \mathbf{m}'_\ell - \mathbf{m}_\ell\|^2)} \widehat{W}(r\mathbf{m}_{\ell+1}) \\
&\quad \times \mathcal{T}_{\ell+1,n-1}(r^{d-1} \boldsymbol{\eta} + \mathbf{m}_0 + \boldsymbol{\alpha} + \mathbf{m}'_\ell - \mathbf{m}_\ell) \\
&\quad \times e^{(\frac{1}{2} s_{\ell+1} \|r^{d-1} \boldsymbol{\eta} + \mathbf{m}_0 + \boldsymbol{\alpha}\|^2)} \widehat{W}(-r\mathbf{m}'_{\ell+1}) \\
&\quad \times \bar{\mathcal{T}}_{\ell+1,n-1}(r^{d-1} \boldsymbol{\eta} + \mathbf{m}_0 + \boldsymbol{\alpha}) \\
&\quad \times \tilde{\tilde{a}}(\boldsymbol{\eta}, h(\mathbf{m}_0 + \boldsymbol{\alpha} + \frac{1}{2} r^{d-1} \boldsymbol{\eta})) d\boldsymbol{\eta} + O(r^\infty). \tag{C-6}
\end{aligned}$$

We then make the substitution $\mathbf{m}_0 \rightarrow \mathbf{m}_0 - \mathbf{m}'_\ell$, followed by the substitutions $\mathbf{m}_j \rightarrow \mathbf{m}_0 - \mathbf{m}_j$ for $j = 1, \dots, \ell$ and $\mathbf{m}_j \rightarrow \mathbf{m}_\ell - \mathbf{m}_j$ for $j = \ell + 1, \dots, n$ as well as the analogous substitutions for the \mathbf{m}'_j . This yields the simpler expression

$$\begin{aligned}
& \text{Tr}_\alpha[K_{1,\ell}(\mathbf{s})^\dagger K_{1,\ell}(\mathbf{s}) \text{Op}_{r,h}(a) K_{\ell+1,n}(\mathbf{s}) K_{\ell+1,n}(\mathbf{s})^\dagger \text{Op}_{r,h}(\bar{a})] \\
&= r^{2nd} h^d \sum_{\substack{\mathbf{m}_0, \mathbf{m}_1, \dots, \mathbf{m}_n \in \mathbb{Z}^d \\ \mathbf{m}'_1, \dots, \mathbf{m}'_n \in \mathbb{Z}^d}} \mathbb{1}[\mathbf{m}'_n - \mathbf{m}_n + \mathbf{m}_\ell - \mathbf{m}'_\ell = 0] \\
&\quad \times \int_{\mathbb{R}^d} \widehat{W}(r(\mathbf{m}_0 - \mathbf{m}_1)) \mathcal{T}_{1,\ell-1}^-(\boldsymbol{\alpha}) e^{(\frac{1}{2} s_\ell (\|\mathbf{m}_\ell + \boldsymbol{\alpha}\|^2 - \|\mathbf{m}'_\ell + \boldsymbol{\alpha}\|^2))} \\
&\quad \times \widehat{W}(r(\mathbf{m}'_1 - \mathbf{m}_0)) \bar{\mathcal{T}}_{1,\ell-1}^-(\boldsymbol{\alpha}) \tilde{a}(-\boldsymbol{\eta}, h(\mathbf{m}_\ell + \boldsymbol{\alpha} + \frac{1}{2} r^{d-1} \boldsymbol{\eta})) \\
&\quad \times \widehat{W}(r(\mathbf{m}_\ell - \mathbf{m}_{\ell+1})) \mathcal{T}_{\ell+1,n-1}^-(r^{d-1} \boldsymbol{\eta} + \boldsymbol{\alpha}) \\
&\quad \times e^{(\frac{1}{2} s_{\ell+1} (\|r^{d-1} \boldsymbol{\eta} + \mathbf{m}'_\ell + \boldsymbol{\alpha}\|^2 - \|r^{d-1} \boldsymbol{\eta} + \boldsymbol{\alpha} + \mathbf{m}_\ell\|^2))} \\
&\quad \times \widehat{W}(r(\mathbf{m}'_{\ell+1} - \mathbf{m}'_\ell)) \bar{\mathcal{T}}_{\ell+1,n-1}^-(r^{d-1} \boldsymbol{\eta} + \boldsymbol{\alpha}) \\
&\quad \times \tilde{\tilde{a}}(\boldsymbol{\eta}, h(\mathbf{m}'_\ell + \boldsymbol{\alpha} + \frac{1}{2} r^{d-1} \boldsymbol{\eta})) d\boldsymbol{\eta} + O(r^\infty). \tag{C-7}
\end{aligned}$$

This yields (11-3) after substituting $\boldsymbol{\eta} \rightarrow -\boldsymbol{\eta}$.

Acknowledgements

We thank László Erdős and Leonid Parnowski for helpful discussions, and the anonymous referee for many valuable comments. We are grateful to the Isaac Newton Institute, Cambridge, for its support and hospitality during the programme ‘‘Periodic and Ergodic Spectral Problems’’. The research leading to these results has received funding from the European Research Council under the European Union’s Seventh Framework Programme (FP/2007-2013) / ERC Grant Agreement no. 291147.

References

- [Allaire and Piatnitski 2005] G. Allaire and A. Piatnitski, “Homogenization of the Schrödinger equation and effective mass theorems”, *Comm. Math. Phys.* **258**:1 (2005), 1–22. MR Zbl
- [Bal et al. 1999] G. Bal, A. Fannjiang, G. Papanicolaou, and L. Ryzhik, “Radiative transport in a periodic structure”, *J. Stat. Phys.* **95**:1-2 (1999), 479–494. MR Zbl
- [Bal et al. 2002] G. Bal, G. Papanicolaou, and L. Ryzhik, “Radiative transport limit for the random Schrödinger equation”, *Nonlinearity* **15**:2 (2002), 513–529. MR Zbl
- [Bal et al. 2011] G. Bal, T. Komorowski, and L. Ryzhik, “Asymptotics of the solutions of the random Schrödinger equation”, *Arch. Ration. Mech. Anal.* **200**:2 (2011), 613–664. MR Zbl
- [Benoit and Gloria 2017] A. Benoit and A. Gloria, “Long-time homogenization and asymptotic ballistic transport of classical waves”, preprint, 2017. To appear in *Ann. Sci. École Norm. Sup.* arXiv
- [Bensoussan et al. 1978] A. Bensoussan, J.-L. Lions, and G. Papanicolaou, *Asymptotic analysis for periodic structures*, Stud. Math. Appl. **5**, North-Holland, Amsterdam, 1978. MR Zbl
- [Berry 1981] M. V. Berry, “Quantizing a classically ergodic system: Sinai’s billiard and the KKR method”, *Ann. Physics* **131**:1 (1981), 163–216. MR
- [Birman and Suslina 2003] M. S. Birman and T. A. Suslina, “Periodic second-order differential operators: threshold properties and averaging”, *Algebra i Analiz* **15**:5 (2003), 1–108. In Russian; translated in *St. Petersburg Math. J.* **15**:5 (2004), 639–714. MR
- [Boldrighini et al. 1983] C. Boldrighini, L. A. Bunimovich, and Y. G. Sinai, “On the Boltzmann equation for the Lorentz gas”, *J. Stat. Phys.* **32**:3 (1983), 477–501. MR Zbl
- [Caglioti and Golse 2010] E. Caglioti and F. Golse, “On the Boltzmann–Grad limit for the two dimensional periodic Lorentz gas”, *J. Stat. Phys.* **141**:2 (2010), 264–317. MR Zbl
- [Castella 1999] F. Castella, “On the derivation of a quantum Boltzmann equation from the periodic von Neumann equation”, *M2AN Math. Model. Numer. Anal.* **33**:2 (1999), 329–349. MR Zbl
- [Castella 2001] F. Castella, “From the von Neumann equation to the quantum Boltzmann equation in a deterministic framework”, *J. Stat. Phys.* **104**:1-2 (2001), 387–447. MR Zbl
- [Castella 2002] F. Castella, “From the von Neumann equation to the quantum Boltzmann equation, II: Identifying the Born series”, *J. Stat. Phys.* **106**:5-6 (2002), 1197–1220. MR Zbl
- [Castella and Plagne 2002] F. Castella and A. Plagne, “Non derivation of the quantum Boltzmann equation from the periodic von Neumann equation”, *Indiana Univ. Math. J.* **51**:4 (2002), 963–1016. MR Zbl
- [Craster et al. 2010] R. V. Craster, J. Kaplunov, and A. V. Pichugin, “High-frequency homogenization for periodic media”, *Proc. R. Soc. Lond. Ser. A Math. Phys. Eng. Sci.* **466**:2120 (2010), 2341–2362. MR Zbl
- [Dahlqvist and Vattay 1998] P. Dahlqvist and G. Vattay, “Periodic orbit quantization of the Sinai billiard in the small scatterer limit”, *J. Phys. A* **31**:30 (1998), 6333–6345. MR Zbl
- [Eng and Erdős 2005] D. Eng and L. Erdős, “The linear Boltzmann equation as the low density limit of a random Schrödinger equation”, *Rev. Math. Phys.* **17**:6 (2005), 669–743. MR Zbl
- [Erdős and Yau 2000] L. Erdős and H.-T. Yau, “Linear Boltzmann equation as the weak coupling limit of a random Schrödinger equation”, *Comm. Pure Appl. Math.* **53**:6 (2000), 667–735. MR Zbl
- [Eskin et al. 2005] A. Eskin, G. Margulis, and S. Mozes, “Quadratic forms of signature (2, 2) and eigenvalue spacings on rectangular 2-tori”, *Ann. of Math. (2)* **161**:2 (2005), 679–725. MR Zbl
- [Folland 1989] G. B. Folland, *Harmonic analysis in phase space*, Ann. of Math. Studies **122**, Princeton Univ. Press, 1989. MR Zbl
- [Gallavotti 1969] G. Gallavotti, “Divergences and the approach to equilibrium in the Lorentz and the wind-tree models”, *Phys. Rev.* **185**:1 (1969), 308–322.
- [Gérard 1991] P. Gérard, “Mesures semi-classiques et ondes de Bloch”, exposé 16 in *Séminaire sur les Équations aux Dérivées Partielles: 1990–1991*, École Polytech., Paris, 1991. MR Zbl

- [Gérard et al. 1997] P. Gérard, P. A. Markowich, N. J. Mauser, and F. Poupaud, “Homogenization limits and Wigner transforms”, *Comm. Pure Appl. Math.* **50**:4 (1997), 323–379. MR Zbl
- [Golse 2008] F. Golse, “On the periodic Lorentz gas and the Lorentz kinetic equation”, *Ann. Fac. Sci. Toulouse Math.* (6) **17**:4 (2008), 735–749. MR Zbl
- [Griffin 2017] J. Griffin, *Quantum dynamics in highly localised periodic potentials*, Ph.D. thesis, University of Bristol, 2017.
- [Griffin and Marklof \geq 2019] J. Griffin and J. Marklof, “Probabilistic model for quantum transport in a low-density periodic potential”, in preparation.
- [Harutyunyan et al. 2016] D. Harutyunyan, G. W. Milton, and R. V. Craster, “High-frequency homogenization for travelling waves in periodic media”, *Proc. A.* **472**:2191 (2016), art. id. 20160066. MR Zbl
- [Lorentz 1905] H. A. Lorentz, “Le mouvement des électrons dans les métaux”, *Arch. Néerl.* **10** (1905), 337–371. Zbl
- [Margulis and Mohammadi 2011] G. Margulis and A. Mohammadi, “Quantitative version of the Oppenheim conjecture for inhomogeneous quadratic forms”, *Duke Math. J.* **158**:1 (2011), 121–160. MR Zbl
- [Marklof 2002] J. Marklof, “Pair correlation densities of inhomogeneous quadratic forms, II”, *Duke Math. J.* **115**:3 (2002), 409–434. Correction in **120**:1 (2003), 227–228. MR Zbl
- [Marklof 2003] J. Marklof, “Pair correlation densities of inhomogeneous quadratic forms”, *Ann. of Math.* (2) **158**:2 (2003), 419–471. MR Zbl
- [Marklof and Strömbergsson 2008] J. Marklof and A. Strömbergsson, “Kinetic transport in the two-dimensional periodic Lorentz gas”, *Nonlinearity* **21**:7 (2008), 1413–1422. MR Zbl
- [Marklof and Strömbergsson 2010] J. Marklof and A. Strömbergsson, “The distribution of free path lengths in the periodic Lorentz gas and related lattice point problems”, *Ann. of Math.* (2) **172**:3 (2010), 1949–2033. MR Zbl
- [Marklof and Strömbergsson 2011a] J. Marklof and A. Strömbergsson, “The Boltzmann–Grad limit of the periodic Lorentz gas”, *Ann. of Math.* (2) **174**:1 (2011), 225–298. MR Zbl
- [Marklof and Strömbergsson 2011b] J. Marklof and A. Strömbergsson, “The periodic Lorentz gas in the Boltzmann–Grad limit: asymptotic estimates”, *Geom. Funct. Anal.* **21**:3 (2011), 560–647. MR Zbl
- [Markowich et al. 1994] P. A. Markowich, N. J. Mauser, and F. Poupaud, “A Wigner-function approach to (semi)classical limits: electrons in a periodic potential”, *J. Math. Phys.* **35**:3 (1994), 1066–1094. MR Zbl
- [Nier 1995] F. Nier, “Asymptotic analysis of a scaled Wigner equation and quantum scattering”, *Transport Theory Statist. Phys.* **24**:4-5 (1995), 591–628. MR Zbl
- [Nier 1996] F. Nier, “A semi-classical picture of quantum scattering”, *Ann. Sci. École Norm. Sup.* (4) **29**:2 (1996), 149–183. MR Zbl
- [Panati et al. 2003] G. Panati, H. Spohn, and S. Teufel, “Effective dynamics for Bloch electrons: Peierls substitution and beyond”, *Comm. Math. Phys.* **242**:3 (2003), 547–578. MR Zbl
- [Spohn 1977] H. Spohn, “Derivation of the transport equation for electrons moving through random impurities”, *J. Stat. Phys.* **17**:6 (1977), 385–412. MR Zbl
- [Spohn 1978] H. Spohn, “The Lorentz process converges to a random flight process”, *Comm. Math. Phys.* **60**:3 (1978), 277–290. MR Zbl
- [Strömbergsson and Vishe 2018] A. Strömbergsson and P. Vishe, “An effective equidistribution result for $SL(2, \mathbb{R}) \times (\mathbb{R}^2)^{\oplus k}$ and application to inhomogeneous quadratic forms”, preprint, 2018. arXiv

Received 2 Nov 2018. Revised 20 Mar 2019. Accepted 6 Jun 2019.

JORY GRIFFIN: j.griffin@queensu.ca

Department of Mathematics and Statistics, Queen’s University, Kingston, ON, Canada

JENS MARKLOF: j.marklof@bristol.ac.uk

School of Mathematics, University of Bristol, Bristol, United Kingdom

PURE and APPLIED ANALYSIS

msp.org/paa

EDITORS-IN-CHIEF

- Charles L. Epstein University of Pennsylvania
cle@math.upenn.edu
- Maciej Zworski University of California at Berkeley
zworski@math.berkeley.edu

EDITORIAL BOARD

- Sir John M. Ball University of Oxford
ball@maths.ox.ac.uk
- Michael P. Brenner Harvard University
brenner@seas.harvard.edu
- Charles Fefferman Princeton University
cf@math.princeton.edu
- Susan Friedlander University of Southern California
susanfri@usc.edu
- Anna Gilbert University of Michigan
annacg@umich.edu
- Leslie F. Greengard Courant Institute, New York University, and
Flatiron Institute, Simons Foundation
greengard@cims.nyu.edu
- Yan Guo Brown University
yan_guo@brown.edu
- Claude Le Bris CERMICS - ENPC
lebris@cermics.enpc.fr
- Robert J. McCann University of Toronto
mccann@math.toronto.edu
- Michael O'Neil Courant Institute, New York University
oneil@cims.nyu.edu
- Jill Pipher Brown University
jill_pipher@brown.edu
- Johannes Sjöstrand Université de Dijon
johannes.sjostrand@u-bourgogne.fr
- Vladimir Šverák University of Minnesota
sverak@math.umn.edu
- Daniel Tataru University of California at Berkeley
tataru@berkeley.edu
- Michael I. Weinstein Columbia University
miw2103@columbia.edu
- Jon Wilkening University of California at Berkeley
wilken@math.berkeley.edu
- Enrique Zuazua DeustoTech-Bilbao, and
Universidad Autónoma de Madrid
enrique.zuazua@deusto.es

PRODUCTION

- Silvio Levy (Scientific Editor)
production@msp.org

Cover image: The figure shows the outgoing scattered field produced by scattering a plane wave, coming from the northwest, off of the (stylized) letters P A A. The total field satisfies the homogeneous Dirichlet condition on the boundary of the letters. It is based on a numerical computation by Mike O'Neil of the Courant Institute.


See inside back cover or msp.org/paa for submission instructions.

The subscription price for 2019 is US \$495/year for the electronic version, and \$555/year (+\$25, if shipping outside the US) for print and electronic. Subscriptions, requests for back issues and changes of subscriber address should be sent to MSP.

Pure and Applied Analysis (ISSN 2578-5885 electronic, 2578-5893 printed) at Mathematical Sciences Publishers, 798 Evans Hall #3840, c/o University of California, Berkeley, CA 94720-3840 is published continuously online. Periodical rate postage paid at Berkeley, CA 94704, and additional mailing offices.

PAA peer review and production are managed by EditFlow® from MSP.

PUBLISHED BY

 **mathematical sciences publishers**
nonprofit scientific publishing

<http://msp.org/>

© 2019 Mathematical Sciences Publishers

PURE and APPLIED ANALYSIS

vol. 1 no. 4 2019

- Connecting dots: from local covariance to empirical intrinsic geometry and locally linear embedding 515
JOHN MALIK, CHAO SHEN, HAU-TIENG WU and NAN WU
- On geometric and analytic mixing scales: comparability and convergence rates for transport problems 543
CHRISTIAN ZILLINGER
- Quantum transport in a low-density periodic potential: homogenisation via homogeneous flows 571
JORY GRIFFIN and JENS MARKLOF
- Cusp universality for random matrices, II: The real symmetric case 615
GIORGIO CIPOLLONI, LÁSZLÓ ERDŐS, TORBEN KRÜGER and DOMINIK SCHRÖDER
- Explicit unconditionally stable methods for the heat equation via potential theory 709
ALEX BARNETT, CHARLES L. EPSTEIN, LESLIE GREENGARD, SHIDONG JIANG and JUN WANG



2578-5893 (2019)1:4;1-3