

PURE and APPLIED ANALYSIS

PAM

WOLFGANG ARENDT AND DANIEL HAUER

MAXIMAL L^2 -REGULARITY IN NONLINEAR GRADIENT
SYSTEMS
AND PERTURBATIONS OF SUBLINEAR GROWTH



vol. 2 no. 1 2020

MAXIMAL L^2 -REGULARITY IN NONLINEAR GRADIENT SYSTEMS AND PERTURBATIONS OF SUBLINEAR GROWTH

WOLFGANG ARENDT AND DANIEL HAUER

The nonlinear semigroup generated by the subdifferential of a convex lower semicontinuous function φ has a smoothing effect, discovered by Haïm Brezis, which implies maximal regularity for the evolution equation. We use this and Schaefer's fixed point theorem to solve the evolution equation perturbed by a Nemyskii operator of sublinear growth. For this, we need that the sublevel sets of φ are not only closed, but even compact. We apply our results to the p -Laplacian and also to the Dirichlet-to-Neumann operator with respect to p -harmonic functions.

1. Introduction

Let H be a real Hilbert space, $\varphi : H \rightarrow (-\infty, +\infty]$ a proper, convex, lower semicontinuous function, $A = \partial\varphi$ the subdifferential of φ , and $D(\varphi) := \{u \in H \mid \varphi(u) < +\infty\}$ the effective domain of φ (see Section 2 for more details). Then A is a maximal monotone (in general, multivalued) operator on H for which the following remarkable well-posedness result holds.

Theorem 1.1 [Brezis 1971]. *Let $u_0 \in \overline{D(\varphi)}$ and $f \in L^2(0, T; H)$. Then, there exists a unique $u \in H_{\text{loc}}^1((0, T]; H) \cap C([0, T]; H)$ such that*

$$\begin{cases} \dot{u}(t) + Au(t) \ni f(t) & \text{a.e. on } (0, T), \\ u(0) = u_0. \end{cases} \quad (1-1)$$

If $u \in D(\varphi)$ then $\dot{u} \in L^2(0, T; H)$.

Our aim in this article is to establish existence of solutions of a perturbed version of (1-1) and to show that these solutions have the same regularity result as in Theorem 1.1. We fix $T > 0$, and denote by \mathcal{H} the space $L^2(0, T; H)$ and by $\|\cdot\|_{\mathcal{H}}$ the norm $\|\cdot\|_{L^2(0, T; H)}$. Then for $f \in \mathcal{H}$ and $u_0 \in H$, we call a function $u : [0, T] \rightarrow H$ a (strong) solution of (1-1) if $u \in H_{\text{loc}}^1((0, T]; H) \cap C([0, T]; H)$, $u(0) = u_0$ and for a.e. $t \in (0, T)$ we have $u(t) \in D(A)$ and $f(t) - \dot{u}(t) \in Au(t)$.

Now, let $G : \mathcal{H} \rightarrow \mathcal{H}$ be a continuous mapping satisfying the *sublinear* growth condition

$$\|Gv(t)\|_H \leq L \|v(t)\|_H + b(t) \quad \text{a.e. on } (0, T) \text{ and for all } v \in \mathcal{H}, \quad (1-2)$$

Hauer is very grateful for the warm hospitality received during his visits at the University of Ulm.

MSC2010: 35K92, 35K58, 47H20, 47H10.

Keywords: nonlinear semigroups, subdifferential, Schaefer's fixed point theorem, existence, smoothing effect, perturbation, compact sublevel sets.

for some $L, b \in L^2(0, T)$ satisfying $b(t) \geq 0$ for a.e. $t \in (0, T)$. Here we let $Gv(t) := (G(v))(t)$ to use less heavy notation. Then we study the evolution problem

$$\begin{cases} \dot{u}(t) + Au(t) \ni Gu(t) & \text{a.e. on } (0, T), \\ u(0) = u_0. \end{cases} \quad (1-3)$$

Note that $Gu \in \mathcal{H}$. Thus, the inclusion in (1-3) means that $Gu(t) - \dot{u}(t) \in Au(t)$ a.e. on $(0, T)$.

For proving existence of solutions to (1-3), we will use a compactness argument in form of Schaefer's fixed point theorem (see Theorem 2.1 in Section 2). Recall that lower semicontinuity of φ is equivalent to saying that the sublevel sets $E_c := \{u \in H \mid \varphi(u) \leq c\}$, $c \in \mathbb{R}$, are closed. We will assume more, namely, compactness of the sublevel sets E_c . In fact, we need this assumption only for the shifted function φ_ω given by $\varphi_\omega(u) = \varphi(u) + \frac{1}{2}\omega\|u\|_H^2$, $u \in H$, which is important for applications. Then our main result says the following.

Theorem 1.2. *Let $\varphi : H \rightarrow (-\infty, +\infty]$ be a proper function such that for some $\omega \geq 0$, φ_ω is convex and has compact sublevel sets. Let $A = \partial\varphi$ and $G : \mathcal{H} \rightarrow \mathcal{H}$ be a continuous mapping satisfying (1-2). Then for every $u_0 \in \overline{D(\varphi)}$ and $f \in \mathcal{H}$, there exists $u \in H_{\text{loc}}^1((0, T]; H) \cap C([0, T]; H)$ solving (1-3). In particular, if $u_0 \in D(\varphi)$, then $u \in H^1(0, T; H)$.*

We show in Example 3.3 that the solution is not unique in general. Further, we have the following regularity result for the composition $\varphi \circ u$ and a uniform estimate.

Remark 1.3. Suppose, the hypotheses of Theorem 1.2 hold. Then every solution u of (1-3) satisfies

$$\varphi \circ u \in W_{\text{loc}}^{1,1}((0, T]) \cap L^1(0, T)$$

and

$$\|u(t)\|_H \leq (\|u_0\|_H^2 + \|b\|_{L^2(0,T)}^2)^{1/2} e^{(2L+1+2\omega)/2t} \quad \text{for all } t \in [0, T]. \quad (1-4)$$

As application, we consider $H = L^2(\Omega)$ and G a Nemytskii operator. The operator A may be the p -Laplacian ($1 \leq p < +\infty$) with possibly lower-order terms and equipped with some boundary conditions (Dirichlet, Neumann, or Robin, see [Coulhon and Hauer 2016]) or a p -version of the Dirichlet-to-Neumann operator considered recently in [Hauer 2015] and via the abstract theory of j -elliptic functions (see [Arendt and ter Elst 2011; 2012; Chill et al. 2016]).

2. Preliminaries

In this section, we define the precise setting used throughout this paper and explain our main tools: Schaefer's fixed point theorem and Brezis' L^2 -maximal regularity result for semiconvex functions.

We begin by recalling that a mapping \mathcal{T} defined on a Banach space X is called *compact* if \mathcal{T} maps bounded sets into relatively compact sets.

Theorem 2.1 (Schaefer's fixed point theorem [1955]). *Let X be a Banach space and $\mathcal{T} : X \rightarrow X$ be continuous and compact. Assume that the "Schaefer set"*

$$\mathcal{S} := \{u \in X \mid \text{there exists } \lambda \in [0, 1] \text{ such that } u = \lambda\mathcal{T}u\}$$

is bounded in X . Then \mathcal{T} has a fixed point.

This result is a special case of *Leray–Schauder* degree theory, but Schaefer [1955] gave a most elegant proof, which also is valid in locally convex spaces; see also [Arendt and Chill 2010; Evans 2010, §9.2.2].

Given a function $\varphi : H \rightarrow (-\infty, +\infty]$, we call the set $D(\varphi) := \{u \in H \mid \varphi(u) < +\infty\}$ the *effective domain* of φ , and φ is said to be *proper* if $D(\varphi)$ is nonempty. Further, we say that φ is *lower semicontinuous* if for every $c \in \mathbb{R}$ the sublevel set

$$E_c := \{u \in D(\varphi) \mid \varphi(u) \leq c\}$$

is closed in H , and φ is *semiconvex* if there exists an $\omega \in \mathbb{R}$ such that the shifted function $\varphi_\omega : H \rightarrow (-\infty, +\infty]$ defined by

$$\varphi_\omega(u) := \varphi(u) + \frac{1}{2}\omega\|u\|_H^2, \quad u \in H,$$

is convex. Then, $\varphi_{\hat{\omega}}$ is convex for all $\hat{\omega} \geq \omega$, and φ_ω is lower semicontinuous if and only if φ is lower semicontinuous.

Given a function $\varphi : H \rightarrow (-\infty, +\infty]$, its *subdifferential* $A = \partial\varphi$ is defined by

$$\partial\varphi = \left\{ (u, h) \in H \times H \mid \liminf_{t \downarrow 0} \frac{\varphi(u + tv) - \varphi(u)}{t} \geq (h, v)_H \text{ for all } v \in D(\varphi) \right\},$$

which, if φ_ω is convex, reduces to

$$\partial\varphi = \{(u, h) \in H \times H \mid \varphi_\omega(u + v) - \varphi_\omega(u) \geq (h + \omega u, v)_H \text{ for all } v \in D(\varphi)\}.$$

It is standard to identify a (possibly multivalued) operator A on H with its graph and for every $u \in H$, one sets $Au := \{v \in H \mid (u, v) \in A\}$ and calls $D(A) := \{u \in H \mid Au \neq \emptyset\}$ the *domain* of A and $\text{Rg}(A) := \bigcup_{u \in D(A)} Au$ the *range* of A .

Now, suppose $\varphi : H \rightarrow (-\infty, +\infty]$ is proper, lower semicontinuous, and semiconvex; more precisely, let us fix $\omega \in \mathbb{R}$ such that φ_ω is convex. Then the subdifferential $\partial\varphi_\omega$ of φ_ω is a simple perturbation of $\partial\varphi$, namely $\partial\varphi_\omega = \partial\varphi + \omega I$. For this reason, Brezis' well-posedness result (Theorem 1.1) remains true; see [Brezis 1973, Proposition 3.12]. In addition, it is not difficult to verify that each solution of (1-1) satisfies (2-2) and the estimates (2-3)–(2-6) below. For later use, we summarize these results in one theorem.

Theorem 2.2 (Brezis' L^2 -maximal regularity for semiconvex φ). *Let $u_0 \in \overline{D(\varphi)}$ and $f \in \mathcal{H}$. Then, there exists a unique $u \in H_{\text{loc}}^1((0, T]; H) \cap C([0, T]; H)$ satisfying*

$$\begin{cases} \dot{u}(t) + Au(t) \ni f(t) & \text{a.e. on } (0, T), \\ u(0) = u_0. \end{cases} \quad (2-1)$$

Moreover,

$$\varphi \circ u \in W_{\text{loc}}^{1,1}((0, T]) \cap L^1(0, T), \quad (2-2)$$

$$\|u(t)\|_H \leq \left(\|u_0\|_H^2 + \int_0^t \|f(s)\|_H^2 ds \right)^{1/2} e^{(1+2\omega)/2t} \quad \text{for every } t \in (0, T], \quad (2-3)$$

$$\int_0^t \varphi(u(s)) ds \leq \frac{1}{2} \|f\|_{\mathcal{H}}^2 + \frac{1}{2}(1 + \omega) \|u\|_{\mathcal{H}}^2 + \frac{1}{2} \|u_0\|_H^2, \quad (2-4)$$

$$t\varphi(u(t)) \leq \int_0^t \varphi(u(s)) ds + \frac{1}{2} \|\sqrt{\cdot} \cdot f\|_{\mathcal{H}}^2 \quad \text{for every } t \in (0, T], \quad (2-5)$$

$$\|\sqrt{\cdot} \dot{u}\|_{\mathcal{H}}^2 \leq 2 \int_0^T \varphi(u(t)) \, dt + \|\sqrt{\cdot} f\|_{\mathcal{H}}^2. \quad (2-6)$$

Finally, if $u_0 \in D(\varphi)$, then $u \in H^1(0, T; H)$.

Remark 2.3 (maximal L^2 -regularity). If $u_0 \in H$ such that $\varphi(u_0)$ is finite, then Theorem 1.1 (or Theorem 2.2) says that for every $f \in L^2(0, T; H)$, the unique solution u of (1-1) has its time derivative $\dot{u} \in L^2(0, T; H)$ and hence by the differential inclusion

$$\dot{u}(t) + Au(t) \ni f(t) \quad \text{a.e. on } (0, T), \quad (2-7)$$

and also $Au \in L^2(0, T; H)$. In other words, for $f \in L^2(0, T; H)$, \dot{u} and $Au \in L^2(0, T; H)$ admit the maximal possible regularity. For this reason, we call this property *maximal L^2 -regularity*, as it is customary for generators of holomorphic semigroups on Hilbert spaces; see [Arendt 2004] for a survey on this subject.

Given $\omega \in \mathbb{R}$, we say that the shifted function $\varphi_\omega : H \rightarrow (-\infty, +\infty]$ has *compact sublevel sets* if

$$E_{\omega,c} := \{u \in D(\varphi) \mid \varphi_\omega(u) \leq c\} \quad \text{is compact in } H \text{ for every } c \in \mathbb{R}. \quad (2-8)$$

Remark 2.4. We emphasize that condition (2-8) does not imply that φ has compact sublevel sets. This becomes more clear if one considers as φ the function associated with the negative *Neumann p -Laplacian* $-\Delta_p^N$ on a bounded, open subset Ω of \mathbb{R}^d with a Lipschitz boundary $\partial\Omega$. For $\max\{1, 2d/(d+2)\} < p < \infty$, $d \geq 1$, let $V = W^{1,p}(\Omega)$, $H = L^2(\Omega)$, and $\varphi : H \rightarrow (-\infty, +\infty]$ be given by

$$\varphi(u) := \begin{cases} \frac{1}{p} \int_\Omega |\nabla u|^p \, dx & \text{if } u \in V, \\ +\infty & \text{if } u \in H \setminus V \end{cases} \quad (2-9)$$

for every $u \in H$. Then, for every $c > 0$, the sublevel set $E_{0,c}$ of φ contains the sequence $(u_n)_{n \geq 0}$ of constant functions $u_n \equiv n$, which does not admit any convergent subsequence in H . On the other hand, for every $\omega > 0$ and $c > 0$, the sublevel set $E_{\omega,c}$ is a bounded set in V and by Rellich–Kondrachov compactness, $V \hookrightarrow H$ by a compact embedding. Thus, for every $\omega > 0$ and $c > 0$, the sublevel set $E_{\omega,c}$ is compact in $L^2(\Omega)$.

3. An example and nonuniqueness

The main example of perturbations G allowed in Theorem 1.2 are Nemytskii operators on the space $\mathcal{H} = L^2(0, T; L^2(\Omega))$. Let $\Omega \subseteq \mathbb{R}^d$ be open and $g : (0, T) \times \Omega \times \mathbb{R} \rightarrow \mathbb{R}$ be a *Carathéodory function*, that is,

- $g(\cdot, \cdot, v) : (0, T) \times \Omega \rightarrow \mathbb{R}$ is measurable for all $v \in \mathbb{R}$,
- $g(t, x, \cdot) : \mathbb{R} \rightarrow \mathbb{R}$ is continuous for a.e. $(t, x) \in (0, T) \times \Omega$.

Assume furthermore g has *sublinear growth*; that is, there exist $L \geq 0$ and $b \in L^2(0, T; L^2(\Omega))$ such that

$$|g(t, x, v)| \leq L |v| + b(t, x) \quad \text{for all } v \in \mathbb{R}, \text{ a.e. } (t, x) \in (0, T) \times \Omega. \quad (3-1)$$

Proposition 3.1. *Let $\mathcal{H} = L^2(0, T; L^2(\Omega))$. Then, the relation*

$$Gv(t, x) := g(t, x, v(t, x)) \quad \text{for a.e. } (t, x) \in (0, T) \times \Omega \text{ and every } v \in \mathcal{H}, \quad (3-2)$$

defines a continuous operator $G : \mathcal{H} \rightarrow \mathcal{H}$ of sublinear growth (1-2).

The proof of Proposition 3.1 is standard (see [Zeidler 1990, Proposition 26.7]) if one uses that $f_n \rightarrow f$ in \mathcal{H} if and only if each subsequence of $(f_n)_{n \geq 1}$ has a dominated subsequence converging to f a.e. (which is well known from the completeness proof of L^2).

For illustrating the theory developed in this paper, we consider the following standard example: the *Dirichlet p -Laplacian* perturbed by a lower-order term.

Example 3.2. Let Ω be an open, bounded subset of \mathbb{R}^d , $d \geq 1$, $H = L^2(\Omega)$, and for $2d/(d+2) \leq p < \infty$, let $V = W_0^{1,p}(\Omega)$ be the closure of $C_c^1(\Omega)$ equipped with respect to the norm $\|u\|_V := \|\nabla u\|_{L^p(\Omega; \mathbb{R}^d)}$. Then, one has that V is continuously embedded into H (see [Brezis 2011, Theorem 9.16]); we write for this $V \hookrightarrow H$.

Further, let $f = \beta + f_1$ be the sum of a maximal monotone graph β of \mathbb{R} satisfying $(0, 0) \in \beta$ and a *Lipschitz–Carathéodory function* $f_1 : \Omega \times \mathbb{R} \rightarrow \mathbb{R}$ satisfying $f_1(x, 0) = 0$; that is, for a.e. $x \in \Omega$, the function $f_1(x, \cdot)$ is Lipschitz continuous (with constant $\omega > 0$) uniformly for a.e. $x \in \Omega$, and $f_1(\cdot, u)$ is measurable on Ω for every $u \in \mathbb{R}$. Then, there is a proper, convex and lower semicontinuous function $j : \mathbb{R} \rightarrow (-\infty, +\infty]$ satisfying $j(0) = 0$ and $\partial j = \beta$ in \mathbb{R} ; see [Barbu 2010, Example 1, p. 53]. We set

$$\begin{aligned} F_1(u) &= \int_0^{u(x)} f_1(\cdot, s) \, ds, & \varphi_2(u) &:= \begin{cases} \int_{\Omega} j(u(x)) \, dx & \text{if } j(u) \in L^1(\Omega), \\ +\infty & \text{if otherwise,} \end{cases} \\ F(u) &= \varphi_2(u) + \int_{\Omega} F_1(u(x)) \, dx \end{aligned} \tag{3-3}$$

for every $u \in H$. Further, let $\varphi_1 : H \rightarrow (-\infty, +\infty]$ be given by

$$\varphi_1(u) = \begin{cases} \frac{1}{p} \int_{\Omega} |\nabla u|^p \, dx + \int_{\Omega} F_1(u) \, dx & \text{if } u \in V, \\ +\infty & \text{if } u \in H \setminus V \end{cases}$$

for every $u \in H$. Then the domain $D(\varphi_1)$ of φ_1 is V . The function φ_1 is lower semicontinuous on H and is proper, $\varphi_{1,\omega}$ is convex, and for every $u \in V$, φ_1 is Gâteaux-differentiable with

$$D_v \varphi_1(u) = \lim_{t \rightarrow 0^+} \frac{\varphi_1(u + tv) - \varphi_1(u)}{t} = \int_{\Omega} |\nabla u|^{p-2} \nabla u \nabla v + f_1(x, u) v \, dx$$

for every $v \in V$. Since V is dense in H , the subdifferential operator $\partial \varphi_1$ is a single-valued operator on H with domain

$$D(\partial \varphi_1) = \left\{ u \in V \mid \text{there exists } h \in H \text{ such that } D_v \varphi_1(u) = \int_{\Omega} h v \, dx \text{ for all } v \in V \right\},$$

$$\partial \varphi_1(u) = h = -\Delta_p u + f_1(x, u) \quad \text{in } \mathcal{D}'(\Omega).$$

The operator $\partial \varphi_1$ is the negative *Dirichlet p -Laplacian* $-\Delta_p^D$ on Ω with a *Lipschitz continuous lower-order term* f_1 . Next, we add the function φ_2 given by (3-3) to φ_1 . For this, note that φ_2 is proper (since for $u_0 \equiv 0$, we have $\varphi_2(u_0) = 0$) with $\text{int}(D(\varphi_2)) \neq \emptyset$, convex (since j is convex), and lower semicontinuous on H . Thus, the function $\varphi : H \rightarrow (-\infty, +\infty]$, given by

$$\varphi(u) = \varphi_1(u) + \varphi_2(u) \quad \text{for every } u \in H, \tag{3-4}$$

is convex, lower semicontinuous, and proper with domain $D(\varphi) = \{u \in V \mid j(u) \in L^1(\Omega)\}$, and the operator $A = \partial\varphi$ is given by

$$\begin{aligned} D(A) &= \{u \in D(\varphi) \mid \text{there exists } h \in H \text{ such that } D_v\varphi(u) = \int_{\Omega} h v \, dx \text{ for all } v \in D(\varphi)\}, \\ Au &= h = -\Delta_p u + \beta(u) + f_1(x, u). \end{aligned}$$

Here, we note that

$$\overline{D(A)} = \overline{D(\varphi)} = \{u \in H \mid j(u(x)) \in \overline{D(\beta)} \text{ for a.e. } x \in \Omega\}.$$

Due to Theorem 2.1, for every $u_0 \in \overline{D(\varphi)}$ and $f \in \mathcal{H}$, there is a unique solution $u \in H_{\text{loc}}^1((0, T]; H) \cap C([0, T]; H)$ of the parabolic boundary-value problem

$$\begin{cases} \partial_t u(t) - \Delta_p u(t) + \beta(u(t)) + f_1(\cdot, u(t)) \ni f(t) & \text{on } (0, T) \times \Omega, \\ u(t) = 0 & \text{on } (0, T) \times \partial\Omega, \\ u(0) = u_0 & \text{on } \Omega. \end{cases}$$

Here, we write $\partial_t u(t)$ instead of $\dot{u}(t)$ since we rewrote the abstract Cauchy problem (1-1) as an explicit parabolic partial differential equation.

If $\max\{1, 2d/(d+2)\} < p < \infty$, then for the Lipschitz constant ω of f_1 , we have φ_{ω} is convex, and for every $c > 0$ the sublevel set $E_{\omega, c}$ is compact in $L^2(\Omega)$. Furthermore, let $g : (0, T) \times \Omega \times \mathbb{R} \rightarrow \mathbb{R}$ be a Carathéodory function with sublinear growth and $u_0 \in \overline{D(\varphi)}$. Then, there is at least one solution $u \in H_{\text{loc}}^1((0, T]; H) \cap C([0, T]; H)$ of the parabolic boundary-value problem

$$\begin{cases} \partial_t u(t, \cdot) - \Delta_p u(t, \cdot) + \beta(u(t, \cdot)) + f_1(\cdot, u(t, \cdot)) \ni g(t, \cdot, u(t, \cdot)) & \text{on } (0, T) \times \Omega, \\ u(t, \cdot) = 0 & \text{on } (0, T) \times \partial\Omega, \\ u(0, \cdot) = u_0 & \text{on } \Omega. \end{cases}$$

In general, the solutions u to the Cauchy problem (1-3) are not unique. We give an example.

Example 3.3 (nonuniqueness). Let $g(u) = \sqrt{|u|}$, $u \in \mathbb{R}$, and Ω be an open and bounded subset of \mathbb{R}^d , $d \geq 1$, with a Lipschitz boundary $\partial\Omega$. Then, there are $L, b > 0$ such that \hat{g} satisfies

$$|g(u)| \leq L|u| + b \quad \text{for every } u \in \mathbb{R}.$$

Thus, for $H = L^2(\Omega)$ and $\mathcal{H} = L^2((0, T) \times \Omega)$, the associated Nemytskii operator $G : \mathcal{H} \rightarrow \mathcal{H}$ defined by (3-2) satisfies the sublinear growth condition (1-2).

Further, for $\max\{1, 2d/(d+2)\} < p < +\infty$, let $\varphi : L^2(\Omega) \rightarrow (-\infty, +\infty]$ be the energy function (2-9) associated with the negative Neumann p -Laplacian $-\Delta_p^N$ on Ω . Then, by Theorem 1.2, for every $u_0 \in L^2(\Omega)$ and every $T > 0$, there is a solution $u \in H_{\text{loc}}^1((0, T]; L^2(\Omega)) \cap C([0, T]; L^2(\Omega))$ of

$$\begin{cases} \partial_t u(t, \cdot) - \Delta_p^N u(t, \cdot) = \sqrt{|u|(t, \cdot)} & \text{in } (0, T) \times \Omega, \\ |\nabla u(t, \cdot)|^{p-2} D_v u(t, \cdot) = 0 & \text{on } (0, T) \times \partial\Omega, \\ u(0) = u_0 & \text{on } \Omega. \end{cases} \quad (3-5)$$

Here, $|\nabla u|^{p-2} D_v u$ denotes the (weak) conormal derivative of u on $\partial\Omega$; see [Coulhon and Hauer 2016].

Now, for the initial value $u_0 \equiv 0$ on Ω , the constant zero function $u \equiv 0$ is certainly a solution of (3-5). For constructing a nontrivial solution of (3-5) with initial value $u_0 \equiv 0$, let $w \in C^1[0, T]$ be a nontrivial solution of the classical ordinary differential equation

$$w' = \sqrt{|w|} \quad \text{on } (0, T), \quad w(0) = 0, \quad (3-6)$$

For instance, one nontrivial solution is $w(t) = \frac{1}{4}t^2$. Since for every constant $c \in \mathbb{R}$ we have $-\Delta_p^N(c\mathbb{1}_\Omega) = 0$, the function $u(t) := w(t)$ is another nontrivial solution of (3-5) with initial value $u_0 \equiv 0$.

4. Proof of the main result

We now give the proof of Theorem 1.2. After possibly replacing φ by a translation, we may always assume without loss of generality that $0 \in D(\partial\varphi_\omega)$ and φ_ω attains a minimum at 0 with $\varphi_\omega(0) = 0$; for further details see [Barbu 2010, p. 159]. By the convexity of φ_ω , this implies $(0, 0) \in \omega I_H + A$, that is,

$$(h + \omega u, u)_H \geq 0 \quad \text{for all } (u, h) \in A. \quad (4-1)$$

For the proof of Theorem 1.2, we need some auxiliary results. The first concerns continuity and is standard; see [Bénilan et al. ca. 1990, (6.5), p. 87] or [Barbu 2010, (4.2), p. 128].

Lemma 4.1. *Let $f_1, f_2 \in \mathcal{H}$ and $u_1, u_2 \in H^1(0, T; H)$ such that*

$$\dot{u}_1 + Au_1 \ni f_1 \quad \text{on } (0, T),$$

$$\dot{u}_2 + Au_2 \ni f_2 \quad \text{on } (0, T).$$

Then,

$$\|u_1(t) - u_2(t)\|_H \leq e^{\omega t} \|u_1(0) - u_2(0)\|_H + \int_0^t e^{\omega(t-s)} \|f_1(s) - f_2(s)\|_H \, ds \quad (4-2)$$

for every $t \in [0, T]$.

Next, we establish the compactness of the *solution operator* P associated with evolution problem (1-1). We recall that the closure $\overline{D(\varphi)}$ in H of the effective domain of a semiconvex function φ is a convex subset of H .

Lemma 4.2. *Let $P : \overline{D(\varphi)} \times \mathcal{H} \rightarrow \mathcal{H}$ be the mapping defined by*

$$P(u_0, f) = \text{solution } u \text{ of (1-1) for every } u_0 \in \overline{D(\varphi)} \text{ and } f \in \mathcal{H}.$$

Then, P is continuous and compact.

Proof. (a) By Lemma 4.1, the map P is continuous from $\overline{D(\varphi)} \times \mathcal{H}$ to \mathcal{H} .

(b) We show that P is compact. Let $(u_n^{(0)})_{n \geq 1} \subseteq \overline{D(\varphi)}$ and $(f_n)_{n \geq 1} \subseteq \mathcal{H}$ such that $\|u_n^{(0)}\|_H + \|f_n\|_{\mathcal{H}} \leq c$ and $u_n = P(u_n^{(0)}, f_n)$ for every $n \geq 1$. Then, by (2-3), (2-4) and by (2-6), for every $\delta \in (0, T)$, there is a $c_\delta > 0$ such that

$$\sup_{n \geq 1} \|u_n\|_{H^1(\delta, T; H)} \leq c_\delta.$$

Since $H^1(\delta, T; H) \hookrightarrow C^{1/2}([\delta, T]; H)$, the sequence $(u_n)_{n \geq 1}$ is equicontinuous on $[\delta, T]$ for each $0 < \delta < T$. Choose a countable dense subset $D := \{t_m \mid m \in \mathbb{N}\}$ of $(0, T]$. Let $m \geq 1$. Then by (2-5),

$$\sup_{n \geq 1} \varphi(u_n(t_m)) \quad \text{is finite}$$

and since by (2-3), $(u_n(t_m))_{n \geq 1}$ is bounded in H , there is a $c' > 0$ such that $(u_n(t_m))_{n \geq 1}$ is in the sublevel set $E_{\omega, c'}$. Thus and by the assumption (2-8), $(u_n(t_m))_{n \geq 1}$ has a convergent subsequence in H . By Cantor's diagonalization argument, we find a subsequence $(u_{n_k})_{k \geq 1}$ of $(u_n)_{n \geq 1}$ such that

$$\lim_{k \rightarrow +\infty} u_{n_k}(t_m) \quad \text{exists in } H \text{ for all } m \in \mathbb{N}.$$

It follows from the equicontinuity of $(u_{n_k})_{k \geq 1}$ that u_{n_k} converges in $C([\delta, T]; H)$ for all $\delta \in (0, T]$. In particular, $(u_{n_k}(t))_{k \geq 1}$ converges in H for every $t \in (0, T)$ and by (2-3), $(u_{n_k})_{k \geq 1}$ is uniformly bounded in $L^\infty(0, T; H)$. Thus, it follows from Lebesgue's dominated convergence theorem that $u_{n_k} = P(u_{n_k}^{(0)}, f_{n_k})$ converges in \mathcal{H} . \square

Remark 4.3. In the previous proof, we have actually shown that P is compact from $\overline{D(\varphi)} \times \mathcal{H}$ into the Fréchet space $C((0, T]; H)$.

With these preliminaries, we can now give the proof of our main result. Here, we were inspired by the linear case [Arendt and Chill 2010].

Proof of Theorem 1.2. First, let $u_0 \in \overline{D(\varphi)}$.

For $v \in \mathcal{H}$, one has $Gv \in \mathcal{H}$ and so, by Brezis' maximal L^2 -regularity result (Theorem 2.2), there is a unique solution $u \in H_{\text{loc}}^1((0, T]; H) \cap C([0, T]; H)$ of the evolution problem

$$\begin{cases} \dot{u}(t) + Au(t) \ni Gv(t) & \text{a.e. on } (0, T), \\ u(0) = u_0. \end{cases}$$

Let $\mathcal{T}v := P(u_0, Gv)$. Then by the continuity and linear growth of G and since $P(u_0, \cdot) : \mathcal{H} \rightarrow \mathcal{H}$ is continuous and compact (Lemma 4.2), the mapping $\mathcal{T} : \mathcal{H} \rightarrow \mathcal{H}$ is continuous and compact.

(a) We consider the Schaefer set

$$\mathcal{S} := \{u \in \mathcal{H} \mid \text{there exists } \lambda \in [0, 1] \text{ such that } u = \lambda \mathcal{T}u\}.$$

We show that \mathcal{S} is bounded in \mathcal{H} . Let $u \in \mathcal{S}$. We may assume that $\lambda \in (0, 1]$; otherwise, $u \equiv 0$. Then, $u \in H_{\text{loc}}^1((0, T]; H) \cap C([0, T]; H)$ and

$$\begin{cases} \dot{u}/\lambda + A(u/\lambda) \ni Gu & \text{on } (0, T), \\ u(0) = u_0. \end{cases}$$

It follows from (4-1) that

$$\left(-\frac{\dot{u}}{\lambda}(t) + Gu(t) + \omega \frac{u}{\lambda}(t), \frac{u}{\lambda} \right)_H \geq 0 \quad \text{for a.e. } t \in (0, T).$$

Thus and by (1-2),

$$\begin{aligned} \frac{d}{dt} \frac{1}{2} \|u(t)\|_H^2 &= (\dot{u}(t), u(t))_H = (\dot{u}(t) - \lambda Gu(t) - \omega \lambda u(t), u(t))_H + (\lambda Gu(t) + \omega \lambda u(t), u(t))_H \\ &\leq (\lambda Gu(t) + \omega \lambda u(t), u(t))_H \\ &\leq \lambda (\|Gu(t)\|_H \|u(t)\|_H + \omega \|u(t)\|_H^2) \\ &\leq \lambda (L \|u(t)\|_H^2 + b(t) \|u(t)\|_H + \omega \|u(t)\|_H^2) \\ &\leq (2L + 1 + 2\omega) \frac{1}{2} \|u(t)\|_H^2 + \frac{1}{2} b^2(t) \end{aligned}$$

for a.e. $t \in (0, T)$. It follows from Gronwall's lemma that (1-4) holds for every $t \in [0, T]$. Thus, \mathcal{S} is bounded in \mathcal{H} . Now, Schaefer's fixed point theorem implies that there exists $u \in \mathcal{H}$ such that $u = \mathcal{T}u$; that is, $u \in H_{\text{loc}}^1((0, T]; H) \cap C([0, T]; H)$ is a solution of the evolution problem (1-3).

(b) Let $u_0 \in D(\varphi)$. Then, by the first part of this proof, there is a solution $u \in H_{\text{loc}}^1((0, T]; H) \cap C([0, T]; H)$ of the evolution problem (1-3). However, by Brezis' maximal regularity result applied to $f = Gu \in \mathcal{H}$, it follows that $u \in H^1(0, T; H)$. \square

5. Application to j -elliptic functions

In Examples 3.2 and 3.3, V is a Banach space injected in H . Recently, in Chill, Hauer and Kennedy [Chill et al. 2016] extended results of [Arendt and ter Elst 2011; 2012] to a nonlinear framework of j -elliptic functions $\varphi : V \rightarrow (-\infty, +\infty]$ generating a quasimaximal monotone operator $\partial_j \varphi$ on H , where $j : V \rightarrow H$ is just a linear operator which is not necessarily injective. This enabled the authors of [Chill et al. 2016] to show that several coupled parabolic-elliptic systems can be realized as a gradient system in a Hilbert space H and to extend the linear variational theory of the Dirichlet-to-Neumann operator to the nonlinear p -Laplace operator; see also [Belhachmi and Chill 2015; 2018] for further applications and extensions of this theory.

The aim of this section is to illustrate that Theorem 1.2 of Section 3 can also be applied to the framework of j -elliptic functions.

Let us briefly recall some basic notions and facts about j -elliptic functions from [Chill et al. 2016]. Let V be a real locally convex topological vector space and $j : V \rightarrow H$ be a linear operator which is merely weak-to-weak continuous (and, in general, not injective). Given a function $\varphi : V \rightarrow (-\infty, +\infty]$, the j -subdifferential is the operator

$$\partial_j \varphi := \left\{ (u, f) \in H \times H \mid \begin{array}{l} \text{there exists } \hat{u} \in D(\varphi) \text{ such that } j(\hat{u}) = u \text{ and for every } \hat{v} \in V \\ \liminf_{t \searrow 0} (\varphi(\hat{u} + t\hat{v}) - \varphi(\hat{u}))/t \geq (f, j(\hat{v}))_H \end{array} \right\}.$$

The function φ is called j -semiconvex if there exists $\omega \in \mathbb{R}$ such that the "shifted" function $\varphi_\omega : V \rightarrow (-\infty, +\infty]$ given by

$$\varphi(\hat{u}) + \frac{1}{2}\omega \|j(\hat{u})\|_H^2 \quad \text{for every } \hat{u} \in V$$

is convex. If $V = H$ and $j = I_H$, then j -semiconvex functions φ are the *semiconvex* ones (see Section 1). The function φ is called j -elliptic if there exists $\omega \geq 0$ such that φ_ω is convex and for every $c \in \mathbb{R}$, the sublevel sets $\{\hat{u} \in V \mid \varphi_\omega(\hat{u}) \leq c\}$ are relatively weakly compact. Finally, we say that the function φ is *lower semicontinuous* if the sublevel sets $\{\varphi \leq c\}$ are closed in the topology of V for every $c \in \mathbb{R}$. It was highlighted in [Chill et al. 2016, Lemma 2.2] that:

(a) If φ is j -semiconvex, then there is an $\omega \in \mathbb{R}$ such that

$$\partial_j \varphi = \left\{ (u, f) \in H \times H \mid \begin{array}{l} \text{there exists } \hat{u} \in D(\varphi) \text{ such that } j(\hat{u}) = u \text{ and for every } \hat{v} \in V \\ \varphi_\omega(\hat{u} + \hat{v}) - \varphi_\omega(\hat{u}) \geq (f + \omega j(\hat{u}), j(\hat{v}))_H \end{array} \right\}.$$

(b) If φ is Gâteaux differentiable with directional derivative $D_{\hat{v}}\varphi$, ($\hat{v} \in V$), then

$$\partial_j \varphi = \left\{ (u, f) \in H \times H \mid \begin{array}{l} \text{there exists } \hat{u} \in D(\varphi) \text{ such that } j(\hat{u}) = u \text{ and for every } \hat{v} \in V \\ D_{\hat{v}}\varphi(\hat{u}) = (f, j(\hat{v}))_H \end{array} \right\}.$$

The main result in [Chill et al. 2016] is that the j -subdifferential $\partial_j \varphi$ of a j -elliptic function φ is already a classical subdifferential. More precisely, the following holds.

Theorem 5.1 [Chill et al. 2016, Corollary 2.7]. *Let $\varphi : V \rightarrow (-\infty, +\infty]$ be proper, lower semicontinuous, and j -elliptic. Then there is a proper, lower semicontinuous, semiconvex function $\varphi^H : H \rightarrow (-\infty, +\infty]$ such that $\partial_j \varphi = \partial \varphi^H$. The function φ^H is unique up to an additive constant.*

Thus the operator $A = \partial_j \varphi$ has the properties of maximal regularity we used before. The following result gives a description of φ^H in the convex case and will be important for our intentions in this paper.

Theorem 5.2 [Chill et al. 2016, Theorem 2.9]. *Assume that $\varphi : V \rightarrow (-\infty, +\infty]$ is convex, proper, lower semicontinuous and j -elliptic, and let $\varphi^H : H \rightarrow (-\infty, +\infty]$ be the function from Theorem 5.1. Then, there is a constant $c \in \mathbb{R}$ such that*

$$\varphi^H(u) = c + \inf_{\hat{u} \in j^{-1}(\{u\})} \varphi(\hat{u}) \quad \text{for every } u \in H,$$

with effective domain $D(\varphi^H) = j(D(\varphi))$.

For our perturbation result, we need the compactness of the sublevel sets of φ^H . With the help of Theorem 5.2 we can establish a criterion in terms of the given φ for this property.

Lemma 5.3. *Let $\varphi : V \rightarrow (-\infty, +\infty]$ be proper, lower semicontinuous j -semiconvex, and j -elliptic. Assume that*

$$j : V \rightarrow H \text{ maps weakly relatively compact sets of } V \text{ into relatively norm-compact sets of } H. \quad (5-1)$$

Then there is an $\omega \geq 0$ such that for every $c \in \mathbb{R}$ the sublevel set

$$E_{\omega, c} = \{u \in H \mid \varphi_{\omega}^H(u) \leq c\} \quad \text{is compact in } H.$$

Remark 5.4. If V is a normed space, then by the Eberlein–Šmulian theorem hypothesis (5-1) is equivalent to j maps weakly convergent sequences in V to norm convergent sequences in H . This in turn is equivalent to j being compact if V is reflexive.

Proof of Lemma 5.3. By hypothesis, there is an $\omega \geq 0$ such that φ_{ω} is convex, lower semicontinuous, and for every $c \in \mathbb{R}$, the sublevel sets $\{\hat{u} \in V \mid \varphi_{\omega}(\hat{u}) \leq c\}$ are weakly relatively compact and closed. By Theorem 5.1, there is a lower semicontinuous, proper function $\varphi^H : H \rightarrow (-\infty, +\infty]$ such that φ_{ω}^H is convex and $\partial \varphi_{\omega}^H = \partial_j \varphi_{\omega}$. Applying Theorem 5.2 to φ_{ω} and φ_{ω}^H , we have

$$\varphi_{\omega}^H(u) = d + \inf_{\hat{u} \in j^{-1}(\{u\})} \varphi_{\omega}(\hat{u}) \quad \text{for every } u \in H \quad (5-2)$$

and some constant $d \in \mathbb{R}$. For $c \in \mathbb{R}$, let $(u_n)_{n \geq 1}$ be an arbitrary sequence in $E_{\omega, c}$. By (5-2), for every $n \in \mathbb{N}$, there is a $\hat{u}_n \in j^{-1}(\{u_n\})$ such that

$$d + \varphi_{\omega}(\hat{u}_n) \leq c + 1.$$

By hypothesis, all sublevel sets of φ_ω are weakly relatively compact in V . Thus, by our hypothesis, the image under j is relatively compact in H . Consequently, there are a subsequence $(u_{n_l})_{l \geq 1}$ of $(u_n)_{n \geq 1}$ and a $u \in H$ such that $u_{n_l} = j(\hat{u}_{n_l}) \rightarrow u$ in H as $l \rightarrow +\infty$. Since $\varphi_\omega^H(u_{n_l}) \leq c$ and since φ^H is lower semicontinuous, it follows that $\varphi^H(u) \leq c$. This shows that $E_{\omega,c}$ is compact. \square

Now, applying Lemma 5.3 to Theorem 1.2, we can state the following existence theorem.

Theorem 5.5. *Let $\varphi : V \rightarrow (-\infty, +\infty]$ be proper, lower semicontinuous j -semiconvex, and j -elliptic. Assume that the mapping j satisfies (5-1) and let $G : \mathcal{H} \rightarrow \mathcal{H}$ be a continuous mapping of sublinear growth (1-2). Then, for $A = \partial_j \varphi$ the nonlinear evolution problem (1-3) admits for every $u_0 \in j(\overline{D(\varphi)})$ and $f \in \mathcal{H}$ at least one solution $u \in H_{\text{loc}}^1((0, T]; H) \cap C([0, T]; H)$. In particular, $\varphi \circ u$ belongs to $W_{\text{loc}}^{1,1}((0, T]) \cap L^1(0, T)$ and inequality (1-4) holds. If $u_0 \in j(D(\varphi))$, then problem (1-3) has a solution $u \in H^1(0, T; H)$.*

We complete this section by considering the following evolution problem involving the *Dirichlet-to-Neumann operator* associated with the p -Laplacian; see [Hauer 2015; Chill et al. 2016].

Example 5.6. Let Ω be a bounded domain with a Lipschitz continuous boundary $\partial\Omega$. Then, for $2d/(d+1) < p < +\infty$, the trace operator $\text{Tr} : W^{1,p}(\Omega) \rightarrow L^2(\partial\Omega)$ is a completely continuous operator (see [Nečas 1967, Théorème 6.2] for the case $p < d$; the other cases $p = d$ and $p > d$ can be deduced from Conséquences 6.2 and 6.3 of the same work). Now, we take

$$V = W^{1,p}(\Omega), \quad H = L^2(\partial\Omega), \quad \text{and} \quad j = \text{Tr}.$$

Then, j is a linear bounded mapping satisfying hypothesis (5-1). In fact, j is a prototype of a noninjective mapping. Furthermore, let $\varphi : V \rightarrow \mathbb{R}$ be the function given by

$$\varphi(\hat{u}) = \frac{1}{p} \int_{\Omega} |\nabla \hat{u}|^p \, dx \quad \text{for every } \hat{u} \in V.$$

Then, φ is continuously differentiable on V and convex. Thus, the Tr -subdifferential operator $\partial_{\text{Tr}}\varphi$ is given by

$$\partial_{\text{Tr}}\varphi = \left\{ (u, f) \in H \times H \mid \begin{array}{l} \text{there exists } \hat{u} \in V \text{ such that } \text{Tr}(\hat{u}) = u \text{ and for every } \hat{v} \in V \\ \int_{\Omega} |\nabla \hat{u}|^{p-2} \nabla \hat{u} \nabla \hat{v} \, dx = (f, j(\hat{v}))_H \end{array} \right\}.$$

Moreover, by [Hauer 2015, inequality (20)], for any $\omega > 0$, the shifted function φ_ω has bounded sublevel sets in V . Since V is reflexive, every sublevel set of φ_ω is weakly compact in V . In addition, by Lemma 2.1 of the same work, $j(D(\varphi))$ is dense in H .

Now, let $g : (0, T) \times \Omega \times \mathbb{R} \rightarrow \mathbb{R}$ be a Carathéodory function with sublinear growth. Then by Theorem 5.5, for every $u_0 \in L^2(\partial\Omega)$, there is at least one solution $u \in H_{\text{loc}}^1((0, T]; L^2(\partial\Omega)) \cap C([0, T]; L^2(\partial\Omega))$ of the elliptic-parabolic boundary-value problem

$$\left\{ \begin{array}{ll} -\Delta_p \hat{u}(t, \cdot) = 0 & \text{on } (0, T) \times \Omega, \\ \partial_t u(t, \cdot) + |\nabla u(t, \cdot)|^{p-2} \frac{\partial}{\partial \nu} u(t, \cdot) = g(t, \cdot, u(t, \cdot)) & \text{on } (0, T) \times \partial\Omega, \\ u(t, \cdot) = \hat{u}(t, \cdot) & \text{on } (0, T) \times \partial\Omega, \\ u(0, \cdot) = u_0 & \text{on } \partial\Omega. \end{array} \right.$$

References

- [Arendt 2004] W. Arendt, “Semigroups and evolution equations: functional calculus, regularity and kernel estimates”, pp. 1–85 in *Evolutionary equations, Vol. I*, edited by C. M. Dafermos and E. Feireisl, North-Holland, Amsterdam, 2004. MR Zbl
- [Arendt and Chill 2010] W. Arendt and R. Chill, “Global existence for quasilinear diffusion equations in isotropic nondivergence form”, *Ann. Sc. Norm. Super. Pisa Cl. Sci. (5)* **9**:3 (2010), 523–539. MR Zbl
- [Arendt and ter Elst 2011] W. Arendt and A. F. M. ter Elst, “The Dirichlet-to-Neumann operator on rough domains”, *J. Differential Equations* **251**:8 (2011), 2100–2124. MR Zbl
- [Arendt and ter Elst 2012] W. Arendt and A. F. M. ter Elst, “Sectorial forms and degenerate differential operators”, *J. Operator Theory* **67**:1 (2012), 33–72. MR Zbl
- [Barbu 2010] V. Barbu, *Nonlinear differential equations of monotone types in Banach spaces*, Springer, 2010. MR Zbl
- [Belhachmi and Chill 2015] Z. Belhachmi and R. Chill, “Application of the j -subgradient in a problem of electropermeabilization”, *J. Elliptic Parabol. Equ.* **1** (2015), 13–29. MR Zbl
- [Belhachmi and Chill 2018] Z. Belhachmi and R. Chill, “The bidomain problem as a gradient system”, 2018. arXiv
- [Bénilan et al. ca. 1990] P. Bénilan, M. G. Crandall, and A. Pazy, “Evolution problems governed by accretive operators”, unpublished manuscript, ca. 1990.
- [Brezis 1971] H. Brézis, “Propriétés régularisantes de certains semi-groupes non linéaires”, *Israel J. Math.* **9** (1971), 513–534. MR Zbl
- [Brezis 1973] H. Brézis, *Opérateurs maximaux monotones et semi-groupes de contractions dans les espaces de Hilbert*, North-Holland Mathematics Studies **5**, North-Holland, Amsterdam, 1973. MR Zbl
- [Brezis 2011] H. Brézis, *Functional analysis, Sobolev spaces and partial differential equations*, Springer, 2011. MR Zbl
- [Chill et al. 2016] R. Chill, D. Hauer, and J. Kennedy, “Nonlinear semigroups generated by j -elliptic functionals”, *J. Math. Pures Appl. (9)* **105**:3 (2016), 415–450. MR Zbl
- [Coulhon and Hauer 2016] T. Coulhon and D. Hauer, “Regularisation effects of nonlinear semigroups”, preprint, 2016. To appear in *BCAM Springer Briefs*. arXiv
- [Evans 2010] L. C. Evans, *Partial differential equations*, 2nd ed., Graduate Studies in Mathematics **19**, American Mathematical Society, Providence, RI, 2010. MR Zbl
- [Hauer 2015] D. Hauer, “The p -Dirichlet-to-Neumann operator with applications to elliptic and parabolic problems”, *J. Differential Equations* **259**:8 (2015), 3615–3655. MR Zbl
- [Nečas 1967] J. Nečas, *Les méthodes directes en théorie des équations elliptiques*, Masson et Cie, Paris, 1967. MR Zbl
- [Schaefer 1955] H. Schaefer, “Über die Methode der a priori-Schranken”, *Math. Ann.* **129** (1955), 415–416. MR Zbl
- [Zeidler 1990] E. Zeidler, *Nonlinear functional analysis and its applications, II/B: Nonlinear monotone operators*, Springer, 1990. MR Zbl

Received 14 Feb 2019. Revised 3 Aug 2019. Accepted 9 Sep 2019.

WOLFGANG ARENDT: wolfgang.arendt@uni-ulm.de
 Institute of Applied Analysis, University of Ulm, Ulm, Germany

DANIEL HAUER: daniel.hauer@sydney.edu.au
 School of Mathematics and Statistics, The University of Sydney, Sydney, Australia

PURE and APPLIED ANALYSIS

msp.org/paa

EDITORS-IN-CHIEF

- Charles L. Epstein University of Pennsylvania
cle@math.upenn.edu
- Maciej Zworski University of California at Berkeley
zworski@math.berkeley.edu

EDITORIAL BOARD

- Sir John M. Ball University of Oxford
ball@maths.ox.ac.uk
- Michael P. Brenner Harvard University
brenner@seas.harvard.edu
- Charles Fefferman Princeton University
cf@math.princeton.edu
- Susan Friedlander University of Southern California
susanfri@usc.edu
- Anna Gilbert University of Michigan
annacg@umich.edu
- Leslie F. Greengard Courant Institute, New York University, and
Flatiron Institute, Simons Foundation
greengard@cims.nyu.edu
- Yan Guo Brown University
yan_guo@brown.edu
- Claude Le Bris CERMICS - ENPC
lebris@cermics.enpc.fr
- Robert J. McCann University of Toronto
mccann@math.toronto.edu
- Michael O'Neil Courant Institute, New York University
oneil@cims.nyu.edu
- Jill Pipher Brown University
jill_pipher@brown.edu
- Johannes Sjöstrand Université de Dijon
johannes.sjostrand@u-bourgogne.fr
- Vladimir Šverák University of Minnesota
sverak@math.umn.edu
- Daniel Tataru University of California at Berkeley
tataru@berkeley.edu
- Michael I. Weinstein Columbia University
miw2103@columbia.edu
- Jon Wilkening University of California at Berkeley
wilken@math.berkeley.edu
- Enrique Zuazua DeustoTech-Bilbao, and
Universidad Autónoma de Madrid
enrique.zuazua@deusto.es

PRODUCTION

- Silvio Levy (Scientific Editor)
production@msp.org

Cover image: The figure shows the outgoing scattered field produced by scattering a plane wave, coming from the northwest, off of the (stylized) letters P A A. The total field satisfies the homogeneous Dirichlet condition on the boundary of the letters. It is based on a numerical computation by Mike O'Neil of the Courant Institute.

See inside back cover or msp.org/paa for submission instructions.

The subscription price for 2020 is US \$505/year for the electronic version, and \$565/year (+\$25, if shipping outside the US) for print and electronic. Subscriptions, requests for back issues and changes of subscriber address should be sent to MSP.

Pure and Applied Analysis (ISSN 2578-5885 electronic, 2578-5893 printed) at Mathematical Sciences Publishers, 798 Evans Hall #3840, c/o University of California, Berkeley, CA 94720-3840 is published continuously online. Periodical rate postage paid at Berkeley, CA 94704, and additional mailing offices.

PAA peer review and production are managed by EditFlow® from MSP.

PUBLISHED BY

 **mathematical sciences publishers**
nonprofit scientific publishing

<http://msp.org/>

© 2020 Mathematical Sciences Publishers

PURE and APPLIED ANALYSIS

vol. 2 no. 1 2020

- Emergence of nontrivial minimizers for the three-dimensional Ohta–Kawasaki energy 1
HANS KNÜPFER, CYRILL B. MURATOV and MATTEO NOVAGA
- Maximal L^2 -regularity in nonlinear gradient systems and perturbations of sublinear growth 23
WOLFGANG ARENDT and DANIEL HAUER
- The local density approximation in density functional theory 35
MATHIEU LEWIN, ELLIOTT H. LIEB and ROBERT SEIRINGER
- Sparse bounds for the discrete spherical maximal functions 75
ROBERT KESLER, MICHAEL T. LACEY and DARÍO MENA
- Characterization by observability inequalities of controllability and stabilization properties 93
EMMANUEL TRÉLAT, GENGSHENG WANG and YASHAN XU
- Stationary compressible Navier–Stokes equations with inflow condition in domains with piecewise analytical boundaries 123
PIOTR B. MUCHA and TOMASZ PIASECKI
- Optimal constants in nontrapping resolvent estimates and applications in numerical analysis 157
JEFFREY GALKOWSKI, EUAN A. SPENCE and JARED WUNSCH



2578-5893 (2020)2:1;1-B