


PURE and APPLIED ANALYSIS

PAM

ROBERT KESLER, MICHAEL T. LACEY AND DARÍO MENA

**SPARSE BOUNDS FOR THE DISCRETE SPHERICAL MAXIMAL
FUNCTIONS**



vol. 2 no. 1 2020

SPARSE BOUNDS FOR THE DISCRETE SPHERICAL MAXIMAL FUNCTIONS

ROBERT KESLER, MICHAEL T. LACEY AND DARÍO MENA

We prove sparse bounds for the spherical maximal operator of Magyar, Stein and Wainger. The bounds are conjecturally sharp, and contain an endpoint estimate. The new method of proof is inspired by ones by Bourgain and Ionescu, is very efficient, and has not been used in the proof of sparse bounds before. The Hardy–Littlewood circle method is used to decompose the multiplier into major and minor arc components. The efficiency arises as one only needs a single estimate on each element of the decomposition.

1. Introduction

Let $\mathcal{A}_\lambda f = d\sigma_\lambda * f$, where $d\sigma_\lambda$ is a uniform unit mass spherical measure on a sphere of radius λ in \mathbb{R}^d for $d \geq 3$. Set the Stein spherical maximal operator to be

$$\mathcal{A}f(x) = \sup_{\lambda > 0} \mathcal{A}_\lambda f,$$

where f is a nonnegative compactly supported and bounded function. We are interested in a sparse bound for the maximal function. In the continuous case, this estimate holds, and is sharp, up to the boundary.

Theorem 1.1 [Lacey 2017]. *Let $d \geq 3$ and set \mathbf{R}_d to be the polygon with vertices*

$$R_0 = \left(\frac{d-1}{d}, \frac{1}{d} \right), \quad R_1 = \left(\frac{d-1}{d}, \frac{d-1}{d} \right), \quad R_2 = \left(\frac{d^2-d}{d^2+1}, \frac{d^2-d+2}{d^2+1} \right), \quad R_3 = (0, 1).$$

(See Figure 1.) *Then, for all $(\frac{1}{p}, \frac{1}{q})$ in the interior of \mathbf{R}_d , we have the sparse bound $\|\mathcal{A}\|_{p,q} < \infty$.*

We set notation for the sparse bounds. Call a collection of cubes \mathcal{S} in \mathbb{R}^n *sparse* if there are sets $\{E_S : S \in \mathcal{S}\}$ which are pairwise disjoint and satisfy $E_S \subset S$ and $|E_S| > \frac{1}{4}|S|$. For any cube Q and $1 \leq r < \infty$, set $\langle f \rangle_{Q,r}^r = |Q|^{-1} \int_Q |f|^r dx$. Then the (r, s) -sparse form $\Lambda_{\mathcal{S},r,s} = \Lambda_{r,s}$ indexed by the sparse collection \mathcal{S} is

$$\Lambda_{\mathcal{S},r,s}(f, g) = \sum_{S \in \mathcal{S}} |S| \langle f \rangle_{S,r} \langle g \rangle_{S,s}.$$

For a sublinear operator T , we set $\|T\|_{r,s}$ to be the best constant C in the inequality

$$\langle Tf, g \rangle < C \sup_{\mathcal{S}} \Lambda_{\mathcal{S},r,s}(f, g).$$

We use the same notation for sublinear operators T acting on functions defined on \mathbb{Z}^d .

Lacey was supported in part by grant from the US National Science Foundation, DMS-1600693 and the Australian Research Council ARC DP160100153. Mena was supported by project 821-B8-287, CIMPA, Escuela de Matemática, UCR.

MSC2010: 11K70, 42B25.

Keywords: sparse bounds, spherical averages, discrete, spherical maximal function.




Figure 1. Sparse bounds hold for points $(\frac{1}{p}, \frac{1}{q})$ in the interior of the four-sided region R_d . The points R_1 and R_2 are as given in Theorem 1.1.

The theorem above refines the well-known L^p -improving properties for the local maximal function $\sup_{1 \leq \lambda \leq 2} \mathcal{A}_\lambda * f$, proved in [Schlag 1997; Schlag and Sogge 1997]; also see [Lee 2003]. The theorem above has as immediate corollaries (a) vector-valued inequalities, and (b) weighted consequences. Both sets of consequences are the strongest known. The method of proof uses the L^p -improving inequalities for the spherical maximal function. That is, the proof is, in some sense, standard, although only recently discovered, and yields the best known information about the mapping properties of the spherical maximal function.

We turn to the setting of discrete spherical averages. Provided λ^2 is an integer, and the dimension d satisfies $d \geq 5$, we can define

$$A_\lambda f(x) = \lambda^{2-d} \sum_{n \in \mathbb{Z}^d : |n| = \lambda} f(x - n)$$

for functions $f \in \ell^2(\mathbb{Z}^d)$. We restrict attention to the case of $d \geq 5$ as in that case $|\{n \in \mathbb{Z}^d : |n| = \lambda\}| \simeq \lambda^{d-2}$ for all $\lambda^2 \in \mathbb{N}$. Let $Af = \sup_\lambda A_\lambda f$, where we will always understand that $\lambda^2 \in \mathbb{N}$. This is the maximal function of [Magyar 1997; Magyar, Stein, and Wainger 2002]. The following theorem is the best known extension of the sparse bounds for the continuous spherical maximal function to the discrete setting.

Theorem 1.2. *Let Z_d be the polygon with vertices*

$$Z_j = \frac{d-4}{d-2} R_j + \frac{2}{d-2} \left(\frac{1}{2}, \frac{1}{2} \right), \quad j = 0, 1, 2, \quad (1.3)$$

and $Z_3 = (0, 1)$. (See Figure 2.) *There holds:*

- (1) *For all $(\frac{1}{p}, \frac{1}{q})$ in the interior Z_d , we have the sparse bound $\|Af\|_{p,q} < \infty$.*
- (2) *When $f = \mathbf{1}_F$ and $g = \mathbf{1}_G$,*

$$\langle A\mathbf{1}_F, \mathbf{1}_G \rangle \lesssim \sup_S \Lambda_{S, d/(d-2), d/(d-2)}(\mathbf{1}_F, \mathbf{1}_G). \quad (1.4)$$




Figure 2. Sparse bounds for the discrete spherical maximal function hold for points $(\frac{1}{p}, \frac{1}{q})$ in the interior of the four-sided figure above. The dotted lines pass through the points $(\frac{1}{2}, \frac{1}{2})$ and the points R_1 and R_2 of Figure 1. Circles along these lines are the points Z_1 and Z_2 . The restricted weak-type sparse bound (1.4) holds at the filled in circle, $Z_1 = (\frac{d}{d-2}, \frac{d}{d-2})$.

By direct computation,

$$Z_0 = \left(\frac{d-2}{d}, \frac{2}{d} \right), \quad Z_1 = \left(\frac{d-2}{d}, \frac{d-2}{d} \right), \quad Z_2 = \left(\frac{d^3-4d^2+4d+1}{d^3-2d^2+d-2}, \frac{d^3-4d^2+6d-7}{d^3-2d^2+d-2} \right). \quad (1.5)$$

The sparse bound near the point $(\frac{d-2}{d}, \frac{2}{d})$ implies the maximal inequality of [Magyar, Stein, and Wainger 2002], namely that $A : \ell^p(\mathbb{Z}^d) \rightarrow \ell^p(\mathbb{Z}^d)$ for $p > \frac{d}{d-2}$. The sparse bound (1.4) requires that both functions be indicator sets, and so is of restricted weak type. It implies the restricted weak-type inequality of [Ionescu 2004]. These inequalities imply a wide range of weighted and vector-valued inequalities, all of which are new. See the applications of the sparse bound in the continuous case in [Lacey 2017].

The discrete spherical maximal function ℓ^p -bounds were established in [Magyar, Stein, and Wainger 2002], with an endpoint restricted weak-type estimate proved in [Ionescu 2004]. The discrete ℓ^p -improving inequalities have only recently been investigated. The case of a fixed radius was addressed, independently, in [Hughes 2018; Kesler and Lacey 2018]. Spherical maximal functions, restricting to lacunary and superlacunary cases, require different techniques [Hughes 2019; Cook 2018; Kesler, Lacey, and Mena 2019]. Robert Kesler [2018a; 2018b] established sparse bounds for the discrete case. This paper extends and simplifies those arguments.

It is very tempting to conjecture that our sparse bounds form the sharp range, up to the endpoints. One would expect that certain kinds of natural examples would demonstrate this. But examples are much harder to come by in the discrete setting. We return to this in Section 5.

The argument in this paper is elegant, especially if one restricts attention to the endpoint estimate (1.4), and much simpler than the arguments in [Kesler 2018a; 2018b]. It proceeds by decomposing the maximal function into a series of terms, guided by the Hardy–Littlewood circle method decomposition developed in [Magyar, Stein, and Wainger 2002]. The decomposition has many parts, as indicated in Figures 3 and 4. But, for each part of the decomposition, we need only one estimate, either an ℓ^2 -estimate, or an endpoint estimate. Roughly speaking, one uses either a “high-frequency” ℓ^2 -estimate, or a “low-frequency” inequality, in which one compares to smoother averages. Interestingly, the notion of “smoother averages” varies. The argument of Ionescu combined with the sparse perspective yields a powerful inequality. Notation and conventions will be established in this section, and used throughout the paper.

2. Proof of the sparse bounds inside the polygon Z_d

A sparse bound is typically proved by recursion. So, the main step is to prove the recursive statement. To do this, we fix a large dyadic cube E and functions $f = \mathbf{1}_F$ and $g = \mathbf{1}_G$ supported on E . We say that $\tau : E \rightarrow \{1, \dots, \ell E\}$ is an *admissible stopping time* if for any subcube $Q \subset E$ with $\langle f \rangle_Q > C \langle f \rangle_E$, for some large constant C to be chosen later, we have $\min_{x \in Q} \tau(x) > \ell Q$.

Lemma 2.1. *Let $(\frac{1}{p}, \frac{1}{q})$ be in the interior of Z_d . For any dyadic cube, functions $f = \mathbf{1}_F$ and $g = \mathbf{1}_G$ supported on E , and any admissible stopping time τ , there holds*

$$|E|^{-1} \langle A_\tau f, g \rangle \lesssim \langle f \rangle_E^{1/p} \langle g \rangle_E^{1/q}. \quad (2.2)$$

We complete the proof of the main result.

Proof of Theorem 1.2(1). We can assume that there is a fixed dyadic cube E so that $f = \mathbf{1}_F$ is supported on the cube $3E$, and $g = \mathbf{1}_G$ is supported on E . Let \mathcal{Q}_E be the maximal dyadic subcubes of E for which $\langle f \rangle_{3Q} > C \langle f \rangle_{3E}$ for a large constant C . Observe that we have, for an appropriate choice of admissible $\tau(x)$,

$$\left\langle \sup_{\lambda \leq \ell(E)} A_\lambda f, g \right\rangle \leq \langle A_\tau f, g \rangle + \sum_{Q \in \mathcal{Q}_E} \left\langle \sup_{\lambda \leq \ell(Q)} A_\lambda (f \mathbf{1}_{3Q}), g \mathbf{1}_Q \right\rangle.$$

The first term is controlled by (2.2). For an appropriate constant $C \simeq 3^d$, we have

$$\sum_{Q \in \mathcal{Q}_E} |Q| \leq \frac{1}{4} |E|.$$

We can clearly recurse on the second term above to construct our sparse bound. This proves a sparse bound for all indicator functions in the interior of Z_d .

Sparse bounds for indicator functions in an open set self-improve to sparse bounds for functions. We give the details in the last section; see Lemma 4.1. \square

We use the corresponding recursive inequality for spherical averages on \mathbb{R}^d . Recall that \mathcal{A}_λ is the continuous spherical average.

Lemma 2.3 [Lacey 2017, Lemma 3.4]. *Let $(\frac{1}{\bar{p}}, \frac{1}{\bar{q}})$ be in the interior of \mathbf{R}_d . For any dyadic cube E , functions $\phi = \mathbf{1}_F$ and $\gamma = \mathbf{1}_G$ supported on E , and any admissible stopping time τ , there holds*

$$|E|^{-1} \langle \mathcal{A}_\tau \phi, \gamma \rangle \lesssim \langle \phi \rangle_E^{1/\bar{p}} \langle \gamma \rangle_E^{1/\bar{q}}. \quad (2.4)$$

We turn to the proof of Lemma 2.1. The restriction to indicator functions will allow us to use interpolation arguments, even though our setting has stopping times, and hence is nonlinear. Let L be the line through $(\frac{1}{2}, \frac{1}{2})$ and $(\frac{1}{\bar{p}}, \frac{1}{\bar{q}})$. Then, let $(\frac{1}{p}, \frac{1}{q})$ be a point on L that is in the interior of \mathbf{R}_d , and very close to the boundary. (The dashed lines in Figure 2 are examples of the lines L we are discussing here.)

This is the point: Fix $(\frac{1}{\bar{p}}, \frac{1}{\bar{q}}) \in \mathbf{R}_d$. For all sufficiently small $0 < \epsilon < 1$ so that $(\frac{1}{\bar{p}+\epsilon}, \frac{1}{\bar{q}+\epsilon}) \in \mathbf{R}_d$ and integers $N \in \mathbb{N}$, we can write $A_\tau f \leq M_1 + M_2$, where

$$|E|^{-1} \langle M_1, g \rangle \lesssim N^{1+\epsilon} \langle f \rangle_E^{1/(\bar{p}+\epsilon)} \langle g \rangle_E^{1/(\bar{q}+\epsilon)}, \quad (2.5)$$

$$|E|^{-1} \langle M_2, g \rangle \lesssim N^{d\epsilon+(4-d)/2} \langle f \rangle_E^{1/2} \langle g \rangle_E^{1/2}. \quad (2.6)$$

Implied constants depend upon \bar{p}, \bar{q} and ϵ , but we do not track the dependence. Once this is proved, one has

$$|E|^{-1} \langle A_\tau f, g \rangle \lesssim N^{1+\epsilon} \langle f \rangle_E^{1/(\bar{p}+\epsilon)} \langle g \rangle_E^{1/(\bar{q}+\epsilon)} + N^{d\epsilon+(4-d)/2} \langle f \rangle_E^{1/2} \langle g \rangle_E^{1/2}.$$

Choosing N to minimize the right-hand side and letting $(\frac{1}{\bar{p}}, \frac{1}{\bar{q}})$ and $0 < \epsilon < 1$ vary completes the proof. Indeed, ignoring ϵ 's, we see that the value of p is given by

$$\frac{1}{p} = \frac{1}{\bar{p}} + \frac{2}{d-2} \left(\frac{1}{2} - \frac{1}{\bar{p}} \right) = \frac{2}{d-2} \cdot \frac{1}{2} + \frac{d-4}{d-2} \cdot \frac{1}{\bar{p}}.$$

Compare this to our description of the extreme points of \mathbf{Z}_d in (1.3). Thus, our lemma follows.

In proving (2.5) and (2.6), it suffices, given $0 < \epsilon < 1$, to prove the statement for sufficiently large N . We will do so for $N > N_0$, for a sufficiently large choice of $N_0 > 0$. Indeed, we find it necessary to use an absorption argument. We show that

$$A_\tau f \leq M_1 + M_2 + \frac{1}{2} A_\tau f, \quad (2.7)$$

where M_1 and M_2 are as in (2.5) and (2.6).

The reader can consult Figure 3 for a guide to the argument. For a technical reason, we assume that $F \subset (2\mathbb{Z}^d) + \delta_f$ and $G \subset (2\mathbb{Z}^d) + \delta_g$. Here, $\delta_f, \delta_g \in \{0, 1\}^d$. This can be assumed without loss of generality.

Small values of τ . The terms M_1 and M_2 have several components. The first contribution to M_1 is the term $M_{1,1} = \mathbf{1}_{\tau \leq N^{1+\epsilon}} A_\tau f$. Our verification that $M_{1,1}$ satisfies (2.5) is our first application of the \mathbb{R}^d inequality (2.4).

We need functions on \mathbb{R}^d . Take $\phi(x) = \sum_{n \in \mathbb{Z}^d} \mathbf{1}_F(n) \mathbf{1}_{n+[-1,1]^d}(x)$, and define γ similarly. By the reduction we made above, these are indicator functions. Moreover, if τ is an admissible stopping time for f , then it is for ϕ as well. The inequality (2.4) holds for these two functions on \mathbb{R}^d . Then, notice that we can compare the discrete and continuous spherical averages as follows:

$$A_\tau f(x) \lesssim \tau A_\tau \phi(x). \quad (2.8)$$

Therefore, if we require that $\tau \leq N^{1+\epsilon}$, we see that (2.4) implies that $M_{1,1}$ satisfies (2.5).




Figure 3. The flow of the proof of (2.2). The nodes of the tree indicate the different elements of the decomposition, and a label on an arrow shows to which of M_1 or M_2 that term contributes. Above, λ represents a fixed choice of radius, and $\tau = \tau(x)$ an admissible choice of radius. For space considerations, several terms of the form $e_q(-\lambda^2 a)$ have been omitted; compare to (2.10).

The decomposition. Below, we assume that $\tau > N^{1+\epsilon}$ pointwise. At this point, we need a decomposition of $A_\lambda f$ into a family of multipliers. We recall this from [Magyar, Stein, and Wainger 2002]. Upper case letters denote a convolution operator, and lower case letters denote the corresponding multiplier. Let $e(x) = e^{2\pi i x}$ and for integers q , $e_q(x) = e(\frac{x}{q})$.

$$A_\lambda f = C_\lambda f + E_\lambda f, \quad (2.9)$$

$$C_\lambda f = \sum_{1 \leq q \leq \lambda} \sum_{a \in \mathbb{Z}_q^\times} e_q(-\lambda^2 a) C_\lambda^{a/q} f, \quad (2.10)$$

$$c_\lambda^{a/q}(\xi) = \hat{C}_\lambda^{a/q}(\xi) = \sum_{\ell \in \mathbb{Z}_q^d} G\left(\frac{a}{q}, \ell\right) \tilde{\psi}_q\left(\xi - \frac{\ell}{q}\right) \widetilde{d\sigma}_\lambda\left(\xi - \frac{\ell}{q}\right), \quad (2.11)$$

$$G\left(\frac{a}{q}, \ell\right) = q^{-d} \sum_{n \in \mathbb{Z}_q^d} e_q(|n|^2 a + n \cdot \ell).$$

The term $G(\frac{a}{q}, \ell)$ is a normalized Gauss sum. In (2.10), the sum over $a \in \mathbb{Z}_q^\times$ means that $(a, q) = 1$. In (2.11), the hat indicates the Fourier transform on \mathbb{Z}^d , and the notation conflates the operator $C_\lambda^{a/q}$ and

the kernel. All our operators are convolution operators or maximal operators formed in the same way. The function ψ is a Schwartz function on \mathbb{R}^d which satisfies

$$\mathbf{1}_{[-1/2, 1/2]}(|\xi|) \leq \tilde{\psi}(\xi) \leq \mathbf{1}_{[-1, 1]}(|\xi|).$$

Above, \tilde{f} denotes the Fourier transform of f on \mathbb{R}^d , and $\tilde{\psi}_q(\xi) = \tilde{\psi}(q\xi)$. The Fourier transform on \mathbb{R}^d of $d\sigma_\lambda$ is $\widehat{d\sigma}_\lambda$. Finally, we will use the notation λ for describing multipliers and so on, and using τ especially when obtaining estimates. In this way, many supremums will be suppressed from the notation.

The error term E_λ . The first contribution to M_2 is $M_{2,1} = |E_\tau f|$. The inequality below is from [Magyar, Stein, and Wainger 2002, Proposition 4.1], and it implies that $M_{2,1}$ satisfies (2.6) since $\tau > N^{1+\epsilon}$:

$$\left\| \sup_{\Lambda \leq \lambda \leq 2\Lambda} |E_\lambda \cdot| \right\|_{2 \rightarrow 2} \lesssim \Lambda^{(4-d)/2}, \quad \Lambda \geq 1. \quad (2.12)$$

Large denominators. The second contribution to M_2 is

$$M_{2,2} = \left| \sum_{N^{1+\epsilon} \leq q \leq \tau} e_q(-\lambda^2 a) C_\tau^{a/q} f \right|. \quad (2.13)$$

The estimate below is a result of [Magyar, Stein, and Wainger 2002, Proposition 3.1], and it verifies that $M_{2,2}$ satisfies (2.6). We need only sum it over $1 \leq a \leq q$, and $q > N^{1+\epsilon}$:

$$\left\| \sup_{\lambda > q} |C_\lambda^{a/q} f| \right\|_2 \lesssim q^{-d/2} \|f\|_2. \quad (2.14)$$

Small denominators: a secondary decomposition. It remains to bound the small denominator case, namely

$$\sum_{1 \leq q \leq N^{1+\epsilon}} \sum_{a \in \mathbb{Z}_q^\times} e_q(-\lambda^2 a) C_\tau^{a/q} f,$$

with further contributions to M_1 and M_2 . Write $C_\tau^{a/q} = C_\tau^{a/q,1} + C_\tau^{a/q,2}$, with this understanding. For an integer $1 \leq Q \leq \frac{1}{2}N$, and $Q \leq q < 2Q$, define

$$\hat{C}_\lambda^{a/q,1}(\xi) = \sum_{\ell \in \mathbb{Z}^d} G(a, \ell, q) \tilde{\psi}_q\left(\xi - \frac{\ell}{q}\right) \tilde{\psi}_{\lambda Q/N}\left(\xi - \frac{\ell}{q}\right) \widehat{d\sigma}_\lambda\left(\xi - \frac{\ell}{q}\right). \quad (2.15)$$

Above, we have adjusted the cutoff around each point $\frac{\ell}{q} \in \mathbb{T}^d$.

Small denominators: the ℓ^2 -part. We complete the construction of the term in M_2 , (2.6), by showing that

$$\left\| \sup_{N^{1+\epsilon} \leq \lambda \leq \ell(E)} |C_\lambda^{a/q,2} f| \right\|_2 \lesssim q^{-1} N^{-(d-2)/2} \|f\|_2. \quad (2.16)$$

It follows that

$$\left\| \sum_{1 \leq q \leq N^{1+\epsilon}} \sum_{a \in \mathbb{Z}_q^\times} e_q(-\lambda^2 a) C_\tau^{a/q,2} f \right\|_2 \lesssim N^{-(d-4)/2+\epsilon} \|f\|_2. \quad (2.17)$$

This is the third and final contribution to M_2 . We remark that the proof detailed below is a quantitative variant of the proof of Magyar, Stein and Wainger's inequality (2.14). The inequality (2.16) is [Ionescu 2004, (2.14)], but we include details here.

Let m be a smooth function supported on $[-\frac{1}{2}, \frac{1}{2}]^d$, and let T_m be the corresponding multiplier operator, either on \mathbb{Z}^d or \mathbb{R}^d , with the notation indicating in which setting we are considering the multiplier.

This is a factorization argument from [Magyar, Stein, and Wainger 2002, p. 200]. Using the notation of (2.23) to define $M_{\psi_{Q/2}}$ below, for $Q \leq q \leq 2Q$, we have

$$\begin{aligned} \hat{C}_\lambda^{a/q,2}(\xi) &= \hat{M}_{\psi_{Q/2},q}(\xi) \cdot \sum_{\ell \in \mathbb{Z}^d} \tilde{\psi}_q\left(\xi - \frac{\ell}{q}\right) \left(1 - \tilde{\psi}_{\lambda Q/N}\left(\xi - \frac{\ell}{q}\right)\right) \widetilde{d\sigma}_\lambda\left(\xi - \frac{\ell}{q}\right) \\ &:= \hat{M}_{\psi_{q/2},q}(\xi) \cdot \hat{C}_\lambda^{a/q,3}(\xi). \end{aligned}$$

That is, the operator in question factors as $C_\lambda^{a/q,2} = C_\lambda^{a/q,3} \circ M_{\psi_{Q/2},q}$. Notice that by the Gauss sum estimate, we have

$$\|M_{\psi_{Q/2},q}\|_{2 \rightarrow 2} \lesssim Q^{-d/2}. \quad (2.18)$$

In controlling the supremum, we need only consider the supremum over $C_\lambda^{a/q,3}$. The transference lemma [Magyar, Stein, and Wainger 2002, Corollary 2.1] allows us to estimate this supremum on $L^2(\mathbb{R}^d)$. We have

$$\|C_\tau^{a/q,3}\|_{\ell^2(\mathbb{Z}^d) \rightarrow \ell^2(\mathbb{Z}^d)} \lesssim \left\| \sup_{\lambda > 0} |\Psi_{\lambda,q} \cdot| \right\|_{L^2(\mathbb{R}^d) \rightarrow L^2(\mathbb{R}^d)}, \quad (2.19)$$

where $\tilde{\Psi}_{\lambda,q} = \tilde{\psi}_q(1 - \tilde{\psi}_{\lambda Q/N}) \widetilde{d\sigma}_\lambda$. To estimate this last norm on $L^2(\mathbb{R}^d)$, we use this lemma of Bourgain:

Lemma 2.20 [Bourgain 1985, Proposition 2]. *Let m be a smooth function on \mathbb{R}^d . We have*

$$\left\| \sup_{r > 0} |T_m(r \cdot) \cdot| \right\|_2 \lesssim \sum_{j \in \mathbb{Z}} \alpha_j^{1/2} (\alpha_j^{1/2} + \beta_j^{1/2}),$$

where

$$\alpha_j = \|\mathbf{1}_{2^j \leq |\xi| \leq 2^{j+1}} m(\xi)\|_\infty \quad \text{and} \quad \beta_j = \|\mathbf{1}_{2^j \leq |\xi| \leq 2^{j+1}} \nabla m(\xi) \cdot \xi\|_\infty. \quad (2.21)$$

We bound the right side of (2.19). Composition with T_{ψ_q} is uniformly bounded on L^2 . The multiplier in question is then $m(\xi) = (1 - \tilde{\psi}_{Q/N})(\xi) \widetilde{d\sigma}_1(\xi)$. This is identically zero for $|\xi| \lesssim \frac{N}{Q}$. That means that for the terms in (2.21), we need only consider $2^j \gtrsim \frac{N}{Q} \geq 100$. Recall the standard stationary phase estimate

$$|\nabla \widetilde{d\sigma}_1(\xi)| + |\widetilde{d\sigma}_1(\xi)| \lesssim |\xi|^{-(d-1)/2}.$$

Hence, the bound for our multiplier is

$$\left\| \sup_{\lambda > 0} |\Psi_{\lambda,q} \cdot| \right\|_{L^2(\mathbb{R}^d) \rightarrow L^2(\mathbb{R}^d)} \lesssim \sum_{j: 2^j \geq N/Q} 2^{-j(d-1)/2} \cdot 2^{-j(d-3)/2} \lesssim \left(\frac{Q}{N}\right)^{(d-2)/2}.$$

This estimate combined with (2.18) and (2.19) completes the proof of (2.16).

Small denominators: the sparse part. Recalling the notation from (2.15), we turn to

$$M_{1,2}f = \sum_{1 \leq q \leq N} \sum_{a \in \mathbb{Z}_q^\times} e_q(-\lambda^2 a) C_\tau^{a/q,1} f$$

and show that this term is as in (2.7). This is the term in which the absorbing term $\frac{1}{2}A_\tau f$ in (2.7) arises. It is also the core of the proof.

Define

$$M_{1,Q}f = \sum_{Q \leq q < 2Q} \sum_{a \in \mathbb{Z}_q^\times} e_q(-\lambda^2 a) C_\tau^{a/q,1} f, \quad 1 \leq Q \leq \frac{1}{2}N,$$

Above, and below, we will treat $M_{1,Q}$ as an operator, as we have yet to tease out some of its additional properties. The main estimates to prove are

$$\begin{aligned} M_{1,Q}f &\leq \overline{M}_{1,Q}f + N^{-\epsilon} A_\tau f, \\ |E|^{-1} \langle \overline{M}_{1,Q}f, g \rangle &\lesssim N^{1+\epsilon/2} \langle f \rangle_E^{1/(\bar{p}+\epsilon)} \langle g \rangle_E^{1/(\bar{q}+\epsilon)}. \end{aligned} \quad (2.22)$$

These are summed over dyadic $1 \leq Q \leq \frac{1}{2}N$ to complete the proof of the absorption inequality (2.7). This step requires that N be sufficiently large, $N > \kappa^{1/\epsilon}$, but that is sufficient for our purposes.

We need the estimate (2.4) on \mathbb{R}^d . We also need kernel estimates for the operators $M_{1,Q}$, and for that we require this preparation, which has been noted before [Magyar, Stein, and Wainger 2002; Ionescu 2004, p. 1415]. For a function ζ with $\tilde{\zeta}$ supported on $[-1, 1]^d$, define a family of Fourier multipliers by

$$\hat{M}_{\tilde{\zeta},q}(\xi) = \sum_{\ell \in \mathbb{Z}_q^d} G\left(\frac{a}{q}, \ell\right) \tilde{\zeta}\left(\xi - \frac{\ell}{q}\right).$$

By inspection, the Gauss sum map $\ell \mapsto G\left(\frac{a}{q}, \ell\right)$ is the Fourier transform of $e_q(|x|^2 a)$ as a function on \mathbb{Z}_q^d . From this, and a routine computation, it follows that

$$M_{\tilde{\zeta},q}(x) = e_q(a|x|^2) \zeta(x). \quad (2.23)$$

(Here we identify the kernel of the convolution operator, and the operator itself.)

It follows that the kernel of $M_{1,Q}f$ is

$$\begin{aligned} M_{1,Q}(n) &= \psi_{\tau Q/N} * d\sigma_\tau(n) \sum_{Q \leq q \leq 2Q} \sum_{a \in \mathbb{Z}_q^\times} e_q(a(|n|^2 - \lambda^2)) \\ &= P_{Q,\tau}(n) \cdot C_Q(|n|^2 - \lambda^2). \end{aligned} \quad (2.24)$$

Note that $P_{Q,\tau}$ is a maximal average over annuli of outer radius τ , and width about $\tau \frac{Q}{N} < \tau$. The second term above is related to Ramanujan sums, defined by

$$c_q(m) = \sum_{a \in \mathbb{Z}_q^\times} e_q(am), \quad m \in \mathbb{Z},$$

so that in (2.24), $C_Q = \sum_{Q \leq q \leq 2Q} c_q$. Ramanujan sums satisfy very good cancellation properties. The properties we will need are summarized in the next result.

Lemma 2.25. *The following estimates hold for any $k \in \mathbb{N}$, and $\epsilon > 0$:*

(1) *For any Q and n , $|C_Q(n)| \leq Q^2$.*

(2) *There holds*

$$\max_{0 < m \leq Q^k} |C_Q(m)| \lesssim Q^{1+\epsilon}. \quad (2.26)$$

(3) *For $M > Q^k$,*

$$\left[\frac{1}{M} \sum_{m \leq M} |C_Q(m)|^k \right]^{1/k} \lesssim Q^{1+\epsilon}. \quad (2.27)$$

The implied constants depend upon k and ϵ .

Proof. The first estimate is trivial, but we include it for the sake of clarity. Note that $C_Q(0) \simeq Q^2$, which will arise in the absorption argument below.

An argument for the second inequality (2.26) begins with the inequality $|c_q(m)| \leq (q, m)$ for $m > 0$. This can be checked by inspection if q is a power of a prime. The general case follows as both sides are multiplicative functions.

Then, of course we have, for any $1 \leq d \leq q$,

$$\sum_{k: dk \leq Q} d \leq Q.$$

It follows that

$$\sum_{q \leq Q} (q, m) \leq Q \sum_{d \leq q: d | m} 1 = Q\delta(m; Q),$$

where $\delta(m; Q)$ is the number of divisors of m that are less than or equal to Q . But, $m \leq Q^k$, so by a well-known logarithmic-type estimate for the divisor function, we have (2.26).

The third property is harder. It is due to [Bourgain 1993]. There are proofs in [Kesler, Lacey, and Mena 2019, Lemma 2.13] and [Kesler 2018b, Lemma 5]. \square

A tertiary decomposition. The preparations are finished. It remains to prove (2.22), and this argument is indicated in Figure 4. There are three cases, namely,

(1) $Q < N^{1/2}$,

(2) $NQ^{k_1-1} < \tau$, where $k_1 = k_1(\bar{p}, \bar{q})$,

(3) $N^{1/2} \leq Q$ and $\tau \leq NQ^{k_1-1}$, implying $N < \tau < Q^{k_2}$, where $k_2 = k_2(\bar{p}, \bar{q})$.

We treat these cases in order, with the core case being the last one.

The first and easiest case concerns $Q < N^{1/2}$. Using the trivial bound $|c_q(n)| \leq q$ and (2.24) we have

$$|M_{1,Q}f| < N \cdot P_{Q,\tau}f.$$

It then follows from the continuous sparse bound (2.4) that we have

$$|E|^{-1} \langle M_{1,Q}f, g \rangle \lesssim N \langle f \rangle_E^{1/\bar{p}} \langle g \rangle^{1/\bar{q}}. \quad (2.28)$$

This is as required in (2.22).




Figure 4. The flow of the proof of (2.22). The integers k_1 and k_2 are large and functions of ϵ , \bar{p} and \bar{q} . The first level of the decomposition is motivated by the estimates for Ramanujan sums in Lemma 2.25. The second level of the diagram is associated with the Ramanujan estimate (2.26). It requires a further decomposition of the kernel $P_{\tau, Q}$ in (2.24).

The second case we restrict to the case that $\tau > NQ^{k_1-1}$ for a sufficiently large integer k_1 that is a function of (\bar{p}, \bar{q}) . This case does not have an absorbing term.

Dominate, using Hölder's inequality with $\ell^{k_1} - \ell^{k'_1}$ duality,

$$\begin{aligned}
 |M_{1, Q} f(n)| &\lesssim [P_{Q, \tau} * f(n)]^{1/k'_1} \left[\sum_{x \in \mathbb{Z}^d} |C_Q(|x|^2 - \lambda^2)|^{k_1} P_{Q, \tau}(x) \right]^{1/k_1} \\
 &\lesssim Q^{1+\epsilon/2} P_{Q, \tau} * f(n)^{1/k'_1}.
 \end{aligned} \tag{2.29}$$

Recall that f is an indicator function. Notice that we are using a bound on the Ramanujan sums that follows from (2.27). Recall that $P_{Q, \tau}$ is an average over an annulus around the sphere of radius λ , of width $\tau \frac{Q}{N}$. In particular, the width is greater than Q^{k_1} by the assumption that $\tau > NQ^{k_1-1}$.

But, then, we are free to conclude our statement; since $(\frac{1}{\bar{p}}, \frac{1}{\bar{q}})$ are in the interior of \mathbf{R}_d , we have, for k_1 sufficiently large, so that k'_1 is sufficiently close to 1, as required in (2.22),

$$\begin{aligned}
 |E|^{-1} \langle M_1 f, g \rangle &\lesssim Q^{1+\epsilon} |E|^{-1} \langle [P_{Q, \tau} * f]^{1/k'_1}, g \rangle \\
 &\lesssim N [|E|^{-1} \langle P_{Q, \tau} * f, g \rangle]^{1/k'_1} \\
 &\lesssim \langle f \rangle_E^{1/(\bar{p}k'_1)} \langle g \rangle_E^{1/(\bar{q}k'_1)}.
 \end{aligned} \tag{2.30}$$

Here, we use (2.29) and then the real-variable inequality (2.4), which we can do if k_1 is sufficiently large, so that $(k'_1 \bar{p}, k'_1 \bar{q}) \in \mathbf{R}_d$. This is our second application of (2.4).

We turn to third case of $N < \tau < Q^{k_2}$. A final, fourth decomposition of $P_{\tau, Q}$ is needed, and the absorption argument appears. Let $\mathbb{S}_\lambda = \{n \in \mathbb{Z}^d : |n| = \lambda\}$ be the integer sphere of radius λ , and set

$$P_{\tau, Q} = \sum_{j=0}^2 P_{\tau, Q}^j,$$

where

$$\begin{aligned} P_{\tau,Q}^0(n) &= P_{\tau,Q}(n) \mathbf{1}_{\mathbb{S}_\tau}(n), \\ P_{\tau,Q}^1(n) &= P_{\tau,Q}(n) \mathbf{1}_{0 < \text{dist}(n, \mathbb{S}_\tau) < \tau Q^{1+\epsilon}/N}, \end{aligned} \quad (2.31)$$

and $P_{\tau,Q}^2$ is then defined. The term $P_{\tau,Q}^0$ is a multiple of the average over the integer sphere of radius λ , and the term $P_{\tau,Q}^1$ is just that part of $P_{\tau,Q}$ that is close to, but not equal to, the sphere of radius \mathbb{S}_τ .

Let us detail the absorbing term. Note that $C_Q(0) \simeq Q^2$. Indeed, we have to single out this case as there is no cancellation in the Ramanujan sum when the argument is zero. Using the definition (2.31), we have

$$\begin{aligned} |P_{\tau,Q}^0(n) \cdot C_Q(|n|^2 - \tau^2)| &\lesssim Q^2 \frac{N}{Q\tau^d} \mathbf{1}_{\mathbb{S}_\tau}(n) \\ &\lesssim \frac{NQ}{\tau^2} \cdot \tau^{2-d} \mathbf{1}_{\mathbb{S}_\tau}(n) \lesssim N^{-2\epsilon} \cdot \tau^{2-d} \mathbf{1}_{\mathbb{S}_\tau}(n). \end{aligned} \quad (2.32)$$

This is as required in (2.7) and (2.22).

The inequality (2.26) on the Ramanujan sums applies in the analysis of $P_{\tau,Q}^1$, due to our assumptions $Q < \tau < Q^{k_2}$. It shows that

$$|P_{Q,\tau}^1(n) \cdot C_Q(|n|^2 - \tau^2)| \lesssim Q^{1+\epsilon} \frac{N}{\tau^d Q} \mathbf{1}_{0 < \text{dist}(n, \mathbb{S}_\tau) < \tau Q^{1+\epsilon}/N}.$$

Keeping normalizations in mind, it follows from our real-variable sparse inequality (2.4) that

$$|E|^{-1} \langle P_{Q,\tau}^1 * f, g \rangle \lesssim Q^{1+2\epsilon} \langle f \rangle_E^{1/\bar{p}} \langle g \rangle_E^{1/\bar{q}}. \quad (2.33)$$

This is as required in (2.22).

The last term is $P_{Q,\tau}^2$. As noted, $|C_Q(m)| \leq Q^2$. The condition $\tau < Q^{k_2}$, and simple Schwartz tail considerations then show that

$$|P_{Q,\tau}^2(n) \cdot C_Q(|n|^2 - \tau^2)| \lesssim \frac{Q}{\tau^d} \left[1 + \frac{|n|}{\tau} \right]^{-2d}. \quad (2.34)$$

Since τ is an admissible stopping time, it follows that

$$|P_{Q,\tau}^1 * f| \lesssim \langle f \rangle_E.$$

This completes the proof of (2.22).

3. The endpoint sparse bound

We need this definition. Given a cube E , we say that collection \mathcal{Q}_E of subcubes $Q \subset E$ are *presparse* if the cubes $\{\frac{1}{3}Q : Q \in \mathcal{Q}_E\}$ are pairwise disjoint. Associated to a presparse collection \mathcal{Q}_E is a family of stopping times. We say that τ is \mathcal{Q}_E admissible (or just admissible) if

$$\ell(E) \geq \tau_{\mathcal{Q}_E}(x) = \tau(x) \geq \max\{1, \ell(Q) \mathbf{1}_{Q/3} : Q \in \mathcal{Q}_E\}, \quad x \in E.$$

The relevant lemma is this:




Figure 5. The flow of the proof of (3.3)–(3.5). The notation is similar to Figure 3.

Lemma 3.1. *For $f = \mathbf{1}_F$ supported on the cube $3E$, there is a presparse collection \mathcal{Q}_E so that for all \mathcal{Q}_E -admissible $\tau = \tau(x)$, and all $g = \mathbf{1}_G$ supported on E , we have*

$$\langle A_\tau f, g \rangle \lesssim \langle f \rangle_{3E, d/(d-2)} \langle g \rangle_{E, d/(d-2)} |E|. \quad (3.2)$$

The lemma follows from this: for integers $N > 1$, there is a decomposition

$$A_\tau f \leq M_1 + M_2, \quad (3.3)$$

$$\langle M_1, g \rangle \lesssim N^2 \langle f \rangle_{3E, 1} \langle g \rangle_{E, 1} |E|, \quad (3.4)$$

$$\langle M_2, g \rangle \lesssim N^{-(d-4)/2} \langle f \rangle_{3E, 2} \langle g \rangle_{E, 2} |E|. \quad (3.5)$$

Recalling that $f = \mathbf{1}_F$ and $g = \mathbf{1}_G$, the right sides above are comparable for $N \simeq [\langle f \rangle_{3E, 1} \langle g \rangle_{E, 1}]^{-1/d}$, and this proves (3.2).

It remains to prove (3.3)–(3.5). Our proof will be much shorter because we do not need to compare to the very rough continuous spherical averages, but to averages over balls. (In particular, we will not need any subtle facts about Ramanujan sums.) A guide to the argument is in Figure 5. The decomposition has several elements. The first begins with the trivial bound $A_\lambda f(x) \lesssim \lambda^2 B_\lambda * f(x)$, where B_λ is the average of over a ball of radius λ . Our first contribution to M_1 is

$$M_{1,1} = \mathbf{1}_{\tau(x) \leq 100N} A_\tau f,$$

which is pointwise bounded by $CN^2\langle f \rangle_{3E,1}$, by choice of \mathcal{Q}_E . Thus (3.4) holds for this term. Below, we are free to assume that $\tau \geq 100N$.

Recall the decomposition of $A_\lambda f$, beginning with (2.9). Our first contribution to M_2 is $M_{2,1} = |E_\tau f|$. This satisfies (3.5) by (2.12).

It remains to bound $C_\tau f$ as defined in (2.10). This requires further contributions to M_1 and M_2 . Apply (2.14) to see that this term obeys (3.5):

$$M_{2,2} = \sum_{N/100 \leq q \leq \lambda} \sum_{a \in \mathbb{Z}_q^\times} |C_\tau^{a/q} f|.$$

The remaining terms are

$$\sum_{1 \leq q \leq N/100} \sum_{a \in \mathbb{Z}_q^\times} e_q(-\lambda^2 a) C_\tau^{a/q} f.$$

Control will consist of an additional contribution to M_1 and M_2 .

Write $C_\lambda^{a/q} = C_\lambda^{a/q,1} + C_\lambda^{a/q,2}$, where a different cut-off in frequency is inserted:

$$\begin{aligned} \hat{C}_\lambda^{a/q,1}(\xi) &= \sum_{\ell \in \mathbb{Z}^d} G\left(\frac{a}{q}, \ell\right) \tilde{\psi}_q\left(\xi - \frac{\ell}{q}\right) \tilde{\psi}_{\lambda q/N}\left(\xi - \frac{\ell}{q}\right) \widetilde{d\sigma}_\lambda\left(\xi - \frac{\ell}{q}\right) \\ &= \sum_{\ell \in \mathbb{Z}^d} G\left(\frac{a}{q}, \ell\right) \tilde{\psi}_{\lambda q/N}\left(\xi - \frac{\ell}{q}\right) \widetilde{d\sigma}_\lambda\left(\xi - \frac{\ell}{q}\right). \end{aligned}$$

This follows from the definition of ψ in (2.11). This is slightly different from (2.15); in particular, the term below satisfies (3.5), just as in the previous section:

$$M_{2,3} = \left| \sum_{1 \leq q \leq N/100} \sum_{a \in \mathbb{Z}_q^\times} e_q(-\lambda^2 a) C_\tau^{a/q,2} f \right|.$$

We claim that

$$\left\| \sum_{1 \leq q \leq N/100} \sum_{a \in \mathbb{Z}_q^\times} e_q(-\lambda^2 a) C_\tau^{a/q,1} f \right\|_\infty \lesssim N^2 \langle f \rangle_{E,1}. \quad (3.6)$$

That is, the term on the left is our second and final contribution to M_1 , as in (3.4).

Recalling (2.23) and (2.24), we have

$$|C_\lambda^{a/q,1}(n)| \lesssim d\sigma_\lambda * \psi_{\lambda q/N}(n)$$

The convolution is with $d\sigma_\lambda$ and $\psi_{\lambda q/N}$, which is a bump function of integral 1, supported on scale $\frac{N}{\lambda q}$, which is much smaller than λ . As a consequence, we have

$$|C_\lambda^{a/q,1}(n)| \lesssim \frac{N}{q} \cdot \lambda^{-d} \left[1 + \frac{|n|}{\lambda} \right]^{-3d}.$$

This is summed over $1 \leq a < q \leq \frac{1}{100}N$. And, one appeals to the admissibility of the stopping time τ to complete the proof of (3.6).

4. Interpolation of sparse bounds

We show that if a sublinear operator satisfies an open range of sparse bounds for indicator sets, then they improve to sparse bounds for functions.

Lemma 4.1. *Suppose that a sublinear operator T satisfies the bound below for $1 < p, q < \infty$. For a fixed function f and all $|g| \leq \mathbf{1}_G$, there is a sparse collection \mathcal{S} so that*

$$|\langle Tf, g \rangle| \lesssim \Lambda_{\mathcal{S}, p, q}(f, \mathbf{1}_G).$$

Then,

$$|\langle Tf, g \rangle| \lesssim \sup_{\mathcal{S}} \Lambda_{\mathcal{S}, p, q}(f, g), \quad q < r < \infty.$$

Proof. In this proof, we will work on \mathbb{R}^d , with the same proof working on \mathbb{Z}^d . We will also assume that (1) T is a positive operator, and (b) all cubes are dyadic. The general case is not much harder than these considerations. Let f be a fixed function, and let g be a bounded compactly supported function. We will show that there is a sparse collection \mathcal{S} so that

$$\langle Tf, g \rangle \lesssim \sum_{Q \in \mathcal{S}} \langle f \rangle_Q^{1/p} \langle g \rangle_{Q, q, 1} |Q|, \quad (4.2)$$

where $\langle g \rangle_{Q, q, 1} = \|g \mathbf{1}_Q\|_{L^{q, 1}(dx/|Q|)}$ is the Lorentz space with normalized measure. Since $\langle g \rangle_{Q, q, 1} < \langle g \rangle_{Q, r}$ for $1 < r < q$, this completes the proof of the lemma.

The argument for (4.2) is a level set argument. Thus, write $g \leq \sum_{k \in \mathbb{Z}} 2^k \mathbf{1}_{G_k}$ for disjoint sets G_k . Apply the assumed sparse bound for indicators for each pair of sets (F, G_k) . We get a sequence of sparse sets \mathcal{S}_k so that

$$\langle Tf, g \rangle \lesssim \sum_{k \in \mathbb{Z}} \Lambda_{\mathcal{S}_k, p, q}(f, \mathbf{1}_{G_k}). \quad (4.3)$$

Let \mathcal{F} be a sequence of stopping cubes for the averages $\langle f \rangle_{Q, p}$. That is, we choose \mathcal{F} so that for any dyadic cube Q there is a dyadic cube in \mathcal{F} that contains Q . And setting Q^a to be the minimal such cube in \mathcal{F} , we have $\langle f \rangle_{Q, p} \lesssim \langle f \rangle_{Q^a, p}$. The sum in (4.3) is organized according to \mathcal{F} . Below, we take $q < \alpha < \infty$, very close to q . For each $P \in \mathcal{F}$ we have

$$\begin{aligned} \Gamma(P) &= \sum_{k \in \mathbb{Z}} 2^{-k} \sum_{\substack{Q \in \mathcal{S}_k \\ Q^a = P}} \langle \mathbf{1}_{G_k} \rangle_Q^{1/q} |Q| \lesssim \sum_{k \in \mathbb{Z}} 2^{-k} |P|^{1/\alpha'} \left[\sum_{\substack{Q \in \mathcal{S}_k \\ Q^a = P}} \langle \mathbf{1}_{G_k} \rangle_Q^{\alpha/q} \right]^{1/\alpha} \\ &\lesssim |P| \sum_{k \in \mathbb{Z}} 2^{-k} \langle \mathbf{1}_{G_k} \rangle_P^{1/q} = \langle g \rangle_{L^{q, 1}(P)} |P|. \end{aligned} \quad (4.4)$$

Here, we have used Hölder's inequality, sparseness of the collections \mathcal{S}_k and the Carleson embedding inequality. The last inequality is one way to define the $L^{q, 1}$ norm.

And, so we have

$$(4.3) \lesssim \sum_{P \in \mathcal{F}} \langle f \rangle_{P, p} \Gamma(P).$$

The bound in (4.4) then implies (4.2). \square

Concerning our Theorem 1.2. For the first assertion, we have proved a restricted weak-type inequality for all $(\frac{1}{p}, \frac{1}{q})$ in the interior of Z_d . We see that we can then replace the indicator functions in both coordinates by functions. That is, we have $\langle A \rangle_{p,q} < \infty$ for all $(\frac{1}{p}, \frac{1}{q})$ in the interior of Z_d .

Concerning the second assertion, we have the restricted weak-type inequality at $(\frac{d}{d-2}, \frac{d}{d-2})$. We see from the theorem above that we have

$$\langle Af, g \rangle \lesssim \sup_S \sum_{Q \in S} \langle f \rangle_{Q, d/(d-2), 1} \langle g \rangle_{Q, q} |Q|, \quad q > \frac{d}{d-2}.$$

5. Counterexamples

We can show the following proposition, showing necessary conditions on (p, q) for the sparse bound to hold. The gap between the sufficient conditions for a sparse bound and these necessary conditions is illustrated in Figure 6.

Proposition 5.1. *If the sparse bound $\|A\|_{p,q} < \infty$ holds, we have*

$$\frac{1}{p_1} \leq \frac{d-2}{d}, \quad \frac{2}{p} + \frac{d-1}{q} \leq d. \quad (5.2)$$

Proof. If a (p, q) -sparse bound held we would conclude that $A : \ell^p \rightarrow \ell^{p,\infty}$. Taking $f = \mathbf{1}_0$, note that $Af(x) \simeq (1 + |x|)^{2-d}$. The latter function is $\ell^{d/(d-2),\infty}$. Hence, $p_1 \geq \frac{d}{d-2}$ is necessary.

For the second inequality in (5.2), set $\mathbb{S}_\lambda = \{n \in \mathbb{Z}^d : |n| = \lambda\}$. We recall that for $d \geq 4$, and any odd choice of $\lambda^2 \in \mathbb{N}$, we have $|\mathbb{S}_\lambda| \simeq \lambda^{d-2}$. For an odd choice of $\lambda^2 \in \mathbb{N}$, let $f = \mathbf{1}_{\mathbb{S}_\lambda}$, and consider $G = \{Af > \frac{c}{\lambda}\}$. This is the set of $x \in \mathbb{Z}^d$ for which the two spheres \mathbb{S}_λ and $x + \mathbb{S}_\mu$, for some choice of $\mu \simeq \lambda$, have about the expected size. Here, necessarily $G \subset E$, a cube centered at the origin of side length about λ .

We claim that $|G| \gtrsim \lambda$. And observe that $Af(x) \gtrsim \lambda^{-1}$ for $x \in G$. Moreover, from the assumed (p, q) -sparse bound,

$$\lambda^{-1} \langle \mathbf{1}_G \rangle_E \lesssim \lambda^{-d} \langle Af, g \rangle \lesssim \langle f \rangle_E^{1/p} \langle \mathbf{1}_G \rangle_E^{1/q} \lesssim \lambda^{-2/p} \langle \mathbf{1}_G \rangle_E^{1/q}. \quad (5.3)$$




Figure 6. The horizontal line is set at $\frac{1}{q} = \frac{d-2}{d}$ for reasons of clarity. Sparse bounds hold below the solid line from $(0, 1)$, to Z_2 to $Z_1 = (\frac{d-2}{d}, \frac{d-2}{d})$. (Recall that Z_2 is defined in (1.5).) They cannot hold to the right of Z_1 , nor above the dotted line. The gray area is unresolved.

For this pair of functions, it is easy to see that the maximal sparse form is the expression on the right. Our lower bound on the size of G proves the proposition.

Note that since $d \geq 5$, and λ^2 is odd, there are about λ^{d-3} choices of vectors $y = (0, y_2, \dots, y_d) \in \mathbb{S}_\lambda$. (We insist on λ^2 being odd to capture the case of $d = 5$ here.) Then, if $|x_1| \leq \frac{1}{2}\lambda$, note that

$$\|(0, y_2, \dots, y_d) - (x_1, 0, \dots, 0)\| = \sqrt{\lambda^2 + x_1^2} = \lambda'.$$

From that, it follows that

$$A_{\lambda'} f(x_1, 0, \dots, 0) \simeq \lambda^{-1}.$$

This shows that $|G| \gtrsim \lambda$. □

Observe that the $(\frac{d}{d-2}, \frac{d}{d-2})$ -sparse bound and (5.3) imply that $|G| \lesssim \lambda^{(d+4)/2}$. Our lower bound is certainly not sharp. What is the correct size of G ?

References

- [Bourgain 1985] J. Bourgain, “Estimations de certaines fonctions maximales”, *C. R. Acad. Sci. Paris Sér. I Math.* **301**:10 (1985), 499–502. MR
- [Bourgain 1993] J. Bourgain, “Fourier transform restriction phenomena for certain lattice subsets and applications to nonlinear evolution equations, I: Schrödinger equations”, *Geom. Funct. Anal.* **3**:2 (1993), 107–156. MR Zbl
- [Cook 2018] B. Cook, “A note on discrete spherical averages over sparse sequences”, 2018. To appear in *Proc. Amer. Math. Soc.* arXiv
- [Hughes 2018] K. Hughes, “ ℓ^p -improving for discrete spherical averages”, preprint, 2018. arXiv
- [Hughes 2019] K. Hughes, “The discrete spherical averages over a family of sparse sequences”, *J. Anal. Math.* (online publication July 2019).
- [Ionescu 2004] A. D. Ionescu, “An endpoint estimate for the discrete spherical maximal function”, *Proc. Amer. Math. Soc.* **132**:5 (2004), 1411–1417. MR Zbl
- [Kesler 2018a] R. Kesler, “ $\ell^p(\mathbb{Z}^d)$ -improving properties and sparse bounds for discrete spherical maximal averages”, 2018. To appear in *J. Anal. Math.* arXiv
- [Kesler 2018b] R. Kesler, “ $\ell^p(\mathbb{Z}^d)$ -improving properties and sparse bounds for discrete spherical maximal means, revisited”, preprint, 2018. arXiv
- [Kesler and Lacey 2018] R. Kesler and M. T. Lacey, “ ℓ^p -improving inequalities for discrete spherical averages”, preprint, 2018. arXiv
- [Kesler, Lacey, and Mena 2019] R. Kesler, M. T. Lacey, and D. Mena Arias, “Lacunary discrete spherical maximal functions”, *New York J. Math.* **25** (2019), 541–557.
- [Lacey 2017] M. T. Lacey, “Sparse bounds for spherical maximal functions”, 2017. To appear in *J. Anal. Math.* arXiv
- [Lee 2003] S. Lee, “Endpoint estimates for the circular maximal function”, *Proc. Amer. Math. Soc.* **131**:5 (2003), 1433–1442. MR Zbl
- [Magyar 1997] A. Magyar, “ L^p -bounds for spherical maximal operators on \mathbb{Z}^n ”, *Rev. Mat. Iberoam.* **13**:2 (1997), 307–317. MR Zbl
- [Magyar, Stein, and Wainger 2002] A. Magyar, E. M. Stein, and S. Wainger, “Discrete analogues in harmonic analysis: spherical averages”, *Ann. of Math. (2)* **155**:1 (2002), 189–208. MR Zbl
- [Schlag 1997] W. Schlag, “A generalization of Bourgain’s circular maximal theorem”, *J. Amer. Math. Soc.* **10**:1 (1997), 103–122. MR Zbl
- [Schlag and Sogge 1997] W. Schlag and C. D. Sogge, “Local smoothing estimates related to the circular maximal theorem”, *Math. Res. Lett.* **4**:1 (1997), 1–15. MR Zbl

Received 8 Apr 2019. Accepted 5 Jul 2019.

ROBERT KESLER: robertmkessler@gmail.com

School of Mathematics, Georgia Institute of Technology, Atlanta, GA, United States

MICHAEL T. LACEY: lacey@math.gatech.edu

School of Mathematics, Georgia Institute of Technology, Atlanta, GA, United States

DARÍO MENA: dario.menaarias@ucr.ac.cr

Escuela de Matemática, Universidad de Costa Rica, San José, Costa Rica

PURE and APPLIED ANALYSIS

msp.org/paa

EDITORS-IN-CHIEF

Charles L. Epstein	University of Pennsylvania cle@math.upenn.edu
Maciej Zworski	University of California at Berkeley zworski@math.berkeley.edu

EDITORIAL BOARD

Sir John M. Ball	University of Oxford ball@maths.ox.ac.uk
Michael P. Brenner	Harvard University brenner@seas.harvard.edu
Charles Fefferman	Princeton University cf@math.princeton.edu
Susan Friedlander	University of Southern California susanfri@usc.edu
Anna Gilbert	University of Michigan annacg@umich.edu
Leslie F. Greengard	Courant Institute, New York University, and Flatiron Institute, Simons Foundation greengard@cims.nyu.edu
Yan Guo	Brown University yan_guo@brown.edu
Claude Le Bris	CERMICS - ENPC lebris@cermics.enpc.fr
Robert J. McCann	University of Toronto mccann@math.toronto.edu
Michael O'Neil	Courant Institute, New York University oneil@cims.nyu.edu
Jill Pipher	Brown University jill_pipher@brown.edu
Johannes Sjöstrand	Université de Dijon johannes.sjostrand@u-bourgogne.fr
Vladimir Šverák	University of Minnesota sverak@math.umn.edu
Daniel Tataru	University of California at Berkeley tataru@berkeley.edu
Michael I. Weinstein	Columbia University miw2103@columbia.edu
Jon Wilkening	University of California at Berkeley wilken@math.berkeley.edu
Enrique Zuazua	DeustoTech-Bilbao, and Universidad Autónoma de Madrid enrique.zuazua@deusto.es

PRODUCTION

Silvio Levy	(Scientific Editor) production@msp.org
-------------	---

Cover image: The figure shows the outgoing scattered field produced by scattering a plane wave, coming from the northwest, off of the (stylized) letters P A A. The total field satisfies the homogeneous Dirichlet condition on the boundary of the letters. It is based on a numerical computation by Mike O'Neil of the Courant Institute.

See inside back cover or msp.org/paa for submission instructions.

The subscription price for 2020 is US \$505/year for the electronic version, and \$565/year (+\$25, if shipping outside the US) for print and electronic. Subscriptions, requests for back issues and changes of subscriber address should be sent to MSP.

Pure and Applied Analysis (ISSN 2578-5885 electronic, 2578-5893 printed) at Mathematical Sciences Publishers, 798 Evans Hall #3840, c/o University of California, Berkeley, CA 94720-3840 is published continuously online. Periodical rate postage paid at Berkeley, CA 94704, and additional mailing offices.

PAA peer review and production are managed by EditFlow® from MSP.

PUBLISHED BY

 **mathematical sciences publishers**
nonprofit scientific publishing
<http://msp.org/>

© 2020 Mathematical Sciences Publishers

PURE and APPLIED ANALYSIS

vol. 2 no. 1 2020

Emergence of nontrivial minimizers for the three-dimensional Ohta–Kawasaki energy	1
HANS KNÜPFER, CYRILL B. MURATOV and MATTEO NOVAGA	
Maximal L^2 -regularity in nonlinear gradient systems and perturbations of sublinear growth	23
WOLFGANG ARENDT and DANIEL HAUER	
The local density approximation in density functional theory	35
MATHIEU LEWIN, ELLIOTT H. LIEB and ROBERT SEIRINGER	
Sparse bounds for the discrete spherical maximal functions	75
ROBERT KESLER, MICHAEL T. LACEY and DARÍO MENA	
Characterization by observability inequalities of controllability and stabilization properties	93
EMMANUEL TRÉLAT, GENGSHENG WANG and YASHAN XU	
Stationary compressible Navier–Stokes equations with inflow condition in domains with piecewise analytical boundaries	123
PIOTR B. MUCHA and TOMASZ PIASECKI	
Optimal constants in nontrapping resolvent estimates and applications in numerical analysis	157
JEFFREY GALKOWSKI, EUAN A. SPENCE and JARED WUNSCH	



2578-5893 (2020)2:1;1-B