

# PURE and APPLIED ANALYSIS

# PAA

RAINER MANDEL

**DISPERSIVE ESTIMATES, BLOW-UP AND FAILURE OF  
STRICHARTZ ESTIMATES FOR THE SCHRÖDINGER EQUATION  
WITH SLOWLY DECAYING INITIAL DATA**

# DISPERSIVE ESTIMATES, BLOW-UP AND FAILURE OF STRICHARTZ ESTIMATES FOR THE SCHRÖDINGER EQUATION WITH SLOWLY DECAYING INITIAL DATA

RAINER MANDEL

The initial value problem for the homogeneous Schrödinger equation is investigated for radially symmetric initial data with slow decay rates and not too wild oscillations. Our global well-posedness results apply to initial data for which Strichartz estimates fail.

## 1. Introduction

In this paper we investigate the initial value problem for the Schrödinger equation

$$i \partial_t \psi + \Delta \psi = 0 \quad \text{in } \mathbb{R}^n, \quad \psi(0) = \phi, \tag{1}$$

for radial initial data  $\phi$  with slow decay at infinity. In particular, we are interested in a solution theory for (1) without assuming  $\phi$  to belong to one of the Lebesgue spaces  $L^r(\mathbb{R}^n)$  with  $r \in [1, 2]$ . In this case Strichartz estimates are not available and local or global well-posedness results for (1) are unknown. Surprisingly, we could not find a complete statement about Strichartz estimates for such initial data in the literature, so we clarify this point here.

**Theorem 1.** *Let  $n \in \mathbb{N}$  and  $p, q \in [1, \infty]$ ,  $r > 2$ . Then there is no Strichartz estimate*

$$\|e^{it\Delta} \phi\|_{L^p_t(\mathbb{R}; L^q(\mathbb{R}^n))} \lesssim \|\phi\|_{L^r(\mathbb{R}^n)}. \tag{2}$$

This theorem partly generalizes the known fact that for any given  $t > 0$  the Schrödinger propagator  $e^{it\Delta}$  is unbounded as a map from  $L^r(\mathbb{R}^n)$  to  $L^q(\mathbb{R}^n)$  for all  $r > 2$ ,  $q \in [1, \infty]$ ; see [Linares and Ponce 2015, p. 63] for the case  $q = r$ . Theorem 1 may seem surprising in view of the fact that the optimal conditions for Strichartz estimates in the most important special case  $r = 2$  do not provide any obvious reason why the estimates should break down completely for  $r > 2$ . Recall that these conditions are given by

$$p, q \geq 2, \quad (p, q, n) \neq (2, \infty, 2), \quad \frac{2}{p} + \frac{n}{q} = \frac{n}{2};$$

see for instance [Cazenave 2003, Theorem 2.3.3]. We refer to [Strichartz 1977; Keel and Tao 1998; Ginibre and Velo 1985] for three milestone contributions related to the discovery of these conditions. At least for  $n \geq 3$ , each of the above conditions has a counterpart in the range  $r > 2$ . The scaling invariance of the Schrödinger equation implies  $2/p + n/q = n/r$  so that  $q \geq r$  would be an immediate consequence that

MSC2010: primary 35Q41; secondary 35B40, 35B44.

Keywords: Schrödinger equation, failure of Strichartz estimates, dispersive blow-up.

replaces the condition  $q \geq 2$ . As we discuss in the [Appendix](#)  $p \geq 2$  generalizes to  $p \geq 2r/(2n - r(n - 1))_+$ . In particular, there is no evident reason for the necessity of  $r \leq 2$  so that [Theorem 1](#) seems to fill a gap in the literature. Its short proof relies on a thorough analysis of a counterexample due to Bona, Ponce, Saut, and Sparber [[Bona et al. 2014](#)]. The main feature of their solution is that the corresponding initial datum oscillates quadratically with respect to the distance to the origin, which produces  $L^\infty$ -blow-up (or dispersive blow-up) of the solution at some prescribable finite time; see [[Bona et al. 2014](#), Lemma 2.1]. We reconsider this self-similar blow-up analysis for partly more general initial data and estimate the blow-up rate in  $L^q(\mathbb{R}^n)$ , which eventually leads to [Theorem 1](#). Accordingly, our proof even reveals that local Strichartz estimates cannot hold either and that no improvement in the radial situation is possible.

Given that Strichartz estimates fail, the question arises of how well-posedness results for the Schrödinger equation can be achieved if the initial datum lies in  $L^r(\mathbb{R}^n)$  only for  $r > 2$ . In view of the above-mentioned counterexample it seems reasonable to impose a condition on the oscillations of the initial datum. In the following we present one possible approach in the radially symmetric case which relies on suitably weighted Sobolev norms of the initial data. For instance we identify a class of initial data lying in  $L^r(\mathbb{R}^n)$  only for  $r > 2n/(n - 1)$  with solutions that are bounded in time and uniformly localized in space; see [Corollary 3](#). In that case dispersion need not occur because there are solutions of the form

$$\psi(x, t) = e^{-i\omega^2 t} \phi(x), \quad \text{where } \phi(x) = |x|^{(2-n)/2} J_{(n-2)/2}(\omega|x|) \text{ for some } \omega \in \mathbb{R}. \tag{3}$$

Here,  $J_{(n-2)/2}$  denotes the Bessel function of the first kind and  $\phi$  solves the linear Helmholtz equation  $\Delta\phi + \omega^2\phi = 0$  in  $\mathbb{R}^n$ . Aiming for a more general result in this direction, we first consider initial profiles of the form  $\phi(x) = e^{i\omega|x|}\phi_\omega(|x|)$ , where  $\phi_\omega$  belongs to the function spaces  $X$  and  $Y_m$  for some  $m \in \{0, \dots, n\}$ , which we introduce now. The space  $X$  is defined to be the completion of  $C_c^\infty(\mathbb{R}_{\geq 0}; \mathbb{C})$  with respect to the norm  $\|\cdot\|_X := \|\cdot\|_{X_1} + \|\cdot\|_{X_2}$  given by

$$\begin{aligned} \|f\|_{X_1} &:= \sup_{z>0} z^{(1-n)/2} \int_0^z (|f(r)|r^{n-2} + |f'(r)|r^{n-1}) dr, \\ \|f\|_{X_2} &:= \int_0^\infty \left| \frac{d}{dr}(f(r)r^{(n-1)/2}) \right| dr + \sup_{z>0} \left( z \int_z^\infty |f(r)|r^{(n-5)/2} dr \right). \end{aligned}$$

Similarly, we define  $Y_m$  to be the completion of  $C_c^\infty(\mathbb{R}_{\geq 0}; \mathbb{C})$  with respect to the norm

$$\begin{aligned} \|f\|_{Y_m} &:= \sum_{k=0}^m \int_0^\infty |f^{(k)}(r)|r^{n-m+k-1} dr \quad \text{if } m \in \{0, \dots, n-1\}, \\ \|f\|_{Y_n} &:= \sum_{k=1}^n \int_0^\infty |f^{(k)}(r)|r^{k-1} dr + \sup_{z>0} z^{-2} \int_0^z f(r)r dr + |f(0)|. \end{aligned}$$

One main feature of these spaces is that slow decay rates of their elements are admissible only provided that their derivatives decay fast enough. For instance, we have that  $r \mapsto (1+r)^{-\alpha}$  lies in  $X$  if and only if  $\alpha \geq (n-1)/2$  and in  $Y_m$  if and only if  $\alpha > n-m$ , whereas  $r \mapsto e^{ir}(1+r)^{-\alpha}$  belongs to  $X$  if and only if  $\alpha > (n+1)/2$  and to  $Y_m$  if and only if  $\alpha > n$ . Using these spaces we find a local well-posedness theory for at most linearly oscillating radial initial data.

**Theorem 2.** Let  $n \in \mathbb{N}$ ,  $m \in \{0, \dots, n\}$ , and  $\phi(x) = \phi_\omega(|x|)e^{i\omega|x|}$  for some  $\omega \in \mathbb{R}$ :

(i) If  $\phi_\omega \in Y_m$  then (1) has a unique global solution  $\psi$  satisfying

$$|\psi(x, t)| \leq C(\sqrt{t})^{m-n} \|\phi_\omega\|_{Y_m}.$$

(ii) If  $\phi_\omega \in X$  then (1) has a unique global solution  $\psi$  satisfying

$$|\psi(x, t)| \leq C|x|^{(1-n)/2} \|\phi_\omega\|_X.$$

In (i) and (ii) the constant  $C$  does not depend on  $\omega$ .

Here we used the term “global solution” to indicate a distributional solution of the Schrödinger equation (1) away from  $t = 0$ , i.e., on  $\mathbb{R} \setminus \{0\} \times \mathbb{R}^n$ . Given that the test functions are dense in both  $X$  and  $Y_m$  and that the above estimates hold, one even gets that such solutions are limits of smooth classical solutions with respect to uniform convergence on all compact sets avoiding  $t = 0$  and  $x = 0$ . By the estimate (i) for  $m = n$  this convergence is even uniform on  $\mathbb{R} \times \mathbb{R}^n$  if we assume  $\phi_\omega \in Y_n$ . Combining the estimates (i) and (ii) we deduce the following.

**Corollary 3.** Let  $n \in \mathbb{N}$  and assume  $\phi(x) = \int_{\mathbb{R}} \phi_\omega(|x|)e^{i\omega|x|} d\mu(\omega)$  for some Borel measure  $\mu$  on  $\mathbb{R}$ . Then (1) has a unique global solution satisfying

$$|\psi(x, t)| \leq C(1 + |x|)^{(1-n)/2} \int_{\mathbb{R}} (\|\phi_\omega\|_X + \|\phi_\omega\|_{Y_n}) d\mu(\omega),$$

provided the right-hand side is finite.

**Corollary 4.** Let  $n \in \mathbb{N}$ ,  $m \in \{0, \dots, n\}$ , and assume  $\phi(x) = \int_{\mathbb{R}} \phi_\omega(|x|)e^{i\omega|x|} d\mu(\omega)$  for some Borel measure  $\mu$  on  $\mathbb{R}$ . Then (1) has a unique global solution satisfying

$$|\psi(x, t)| \leq C(1 + t)^{-m/2} \int_{\mathbb{R}} (\|\phi_\omega\|_{Y_{n-m}} + \|\phi_\omega\|_{Y_n}) d\mu(\omega),$$

provided the right-hand side is finite.

**Remark 1.** (a) In the case  $n \geq 2$  we can apply Corollary 3 to initial conditions that are sufficiently regular superpositions of radially symmetric Herglotz waves. The corresponding densities  $a$  are assumed to be Lebesgue measurable and to satisfy  $\int_{\mathbb{R}} |a(\omega)|(|\omega|^{-1/2} + |\omega|^{(n-2)/2}) < \infty$ . Using the asymptotic expansions of the Bessel functions at infinity one finds

$$\phi(x) := \int_0^\infty a(\omega)|x|^{(2-n)/2} J_{(n-2)/2}(\omega|x|) d\omega = \int_{\mathbb{R}} \phi_\omega(|x|)e^{i\omega|x|} d\mu(\omega),$$

where  $d\mu(\omega) = d\delta_0(\omega) + d\omega$  ( $\delta_0$  is the Dirac delta distribution) and

$$\phi_\omega(r) = r^{1-n} \cdot \begin{cases} a(\omega)\omega^{-n/2} B_n(\omega r) & \text{if } \omega > 0, \\ \int_0^\infty a(\xi)\xi^{-n/2} A_n(\xi r) d\xi & \text{if } \omega = 0, \\ a(-\omega)|\omega|^{-n/2} \overline{B_n(-\omega r)} & \text{if } \omega < 0. \end{cases}$$

Here we anticipated the notation from (6). So Proposition 6 and a few computations yield that  $\omega \mapsto \|\phi_\omega\|_X + \|\phi_\omega\|_{Y_n}$  is  $\mu$ -integrable because of

$$\begin{aligned} \|\phi_0\|_X + \int_{\mathbb{R}} \|\phi_\omega\|_X d\omega &\lesssim \int_{\mathbb{R}} |a(\omega)| |\omega|^{-1/2} d\omega, \\ \|\phi_0\|_{Y_n} + \int_{\mathbb{R}} \|\phi_\omega\|_{Y_n} d\omega &\lesssim \int_{\mathbb{R}} |a(\omega)| |\omega|^{(n-2)/2} d\omega. \end{aligned}$$

In particular, Corollary 3 provides an abstract framework for the observation that the solutions of the Schrödinger equation with initial data given by sufficiently regular superpositions of radially symmetric Herglotz waves as in (3) remain bounded in time and uniformly localized in space.

(b) Theorem 2(i) is a generalized version of the fact that integrable initial data yield bounded solutions. Indeed, the latter statement corresponds to  $m = 0$  in the theorem. So we get that less integrability of the initial datum is still sufficient for the absence of finite time  $L^\infty$ -blow-up provided that the derivatives decay sufficiently fast. Notice that some kind of control on the derivatives seems necessary given that there are initial data in  $L^s(\mathbb{R}^n)$  for any given  $s > 1$ , the corresponding solutions of which become unbounded in finite time; see [Bona et al. 2014, Lemma 2.1 and Remark 2.2].

(c) The decay rates from Corollary 3 improve once we add regularity assumptions on  $\mu$ . In the simplest situation  $d\mu(\omega) = b(\omega) d\omega$  for  $b \in W_0^{k,1}(\mathbb{R})$  and  $\omega \mapsto \phi_\omega \in W^{k,\infty}(\mathbb{R}; X \cap Y_n)$ , the decay rate improves to  $(1 + |x|)^{(1-n)/2-k}$ . This is proved using integration by parts as in the method of stationary phase. In the Appendix we discuss the densities  $b(\omega) = (\omega - 1)^{-\delta} \mathbb{1}_{[1,2]}(\omega)$  with  $\delta \in (0, 1)$  and find the intermediate decay rates  $(1 + |x|)^{(1-n)/2-(1-\delta)}$ .

(d) Theorem 2(i) tells us that nondispersive solutions of the Schrödinger equation (1) can only occur for radial initial data  $\phi(x) = \phi_{\text{rad}}(|x|)$  that satisfy  $\|\phi_{\text{rad}}\|_{Y_m} = \infty$  for all  $m \in \{0, \dots, n - 1\}$ . For smooth initial data this essentially means that for some  $k \in \{0, \dots, n - 1\}$  the function  $|\phi_{\text{rad}}^{(k)}(r)|$  does not decay faster than  $(1 + r)^{-1-k}$  as  $r \rightarrow \infty$ . We conclude that the lack of dispersion is a phenomenon related to slowly decaying or heavily oscillating initial data.

## 2. Proof of Theorem 2

In the following let  $\psi$  denote the unique solution of the Schrödinger equation (1) with initial datum  $\phi$  given by  $\phi(x) = \phi_{\text{rad}}(|x|) = \phi_\omega(|x|)e^{i\omega|x|}$ . By density, it suffices to prove the estimates for  $\phi_\omega \in C_c^\infty(\mathbb{R}_{\geq 0}; \mathbb{C})$ . In the following we use the abbreviations

$$f(r) := \phi_{\text{rad}}(r)r^{n/2}J_{(n-2)/2}\left(\frac{r|x|}{2t}\right) \quad \text{and} \quad g(\rho) := f(2\sqrt{t}\rho). \tag{4}$$

We first recall the representation formula of the solution for radial initial data.

**Proposition 5.** *We have for all  $x \in \mathbb{R}^n$ ,  $t > 0$ ,*

$$\psi(x, t) = |x|^{(2-n)/2}(\sqrt{t})^{-1}e^{i(|x|^2/(4t)-n\pi/4)} \int_0^\infty g(\rho)e^{i\rho^2} d\rho. \tag{5}$$

*Proof.* From (4.2) in [Linares and Ponce 2015] or (2.2.5) in [Cazenave 2003] we get

$$\begin{aligned}
 \psi(x, t) &= \frac{1}{(4i\pi t)^{n/2}} \int_{\mathbb{R}^n} e^{i|x-y|^2/(4t)} \phi(y) dy \\
 &= \frac{1}{(4i\pi t)^{n/2}} \int_0^\infty \phi_{\text{rad}}(r) r^{n-1} e^{i(|x|^2+r^2)/(4t)} \left( \int_{\partial B_1(0)} e^{ir(x,\omega)/(2t)} d\sigma(\omega) \right) dr \\
 &= \frac{1}{(4i\pi t)^{n/2}} \int_0^\infty \phi_{\text{rad}}(r) r^{n-1} e^{i(|x|^2+r^2)/(4t)} \left( \int_{\partial B_1(0)} e^{ir|x|\omega_1/(2t)} d\sigma(\omega) \right) dr \\
 &= \frac{1}{(4i\pi t)^{n/2}} \int_0^\infty \phi_{\text{rad}}(r) r^{n-1} e^{i(|x|^2+r^2)/(4t)} |\partial B_1(0)| \Gamma\left(\frac{n}{2}\right) 2^{(n-2)/2} \left(\frac{r|x|}{2t}\right)^{(2-n)/2} J_{(n-2)/2}\left(\frac{r|x|}{2t}\right) dr \\
 &= |\partial B_1(0)| \Gamma\left(\frac{n}{2}\right) (4\pi i)^{-n/2} 2^{n-2} |x|^{(2-n)/2} t^{-1} e^{i|x|^2/(4t)} \int_0^\infty f(r) e^{ir^2/(4t)} dr \\
 &= |\partial B_1(0)| \Gamma\left(\frac{n}{2}\right) (4\pi i)^{-n/2} 2^{n-1} |x|^{(2-n)/2} (\sqrt{t})^{-1} e^{i|x|^2/(4t)} \int_0^\infty f(2\sqrt{t}\rho) e^{i\rho^2} d\rho.
 \end{aligned}$$

So the claim follows from  $|\partial B_1(0)| = 2\pi^{n/2} / \Gamma(n/2)$ .  $\square$

It will be convenient to split the integrand in (5) into three parts,  $g = g_1 + g_2 + g_3$ . The function  $g_1$  will be identical to  $g$  for small arguments and the corresponding estimates rely on the behavior of the Bessel function  $J_{(n-2)/2}$  on the interval  $[0, 1]$ . The sum  $g_2 + g_3$  represents  $g$  for large arguments and their definitions are based on the asymptotic expansion of the Bessel function at infinity. To be more precise, we fix some cut-off function  $\chi \in C_0^\infty(\mathbb{R})$  such that  $\chi \equiv 1$  on  $[0, \frac{1}{2}]$  and  $\chi \equiv 0$  on  $[1, \infty]$ . Then, much as in [Watson 1944, p. 202], we write

$$z^{n/2} J_{(n-2)/2}(z) = A_n(z) + e^{iz} B_n(z) + e^{-iz} \overline{B_n(z)}, \quad (6)$$

where the functions  $A_n, B_n$  are given by

$$A_n(z) := \chi(z) z^{n/2} J_{(n-2)/2}(z), \quad B_n(z) := (1 - \chi(z)) e^{-i(n-1)\pi/4} \sum_{k=0}^\infty \alpha_k z^{(n-1)/2-k} \quad (7)$$

and the coefficients  $\alpha_k \in \mathbb{C}$  are  $\alpha_0 := 1/\sqrt{2\pi}$  and for  $k \in \mathbb{N}$

$$\alpha_k := \frac{1}{\sqrt{2\pi}} \binom{n-2}{2} \binom{i}{2}^k, \quad \text{where } (v, k) := \frac{(4v^2 - 1^2)(4v^2 - 3^2) \cdots (4v^2 - (2k-1)^2)}{4^k k!}; \quad (8)$$

see [Watson 1944, p. 199]. The motivation for this decomposition is that  $A_n, B_n$  and its derivatives satisfy useful uniform estimates that we provide next.

**Proposition 6.** *We have  $\text{supp}(A_n) \subset [0, 1]$ ,  $\text{supp}(B_n) \subset [\frac{1}{2}, \infty)$ , and for all  $j \in \mathbb{N}_0$ ,  $z \in \mathbb{R}$ ,*

$$|A_n^{(j)}(z)| \lesssim \begin{cases} |z|^{n-1-j} & \text{if } j \in \{0, \dots, n-1\}, \\ |z| & \text{if } j \in \{n, n+2, n+4, \dots\}, \\ 1 & \text{if } j \in \{n+1, n+3, \dots\}, \end{cases}$$

$$\left| \frac{d^j}{dz^j} (B_n(z) z^{(1-n)/2}) \right| \lesssim \begin{cases} 1 & \text{if } j = 0, \\ |z|^{-1-j} & \text{if } j \geq 1. \end{cases}$$

*Proof.* The estimate for  $A_n$  follows from

$$\begin{aligned} A_n(z) &= \chi(z)z^{n/2}J_{(n-2)/2}(z) \\ &= \chi(z)z^{n/2}\sum_{m=0}^{\infty}\frac{(-1)^m}{m!\Gamma(m+n/2)}\left(\frac{z}{2}\right)^{(n-2)/2+2m} \\ &= \chi(z)\sum_{m=0}^{\infty}\frac{(-1)^m}{2^{(n-2)/2+2m}m!\Gamma(m+n/2)}z^{n-1+2m}. \end{aligned} \tag{9}$$

The estimate for  $B_n$  follows from its series representation (7); see also [Watson 1944, p. 206].  $\square$

Given the definition of  $g$  in (4) and the splitting (6) of the Bessel function, we decompose the integrand  $g$  according to  $g = g_1 + g_2 + g_3$ , where, for  $r := 2\sqrt{t}\rho$ ,

$$\begin{aligned} g_1(\rho) &:= \phi_{\text{rad}}(r)A_n\left(\frac{r|x|}{2t}\right)\left(\frac{|x|}{2t}\right)^{-n/2} = e^{2i\sqrt{t}\omega\rho}\phi_{\omega}(r)A_n\left(\frac{r|x|}{2t}\right)\left(\frac{|x|}{2t}\right)^{-n/2}, \\ g_2(\rho) &:= e^{ir|x|/(2t)}\phi_{\text{rad}}(r)B_n\left(\frac{r|x|}{2t}\right)\left(\frac{|x|}{2t}\right)^{-n/2} = e^{2i\sqrt{t}(\omega+|x|/(2t))\rho}\phi_{\omega}(r)B_n\left(\frac{r|x|}{2t}\right)\left(\frac{|x|}{2t}\right)^{-n/2}, \\ g_3(\rho) &:= e^{-ir|x|/(2t)}\phi_{\text{rad}}(r)B_n\left(\frac{r|x|}{2t}\right)\left(\frac{|x|}{2t}\right)^{-n/2} = e^{2i\sqrt{t}(\omega-|x|/(2t))\rho}\phi_{\omega}(r)B_n\left(\frac{r|x|}{2t}\right)\left(\frac{|x|}{2t}\right)^{-n/2}. \end{aligned}$$

We now remove the linear phase factors by putting  $g_{j,a_j}(\rho) := g_j(\rho)e^{-ia_j\rho}$  for

$$a_1 := 2\sqrt{t}\omega, \quad a_2 := 2\sqrt{t}\left(\omega + \frac{|x|}{2t}\right), \quad a_3 := 2\sqrt{t}\left(\omega - \frac{|x|}{2t}\right). \tag{10}$$

This implies, again for  $r := 2\sqrt{t}\rho$ ,

$$\begin{aligned} g_{1,a_1}(\rho) &= \phi_{\omega}(r)A_n\left(\frac{r|x|}{2t}\right)\left(\frac{|x|}{2t}\right)^{-n/2}, \\ g_{2,a_2}(\rho) &= \phi_{\omega}(r)B_n\left(\frac{r|x|}{2t}\right)\left(\frac{|x|}{2t}\right)^{-n/2}, \\ g_{3,a_3}(\rho) &= \phi_{\omega}(r)B_n\left(\frac{r|x|}{2t}\right)\left(\frac{|x|}{2t}\right)^{-n/2}. \end{aligned} \tag{11}$$

So we infer from Proposition 5

$$\psi(x, t)|x|^{(n-2)/2}\sqrt{t}e^{-i(|x|^2/(4t)-n\pi/4)} = \int_0^{\infty} g(\rho)e^{i\rho^2} d\rho = \sum_{j=1}^3 \int_0^{\infty} g_{j,a_j}(\rho)e^{i(\rho^2+a_j\rho)} d\rho. \tag{12}$$

In order to estimate these terms, we make use of the following auxiliary result.

**Proposition 7.** *Let  $a \in \mathbb{R}$  and  $\Xi_a^m \in C^\infty(\mathbb{R}; \mathbb{C})$  for  $m \in \mathbb{N}_0$  be inductively defined by*

$$\Xi_a^0(s) := \int_s^{\infty} e^{i(\rho^2+a\rho)} d\rho, \quad \Xi_a^m(s) := \int_s^{\infty} \Xi_a^{m-1}(\rho) d\rho.$$

*Then  $|\Xi_a^m(s)| \leq C_m$  for all  $a \in \mathbb{R}$ ,  $s \geq 0$ .*

*Proof.* This follows from  $\Xi_a^m(s) = e^{-ia^2/4} \Xi_0^m(s + a/2)$  once we have proved the estimate

$$|\Xi^m(s)| \leq C_m(1 + s_+)^{-m-1} \quad \text{for all } s \in \mathbb{R},$$

where  $\Xi^m := \Xi_0^m$ . The existence of the improper Fresnel integral  $\Xi^0(s)$  is a well-known consequence of the residue theorem. Moreover, l'Hôpital's rule gives

$$\Xi^0(s)(-2is)e^{-is^2} \rightarrow 1 \quad \text{and} \quad s^2(1 + 2is\Xi^0(s)e^{-is^2}) \rightarrow \frac{i}{2} \quad \text{as } s \rightarrow \infty.$$

(For the second limit one may proceed as we do below in the computation of  $z_k$ .) Since the improper integral  $\int_s^\infty (1/\rho)e^{i\rho^2} d\rho$  exists for  $s > 1$  (again by the residue theorem), we obtain from the previous statement that the integral  $\Xi^1(s)$  exists and

$$\Xi^1(s)(-2is)^2e^{-is^2} \rightarrow 1 \quad \text{and} \quad s^2(1 - (-2is)^2\Xi^1(s)e^{-is^2}) \rightarrow \frac{3i}{2} \quad \text{as } s \rightarrow \infty.$$

By induction, we find that for all  $k \in \mathbb{N}$ ,  $k \geq 2$  the (proper) integral  $\Xi^k(s)$  exists and

$$\begin{aligned} z_k &:= \lim_{s \rightarrow \infty} s^2(1 - (-2is)^{k+1}\Xi^k(s)e^{-is^2}) \\ &= \lim_{s \rightarrow \infty} \frac{e^{is^2}(-2is)^{-k-1} - \Xi^k(s)}{e^{is^2}s^{-2}(-2is)^{-k-1}} \\ &= \lim_{s \rightarrow \infty} \frac{-e^{is^2}(-2is)^{-k} + 2(k+1)ie^{is^2}(-2is)^{-k-2} - (\Xi^k)'(s)}{-e^{is^2}s^{-2}(-2is)^{-k} - 2e^{is^2}s^{-3}(-2is)^{-k-1} + 2(k+1)ie^{is^2}s^{-2}(-2is)^{-k-2}} \\ &= \lim_{s \rightarrow \infty} \frac{-(-2is)^{-k} + 2(k+1)i(-2is)^{-k-2} + \Xi^{k-1}(s)e^{-is^2}}{-s^{-2}(-2is)^{-k} - 2s^{-3}(-2is)^{-k-1} + 2(k+1)is^{-2}(-2is)^{-k-2}} \\ &= \lim_{s \rightarrow \infty} \frac{-(k+1)is^{-2} - 2 + 2(-2is)^k\Xi^{k-1}(s)e^{-is^2}}{-2s^{-2} - 2is^{-4} - (k+1)is^{-4}} \\ &= \lim_{s \rightarrow \infty} \frac{(k+1)i + 2s^2(1 - (-2is)^k\Xi^{k-1}(s)e^{-is^2})}{2 + 2is^{-2} + (k+1)is^{-2}} = \frac{(k+1)i}{2} + z_{k-1}, \end{aligned}$$

implying

$$z_k = \lim_{s \rightarrow \infty} s^2(1 - (-2is)^{k+1}\Xi^k(s)e^{-is^2}) = \frac{(k+2)(k+1)i}{4}.$$

This yields the bounds for  $\Xi^m(s)$  and we are done.  $\square$

Let us remark that the proof actually yields the stronger estimate  $|\Xi_a^m(s)| \leq C_m$  for  $0 \leq s \leq a_-$  and  $|\Xi_a^m(s)| \leq 2^{m+1}C_m(1+s)^{-m-1}$  for  $s \geq a_-$ . However, given that these estimates depend on  $a$ , it seems difficult to make use of them. Moreover, the independence of  $a$  guarantees that our estimates below do not depend on  $\omega$  since the latter is completely absorbed in the definitions of  $a_1, a_2, a_3$  from (10). From (12) and Proposition 7 we deduce the following estimate for the solution  $\psi$ .

**Proposition 8.** *For all  $x \in \mathbb{R}^n$ ,  $t \in \mathbb{R}$ ,  $m \in \{0, \dots, n\}$ , we have*

$$|\psi(x, t)| \leq C_{m-1}|x|^{(2-n)/2}t^{-1/2} \left( |\delta_{m,n}|g_{1,a_1}^{(n-1)}(0)| + \int_0^\infty |g_{1,a_1}^{(m)}(\rho)| + |g_{2,a_2}^{(m)}(\rho)| + |g_{3,a_3}^{(m)}(\rho)| d\rho \right), \quad (13)$$

where  $g_{1,a_1}, g_{2,a_2}, g_{3,a_3}$  are given by (11).

*Proof.* In the following integration-by-parts scheme we use  $(\Xi_{a_j}^m)' = -\Xi_{a_j}^{m-1}$  as well as

$$g_{j,a_j}(0) = g'_{j,a_j}(0) = \dots = g_{j,a_j}^{(n-2)}(0) = 0, \quad g_{2,a_2}^{(n-1)}(0) = g_{3,a_3}^{(n-1)}(0) = 0, \tag{14}$$

which follows from (11) and Proposition 6. Recall that the support of  $B_n$  is contained in  $[\frac{1}{2}, \infty)$  by choice of the cut-off function  $\chi$  so that the above estimate is actually trivial for  $j \in \{2, 3\}$ . So we have for  $m \in \{0, \dots, n-1\}$

$$\begin{aligned} \int_0^\infty g_{j,a_j}(\rho) e^{i(\rho^2+a_j\rho)} d\rho &\stackrel{(14)}{=} \lim_{M \rightarrow \infty} \int_0^M \left( \int_0^\rho g'_{j,a_j}(t) dt \right) e^{i(\rho^2+a_j\rho)} d\rho \\ &= \lim_{M \rightarrow \infty} \int_0^M g'_{j,a_j}(t) \left( \int_t^M e^{i(\rho^2+a_j\rho)} d\rho \right) dt \\ &= \int_0^\infty g'_{j,a_j}(t) \Xi_{a_j}^0(t) dt \\ &\stackrel{(14)}{=} \int_0^\infty \left( \int_0^t g''_{j,a_j}(s) ds \right) \Xi_{a_j}^0(t) dt \\ &= \int_0^\infty g''_{j,a_j}(s) \Xi_{a_j}^1(s) ds \\ &\vdots \\ &= \int_0^\infty g_{j,a_j}^{(m)}(\rho) \Xi_{a_j}^{m-1}(\rho) d\rho. \end{aligned} \tag{15}$$

Notice that the limit  $M \rightarrow \infty$  passes under the integral because  $g_{j,a_j}$  has compact support and the  $\Xi_{a_j}^k$  are bounded by Proposition 7. So we obtain for  $m \in \{0, \dots, n-1\}$

$$\begin{aligned} |\psi(x, t)| &\stackrel{(12)}{\leq} |x|^{(2-n)/2} (\sqrt{t})^{-1} \sum_{j=1}^3 \left| \int_0^\infty g_{j,a_j}^{(m)}(\rho) \Xi_{a_j}^{m-1}(\rho) d\rho \right| \\ &\leq C_{m-1} |x|^{(2-n)/2} (\sqrt{t})^{-1} \sum_{j=1}^3 \int_0^\infty |g_{j,a_j}^{(m)}(\rho)| d\rho. \end{aligned}$$

Moreover, using (15) for  $m = n-1$  we get

$$\begin{aligned} |\psi(x, t)| &= |x|^{(2-n)/2} (\sqrt{t})^{-1} \left| \sum_{j=1}^3 \int_0^\infty g_{j,a_j}^{(n-1)}(\rho) \Xi_{a_j}^{n-2}(\rho) d\rho \right| \\ &= |x|^{(2-n)/2} (\sqrt{t})^{-1} \left| \sum_{j=1}^3 \left( g_{j,a_j}^{(n-1)}(0) \int_0^\infty \Xi_{a_j}^{n-2}(\rho) d\rho + \int_0^\infty \left( \int_0^t g_{j,a_j}^{(n)}(\rho) d\rho \right) \Xi_{a_j}^{n-2}(t) dt \right) \right| \\ &= |x|^{(2-n)/2} (\sqrt{t})^{-1} \left| \sum_{j=1}^3 \left( g_{j,a_j}^{(n-1)}(0) \Xi_{a_j}^{n-1}(0) + \int_0^\infty g_{j,a_j}^{(n)}(\rho) \Xi_{a_j}^{n-1}(\rho) d\rho \right) \right| \\ &\stackrel{(14)}{\leq} C_{n-1} |x|^{(2-n)/2} (\sqrt{t})^{-1} \left( |g_{1,a_1}^{(n-1)}(0)| + \sum_{j=1}^3 \int_0^\infty |g_{j,a_j}^{(n)}(\rho)| d\rho \right). \end{aligned} \quad \square$$

For notational convenience we write  $x \lesssim y$ , respectively  $x \gtrsim y$ , instead of  $x \leq cy$ , respectively  $x \geq cy$ , for positive numbers  $c$  that are independent of  $\omega$ ,  $|x|$ ,  $t$ ,  $r$  but may depend on  $m \in \{0, \dots, n\}$  or the space dimension  $n \in \mathbb{N}$ .

**Proposition 9.** *Let  $m \in \{0, \dots, n\}$ . Then the functions  $g_{1,a_1}$ ,  $g_{2,a_2}$ ,  $g_{3,a_3}$  from (11) satisfy the following estimates for all  $\rho \geq 0$  and  $r = 2\sqrt{t}\rho$ :*

$$\begin{aligned} |g_{1,a_1}^{(m)}(\rho)| &\lesssim |x|^{(n-2)/2} (\sqrt{t})^{m-n+2} \sum_{k=0}^m |\phi_\omega^{(k)}(r)| r^{n-m+k-1} \cdot \mathbb{1}_{[0,1]} \left( \frac{r|x|}{2t} \right) \quad \text{if } m < n, \\ |g_{1,a_1}^{(n)}(\rho)| &\lesssim \left( |x|^{(n-2)/2} (\sqrt{t})^2 \sum_{k=1}^n |\phi_\omega^{(k)}(r)| r^{k-1} + |x|^{(n+2)/2} (\sqrt{t})^{-2} |\phi_\omega(r)| r \right) \cdot \mathbb{1}_{[0,1]} \left( \frac{r|x|}{2t} \right), \\ |g_{2,a_2}^{(m)}(\rho)| + |g_{3,a_3}^{(m)}(\rho)| &\lesssim |x|^{-1/2} (\sqrt{t})^{1+m} \sum_{k=0}^m |\phi_\omega^{(k)}(r)| r^{(n-1)/2-m+k} \cdot \mathbb{1}_{[1/2,\infty)} \left( \frac{r|x|}{2t} \right). \end{aligned}$$

*Proof.* We get for  $r = 2\sqrt{t}\rho$  and  $m \in \{0, \dots, n-1\}$

$$\begin{aligned} |g_{1,a_1}^{(m)}(\rho)| &\stackrel{(11)}{=} (2\sqrt{t})^m \left| \frac{d^m}{dr^m} \left( \phi_\omega(r) A_n \left( \frac{r|x|}{2t} \right) \right) \right| \left( \frac{|x|}{2t} \right)^{-n/2} \\ &\lesssim (\sqrt{t})^m \sum_{k=0}^m \left| \phi_\omega^{(k)}(r) A_n^{(m-k)} \left( \frac{r|x|}{2t} \right) \right| \left( \frac{|x|}{2t} \right)^{m-k-n/2} \\ &\stackrel{\text{Prop. 6}}{\lesssim} (\sqrt{t})^m \sum_{k=0}^m |\phi_\omega^{(k)}(r)| \left( \frac{r|x|}{2t} \right)^{n-1-m+k} \left( \frac{|x|}{2t} \right)^{m-k-n/2} \cdot \mathbb{1}_{[0,1]} \left( \frac{r|x|}{2t} \right) \\ &\lesssim |x|^{(n-2)/2} (\sqrt{t})^{m-n+2} \sum_{k=0}^m |\phi_\omega^{(k)}(r)| r^{n-1-m+k} \cdot \mathbb{1}_{[0,1]} \left( \frac{r|x|}{2t} \right). \end{aligned}$$

This implies the first estimate. For  $m = n$  we use the estimate for  $A_n^{(j)}$  from Proposition 6 for  $j \in \{0, \dots, n\}$  and obtain

$$\begin{aligned} |g_{1,a_1}^{(n)}(\rho)| &\stackrel{(11)}{=} (2\sqrt{t})^n \left| \frac{d^n}{dr^n} \left( \phi_\omega(r) A_n \left( \frac{r|x|}{2t} \right) \right) \right| \left( \frac{|x|}{2t} \right)^{-n/2} \\ &\lesssim (\sqrt{t})^n \sum_{k=0}^n \left| \phi_\omega^{(k)}(r) A_n^{(n-k)} \left( \frac{r|x|}{2t} \right) \right| \left( \frac{|x|}{2t} \right)^{n/2-k} \\ &\lesssim (\sqrt{t})^n \left( \sum_{k=1}^n |\phi_\omega^{(k)}(r)| \left( \frac{r|x|}{2t} \right)^{-1+k} \left( \frac{|x|}{2t} \right)^{n/2-k} + |\phi_\omega(r)| \frac{r|x|}{2t} \left( \frac{|x|}{2t} \right)^{n/2} \right) \cdot \mathbb{1}_{[0,1]} \left( \frac{r|x|}{2t} \right) \\ &\lesssim \left( |x|^{(n-2)/2} (\sqrt{t})^2 \sum_{k=1}^n |\phi_\omega^{(k)}(r)| r^{k-1} + |x|^{(n+2)/2} (\sqrt{t})^{-2} |\phi_\omega(r)| r \right) \cdot \mathbb{1}_{[0,1]} \left( \frac{r|x|}{2t} \right). \end{aligned}$$

This yields the second estimate. The third estimate results from

$$|B_n^{(j)}(z)| \lesssim |z|^{(n-1)/2-j}, \quad (16)$$

which is a consequence of Proposition 6. Thus

$$\begin{aligned}
 |g_{2,a_2}^{(m)}(\rho)| + |g_{2,a_3}^{(m)}(\rho)| &\stackrel{(11)}{=} 2(2\sqrt{t})^m \left| \frac{d^m}{dr^m} \left( \phi_\omega(r) B_n \left( \frac{r|x|}{2t} \right) \right) \right| \left( \frac{|x|}{2t} \right)^{-n/2} \\
 &\lesssim (\sqrt{t})^m \sum_{k=0}^m |\phi_\omega^{(k)}(r)| \left| B_n^{(m-k)} \left( \frac{r|x|}{2t} \right) \right| \left( \frac{|x|}{2t} \right)^{m-k-n/2} \\
 &\stackrel{(16)}{\lesssim} (\sqrt{t})^m \sum_{k=0}^m |\phi_\omega^{(k)}(r)| \left( \frac{r|x|}{2t} \right)^{(n-1)/2-m+k} \left( \frac{|x|}{2t} \right)^{m-k-n/2} \cdot \mathbb{1}_{[1/2, \infty)} \left( \frac{r|x|}{2t} \right) \\
 &\lesssim |x|^{-1/2} (\sqrt{t})^{m+1} \sum_{k=0}^m |\phi_\omega^{(k)}(r)| r^{(n-1)/2-m+k} \cdot \mathbb{1}_{[1/2, \infty)} \left( \frac{r|x|}{2t} \right). \quad \square
 \end{aligned}$$

*Proof of Theorem 2(i).* We combine the estimates from Propositions 8 and 9. Under the assumption  $\phi_\omega \in C_c^\infty(\mathbb{R}_{\geq 0}; \mathbb{C})$  we get for all  $m \in \{0, \dots, n-1\}$

$$\begin{aligned}
 |\psi(x, t)| &\stackrel{\text{Prop. 8}}{\lesssim} |x|^{(2-n)/2} (\sqrt{t})^{-1} \int_0^\infty |g_{1,a_1}^{(m)}(\rho)| + |g_{2,a_2}^{(m)}(\rho)| + |g_{3,a_3}^{(m)}(\rho)| d\rho \\
 &\stackrel{\text{Prop. 9}}{\lesssim} (\sqrt{t})^{m-n+1} \sum_{k=0}^m \int_0^{\sqrt{t}/|x|} |\phi_\omega^{(k)}(2\sqrt{t}\rho)| (2\sqrt{t}\rho)^{n-m+k-1} d\rho \\
 &\quad + |x|^{(1-n)/2} (\sqrt{t})^m \sum_{k=0}^m \int_{\sqrt{t}/(2|x|)}^\infty |\phi_\omega^{(k)}(2\sqrt{t}\rho)| (2\sqrt{t}\rho)^{(n-1)/2-m+k} d\rho \\
 &\lesssim (\sqrt{t})^{m-n} \sum_{k=0}^m \int_0^{2t/|x|} |\phi_\omega^{(k)}(r)| r^{n-m+k-1} dr \\
 &\quad + (\sqrt{t})^{m-n} \left( \frac{t}{|x|} \right)^{(n-1)/2} \sum_{k=0}^m \int_{t/|x|}^\infty |\phi_\omega^{(k)}(r)| r^{(n-1)/2-m+k} dr \\
 &\lesssim (\sqrt{t})^{m-n} \sum_{k=0}^m \left( \int_0^{2t/|x|} |\phi_\omega^{(k)}(r)| r^{n-m+k-1} dr + \int_{t/|x|}^\infty |\phi_\omega^{(k)}(r)| r^{n-m+k-1} dr \right) \\
 &\lesssim (\sqrt{t})^{m-n} \|\phi_\omega\|_{Y_m}.
 \end{aligned}$$

In the case  $m = n$  we use the second estimate in Proposition 9 instead of the first one. By density of  $C_c^\infty(\mathbb{R}_{\geq 0}; \mathbb{C})$  in  $Y_m$  the result follows.  $\square$

*Proof of Theorem 2(ii).* For  $r = 2\sqrt{t}\rho$  we use

$$|g'_{1,a_1}(\rho)| \lesssim |x|^{(n-2)/2} (\sqrt{t})^{3-n} (|\phi_\omega(r)| r^{n-2} + |\phi'_\omega(r)| r^{n-1}) \cdot \mathbb{1}_{[0,1]} \left( \frac{r|x|}{2t} \right),$$

as well as

$$\begin{aligned}
 |g'_{2,a_2}(\rho)| + |g'_{2,a_3}(\rho)| &\stackrel{(11)}{=} 4\sqrt{t} \left| \frac{d}{dr} \left( \phi_\omega(r) B_n \left( \frac{r|x|}{2t} \right) \right) \right| \left( \frac{|x|}{2t} \right)^{-n/2} \\
 &\lesssim \sqrt{t} \left| \frac{d}{dr} \left( \phi_\omega(r) r^{(n-1)/2} \right) \right| \left| r^{(1-n)/2} B_n \left( \frac{r|x|}{2t} \right) \right| \left( \frac{|x|}{2t} \right)^{-n/2} \\
 &\quad + \sqrt{t} |\phi_\omega(r)| r^{(n-1)/2} \left| \frac{d}{dr} \left( r^{(1-n)/2} B_n \left( \frac{r|x|}{2t} \right) \right) \right| \left( \frac{|x|}{2t} \right)^{-n/2}
 \end{aligned}$$

$$\begin{aligned}
&\stackrel{\text{Prop. 6}}{\lesssim} \sqrt{t} \left| \frac{d}{dr} (\phi_\omega(r) r^{(n-1)/2}) \right| r^{(1-n)/2} \left( \frac{r|x|}{2t} \right)^{(n-1)/2} \left( \frac{|x|}{2t} \right)^{-n/2} \cdot \mathbb{1}_{[1/2, \infty)} \left( \frac{r|x|}{2t} \right) \\
&\quad + \sqrt{t} |\phi_\omega(r)| r^{(n-1)/2} \left( \frac{r|x|}{2t} \right)^{-2} \left( \frac{|x|}{2t} \right)^{1/2} \cdot \mathbb{1}_{[1/2, \infty)} \left( \frac{r|x|}{2t} \right) \\
&\lesssim \left( |x|^{-1/2} (\sqrt{t})^2 \left| \frac{d}{dr} (\phi_\omega(r) r^{(n-1)/2}) \right| + |x|^{-3/2} (\sqrt{t})^4 |\phi_\omega(r)| r^{(n-5)/2} \right) \cdot \mathbb{1}_{[1/2, \infty)} \left( \frac{r|x|}{2t} \right).
\end{aligned}$$

This implies

$$\begin{aligned}
|\psi(x, t)| &\stackrel{\text{Prop. 8}}{\lesssim} |x|^{(2-n)/2} (\sqrt{t})^{-1} \int_0^\infty |g_{1,a_1}^{(m)}(\rho)| + |g_{2,a_2}^{(m)}(\rho)| + |g_{3,a_3}^{(m)}(\rho)| d\rho \\
&\stackrel{\text{Prop. 9}}{\lesssim} (\sqrt{t})^{2-n} \int_0^{\sqrt{t}/|x|} \left( |\phi_\omega(2\sqrt{t}\rho)| (2\sqrt{t}\rho)^{n-2} + |\phi'_\omega(2\sqrt{t}\rho)| (2\sqrt{t}\rho)^{n-1} \right) d\rho \\
&\quad + |x|^{(1-n)/2} \sqrt{t} \int_{\sqrt{t}/(2|x|)}^\infty \left| \frac{d}{dr} (\phi_\omega(r) r^{(n-1)/2}) \right|_{r=2\sqrt{t}\rho} d\rho \\
&\quad + |x|^{(-1-n)/2} (\sqrt{t})^3 \int_{\sqrt{t}/(2|x|)}^\infty |\phi_\omega(2\sqrt{t}\rho)| (2\sqrt{t}\rho)^{(n-5)/2} d\rho \\
&\lesssim |x|^{(1-n)/2} \left( \frac{t}{|x|} \right)^{(1-n)/2} \int_0^{2t/|x|} (|\phi_\omega(r)| r^{n-2} + |\phi'_\omega(r)| r^{n-1}) dr \\
&\quad + |x|^{(1-n)/2} \left( \int_{t/|x|}^\infty \left| \frac{d}{dr} (\phi_\omega(r) r^{(n-1)/2}) \right| dr + \frac{t}{|x|} \int_{t/|x|}^\infty |\phi_\omega(r)| r^{(n-5)/2} dr \right) \\
&\lesssim |x|^{(1-n)/2} \|\phi_\omega\|_X.
\end{aligned}$$

So we get the result by density of  $C_c^\infty(\mathbb{R}_{\geq 0}; \mathbb{C})$  in  $X$ . □

### 3. Proof of Theorem 1

We estimate the solution of the Schrödinger equation for the initial datum

$$\phi(x) = \phi_{\text{rad}}(|x|), \quad \text{where } \phi_{\text{rad}}(\rho) := e^{-i\rho^2/4} \mathbb{1}_{\rho \geq 1} \rho^{-\sigma}$$

and  $\sigma$  is chosen according to  $(n-3)/2 < \sigma < n$ . In this case, the formula (5) is well-defined and provides a solution of the initial value problem (1). In [Bona et al. 2014, Section 2.1] it was shown that the solution  $\psi$  blows up in  $L^\infty(\mathbb{R}^n)$  as  $t \rightarrow 1$  provided that  $n/2 < \sigma < n$  holds. In fact, in this case the function

$$a(x) := \mathbb{1}_{|x| \geq 1} |x|^{-\sigma}$$

lies in  $L^2(\mathbb{R}^n)$  but not in  $L^1(\mathbb{R}^n)$  so that [Bona et al. 2014, Remark 2.2] applies. We now generalize this analysis to the range  $(n-3)/2 < \sigma < n$  and detect a self-similar blow-up in  $L^q(\mathbb{R}^n)$  for all  $q > n/(n-\sigma)$ , and a lower estimate for the corresponding blow-up rate then implies  $\|\psi\|_{L^p(\mathbb{R}, L^q(\mathbb{R}^n))} = \infty$  for  $p \geq 2q/((n-\sigma)q - n)_+$ . From this we will finally deduce the nonvalidity of Strichartz estimates for initial data  $\phi \in L^r(\mathbb{R}^n)$ , where  $r > 2$ .

We set  $k_t := \sqrt{1/(4t) - 1/4}$  for  $0 \leq t < 1$  and write  $\psi(x) = \psi_{\text{rad}}(|x|)$ . We get for  $|x| = 2tk_t z$

$$\begin{aligned} 2|\psi(x, t)|k_t^{n-\sigma} &\stackrel{(5)}{=} |x|^{(2-n)/2}t^{-1}k_t^{n-\sigma} \left| \int_0^\infty J_{(n-2)/2}\left(\frac{\rho|x|}{2t}\right)\phi_{\text{rad}}(\rho)\rho^{n/2}e^{i\rho^2/(4t)}d\rho \right| \\ &= |x|^{(2-n)/2}t^{-1}k_t^{n-\sigma} \left| \int_1^\infty J_{(n-2)/2}\left(\frac{\rho|x|}{2t}\right)\rho^{n/2-\sigma}e^{i\rho^2k_t^2}d\rho \right| \\ &= (2tk_t z)^{(2-n)/2}t^{-1}k_t^{n-\sigma} \left| \int_1^\infty J_{(n-2)/2}(\rho k_t z)\rho^{n/2-\sigma}e^{i\rho^2k_t^2}d\rho \right| \\ &= (2tk_t z)^{(2-n)/2}t^{-1}k_t^{(n-2)/2} \left| \int_{k_t}^\infty J_{(n-2)/2}(sz)s^{n/2-\sigma}e^{is^2}ds \right| \\ &\rightarrow (2z)^{(2-n)/2} \left| \int_0^\infty J_{(n-2)/2}(sz)s^{n/2-\sigma}e^{is^2}ds \right| \quad \text{as } t \rightarrow 1 \end{aligned}$$

and the convergence is locally uniform in  $z \in (0, \infty)$  due to  $(n - 3)/2 < \sigma < n$ . Since the right-hand side is not identically zero we may find  $\delta > 0$  and radii  $0 < R_1 < R_2$  such that

$$|\psi(x, t)| \gtrsim k_t^{\sigma-n} \quad \text{for } R_1k_t \leq |x| \leq R_2k_t \text{ and } 1 - \delta < t < 1.$$

Hence we get

$$\begin{aligned} \int_{1-\delta}^1 \left( \int_{B_{R_2k_t}(0) \setminus B_{R_1k_t}(0)} |\psi(x, t)|^q dx \right)^{p/q} dt &\gtrsim \int_{1-\delta}^1 \left( \int_{R_1k_t}^{R_2k_t} r^{n-1}k_t^{(\sigma-n)q} dr \right)^{p/q} dt \\ &\gtrsim \int_{1-\delta}^1 (k_t^{n+(\sigma-n)q})^{p/q} dt \\ &\gtrsim \int_{1-\delta}^1 (1-t)^{p(n+(\sigma-n)q)/(2q)} dt. \end{aligned}$$

This integral is finite if and only if  $(p/(2q))(n + (\sigma - n)q) > -1$ . So  $\psi \in L^p(\mathbb{R}; L^q(\mathbb{R}^n))$  can only hold for  $p < 2q/((n - \sigma)q - n)_+$ . Moreover, the initial datum lies in  $L^r(\mathbb{R}^n)$  if and only if  $\sigma > n/r$ . So, for any  $r > 1$  we can consider the limit  $\sigma \searrow \max\{n/r, (n - 3)/2\}$ , and we find that the validity of the Strichartz estimate (2) with initial datum in  $L^r(\mathbb{R}^n)$ ,  $r > 1$ , implies

$$p \leq \frac{2q}{((n - \max\{n/r, (n - 3)/2\})q - n)_+} = \max \left\{ \frac{2qr}{n((r - 1)q - r)_+}, \frac{4q}{((n + 3)q - 2n)_+} \right\}. \tag{17}$$

On the other hand, the scaling invariance of the Schrödinger equation implies  $2/p + n/q = n/r$  and thus  $p = 2qr/(n(q - r))$ . Plugging this into (17) we obtain  $r \leq 2$ . Hence, the Strichartz estimate (2) cannot hold for any  $r > 2$ , which finishes the proof. □

### Appendix

In this appendix we briefly discuss the restriction  $p \geq 2$  in the context of Strichartz estimates of the form

$$\|e^{it\Delta}\phi\|_{L_t^p(\mathbb{R}; L^q(\mathbb{R}^n))} \lesssim \|\phi\|_{L^2(\mathbb{R}^n)},$$

which results from an abstract reasoning involving translation-invariant operators due to [Hörmander 1960]; see [Keel and Tao 1998, p. 970–971]. Here we provide a family of explicit counterexamples that not only implies  $p \geq 2$  for square integrable initial data, but even shows  $p \geq 2r/(2n - r(n - 1))_+$  for initial data in  $L^r(\mathbb{R}^n)$  with  $r > 2n/(n + 1)$ . In order to avoid lengthy computations involving oscillatory integrals, we only sketch the proofs. The starting point is a reasonable choice of an initial condition. We choose

$$\phi(x) = |x|^{(2-n)/2} \int_1^2 (\omega - 1)^{-\delta} J_{(n-2)/2}(\omega|x|) d\omega$$

for  $\delta \in (0, 1)$ . This function corresponds to a singular superposition of Herglotz waves; see Remark 1(c). It is smooth and lengthy computations involving the van der Corput lemma [Stein 1993, p. 334] reveal

$$|\phi(x)| \sim 2|x|^{(1-n)/2-(1-\delta)} \int_0^\infty \operatorname{Re}(e^{i\rho} \alpha_0 e^{i(|x| - ((n-1)\pi/4))\rho}) \rho^{-\delta} d\rho \quad \text{as } |x| \rightarrow \infty,$$

where  $\alpha_0 > 0$  is the dominant term in the series expansion of the Bessel function near infinity; see (8). In particular we get  $\phi \in L^r(\mathbb{R}^n)$  if and only if  $\delta < (n + 1)/2 - n/r$ .

The above choice for the initial datum allows to write down the corresponding solution of the Schrödinger equation semiexplicitly via

$$\widehat{\psi(\cdot, t)}(\xi) = e^{-it|\xi|^2} \widehat{\phi}(\xi) = e^{-it|\xi|^2} |\xi|^{-n/2} (|\xi| - 1)^{-\delta} \mathbb{1}_{[1,2]}(|\xi|).$$

Hence, one gets

$$\psi(x, t) = |x|^{(2-n)/2} \int_1^2 e^{-it\omega^2} (\omega - 1)^{-\delta} J_{(n-2)/2}(\omega|x|) d\omega,$$

and the van der Corput Lemma implies  $|\psi(x, t)| \gtrsim ct^{\delta-1}$  for small  $|x|$  and large  $t$ . In particular,  $\psi \in L^p(\mathbb{R}; L^q(\mathbb{R}^n))$  implies  $p(1-\delta) > 1$ . So we conclude that for any  $r \in (2n/(n+1), \infty]$  we can consider the limit  $\delta \nearrow \min\{1, (n+1)/2 - n/r\}$  in the above computations and obtain that  $\psi \in L^p(\mathbb{R}; L^q(\mathbb{R}^n))$  implies

$$p \geq \frac{2r}{(2n - r(n - 1))_+} \quad \text{provided } r > \frac{2n}{n + 1}. \quad (18)$$

For  $r = 2$ , i.e., for square integrable initial data, this implies  $p \geq 2$ , which is all we wanted to demonstrate.

Let us mention that a detailed analysis of  $\psi$  reveals  $\psi \in L^p(\mathbb{R}; L^q(\mathbb{R}^n))$  if and only if

$$q > \max \left\{ \frac{2n - 1}{n - \delta}, \frac{2n}{n + 1 - 2\delta} \right\},$$

$$p > \max \left\{ \frac{1}{1 - \delta}, \frac{2q}{q(n - \delta) + 1 - 2n}, \frac{2q}{q(n + 1 - 2\delta) - 2n} \right\}.$$

Keeping the scaling condition  $2/p + n/q = n/r$  in mind, these a priori more restrictive conditions do however not result in stronger necessary conditions than (18), so that further necessary conditions cannot be deduced from this example.

## Acknowledgements

This work was funded by the Deutsche Forschungsgemeinschaft (DFG, German Research Foundation) Project-ID 258734477-SFB 1173. The author thanks Dominic Scheider (Karlsruhe Institute of Technology) and Martin Spitz (University of Bielefeld) for valuable remarks and discussions leading to an improvement of the manuscript.

## References

- [Bona et al. 2014] J. L. Bona, G. Ponce, J.-C. Saut, and C. Sparber, “Dispersive blow-up for nonlinear Schrödinger equations revisited”, *J. Math. Pures Appl.* (9) **102**:4 (2014), 782–811. [MR](#) [Zbl](#)
- [Cazenave 2003] T. Cazenave, *Semilinear Schrödinger equations*, Courant Lecture Notes in Mathematics **10**, Courant Institute of Mathematical Sciences, New York, 2003. [MR](#) [Zbl](#)
- [Ginibre and Velo 1985] J. Ginibre and G. Velo, “The global Cauchy problem for the nonlinear Schrödinger equation revisited”, *Ann. Inst. H. Poincaré Anal. Non Linéaire* **2**:4 (1985), 309–327. [MR](#) [Zbl](#)
- [Hörmander 1960] L. Hörmander, “Estimates for translation invariant operators in  $L^p$  spaces”, *Acta Math.* **104** (1960), 93–140. [MR](#) [Zbl](#)
- [Keel and Tao 1998] M. Keel and T. Tao, “Endpoint Strichartz estimates”, *Amer. J. Math.* **120**:5 (1998), 955–980. [MR](#) [Zbl](#)
- [Linares and Ponce 2015] F. Linares and G. Ponce, *Introduction to nonlinear dispersive equations*, 2nd ed., Springer, 2015. [MR](#) [Zbl](#)
- [Stein 1993] E. M. Stein, *Harmonic analysis: real-variable methods, orthogonality, and oscillatory integrals*, Princeton Mathematical Series **43**, Princeton University Press, 1993. [MR](#) [Zbl](#)
- [Strichartz 1977] R. S. Strichartz, “Restrictions of Fourier transforms to quadratic surfaces and decay of solutions of wave equations”, *Duke Math. J.* **44**:3 (1977), 705–714. [MR](#) [Zbl](#)
- [Watson 1944] G. N. Watson, *A treatise on the theory of Bessel functions*, Cambridge University Press, 1944. [MR](#) [Zbl](#)

Received 3 Nov 2019. Revised 21 Jan 2020. Accepted 21 Feb 2020.

RAINER MANDEL: [rainer.mandel@kit.edu](mailto:rainer.mandel@kit.edu)

Karlsruhe Institute of Technology, Institute for Analysis, Karlsruhe, Germany

## EDITORS-IN-CHIEF

- Charles L. Epstein University of Pennsylvania  
[cle@math.upenn.edu](mailto:cle@math.upenn.edu)
- Maciej Zworski University of California at Berkeley  
[zworski@math.berkeley.edu](mailto:zworski@math.berkeley.edu)

## EDITORIAL BOARD

- Sir John M. Ball University of Oxford  
[ball@maths.ox.ac.uk](mailto:ball@maths.ox.ac.uk)
- Michael P. Brenner Harvard University  
[brenner@seas.harvard.edu](mailto:brenner@seas.harvard.edu)
- Charles Fefferman Princeton University  
[cf@math.princeton.edu](mailto:cf@math.princeton.edu)
- Susan Friedlander University of Southern California  
[susanfri@usc.edu](mailto:susanfri@usc.edu)
- Anna Gilbert University of Michigan  
[annacg@umich.edu](mailto:annacg@umich.edu)
- Leslie F. Greengard Courant Institute, New York University, and  
Flatiron Institute, Simons Foundation  
[greengard@cims.nyu.edu](mailto:greengard@cims.nyu.edu)
- Yan Guo Brown University  
[yan\\_guo@brown.edu](mailto:yan_guo@brown.edu)
- Claude Le Bris CERMICS - ENPC  
[lebris@cermics.enpc.fr](mailto:lebris@cermics.enpc.fr)
- Robert J. McCann University of Toronto  
[mccann@math.toronto.edu](mailto:mccann@math.toronto.edu)
- Michael O'Neil Courant Institute, New York University  
[oneil@cims.nyu.edu](mailto:oneil@cims.nyu.edu)
- Galina Perelman Université Paris-Est Créteil  
[galina.perelman@u-pec.fr](mailto:galina.perelman@u-pec.fr)
- Jill Pipher Brown University  
[jill\\_pipher@brown.edu](mailto:jill_pipher@brown.edu)
- Johannes Sjöstrand Université de Dijon  
[johannes.sjostrand@u-bourgogne.fr](mailto:johannes.sjostrand@u-bourgogne.fr)
- Vladimir Šverák University of Minnesota  
[sverak@math.umn.edu](mailto:sverak@math.umn.edu)
- Daniel Tataru University of California at Berkeley  
[tataru@berkeley.edu](mailto:tataru@berkeley.edu)
- Michael I. Weinstein Columbia University  
[miw2103@columbia.edu](mailto:miw2103@columbia.edu)
- Jon Wilkening University of California at Berkeley  
[wilken@math.berkeley.edu](mailto:wilken@math.berkeley.edu)
- Enrique Zuazua DeustoTech-Bilbao, and  
Universidad Autónoma de Madrid  
[enrique.zuazua@deusto.es](mailto:enrique.zuazua@deusto.es)

## PRODUCTION

- Silvio Levy (Scientific Editor)  
[production@msp.org](mailto:production@msp.org)

**Cover image:** The figure shows the outgoing scattered field produced by scattering a plane wave, coming from the northwest, off of the (stylized) letters P A A. The total field satisfies the homogeneous Dirichlet condition on the boundary of the letters. It is based on a numerical computation by Mike O'Neil of the Courant Institute.

---

See inside back cover or [msp.org/paa](http://msp.org/paa) for submission instructions.

---

The subscription price for 2020 is US \$505/year for the electronic version, and \$565/year (+\$25, if shipping outside the US) for print and electronic. Subscriptions, requests for back issues and changes of subscriber address should be sent to MSP.

---

Pure and Applied Analysis (ISSN 2578-5885 electronic, 2578-5893 printed) at Mathematical Sciences Publishers, 798 Evans Hall #3840, c/o University of California, Berkeley, CA 94720-3840 is published continuously online. Periodical rate postage paid at Berkeley, CA 94704, and additional mailing offices.

---

PAA peer review and production are managed by EditFlow<sup>®</sup> from MSP.

PUBLISHED BY

# PURE and APPLIED ANALYSIS

vol. 2 no. 2 2020

- Hypocoercivity without confinement 203  
EMERIC BOUIN, JEAN DOLBEAULT, STÉPHANE MISCHLER, CLÉMENT MOUHOT  
and CHRISTIAN SCHMEISER
- The hyperbolic Yang–Mills equation in the caloric gauge: local well-posedness and control of energy-dispersed solutions 233  
SUNG-JIN OH and DANIEL TATARU
- Scattering resonances on truncated cones 385  
DEAN BASKIN and MENGXUAN YANG
- The Kähler geometry of certain optimal transport problems 397  
GABRIEL KHAN and JUN ZHANG
- Semiclassical asymptotics for nonselfadjoint harmonic oscillators 427  
VÍCTOR ARNAIZ and GABRIEL RIVIÈRE
- On the solution of Laplace’s equation in the vicinity of triple junctions 447  
JEREMY HOSKINS and MANAS RACHH
- Enhanced convergence rates and asymptotics for a dispersive Boussinesq-type system with large ill-prepared data 477  
FRÉDÉRIC CHARVE
- Dispersive estimates, blow-up and failure of Strichartz estimates for the Schrödinger equation with slowly decaying initial data 519  
RAINER MANDEL