

Pacific Journal of Mathematics

LOOPS WITH TRANSITIVE AUTOMORPHISM GROUPS

R. H. BRUCK

LOOPS WITH TRANSITIVE AUTOMORPHISM GROUPS

R. H. BRUCK

1. Introduction. Every automorphism of an additive loop¹ L maps 0 upon 0. The automorphism group $A(L)$ of L will be called *transitive* if $A(L)$ is transitive on the nonzero elements of L . It is readily deduced from work of L. J. Paige [4] and P. T. Bateman [3] that, for every cardinal number n , there exists a loop L of cardinal number n with a transitive automorphism group. We shall demonstrate that (aside from the obvious exceptions) such a loop L must be simple, that is, its only normal subloops must be 0 and L , if it satisfies the following ascending chain condition:

(C) If $N_1 \subset N_2 \subset N_3 \subset \dots$ is an ascending chain of normal subloops of the loop L , there exists an integer i such that $N_i = N_{i+1}$.

2. Theorem. We shall establish the following result.

THEOREM 1. *An additive loop L which satisfies (C) and has a transitive automorphism group is either (i) a simple loop or (ii) a finite abelian p -group of type (p, ρ, \dots, p) .*

Proof. For each nonzero a of L , denote by $M(a)$ the smallest normal subloop of L which contains a .

(1) *The subloop $M(a)$ has a transitive automorphism group and is a minimal normal subloop of L .* If $b \neq 0$ is in $M(a)$, then there exists $\theta \in A(L)$ such that $a\theta = b$. Since θ maps normal subloops upon normal subloops, we have $M(a)\theta = M(b)$. Since $b \in M(a)$, it follows that $M(b) \subset M(a)$. If $\phi = \theta^{-1}$, then $M(a) = M(b)\phi \subset M(a)\phi$, and, by induction, $M(a) \subset M(a)\phi \subset M(a)\phi^2 \subset \dots$. In view of (C), we have $M(a)\phi^i = M(a)\phi^{i+1}$ for some integer i . Since ϕ is an automorphism of L , it follows that $M(a) = M(a)\phi^{-1} = M(a)\theta = M(b)$. Hence θ induces an automorphism of $M(a)$. This is enough to prove (1).

¹ Readers unfamiliar with loop theory will get the sense of the paper if they read *group* in place of *loop*. The necessary loop theory will be found in Baer [1, 2].

Pacific J. Math. 1 (1951), 481-483.

(2) If N is a normal subloop of L , then $N \cap M(a) = 0$ or $M(a)$. This follows from the minimality of $M(a)$.

(3) The loop L is a direct sum of a finite number r of isomorphic simple subloops M_i ; that is, $L = M_1 \oplus M_2 \oplus \cdots \oplus M_r$. If a, b are nonzero elements of L , there exists $\theta \in A(L)$ such that $a\theta = b$. Then $M(a)\theta = M(b)$, showing that all the minimal normal subloops $M(a)$ are isomorphic. If a_1 is an arbitrary nonzero element of L , define $M_1 = M(a_1)$. Now suppose that $M_i = M(a_i)$ has been defined, for $i = 1, 2, \cdots, s$, such that the (normal) subloop N_s generated by the M_i is the direct sum $N_s = M_1 \oplus \cdots \oplus M_s$. Write $t = s + 1$. If there exists a nonzero element a_t of L which is not in N_s , define $M_t = M(a_t)$. Then $N_s \cap M_t = 0$, by (2), and hence $N_t = N_s \oplus M_t = M_1 \oplus \cdots \oplus M_t$. In view of (C), the strictly increasing chain $N_1 \subset N_2 \subset \cdots$ must be finite. Therefore $L = N_r$ for some integer r . If M' is a normal subloop of M_1 , then M' is normal in L , by virtue of the direct decomposition. Hence, by (1), each M_i is simple. This proves (3).

The center $Z(L)$ of a loop L is a characteristic subloop and an abelian group. In view of (1), either $Z(M_i) = 0$ or $Z(M_i) = M_i$. Hence, by (3), either (i') $Z(L) = 0$ or (ii') L is a direct sum of isomorphic simple abelian groups. Since a simple abelian group is cyclic of prime order p , (ii') implies (ii) of Theorem 1. (Conversely, every finite abelian p -group of type (p, p, \cdots, p) satisfies the hypotheses of the theorem.) In the case (i'), assume $r > 1$ in (3). Since $Z(L) = 0$, the decomposition (3) is unique. However, the nonzero element $c = a_1 + a_2$ is in $M_1 \oplus M_2$ but not in any of the M_i . Yet the proof of (3) shows that $M(c)$ could be chosen as the first factor in the direct decomposition of L , a contradiction. Therefore $r = 1$, and we have (i). This completes the proof of Theorem 1.

As the following (trivial) theorem shows, simple loops need not have transitive automorphism groups:

THEOREM 2. *A finite simple group $G \neq 0$ with a transitive automorphism group is necessarily cyclic of prime order.*

Proof. Every nonzero element of G has the same order p , necessarily prime. Thus G is a p -group, $Z(G) \neq 0$, $Z(G) = G$, and G is cyclic of order p .

3. Remarks. The author does not know whether finiteness is necessary for the conclusion of Theorem 2.

The following is the nonabelian loop L of lowest order with a transitive automorphism group; it is readily verified that $A(L)$ is the (alternating) group of order 12 generated by (12)(34) and (123):

+	0	1	2	3	4
0	0	1	2	3	4
1	1	0	3	4	2
2	2	4	0	1	3
3	3	2	4	0	1
4	4	3	1	2	0

REFERENCES

1. Reinhold Baer, *The homomorphism theorems for loops*, Amer. J. Math. 67 (1945), 450-460.
2. ———, *Direct decompositions*, Trans. Amer. Math. Soc. 62 (1947), 62-98.
3. P. T. Bateman, *A remark on infinite groups*, Amer. Math. Monthly 57 (1950), 623-624.
4. L. J. Paige, *Neofields*, Duke Math. J. 16 (1949), 39-60.

UNIVERSITY OF WISCONSIN

PACIFIC JOURNAL OF MATHEMATICS

EDITORS

M.M. SCHIFFER*

Stanford University
Stanford, California

E. HEWITT

University of Washington
Seattle 5, Washington

R. P. DILWORTH

California Institute of Technology
Pasadena 4, California

E.F. BECKENBACH**

University of California
Los Angeles 24, California

ASSOCIATE EDITORS

H. BUSEMANN

P. R. HALMOS

BØRGE JESSEN

J. J. STOKER

HERBERT FEDERER

HEINZ HOPF

PAUL LÉVY

E. G. STRAUS

MARSHALL HALL

R. D. JAMES

GEORGE PÓLYA

KÔSAKU YOSIDA

SPONSORS

UNIVERSITY OF BRITISH COLUMBIA

CALIFORNIA INSTITUTE OF TECHNOLOGY

UNIVERSITY OF CALIFORNIA, BERKELEY

UNIVERSITY OF CALIFORNIA, DAVIS

UNIVERSITY OF CALIFORNIA, LOS ANGELES

UNIVERSITY OF CALIFORNIA, SANTA BARBARA

UNIVERSITY OF NEVADA

OREGON STATE COLLEGE

UNIVERSITY OF OREGON

UNIVERSITY OF SOUTHERN CALIFORNIA

STANFORD RESEARCH INSTITUTE

STANFORD UNIVERSITY

WASHINGTON STATE COLLEGE

UNIVERSITY OF WASHINGTON

* * *

AMERICAN MATHEMATICAL SOCIETY

HUGHES AIRCRAFT COMPANY

Mathematical papers intended for publication in the *Pacific Journal of Mathematics* should be typewritten (double spaced), and the author should keep a complete copy. Manuscripts may be sent to any of the editors. Manuscripts intended for the outgoing editors should be sent to their successors. All other communications to the editors should be addressed to the managing editor, E.G. Straus, at the University of California Los Angeles 24, California.

50 reprints of each article are furnished free of charge; additional copies may be obtained at cost in multiples of 50.

The *Pacific Journal of Mathematics* is published quarterly, in March, June, September, and December. The price per volume (4 numbers) is \$12.00; single issues, \$3.50; back numbers (Volumes 1, 2, 3) are available at \$2.50 per copy. Special price to individual faculty members of supporting institutions and to individual members of the American Mathematical Society: \$4.00 per volume; single issues, \$1.25.

Subscriptions, orders for back numbers, and changes of address should be sent to the publishers, University of California Press, Berkeley 4, California.

Printed at Ann Arbor, Michigan. Entered as second class matter at the Post Office, Berkeley, California.

* To be succeeded in 1955, by H.L. Royden, Stanford University, Stanford, California.

** To be succeeded in 1955, by E.G. Straus, University of California, Los Angeles 24, Calif.

UNIVERSITY OF CALIFORNIA PRESS · BERKELEY AND LOS ANGELES

COPYRIGHT 1954 BY PACIFIC JOURNAL OF MATHEMATICS

R. H. Bruck, <i>Loops with transitive automorphism groups</i>	481
Paul R. Garabedian, <i>A partial differential equation arising in conformal mapping</i>	485
Magnus R. Hestenes, <i>Applications of the theory of quadratic forms in Hilbert space to the calculus of variations</i>	525
Sze-Tsen Hu, <i>On the realizability of homotopy groups and their operations</i>	583