

Pacific Journal of Mathematics

**THE NORMAL COMPLETION OF A SUBSET OF A COMPLETE
LATTICE AND LATTICES OF CONTINUOUS FUNCTIONS**

ALFRED HORN

THE NORMAL COMPLETION OF A SUBSET OF A COMPLETE LATTICE AND LATTICES OF CONTINUOUS FUNCTIONS

ALFRED HORN

Introduction. The normal completion of a partially ordered set C is a complete lattice \tilde{C} whose elements are subsets of C . However, in practice a partially ordered set C is usually given as a subset of a complete lattice B , and it seems desirable to be able to describe directly a subset of B which contains C and which is isomorphic with \tilde{C} . Such a construction is given in § 1 below. The idea of the construction is suggested by the method of formation of the normal completion itself, and is undoubtedly known. However, it does not seem to appear explicitly in the literature. In § 2 we treat the case where C is the lattice of real-valued continuous functions on a topologic space X , and B is the lattice of all real-valued functions on X . This leads to a simple proof of a result of Dilworth [2]. In § 3 we extend another result of Dilworth in proving that for any two topologic spaces, the lattices of regular open sets are isomorphic if and only if the lattices of normal lower semicontinuous functions are isomorphic.

Notations and definitions. If S is a subset of a partially ordered set P , then $\sum_P(S)$ and $\prod_P(S)$ will be used to denote the least upper bound and the greatest lower bound of S , respectively. If S is empty, $\sum_P(S)$ is the least element of P , and $\prod_P(S)$ is the greatest element of P . A subset of P will always be given the order induced by that of P . If $x \in P$, $S \subset P$, we shall write $x \geq S$ whenever $x \geq y$ for all y in S . Two partially ordered sets P_1 and P_2 are said to be isomorphic, in symbols $P_1 \cong P_2$, if there exists a function f on P_1 onto P_2 such that

$$x \leq y \leftrightarrow f(x) \leq f(y).$$

This implies that f is one-to-one.

If A is a subset of a topologic space, we use $\mathcal{I}A$, and $\mathcal{C}A$ for the interior and the closure of A . A set S is called *regular open* if $S = \mathcal{I}\mathcal{C}S$, and *regular closed* if $S = \mathcal{C}\mathcal{I}S$. A zero-dimensional space is one whose open sets have a basis each member of which is both open and closed. A space shall be called

Received December 2, 1951.

Pacific J. Math. 3 (1953), 137-152

extremally disconnected if every regular open set is closed, that is, if the closure of every open set is open. Contrary to the usual custom, we do not require a completely regular space to satisfy any of the separation axioms. If A is a set, k_A will stand for the characteristic function of A . If f is a function, we use $E(f > \lambda)$ to denote $E_x(f(x) > \lambda)$.

1. The normal completion of a subset of a complete lattice. Let B be a complete lattice on which are defined a “closure operation” x^* and a “dual closure operation” x_* with the following properties:

- i) $x_* \leq x \leq x^*$,
- ii) $(x_*)_* = x_*$, $(x^*)^* = x^*$,
- iii) $x \leq y \implies x_* \leq y_*$ and $x^* \leq y^*$.

As a simple consequence, we have

- iv) $((x_*)^*)_*$ = $(x_*)^*$ and $((x^*)_*)^*$ = $(x^*)_*$.

Throughout this section, $\sum(S)$ and $\prod(S)$ will be used to denote $\sum_B(S)$ and $\prod_B(S)$. Let

$$L = E_x(x = x_*), \text{ and } U = E_x(x = x^*).$$

A proof of the following theorem is given in [1, p.49].

THEOREM 1. *L and U are complete lattices in which*

$$\begin{aligned} \sum_L(S) &= \sum(S), & \prod_L(S) &= \prod(S), \\ \sum_U(S) &= \sum(S)^*, & \prod_U(S) &= \prod(S). \end{aligned}$$

Now let

$$NL = E_x(x = (x^*)_*), \text{ and } NU = E_x(x = (x_*)^*).$$

We note that $NL \subset L$ and $NU \subset U$, by ii).

THEOREM 2. *NL and NU are isomorphic complete lattices in which*

$$\begin{aligned} \sum_{NL}(S) &= (\sum(S)^*)_*, & \prod_{NL}(S) &= \prod(S)_*, \\ \sum_{NU}(S) &= \sum(S)^*, & \prod_{NU}(S) &= (\prod(S)_*)^*. \end{aligned}$$

Proof. It is easy to verify by i), ii), iii), and iv) that $(x^*)_*$ is a closure operation in L . By Theorem 1 it follows that NL is a complete lattice in which

$$\sum_{NL}(S) = (\sum_L(S)^*)_* = (\sum(S)^*)_*, \prod_{NL}(S) = \prod_L(S) = \prod(S)_*.$$

For the other half of the proof, we define the dual closure operation $(x_*)^*$ in U , and apply Theorem 1 again. The mapping $f(x) = x^*$ is easily seen to be an isomorphism of NL onto NU whose inverse is $f^{-1}(x) = x_*$.

As an example, if B is the lattice of all subsets of a topologic space, and

$$x_* = \mathcal{I}x, \quad x^* = \mathcal{C}x,$$

then L and U are the lattices of open sets and closed sets, while NL and NU are the lattices of regular open sets and regular closed sets respectively.

The normal completion (completion by cuts) of a partially ordered set C is a canonical embedding of C in a complete lattice \tilde{C} which preserves all existing sums and products. See [1, p.58] for details. Now \tilde{C} may be constructed as follows. If $S \subset C$, let

$$S^\wedge = E_x(x \in C, x \geq S) \quad \text{and} \quad \bar{S} = E_x(x \in C, x \leq S^\wedge).$$

Then \bar{S} is a closure operation in the lattice of all subsets of C . The class

$$\tilde{C} = E_S(S = \bar{S})$$

is the normal completion of C ; C is embedded in \tilde{C} by the mapping

$$f(x) = \bar{S}_x,$$

where S_x is the class whose only member is x .

Now suppose C is an arbitrary subset of B . Our aim is to describe a subset of B which is isomorphic with \tilde{C} . If $x \in B$, we define

$$x_0 = E_y(y \in C, y \leq x) \quad \text{and} \quad x^0 = E_y(y \in C, y \geq x).$$

Now let

$$x_* = \sum(x_0), \quad x^* = \prod(x^0).$$

It is easily verified that x^* and x_* are closure and dual closure operations on B . To check property ii), we need only note that if $y \in C$, then

$$y \leq x_* \leftrightarrow y \leq x.$$

We also remark that

$$x \in C \rightarrow x^* = x = x_*.$$

We now consider the lattices L , U , NL , and NU obtained from this particular definition of x^* , x_* . Obviously $C \subset NL$, $C \subset NU$, and the following theorem shows that NL and NU are extensions of C .

THEOREM 3. *If*

$$x = \sum_C(S), \quad y = \prod_C(S),$$

then

$$x = \sum_{NL}(S) = \sum_{NU}(S) \text{ and } y = \prod_{NL}(S) = \prod_{NU}(S).$$

Proof. First

$$w \in C, \quad w \geq \sum(S) \rightarrow w \in C, \quad w \geq S \rightarrow w \geq \sum_C(S) = x.$$

hence

$$\sum(S)^0 \geq x,$$

and therefore

$$\sum(S)^* \geq x.$$

Also

$$\sum(S)^* \leq x$$

since $x \in C$ and $x \geq \sum(S)$. Thus

$$x = \sum(S)^* = \sum_{NU}(S).$$

Also, since $x \in C$,

$$x = x_* = (\sum(S)^*)_* = \sum_{NL}(S).$$

We omit the proof of the dual statement.

THEOREM 4. NL (and hence NU) is isomorphic with \tilde{C} .

Proof. Let S be any subset of C . We first note that $S^\wedge = \sum(S)^0$ and that $x \leq \sum(S)^0$ if and only if $x \leq \sum(S)^*$. Hence

$$(1) \quad \bar{S} = E_x(x \in C, x \leq \sum(S)^*).$$

It follows that

$$(2) \quad \sum(\bar{S}) = (\sum(S)^*)_*.$$

Thus if $S = \bar{S}$, then $\sum(S) \in NL$.

We now define a mapping ϕ of \tilde{C} onto NL by

$$\phi(S) = \sum(S).$$

ϕ is onto NL because

$$x \in NL \rightarrow x \in L \rightarrow x = \sum(x_0),$$

and by (2),

$$\phi(\bar{x}_0) = \sum(\bar{x}_0) = (x^*)_* = x.$$

Finally if $S \subset T \in \tilde{C}$, then $\phi(S) \leq \phi(T)$, while by (1),

$$\phi(S) \leq \phi(T) \rightarrow S = E_x(x \in C, x \leq \phi(S)^*) \subset E_x(x \in C, x \leq \phi(T)^*) = T.$$

Using Theorem 4, we can show that C may be characterized as follows: Let A be a complete lattice which contains a subset D isomorphic with C and such that, for any $S \subset D$, whenever $\sum_D(S)$ or $\prod_D(S)$ exists then it is equal to $\sum_A(S)$ or $\prod_A(S)$. If no proper subset of A which contains D is a complete lattice, then A is isomorphic with \tilde{C} .

As applications of Theorem 4 we have the following examples. If C is the class of open and closed subsets of a zero-dimensional space Z , then \tilde{C} is isomorphic with the lattice of regular open sets of Z . For it is easily seen that if x is any subset of Z , then

$$x^* = Cx, \text{ and } x_* = \partial x.$$

If C is a class of subsets of a set A which contains all one-point subsets and their complements, then \tilde{C} is isomorphic with the lattice of all subsets of A . For $x = x_* = x^*$ whenever $x \subset A$. Our main application is given in the next section.

A special result which we shall use later is the following.

THEOREM 5. *If B has the property that*

$$x \cap \sum x_\alpha = \sum (x \cap x_\alpha)$$

for any family $\{x_\alpha\}$, and

$$\prod_C(S) = \prod(S)$$

for any finite subset S of C , then

$$\prod_L(S) = \prod(S) \text{ and } \prod_{NL}(S) = \prod(S)$$

for any finite subset S of L and NL respectively.

Proof. Let x_1, \dots, x_n be elements of L . Then

$$x_i = \sum_{\alpha \in I_i} y_{\alpha, i},$$

where the $y_{\alpha, i}$ are elements of $(x_i)_0$. By [1, Theorem 14, p. 146], we have

$$\prod x_i = \sum_f \prod_i y_{f(i), i},$$

where f ranges over all functions such that $f(i) \in I_i$. Since $y_{f(i), i} \in C$, it follows that

$$\prod x_i = \sum_f y_f,$$

where $y_f \in C$. Hence $\prod x_i \in L$, since L is closed under \sum , and therefore

$$\prod_L(x_i) = \prod x_i.$$

If $x_i \in NL$, then

$$\prod_{NL} x_i = \prod_L x_i = \prod x_i.$$

2. The normal completion of lattices of continuous functions. Hereafter B will stand for the set of all real-valued functions on a topologic space X . We allow the functions in B to assume ∞ and $-\infty$ as values; B is then a complete lattice in which

$$\sum(S)(x) = \sup_{f \in S} f(x).$$

We define in B the operations f_- , f^- , where

$$f_-(x) = \sup_{A \in N(x)} \inf_{y \in A} f(y),$$

$$f^-(x) = \inf_{A \in N(x)} \sup_{y \in A} f(y),$$

$N(x)$ standing for the class of neighborhoods of x . It is well known that f^- and f_- are closure and dual closure operations on B . In accordance with Theorem 1, define

$$LS = E_f(f = f_-), \quad US = E_f(f = f^-).$$

These are the classes of lower and upper semicontinuous functions on X . We also define the complete lattices

$$NLS = E_f(f = (f^-)_-), \quad NUS = E_f(f = (f_-)^-),$$

and call their members normal lower and upper semicontinuous. It is well known that $f \in LS$ ($f \in US$) if and only if $E(f > \lambda)$ ($E(f < \lambda)$) is open for each finite λ . We now let $C = LS \cap US$, the class of continuous, real-valued functions on X . If A is a subset of X , then

$$(k_A)^- = k_{C_A}, \quad (k_A)_- = k_{\mathcal{Q}_A}.$$

Hence k_A is in C , LS , US , NLS , or NUS if and only if A is open and closed, open, closed, regular open, or regular closed, respectively.

In terms of the present definitions of B and C , let us now define the operations f^* , f_* and the classes L , U , NL , and NU as in § 1. Theorem 4 specializes to:

THEOREM 6. *The normal completion of the class of continuous, real-valued functions on an arbitrary topologic space X is isomorphic with the class of NL (or NU) functions on X .*

The lattice operations in the class NL are described in Theorem 3. It is easily seen that the hypotheses of Theorem 5 are satisfied in the present case. It follows that the greatest lower bound of a finite family of L functions is also an L function, and that the same is true of NL .

We now investigate the relations between the classes LS , NLS , L , and NL . We shall not state, but shall feel free to use, the duals of the next theorems.

THEOREM 7. If $f \in B$, then $f_* \leq f_-$. Hence $L \subset LS$.

Proof. If $g \in C$, $g \leq f$, then $g = g_- \leq f_-$. Hence

$$f_* = \sum(f_0) \leq f_-.$$

COROLLARY. The relation

$$(f_*)_- = (f_-)_* = f_*$$

holds for every $f \in B$.

Proof. We have

$$f_* \leq f_- \leq f \rightarrow f_* = (f_*)_* \leq (f_-)_* \leq f_*;$$

and applying the theorem to f_* , we have

$$f_* = (f_*)_* \leq (f_*)_- \leq f_*.$$

THEOREM 8. A necessary and sufficient condition that $f_* = f_-$ for every $f \in B$ is that X be completely regular.

Proof. Suppose $f_* = f_-$ for every $f \in B$. Let x be any point of X . Let A be any neighborhood of x , and set $f = k_A$. Then $f \in LS$, and hence

$$f = f_- = f_*.$$

Therefore there is a continuous function g with $g \leq f$, $g(x) > 0$. This establishes the complete regularity of X .

Conversely, if X is completely regular, suppose that for some $f \in B$ there is an x with

$$f_*(x) < f_-(x).$$

Then there is a neighborhood A of x with

$$\inf_{y \in A} f(y) > \lambda > f_*(x).$$

Let $g \in C$ with $g(x) = 1$, $g(y) = 0$ for y outside A , and $0 \leq g(z) \leq 1$ for all $z \in X$. Define

$$h(y) = \lambda + \log g(y),$$

with the convention that $\log 0 = -\infty$. Then

$$h \leq f, h \in C, \text{ and } h(x) > f_*(x).$$

This contradiction completes the proof.

COROLLARY. *If X is completely regular then $\tilde{C} \cong NLS$.*

We now obtain two characterizations of the class NLS .

LEMMA. *For any $f \in B$ and any finite λ ,*

$$E(f_- \geq \lambda) = \bigcap_n \mathcal{J}E(f \geq \lambda - 1/n).$$

Proof. We have

$$f_-(x) \geq \lambda \leftrightarrow \text{for every } n, x \in \mathcal{J}E(f \geq \lambda - 1/n).$$

THEOREM 9. *If $f \in LS$, then $f \in NLS$ if and only if $E(f \geq \lambda)$ is an intersection of regular open sets for each finite λ [2].*

Proof. If $f \in NLS$, then

$$E(f \geq \lambda) = E((f^-)_- \geq \lambda) = \bigcap_n \mathcal{J}E(f^- \geq \lambda - 1/n),$$

by the Lemma. The result now follows from the fact that $f^- \in US$.

Conversely, if $f \in LS$, and $E(f \geq \lambda)$ is an intersection of regular open sets for each finite λ , we have

$$f = f_- \leq (f^-)_-.$$

Suppose

$$f(x) < \lambda < (f^-)_-(x)$$

for some x . Then for some neighborhood A of x we have

$$\inf_{y \in A} f^-(y) > \lambda.$$

But $E(f < \lambda)$ is a union of regular closed sets. Hence there is an open set B with $x \in \overset{\circ}{C} B$ and $f(y) < \lambda$ for each $y \in \overset{\circ}{C} B$. Thus $f^-(y) \leq \lambda$ for $y \in B$. But $B \cap A \neq \emptyset$, since $x \in \overset{\circ}{C} B$. This contradiction completes the proof.

THEOREM 10. *If $f \in LS$, then $f \in NLS$ if and only if $\mathfrak{I}E(f \geq \lambda)$ is a regular open set for each finite λ .*

Proof. If $f \in NLS$, then, by Theorem 8, $E(f \geq \lambda)$ is an intersection of regular open sets, and hence its interior is regular open. Conversely, suppose $\mathfrak{I}E(f \geq \lambda)$ is regular open for each finite λ . We have

$$E(f > \mu) \subset \mathfrak{I}E(f \geq \mu) \subset E(f \geq \mu)$$

for each finite μ . Hence

$$E(f \geq \lambda) = \bigcap_{\mu < \lambda} E(f \geq \mu) = \bigcap_{\mu < \lambda} E(f > \mu) = \bigcap_{\mu < \lambda} \mathfrak{I}E(f \geq \mu),$$

and the last is an intersection of regular open sets.

Other characterizations of NLS are given in [2, Theorem 3.1], and in Theorem 14 below.

We now define C_1 as the class of finite real-valued continuous functions on X , and C_2 as the class of bounded real-valued continuous functions on X , and investigate the normal completions of these lattices. We shall use ∞ to denote the function which is identically ∞ , and similarly for $-\infty$.

THEOREM 11. *Let M be the lattice consisting of all NLS functions which are bounded above and below by members of C_1 together with the functions ∞ , $-\infty$. If X is completely regular, then $\tilde{C}_1 \cong M$.*

Proof. We shall say that a function is *dominated* above or below if it is bounded above or below by a member of C_1 . Let f^0, f^* be defined as in § 1, where C_1 is used instead of C . We first note that if f is dominated above, then $f^* = f^-$ and both are dominated above. Indeed, if $f \leq h \in C_1$, then

$$\begin{aligned} f^* &= \prod_{g \geq f, g \in C_1} g = \prod_{h \geq g \geq f, g \in C_1} g \\ &= \prod_{h \geq g \geq f, g \in C} g = \prod_{g \geq f, g \in C} g = f^-, \end{aligned}$$

by the dual of Theorem 8. On the other hand, if $f^0 = 0$, then $f^* = \infty$.

Now the normal completion of C_1 is isomorphic with

$$NL_1 = E_f(f = (f^*)_*).$$

If f is a finite-valued member of M , then $f^* = f^-$ is dominated above and below.

Hence

$$(f^*)_* = (f^*)_* = (f^-)_- = f,$$

and $f \in NL_1$. Conversely, if $f \in NL_1$ and $f \neq \infty, f \neq -\infty$, then f must be dominated above. For otherwise $f^* = \infty$, and therefore

$$(f^*)_* = f = \infty.$$

Hence $f^* = f^-$, and f^* is dominated above. Also f^- is dominated below, for otherwise

$$f = (f^-)_* = -\infty.$$

Therefore

$$f = (f^*)_* = (f^-)_* = (f^-)_-,$$

and $f \in M$.

In a similar manner we can prove:

THEOREM 12. *If X is completely regular, then \tilde{C}_2 is isomorphic with the lattice consisting of all bounded NLS functions together with the functions $\infty, -\infty$ [2].*

We now prove a theorem of Stone [5] and Nakano [4].

THEOREM 13. *If X is extremally disconnected, then C is complete. If C is complete and X is completely regular, then X is extremally disconnected.*

Proof. Suppose X is extremally disconnected. Then by the lemma preceding Theorem 9,

$$f \in US \rightarrow f_- \in US \rightarrow f_- \in C.$$

For any $f \in B$, we have $f^* \in U \subset US$. Therefore $(f^*)_- \in C$. Consequently, for any f in NL ,

$$f = (f^*)_* = ((f^*)_-)_* = (f^*)_-,$$

by the corollary of Theorem 9. Thus $NL = C$, and C is complete.

Conversely, let A be any regular open set. Then by Theorem 8,

$$k_A \in NLS = NL = C.$$

Therefore A is closed.

3. Regular open sets and NLS functions. Let $R(X)$, or simply R , denote the lattice of regular open sets of X . According to Theorem 3, R is a complete lattice in which

$$\prod_R(S) = \mathfrak{I} \cap (S), \quad \sum_R(S) = \mathfrak{I} C \cup (S).$$

It is well known that R is in fact a complete Boolean algebra. By Stone's Theorem, R is isomorphic with the open and closed sets of an extremally disconnected, zero-dimensional, compact Hausdorff space, which we shall call the representation space of R .

THEOREM 14. *A necessary and sufficient condition that f be normal lower semicontinuous is the existence of a family $\{x_\lambda\}$, $-\infty < \lambda < \infty$, of regular open sets with the properties*

- i) $x_\lambda \subset x_\mu$ for $\lambda > \mu$,
- ii) $f(p) = \sup E_\lambda(p \in x_\lambda)$.

In this case we have

$$\text{iii) } \mathfrak{I} E(f \geq \lambda) = \prod_{R, \mu < \lambda} x_\mu.$$

If $f \in NLS$, we may choose $x_\lambda = \mathfrak{I} E(f \geq \lambda)$.

Proof. Suppose $f \in NLS$. Let

$$x_\lambda = \mathfrak{I} E(f \geq \lambda).$$

Theorem 10 shows that $x_\lambda \in R$, and the proof of that theorem shows that iii) holds. Since i) is obvious, it remains to prove ii). Suppose $\lambda < f(p)$. Then

$$p \in E(f > \lambda) \subset x_\lambda.$$

Hence

$$\sup E_\lambda(p \in x_\lambda) \geq f(p).$$

On the other hand, if $\lambda > f(p)$, then

$$p \notin E(f \geq \lambda) \supset x_\lambda.$$

Therefore

$$\sup E_\lambda(p \in x_\lambda) \leq f(p).$$

Now suppose $\{x_\lambda\}$ is a family of regular open sets satisfying i), and let f be defined by ii). If $f(p) > \lambda$, then $p \in x_\mu$ for some $\mu > \lambda$. But

$$f(q) \geq \mu > \lambda \quad \text{for } q \in x_\mu.$$

Therefore $E(f > \lambda)$ is open and $f \in LS$. Also

$$E(f \geq \lambda) = \bigcap_{\mu < \lambda} x_\mu,$$

which implies iii). Theorem 10 now shows that $f \in NLS$.

THEOREM 15. *If X and Y are any topologic spaces, then $NLS(X) \cong NLS(Y)$ if and only if $R(X) \cong R(Y)$. In particular ([2] when X is completely regular), if Z is the representation space of $R(X)$, then $NLS(X) \cong C(Z)$.*

Proof. Suppose $R(X) \cong R(Y)$. Let ϕ be a function on $R(X)$ to $R(Y)$ which establishes this isomorphism. If $f \in NLS(X)$, let

$$x_\lambda = \mathcal{J}E(f \geq \lambda),$$

Define Tf to be the function on Y defined by

$$Tf(p) = \sup E_\lambda(p \in \phi(x_\lambda)).$$

By Theorem 14, $Tf \in NLS(Y)$. We now show that T is an isomorphic mapping of $NLS(X)$ onto $NLS(Y)$. If $f, g \in NLS(X)$, let

$$x_\lambda = \mathcal{J}E(f \geq \lambda), \quad y_\lambda = \mathcal{J}E(g \geq \lambda).$$

Clearly,

$$f \geq g \rightarrow x_\lambda \supset y_\lambda \rightarrow \phi(x_\lambda) \supset \phi(y_\lambda)$$

for every λ , and therefore $Tf \geq Tg$. Using Theorem 14, iii) twice, we have

$$x_\lambda = \prod_{R(X), \mu < \lambda} x_\mu,$$

which implies

$$\phi(x_\lambda) = \prod_{R(Y), \mu < \lambda} \phi(x_\mu) = \mathfrak{J}E(Tf \geq \lambda).$$

From this it follows that

$$Tf \geq Tg \longrightarrow \mathfrak{J}E(Tf \geq \lambda) \supset \mathfrak{J}E(Tg \geq \lambda) \longrightarrow \phi(x_\lambda) \supset \phi(y_\lambda) \longrightarrow x_\lambda \supset y_\lambda$$

for every λ . This implies $f \geq g$ by Theorem 14, ii). Finally, if $F \in NLS(Y)$, choose $x_\lambda \in R(X)$ so that

$$\phi(x_\lambda) = \mathfrak{J}E(F \geq \lambda),$$

and define f as in Theorem 14, ii). Then $f \in NLS(X)$ and

$$\mathfrak{J}E(f \geq \lambda) = \prod_{R(X), \mu < \lambda} x_\mu,$$

and this last is x_λ because

$$\phi(x_\lambda) = \prod_{R(Y), \mu < \lambda} \phi(x_\mu).$$

This shows that $Tf = F$.

If Z is the representation space of $R(X)$, then $NLS(Z) = C(Z)$ by Theorem 13 and the corollary to Theorem 8.

Now suppose $NLS(X) \cong NLS(Y)$. Let Z_1 and Z_2 be the representation spaces of $R(X)$ and $R(Y)$, respectively. Then

$$C(Z_1) \cong C(Z_2).$$

A theorem of Kaplansky [3] shows that Z_1 is homeomorphic with Z_2 . This implies that

$$R(X) \cong R(Y),$$

and the proof is complete.

In Theorem 14, it is easily seen that f is bounded if and only if $x_\lambda = 0$ for all λ outside some finite interval. It follows from the proof of Theorem 15 that the bounded NLS functions on X are isomorphic with those on Y if and only if

$$NLS(X) \cong NLS(Y).$$

The same is not true of the finite valued NLS functions. If

$$R(X) \cong R(Y),$$

all we can say is that the classes of *NLS* functions f on X and Y , respectively, which are such that $E(|f| = \infty)$ is nowhere dense, are isomorphic. This follows from the fact that $E(f = \infty)$ is nowhere dense if and only if

$$\prod_R x_\lambda = 0,$$

and $E(f = -\infty)$ is nowhere dense if and only if

$$\sum_R x_\lambda = X \text{ or } Y.$$

The following theorem gives a corresponding result for the lattices of continuous functions. We recall that $C_2(X)$ denotes the lattice of bounded real-valued continuous functions on X .

THEOREM 16. *If X and Y are any topologic spaces, then*

$$C_2(X) \cong C_2(Y)$$

if and only if

$$C(X) \cong C(Y).$$

Proof. There exist completely regular Hausdorff spaces X', Y' such that

$$C_2(X) \cong C_2(X'), C_2(Y) \cong C_2(Y'), C(X) \cong C(X'), \text{ and } C(Y) \cong C(Y')$$

(one modifies Dilworth's proof [2, § 2] by defining $x \sim y$ if $f(x) = f(y)$ for all $f \in C(X)$). Now let $\beta X', \beta Y'$ be the Stone-Ćech compactifications of X', Y' . Then

$$C_2(\beta X') \cong C_2(X').$$

Now the mapping

$$\phi(f) = f/(1 + |f|)$$

carries $C(X')$ into $C_2(X')$. We use this to extend members of $C(X')$. Thus

$$C(\beta X') \cong C(X').$$

Combining these facts, we have

$$\begin{aligned} C(X) \cong C(Y) &\leftrightarrow C(X') \cong C(Y') \leftrightarrow C(\beta X') \cong C(\beta Y') \\ &\leftrightarrow \beta X' \text{ is homeomorphic with } \beta Y' \end{aligned}$$

(by Kaplansky's Theorem [3]), \leftrightarrow

$$C_2(\beta X') \cong C_2(\beta Y') \text{ (again by [3])} \leftrightarrow C_2(X') \cong C_2(Y') \leftrightarrow C_2(X) \cong C_2(Y).$$

It would be of interest to give a more elementary proof of this theorem.

REFERENCES

1. G. Birkhoff, *Lattice theory*, Amer. Math. Soc. Colloquium Publications, vol. XXV, Revised Edition, New York, 1948.
2. R. P. Dilworth, *The normal completion of the lattice of continuous functions*, Trans. Amer. Math. Soc. 68 (1950), 427-438.
3. I. Kaplansky, *Lattices of continuous functions*, Bull. Amer. Math. Soc. 53 (1947), 617-622.
4. H. Nakano, *Über das System aller stetigen Funktionen auf einem topologischen Raum*, Proc. Imp. Acad. Tokyo 17 (1941), 308-310.
5. M. H. Stone, *Boundedness properties in function lattices*, Canadian J. Math. 1 (1949), 176-186.

UNIVERSITY OF CALIFORNIA, LOS ANGELES, CALIFORNIA

PACIFIC JOURNAL OF MATHEMATICS

EDITORS

R. M. ROBINSON
University of California
Berkeley 4, California

E. HEWITT
University of Washington
Seattle 5, Washington

R. P. DILWORTH
California Institute of Technology
Pasadena 4, California

E. F. BECKENBACH
University of California
Los Angeles 24, California

ASSOCIATE EDITORS

H. BUSEMANN

HERBERT FEDERER

MARSHALL HALL

P. R. HALMOS

HEINZ HOPF

R. D. JAMES

BØRGE JESSEN

PAUL LÉVY

GEORGE PÓLYA

J. J. STOKER

E. G. STRAUS

KÔSAKU YOSIDA

SPONSORS

UNIVERSITY OF BRITISH COLUMBIA

CALIFORNIA INSTITUTE OF TECHNOLOGY

UNIVERSITY OF CALIFORNIA, BERKELEY

UNIVERSITY OF CALIFORNIA, DAVIS

UNIVERSITY OF CALIFORNIA, LOS ANGELES

UNIVERSITY OF CALIFORNIA, SANTA BARBARA

UNIVERSITY OF NEVADA

OREGON STATE COLLEGE

UNIVERSITY OF OREGON

UNIVERSITY OF SOUTHERN CALIFORNIA

STANFORD RESEARCH INSTITUTE

STANFORD UNIVERSITY

WASHINGTON STATE COLLEGE

UNIVERSITY OF WASHINGTON

* * *

AMERICAN MATHEMATICAL SOCIETY

NATIONAL BUREAU OF STANDARDS,

INSTITUTE FOR NUMERICAL ANALYSIS

Vari-Type Composition by

Elaine Barth

Delores Wierman

With the cooperation of

E. F. Beckenbach

E. G. Straus

Printed in the United States of America by
Edwards Brothers, Inc., Ann Arbor, Michigan

UNIVERSITY OF CALIFORNIA PRESS * BERKELEY AND LOS ANGELES

COPYRIGHT 1953 BY PACIFIC JOURNAL OF MATHEMATICS

Herbert Busemann, <i>Volume in terms of concurrent cross-sections</i>	1
L. Carlitz, <i>Some special equations in a finite field</i>	13
Homer V. Craig and Billie Braden Townsend, <i>On certain metric extensors</i>	25
Philip J. Davis and Henry Pollak, <i>Linear functionals and analytic continuation problems</i>	47
Jacob C. E. Dekker, <i>The constructivity of maximal dual ideals in certain Boolean algebras</i>	73
Harley M. Flanders, <i>The norm function of an algebraic field extension</i>	103
Marshall Hall, <i>Subgroups of free products</i>	115
Israel (Yitzchak) Nathan Herstein, <i>Finite multiplicative subgroups in division rings</i>	121
Joseph Lawson Hodges, Jr. and Murray Rosenblatt, <i>Recurrence-time moments in random walks</i>	127
Alfred Horn, <i>The normal completion of a subset of a complete lattice and lattices of continuous functions</i>	137
Fulton Koehler, <i>Estimates for the errors in the Rayleigh-Ritz method</i>	153
M. H. Martin, <i>The Monge-Ampère partial differential equation $rt - s^2 + \lambda^2 = 0$</i>	165
John E. Maxfield, <i>Normal k-tuples</i>	189
Jack E. McLaughlin, <i>Structured theorems for relatively complemented lattices</i>	197
William H. Mills, <i>A system of quadratic Diophantine equations</i>	209
T. S. Motzkin, Ernst Gabor Straus and F. A. Valentine, <i>The number of farthest points</i>	221
G. Power, <i>Forces on the boundary of a dielectric</i>	233
Ralph Gordon Selfridge, <i>Approximations with least maximum error</i>	247