

Pacific Journal of Mathematics

AN ISOPERIMETRIC MINIMAX

WILLIAM GUSTIN

AN ISOPERIMETRIC MINIMAX

WILLIAM GUSTIN

Introduction. In the preceding paper J. W. Green considers for a given convex body K in the euclidean plane the minimum of the isoperimetric ratio r (ratio of squared perimeter l^2 to area a) taken over all affine transforms k of K . He then investigates the maximum value taken over all K of this minimum ratio, shows by variational methods that such a maximum is attained by some polygon of five or fewer sides, and conjectures that it is, in fact, attained by a triangle with $12\sqrt{3}$, the isoperimetric ratio of an equilateral triangle, as the minimax ratio. I shall prove this conjecture directly by refining an estimation used by Green, the precise statement of results being as follows:

I. Let K be an nontriangular plane convex body; there then exists an affine transform k of K with $r(k) < 12\sqrt{3}$.

II. Let T be a nonequilateral triangle; then $r(T) > 12\sqrt{3}$.

Before taking up the proof of these results we dispose of a lemma.

III. Let k be a possibly degenerate convex body with $s \subset k \subset t$, wherein t is an equilateral triangle, and s a side of t ; there then exists a number x with $0 \leq x \leq 1$ such that

$$l(k) \leq (2/3 + x/3) l(t)$$

$$a(k) \geq x a(t),$$

simultaneous equality occurring if and only if either $x = 0$, $k = s$ or $x = 1$, $k = t$.

Proof of III. Let p be that supporting strip of k parallel to the line-segment s ; and let x be the ratio of the width of p to the width or altitude of t . Thus $0 \leq x \leq 1$, with $x = 0$ or $x = 1$ according as $k = s$ or $k = t$. Choose a point at which k touches the side of p opposite s , and define k_* to be the triangle with this point as apex and s as base. Define k^* to be the trapezoid formed by intersection of p and t . Clearly $s \subset k_* \subset k \subset k^* \subset t$; and $k_* = k = k^*$ if and only if $k = s$ or $k = t$.

Received October 16, 1952.

Pacific J. Math. 3 (1953), 403-405

Since $k \supset k_*$, it follows that $a(k) \geq a(k_*)$, with equality if and only if $k = k_*$. And since $k \subset k^*$, it follows that $l(k) \leq l(k^*)$ with equality if and only if $k = k^*$. These inequalities become, upon the easy computation of $a(k_*)$ and $l(k^*)$, the asserted inequalities of III.

Proof of I. Let K be the given nontriangular convex body. Since the area functional is continuous, it easily follows from a compactness argument that a triangle T of maximal area can be inscribed in K . Let the three sides of T be labelled S_i ($i = 1, 2, 3$), and let V_i be that vertex of T opposite S_i . Because the area of T is maximal, the line L_i through V_i and parallel to S_i is a line of support of K . The triangle formed by the three lines L_i then circumscribes K and also T ; it is composed of four nonoverlapping congruent triangles T and T_i , where T_i is labelled so as to have S_i as a side. That part K_i of K in T_i is a possibly degenerate convex body with $S_i \subset K_i \subset T_i$. Now any triangle can be affinely transformed into any other triangle. In particular, T can be affinely transformed into an equilateral triangle t , with T_i going into t_i , S_i into s_i , K_i into k_i , and K into k . Therefore $s_i \subset k_i \subset t_i$, and t_i is congruent to t . According to III, ratios x_i exist giving inequalities on $l(k_i)$ and $a(k_i)$. Furthermore, since K and hence k is nontriangular, not all $x_i = 0$ and not all $x_i = 1$. Therefore $0 < x < 1$, where $x = \sum x_i/3$. Evidently k is composed of the four nonoverlapping sets t and k_i in such a way that

$$l(k) = \sum l(k_i) - l(t) \leq (1+x) l(t),$$

$$a(k) = \sum a(k_i) + a(t) \geq (1+3x) a(t),$$

whereupon

$$r(k) \leq \frac{(1+x)^2}{1+3x} r(t) = \left[1 - \frac{x(1-x)}{1+3x} \right] 12\sqrt{3} < 12\sqrt{3},$$

as was to be shown.

Proof of II. Through II is merely a matter of trigonometry, and very likely can be verified by exhibiting a neat but perhaps unobtrusive trigonometric identity, I shall here prove it by the sort of methods used above.

Let T be a nonequilateral triangle. Define S_i , V_i , L_i as above. Since T is nonequilateral, some two of its sides, say S_1 and S_2 , are unequal. Let v_3 be that point on the line L_3 , regarded as a linear mirror, at which $v_1 = V_1$ is reflected when viewed from $v_2 = V_2$; and let t be the so symmetrized isosceles triangle

with vertices v_i and sides s_i . Then the path $s_1 s_2$ is shorter than $S_1 S_2$, so $l(t) < l(T)$; and, since both triangles have the same base and altitude, $a(t) = a(T)$. Therefore $r(t) < r(T)$. Consequently if the minimum isoperimetric ratio among triangles is attained, it is attained by an equilateral triangle only; whereupon it would follow that $r(T) > 12\sqrt{3}$, as was to be shown. Now all possible triangle isoperimetric ratios are realized by triangles of fixed perimeter containing a fixed point. By a compactness argument, some such triangle achieves a maximum area and hence a minimum isoperimetric ratio. This completes the proof.

INDIANA UNIVERSITY

PACIFIC JOURNAL OF MATHEMATICS

EDITORS

R. M. ROBINSON
University of California
Berkeley 4, California

R. F. DILWORTH
California Institute of Technology
Pasadena 4, California

E. HEWITT
University of Washington
Seattle 5, Washington

E. F. BECKENBACH
University of California
Los Angeles 24, California

ASSOCIATE EDITORS

H. BUSEMANN P. R. HALMOS
HERBERT FEDERER HEINZ HOPF
MARSHALL HALL R. D. JAMES

BØRGE JESSEN J. J. STOKER
PAUL LÉVY E. G. STRAUS
GEORGE PÓLYA KÔSAKU YOSIDA

SPONSORS

UNIVERSITY OF BRITISH COLUMBIA
CALIFORNIA INSTITUTE OF TECHNOLOGY
UNIVERSITY OF CALIFORNIA, BERKELEY
UNIVERSITY OF CALIFORNIA, DAVIS
UNIVERSITY OF CALIFORNIA, LOS ANGELES
UNIVERSITY OF CALIFORNIA, SANTA BARBARA
UNIVERSITY OF NEVADA
OREGON STATE COLLEGE
UNIVERSITY OF OREGON

UNIVERSITY OF SOUTHERN CALIFORNIA
STANFORD RESEARCH INSTITUTE
STANFORD UNIVERSITY
WASHINGTON STATE COLLEGE
UNIVERSITY OF WASHINGTON
* * *
AMERICAN MATHEMATICAL SOCIETY
NATIONAL BUREAU OF STANDARDS,
INSTITUTE FOR NUMERICAL ANALYSIS

Mathematical papers intended for publication in the *Pacific Journal of Mathematics* should be typewritten (double spaced), and the author should keep a complete copy. Manuscripts may be sent to any of the editors except Robinson, whose term expires with the completion of the present volume; they might also be sent to M. M. Schiffer, Stanford University, Stanford, California, who is succeeding Robinson. All other communications to the editors should be addressed to the managing editor, E. F. Beckenbach, at the address given above.

Authors are entitled to receive 100 free reprints of their published papers and may obtain additional copies at cost.

The *Pacific Journal of Mathematics* is published quarterly, in March, June, September, and December. The price per volume (4 numbers) is \$8.00; single issues, \$2.50. Special price to individual faculty members of supporting institutions and to individual members of the American Mathematical Society: \$4.00 per volume; single issues, \$1.25.

Subscriptions, orders for back numbers, and changes of address should be sent to the publishers, University of California Press, Berkeley 4, California.

Printed at Ann Arbor, Michigan. Entered as second class matter at the Post Office, Berkeley, California.

UNIVERSITY OF CALIFORNIA PRESS · BERKELEY AND LOS ANGELES

COPYRIGHT 1953 BY PACIFIC JOURNAL OF MATHEMATICS

William George Bade, <i>An operational calculus for operators with spectrum in a strip</i>	257
E. F. Beckenbach and Lloyd Kenneth Jackson, <i>Subfunctions of several variables</i>	291
David Blackwell, <i>Extension of a renewal theorem</i>	315
L. Carlitz, <i>Some theorems on the Schur derivative</i>	321
Paul Arnold Clement, <i>Generalized convexity and surfaces of negative curvature</i>	333
Merrill M. Flood, <i>On the Hitchcock distribution problem</i>	369
Watson Bryan Fulks, <i>On the unique determination of solutions of the heat equation</i>	387
John W. Green, <i>Length and area of a convex curve under affine transformation</i>	393
William Gustin, <i>An isoperimetric minimax</i>	403
Arthur Eugene Livingston, <i>Some Hausdorff means which exhibit the Gibbs' phenomenon</i>	407
Charles Loewner, <i>On generation of solutions of the biharmonic equation in the plane by conformal mappings</i>	417
Gábor Szegő, <i>Remark on the preceding paper of Charles Loewner</i>	437
Immanuel Marx and G. Piranian, <i>Lipschitz functions of continuous functions</i>	447
Ting-Kwan Pan, <i>The spherical curvature of a hypersurface in Euclidean space</i>	461
Ruth Lind Potter, <i>On self-adjoint differential equations of second order</i>	467
E. H. Rothe, <i>A note on the Banach spaces of Calkin and Morrey</i>	493
Eugene Schenkman, <i>A generalization of the central elements of a group</i>	501
A. Seidenberg, <i>A note on the dimension theory of rings</i>	505