

# Pacific Journal of Mathematics

**IDENTIFICATIONS IN SINGULAR HOMOLOGY THEORY**

EDWARD RICHARD FADELL

# IDENTIFICATIONS IN SINGULAR HOMOLOGY THEORY

EDWARD R. FADELL

## INTRODUCTION

0.1. Given a Mayer complex  $M$ , a subcomplex  $M'$  is termed an *unessential identifier* for  $M$  if the natural projections from  $M$  onto the factor complex  $M/M'$  induce isomorphisms-onto on the homology level (see [1, §1.2]). The present paper is a continuation and improvement of certain results obtained by Rado' and Reichelderfer (see [1] and [3]) concerning unessential identifiers for the singular complex  $R$  of Rado' (see [1, §0.1]). We shall make use of the results, terminology, and notation in [1] and [3] with one exception. Because of a conflict in notation in [1] and [3], we shall use the notation  $\eta_p$  for the homomorphisms

$$\eta_p : C_p^S \longrightarrow C_p^R,$$

defined as the trivial homomorphism for  $p < 0$ , and for  $p \geq 0$  as follows:

$$\eta_p(d_0, \dots, d_p, T)^S = (d_0, \dots, d_p, T)^R$$

(see [1, §0.3]).

0.2. The principal results of the present paper may be described as follows. Let  $N(\sigma_p \beta_p^R)$  denote the nucleus of the product homomorphism

$$\sigma_p \beta_p^R : C_p^R \longrightarrow C_p^S.$$

**THEOREM.** *The system  $\{N(\sigma_p \beta_p^R)\}$  is an unessential identifier for  $R$ .*

Furthermore, for each  $p$  we have

$$N(\sigma_p \beta_p^R) \supset \hat{\Delta}_p^R \supset \hat{\Gamma}_p^R,$$

where  $\{\hat{\Delta}_p^R\}$  and  $\{\hat{\Gamma}_p^R\}$  are the largest unessential identifiers for  $R$  obtained by Reichelderfer [3, §3.6] and Radó [1, §4.7], respectively. Thus  $\{N(\sigma_p \beta_p^R)\}$  is the largest unessential identifier presently known for  $R$  and imposes all the classical identifications in  $R$ .

Let  $N(\beta_p^S)$  denote the nucleus of the barycentric homomorphism

$$\beta_p^S : C_p^S \rightarrow C_p^S.$$

**THEOREM.** *The system  $\{N(\beta_p^S)\}$  is an unessential identifier for  $S$ .*

It is interesting to note that the foregoing theorem gives for the Eilenberg complex  $S$  the result corresponding to that of Reichelderfer for the Radó complex  $R$  (see [3, §3.2]).

## I. PRELIMINARIES

1.1. Let  $v_0, \dots, v_p$  denote  $p+1$  points in Hilbert space  $E_\infty$ . The barycenter  $b = b(v_0, \dots, v_p)$  of these points is given by

$$b = (v_0 + \dots + v_p)/(p+1).$$

The following lemmas are easily verified.

1.2. **LEMMA.** *Let  $v_j$  ( $j = 0, \dots, p$ ) denote  $p+1$  points in  $E_\infty$ , and*

$$x = \sum_{j=0}^p \mu_j b(v_0, \dots, v_j),$$

where  $\mu_j$  is real for  $j = 0, \dots, p$ . Then

$$x = \sum_{j=0}^p \sum_{l=j}^p \frac{\mu_l}{l+1} v_j, \quad \text{with} \quad \sum_{j=0}^p \sum_{l=j}^p \frac{\mu_l}{l+1} = \sum_{j=0}^p \mu_j.$$

1.3. **LEMMA.** *Let  $v_j$  ( $j = 0, \dots, p$ ) denote  $p+1$  points in  $E_\infty$ , and*

$$x = \sum_{j=0}^p \mu_j v_j,$$

with  $\mu_j$  ( $j = 0, \dots, p$ ) real and satisfying

$$\mu_0 \geq \mu_1 \geq \cdots \geq \mu_p \geq 0.$$

Then

$$x = \sum_{j=0}^p \lambda_j b(v_0 \cdots v_j),$$

with

$$\lambda_j = (j+1)(\mu_j - \mu_{j+1}) \text{ for } j = 0, \dots, p-1 \text{ (provided } p-1 \geq 0),$$

$$\lambda_p = (p+1)\mu_p,$$

and

$$\sum_{j=0}^p \lambda_j = \sum_{j=0}^p \mu_j.$$

1.4. As in [1], let  $d_0, d_1, d_2, \dots$  denote the sequence of points  $(1, 0, 0, 0, \dots)$ ,  $(0, 1, 0, 0, \dots)$ ,  $(0, 0, 1, 0, \dots)$ ,  $\dots$  in  $E_\infty$ . For integers  $p, q$  such that  $p \geq 0, 0 \leq q \leq p+1$ , the homomorphism

$$q_{*p} : C_p \longrightarrow C_{p+1}$$

in the formal complex  $K$  of  $E_\infty$  is defined by the relation

$$q_{*p}(v_0, \dots, v_p) = \begin{cases} (d_{p+1}, v_0, \dots, v_p) & \text{for } q = 0, \\ (-1)^q (v_0, \dots, v_{q-1}, d_{p+1}, v_q, \dots, v_p) & \text{for } 1 \leq q \leq p, \\ (-1)^{p+1} (v_0, \dots, v_p, d_{p+1}) & \text{for } q = p+1. \end{cases}$$

1.5. For  $p \geq 0$ , let  $\tau_p$  denote an element of  $T_{p0}$  (see [3, §1.9]), and let  $(i_0, \dots, i_p)$  denote the permutation of  $0, \dots, p$  which gives rise to  $\tau_p$ . Then we let  $\text{sgn } \tau_p$  denote the sign of the permutation  $(i_0, \dots, i_p)$ : i.e.,  $\text{sgn } \tau_p$  is  $+1$  or  $-1$  according as an even or odd number of transpositions is required to obtain  $(i_0, \dots, i_p)$ .

The following lemmas are then obvious.

1.6. LEMMA. For  $p \geq 0$  and  $\tau_{p+1} \in T_{p+10}$ , there exists a unique  $\pi_p \in T_{p0}$ ,

and a unique  $q$ ,  $0 \leq q \leq p + 1$ , such that

$$\tau_{p+1}(d_0, \dots, d_{p+1}) = q_{*p} \pi_p(p+1)_{p+1}(d_0, \dots, d_{p+1}).$$

1.7. LEMMA. For  $p \geq 0$ , let  $E_{p+1}$  denote the set of ordered pairs  $(q, \pi_p)$ ,  $0 \leq q \leq p + 1$ ,  $\pi_p \in T_{p0}$ . There exists a biunique correspondence

$$\xi : T_{p+10} \longrightarrow E_{p+1}$$

with

$$\xi \tau_{p+1} = (q, \pi_p),$$

such that

$$\tau_{p+1}(d_0, \dots, d_{p+1}) = q_{*p} \pi_p(p+1)_{p+1}(d_0, \dots, d_{p+1})$$

and

$$\text{sgn } \tau_{p+1} = (-1)^{p+q+1} \text{sgn } \pi_p.$$

1.8. Let

$$h_p : C_p \longrightarrow C_q$$

denote a homomorphism in  $K$  such that

$$h_p(d_0 \dots d_p) = \pm (w_0, \dots, w_q).$$

Then  $[h_p]$  will denote the usual affine mapping from the convex hull  $|d_0, \dots, d_q|$  of the points  $d_0, \dots, d_q$  onto the convex hull  $|w_0, \dots, w_q|$  of the points  $w_0, \dots, w_q$  such that  $[h_p](d_i) = w_i$  for  $i = 0, \dots, q$ .

1.9. Let  $\beta_p^R$  denote the barycentric homomorphism in  $R$ , and  $\rho_{*p}^R$  the barycentric homotopy operator in  $R$  of Reichelderfer (see [3, § 2.1]). The barycentric homomorphism

$$\beta_p^S : C_p^S \longrightarrow C_p^S$$

in  $S$  may be given by

$$\beta_p^S = \sigma_p \beta_p^R \eta_p \quad (\text{see [2, § 3.7]}).$$

The corresponding homotopy operator

$$\rho_{*p}^S : C_p^S \longrightarrow C_{p+1}^S$$

is given by

$$\rho_{*p}^S = \sigma_{p+1} \rho_{*p}^R \eta_p,$$

1.10. Employing the structure theorems for  $\beta_p^R$ ,  $\rho_{*p}^R$  (see [3, § 2.2]) we obtain the following:

LEMMA. For  $p \geq 0$ ,

$$\beta_p^S(d_0, \dots, d_p, T)^S = \sum_{\tau_p \in T_{p0}} \operatorname{sgn} \tau_p(d_0, \dots, d_p, T[0_{p+1} b_{p0} \tau_p])^S,$$

$$\rho_{*p}^S(d_0, \dots, d_p, T)^S = \sum_{k=0}^p \sum_{\tau_p \in T_{pk}} (-1)^k \operatorname{sgn} \tau_p(d_0, \dots, d_{p+1}, T[b_{pk} \tau_p])^S.$$

*Proof.* We have

$$\begin{aligned} \beta_p^S(d_0, \dots, d_p, T)^S &= \sigma_p \beta_p^R(d_0, \dots, d_p, T)^R \\ &= \sigma_p \sum_{\tau_p \in T_{p0}} (0_{p+1} b_{p0} \tau_p(d_0, \dots, d_p), T)^R \\ &= \sum_{\tau_p \in T_{p0}} \operatorname{sgn} \tau_p(d_0, \dots, d_p, T[0_{p+1} b_{p0} \tau_p])^S, \end{aligned}$$

and

$$\begin{aligned} \rho_{*p}^S(d_0, \dots, d_p, T)^S &= \sigma_{p+1} \rho_{*p}^R(d_0, \dots, d_p, T)^R \\ &= \sigma_{p+1} \sum_{k=0}^p \sum_{\tau_p \in T_{pk}} (b_{pk} \tau_p(d_0, \dots, d_p), T)^R \\ &= \sum_{k=0}^p \sum_{\tau_p \in T_{pk}} (-1)^k \operatorname{sgn} \tau_p(d_0, \dots, d_{p+1}, T[b_{pk} \tau_p])^S. \end{aligned}$$

1.11. In [2], Rado' makes use of the following identities which we state in terms of  $\rho_{*p}^R$  :

$$(1) \quad \sigma_{p+1} \rho_{*p}^R \eta_p \sigma_p = \sigma_{p+1} \rho_{*p}^R, \quad -\infty < p < \infty,$$

$$(2) \quad \sigma_p \beta_p^R \eta_p \sigma_p = \sigma_p \beta_p^R, \quad -\infty < p < \infty.$$

The proof of (1) may be modeled after the proof for the corresponding identity stated in terms of the classical homotopy operator  $\rho_p^R$  (see [2, § 3.5]). From identities (1) and (2), we have

$$(3) \quad \beta_p^S \sigma_p = \sigma_p \beta_p^R,$$

$$(4) \quad \rho_{*p}^S \sigma_p = \sigma_{p+1} \rho_{*p}^R,$$

$$(5) \quad \beta_{p+1}^S \rho_{*p}^S \sigma_p = \sigma_{p+1} \beta_{p+1}^R \rho_{*p}^R$$

for all integers  $p$ .

1.12. Let  $P_1$  and  $P_2$  denote the following propositions:

$P_1$ . Let  $c_p^S$  denote a  $p$ -chain of  $S$  such that

$$\beta_p^S c_p^S = 0.$$

Then

$$\beta_{p+1}^S \rho_{*p}^S c_p^S = 0.$$

$P_2$ . Let  $c_p^R$  denote a  $p$ -chain of  $R$  such that

$$\sigma_p \beta_p^R c_p^R = 0.$$

Then

$$\sigma_{p+1} \beta_{p+1}^R \rho_{*p}^R c_p^R = 0.$$

**THEOREM.**  $P_1 \equiv P_2$ ; i.e.,  $P_1$  is true if and only if  $P_2$  is true.

*Proof.* Assume  $P_1$ , and let  $c_p^R$  denote a  $p$ -chain of  $R$  such that

$$\sigma_p \beta_p^R c_p^R = 0.$$

Then via identity (3) we have

$$\beta_p^S \sigma_p c_p^R = 0.$$

Therefore

$$\beta_{*p+1}^S \rho_{*p}^S \sigma_p c_p^R = 0.$$

But via identity (5), we have

$$\sigma_{p+1} \beta_{p+1}^R \rho_{*p}^R c_p^R = 0,$$

and  $P_2$  follows.

Now assume  $P_2$ , and let  $c_p^S$  denote a  $p$ -chain of  $S$  such that

$$\beta_p^S c_p^S = 0.$$

Then since

$$\beta_p^S = \sigma_p \beta_p^R \eta_p,$$

we have

$$\sigma_p \beta_p^R \eta_p c_p^S = 0.$$

Therefore, via  $P_2$ , we have

$$\sigma_{p+1} \beta_{p+1}^R \rho_{*p}^R \eta_p c_p^S = 0.$$

But via (5) and the fact that  $\sigma_p \eta_p = 1$ , we have

$$\sigma_{p+1} \beta_{p+1}^R \rho_{*p}^R \eta_p c_p^S = \beta_{p+1}^S \rho_{*p}^S \sigma_p \eta_p c_p^S = \beta_{p+1}^S \rho_{*p}^S c_p^S = 0,$$

and  $P_1$  follows.

## II. THE PROOF OF $P_1$

2.1. We shall use throughout this section the notation  $T$  for the  $p$ -cell



$(d_0, \dots, d_p, T)^S$  when there is little chance for ambiguity. Under this convention a chain  $c_p^S$  having the representation

$$c_p^S = \sum_{j=1}^n \lambda_j (d_0, \dots, d_p, T_j)^S$$

may be written  $\sum_{j=1}^n \lambda_j T_j$ . Thus  $T$  represents both a transformation from the convex hull  $|d_0, \dots, d_p|$  into the topological space  $X$  and the  $p$ -cell  $(d_0, \dots, d_p, T)^S$ .

2.2. For  $p < 0$ , the proposition  $P_1$  is trivial. For  $p = 0$ ,  $P_1$  is also trivial. For since  $\beta_0^R = 1$  and  $\sigma_0 \eta_0 = 1$ , we have

$$\beta_0^S c_0^S = 0$$

implying

$$\sigma_0 \beta_0^R \eta_0 c_0^S = \sigma_0 \eta_0 c_0^S = c_0^S = 0,$$

whence clearly

$$\beta_1^S \rho_{*0}^S c_0^S = 0.$$

Now, take a fixed  $p \geq 1$ . Let

$$c_p^S = \sum_{j=1}^n \lambda_j T_j \tag{\lambda_j \neq 0}$$

denote a  $p$ -chain of  $S$  such that

$$\beta_p^S c_p^S = 0.$$

Via § 1.10,

$$(1) \quad \beta_p^S c_p^S = \sum_{j=1}^n \sum_{\tau_p \in T_{p0}} \lambda_j \operatorname{sgn} \tau_p T_j [0_{p+1} b_{p0} \tau_p].$$

Let  $E$  denote the set of ordered pairs  $(j, \tau_p)$ ,  $1 \leq j \leq n$ ,  $\tau_p \in T_{p0}$ . Then

$$(2) \quad \beta_p^S c_p^S = \sum_{(j, \tau_p) \in E} \lambda_j \operatorname{sgn} \tau_p T_j [0_{p+1} b_{p0} \tau_p].$$

We now define a binary relation “ $\equiv$ ” on  $E$  as follows:

$$(j, \tau_p) \equiv (j', \tau_p')$$

if and only if  $T_j [0_{p+1} b_{p0} \tau_p]$ ,  $T_{j'} [0_{p+1} b_{p0} \tau_p']$  are identical  $p$ -cells. Then “ $\equiv$ ” as defined is obviously a true equivalence relation and induces a partitioning of  $E$  into nonempty, mutually disjoint sets  $E_s$  ( $s = 1, \dots, t$ ) with

$$E = \bigcup_{s=1}^t E_s.$$

Therefore, via (2), we have

$$(3) \quad \beta_p^S c_p^S = \sum_{s=1}^t \sum_{(j, \tau_p) \in E_s} \lambda_j \operatorname{sgn} \tau_p T_j [0_{p+1} b_{p0} \tau_p].$$

Take  $1 \leq s < s' \leq t$ . Then for  $(j, T_p) \in E_s$ ,  $(j', T_p') \in E_{s'}$ , the  $p$ -cells  $T_j [0_{p+1} b_{p0} \tau_p]$ ,  $T_{j'} [0_{p+1} b_{p0} \tau_p']$  are distinct. Therefore, since

$$\beta_p^S c_p^S = 0,$$

we must have for each  $s$ ,  $1 \leq s \leq t$ ,

$$(4) \quad \sum_{(j, \tau_p) \in E_s} \lambda_j \operatorname{sgn} \tau_p T_j [0_{p+1} b_{p0} \tau_p] = 0,$$

and hence

$$(5) \quad \sum_{(j, \tau_p) \in E_s} \lambda_j \operatorname{sgn} \tau_p = 0,$$

since all  $p$ -cells occurring in (4) are identical.

2.3. Again via §1.10,

$$(6) \quad \beta_{p+1}^S \rho_{*p}^S c_p^S = \sum_{j=1}^n \sum_{k=0}^p \sum_{\tau_p \in T_{pk}} \sum_{\tau_{p+1} \in T_{p+10}} (-1)^k \operatorname{sgn} \tau_p \operatorname{sgn} \tau_{p+1} \lambda_j T_j [b_{pk} \tau_p] [0_{p+2} b_{p+10} \tau_{p+1}].$$

Applying the lemma of § 1.7, we obtain

$$(7) \quad \beta_{p+1}^S \rho_{*p}^S c_p^S = \sum_{k=0}^p \sum_{q=0}^{p+1} (-1)^{p+q+k+1} \left\{ \sum_{j=1}^n \sum_{\tau_p \in T_{pk}} \sum_{\pi_p \in T_{p0}} \lambda_j \operatorname{sgn} \tau_p \operatorname{sgn} \pi_p T_j [b_{pk} \tau_p] [0_{p+2} b_{p+10} q_{*p} \pi_p (p+1)_{p+1}] \right\}.$$

Thus, to prove that

$$\beta_{p+1}^S \rho_{*p}^S c_p^S = 0,$$

we are led to consider for a fixed  $k$  and  $q$ ,  $0 \leq k \leq p$ ,  $0 \leq q \leq p+1$ , the expression

$$(8) \quad Y_{kq} = \sum_{j=1}^n \sum_{\tau_p \in T_{pk}} \sum_{\pi_p \in T_{p0}} \lambda_j \operatorname{sgn} \tau_p \operatorname{sgn} \pi_p T_j [b_{pk} \tau_p] [0_{p+2} b_{p+10} q_{*p} \pi_p (p+1)_{p+1}].$$

Now to prove  $P_1$  we need only show that  $Y_{kq} = 0$ . Therefore  $k$  and  $q$  will remain fixed throughout the remainder of this section; and even though subsequent definitions will depend upon  $k$  and  $q$ , they will not be displayed in the notation.

2.4. For

$$\tau_p = \tau_p(i_0, \dots, i_p) \in T_{p0}$$

(see [3, § 1.9]) there exists a unique permutation  $(n_0, \dots, n_k)$  of  $0, \dots, k$  such that  $i_{n_0} < \dots < i_{n_k}$ . Let

$$\bar{\tau}_p = \bar{\tau}_p(j_0, \dots, j_p),$$

where  $j_l = i_{n_l}$  for  $l = 0, \dots, k$ , and  $j_l = i_l$  for  $k+1 \leq l \leq p$ . Then there exists

a unique permutation  $(m_0, \dots, m_k)$  of  $0, \dots, k$ , namely  $(n_0, \dots, n_k)^{-1}$ , such that

$$\tau_p = \tau_p(j_{m_0}, \dots, j_{m_k}, j_{k+1}, \dots, j_p).$$

Furthermore, let  $A(\tau_p)$  denote the set of  $\pi_p \in T_{p0}$  defined as follows. For

$$\pi_p = \pi_p(u_0, \dots, u_p) \in T_{p0}$$

we have a unique set of integers  $l_0, \dots, l_k$ ,  $0 \leq l_0 < \dots < l_k \leq p$  such that  $(u_{l_0}, \dots, u_{l_k})$  is a permutation of  $0, \dots, k$ . Set  $\pi_p \in A(\tau_p)$  if and only if  $m_0 = u_{l_0}, \dots, m_k = u_{l_k}$ .

2.5. Let  $B$  denote the set of ordered pairs  $(\tau_p, \pi_p)$ ,  $\tau_p \in T_{p0}$ ,  $\pi_p \in A(\tau_p)$ , and  $B'$  the set of ordered pairs  $(\tau'_p, \pi'_p)$ ,  $\tau'_p \in T_{pk}$ ,  $\pi'_p \in T_{p0}$ . We define a mapping

$$\gamma : B \rightarrow B'$$

as follows:

$$\gamma(\tau_p, \pi_p) = (\tau'_p, \pi'_p)$$

where  $\tau'_p = \overline{\tau_p}$  and  $\pi'_p = \pi_p$ . One shows with little difficulty that  $\gamma$  is biunique. Therefore

$$(9) \quad Y_{kq} = \sum_{j=1}^n \sum_{\tau_p \in T_{p0}} \sum_{\pi_p \in A(\tau_p)} \lambda_j \operatorname{sgn} \overline{\tau_p} \operatorname{sgn} \pi_p I_j [b_{pk} \overline{\tau_p}]$$

$$[0_{p+2} b_{p+10} q_{*p} \pi_p (p+1)_{p+1}].$$

2.6. Let  $A = A(\tau_p(0, \dots, p))$ . For  $\tau_p \in T_{p0}$  we define

$$f_{\tau_p} : A \rightarrow A(\tau_p)$$

as follows. For  $\pi_p(u_0, \dots, u_p) \in A$ , there exist integers  $l_0, \dots, l_k$ ,  $0 \leq l_0 < \dots < l_k \leq p$ , such that  $u_{l_0} = 0, \dots, u_{l_k} = k$ . Define

$$f_{\tau_p} \pi_p = \pi'_p(u'_0, \dots, u'_p)$$

as follows. Let

$$\bar{\tau}_p = \bar{\tau}_p(j_0, \dots, j_p) \text{ and } \tau_p = \tau_p(j_{m_0}, \dots, j_{m_k}, j_{k+1}, \dots, j_p),$$

where  $(m_0, \dots, m_k)$  is a permutation of  $0, \dots, k$ . Set  $u'_{l_0} = m_0, \dots, u'_{l_k} = m_k$ , and  $u'_r = u_r$  for  $r \neq l_0, \dots, l_k$ . Here again it is easy to show that  $f_{\tau_p}$  is bi-unique. We have then

$$(10) \quad Y_{kq} = \sum_{j=1}^n \sum_{\tau_p \in T_{p0}} \sum_{\pi_p \in A} \lambda_j \operatorname{sgn} \bar{\tau}_p \operatorname{sgn} f_{\tau_p} \pi_p T_j [b_{pk} \bar{\tau}_p] \\ [0_{p+2} b_{p+1} 0 q_{*p} f_{\tau_p} \pi_p (p+1)_{p+1}],$$

and hence

$$(11) \quad Y_{kq} = \sum_{s=1}^t \sum_{\pi_p \in A} \sum_{(j, \tau_p) \in E_s} \lambda_j \operatorname{sgn} \bar{\tau}_p \operatorname{sgn} f_{\tau_p} \pi_p T_j [b_{pk} \bar{\tau}_p] \\ [0_{p+2} b_{p+1} 0 q_{*p} f_{\tau_p} \pi_p (p+1)_{p+1}]$$

(see § 2.2).

2.7. LEMMA. Take  $\pi_p(u_0, \dots, u_p) \in T_{p0}$  and let

$$\alpha = [0_{p+2} b_{p+1} 0 q_{*p} \pi_p (p+1)_{p+1}].$$

Let

$$x = \sum_{j=0}^{p+1} \mu_j d_j,$$

with

$$\mu_j \geq 0, \quad j = 0, \dots, p+1, \text{ and } \sum_{j=0}^{p+1} \mu_j = 1,$$

denote a point of  $|d_0, \dots, d_{p+1}|$ . Then

$$\alpha(x) = \sum_{j=0}^{p+1} a_j d_j,$$

where

$$(i) \quad a_j \geq 0, \quad j = 0, \dots, p+1;$$

$$(ii) \quad \sum_{j=0}^{p+1} a_j = 1;$$

$$(iii) \quad a_{u_0} \geq a_{u_1} \geq \dots \geq a_{u_p};$$

(iv)  $a_{u_0}, \dots, a_{u_p}, a_{p+1}$  are independent of  $\pi_p$ ; i.e., if  $\pi'_p = \pi'_p(u'_0, \dots, u'_p) \in T_{p0}$  and

$$\alpha' = [0_{p+2} \ b_{p+1} \ 0 \ q_{*p} \ \pi'_p(p+1)_{p+1}],$$

then

$$\alpha'(x) = \sum_{j=0}^{p+1} a'_j d_j$$

with

$$a_{u_0} = a'_{u'_0}, \dots, a_{u_p} = a'_{u'_p}, \quad a_{p+1} = a'_{p+1}.$$

*Proof.* We consider only the case  $1 \leq q \leq p$  since the fringe cases  $q = 0$ ,  $p+1$  follow in a completely analogous manner. In case  $1 \leq q \leq p$  we have

$$\alpha = [b(w_0) b(w_0, w_1) \dots b(w_0, \dots, w_{p+1})],$$

where

$$w_l = d_{u_l}, \quad l = 0, \dots, q-1, \quad w_q = d_{p+1}, \quad w_l = d_{u_{l-1}}, \quad l = q+1, \dots, p+1.$$

Therefore,

$$\alpha(x) = \sum_{j=0}^{p+1} \mu_j b(w_0, \dots, w_j) = \sum_{j=0}^{p+1} \left( \sum_{l=j}^{p+1} \frac{\mu_l}{l+1} \right) w_j$$

(see § 1.2). Let

$$a_{p+1} = \sum_{l=q}^{p+1} \frac{\mu_l}{l+1}, \quad a_{u_r} = \sum_{l=r}^{p+1} \frac{\mu_l}{l+1} \quad \text{for } r = 0, \dots, q-1$$

and

$$a_{u_r} = \sum_{l=r+1}^{p+1} \frac{\mu_l}{l+1} \quad \text{for } r = q, \dots, p.$$

Clearly,  $a_{u_0}, \dots, a_{u_p}, a_{p+1}$  are independent of  $\pi_p$  in the sense of (iv), and  $a_{u_0} \geq \dots \geq a_{u_p}$ . Furthermore,  $a_j \geq 0$  ( $j = 0, \dots, p+1$ ), and

$$\sum_{j=0}^{p+1} a_j = \sum_{j=0}^{p+1} \mu_j = 1.$$

Also,

$$\alpha(x) = \sum_{j=0}^{q-1} a_{u_j} d_{u_j} + a_{p+1} d_{p+1} + \sum_{j=q}^p a_{u_j} d_{u_j} = \sum_{j=0}^{p+1} a_j d_j,$$

and the lemma follows.

2.8. LEMMA. Take  $(j, \tau_p)$  and  $(j', \tau'_p) \in E_s$  (see §2.2),  $1 \leq s \leq t$ , and  $\pi_p^* \in A$ . Then

$$\begin{aligned} & T_j [b_{pk} \bar{\tau}_p] [0_{p+2} b_{p+1} 0 q_{*p} f_{\tau_p} \pi_p^*(p+1)_{p+1}] \\ &= T_{j'} [b_{pk} \bar{\tau}'_p] [0_{p+2} b_{p+1} 0 q_{*p} f_{\tau'_p} \pi_p^*(p+1)_{p+1}]. \end{aligned}$$

*Proof.* Since  $(j, \tau_p), (j', \tau'_p)$  lie in  $E_s$ , we have

$$T_j [0_{p+1} b_{p0} \tau_p] = T_{j'} [0_{p+1} b_{p0} \tau'_p],$$

Let

$$\pi_p = f_{\tau_p} \pi_p^* = \pi_p(u_0, \dots, u_p), \quad \pi'_p = f_{\tau'_p} \pi_p^* = \pi'_p(u'_0, \dots, u'_p),$$

$$\alpha = [0_{p+2} b_{p+1} 0 q_{*p} \pi_p(p+1)_{p+1}], \quad \alpha' = [0_{p+2} b_{p+1} 0 q_{*p} \pi'_p(p+1)_{p+1}],$$

$$\gamma = [b_{pk} \bar{\tau}_p], \quad \text{and } \gamma' = [b_{pk} \bar{\tau}'_p].$$

Furthermore, let

$$\begin{aligned}\tau_p &= \tau_p(i_0, \dots, i_p), \quad \bar{\tau}_p = \bar{\tau}_p(j_0, \dots, j_p), \\ \tau'_p &= \tau'_p(i'_0, \dots, i'_p), \quad \bar{\tau}'_p = \bar{\tau}'_p(j'_0, \dots, j'_p).\end{aligned}$$

We have permutations  $(m_0, \dots, m_k), (n_0, \dots, n_k)$  of  $0, \dots, k$  such that

$$\begin{aligned}\tau_p &= \tau_p(j_{m_0}, \dots, j_{m_k}, j_{k+1}, \dots, j_p), \\ \tau'_p &= \tau'_p(j'_{n_0}, \dots, j'_{n_k}, j'_{k+1}, \dots, j'_p)\end{aligned}$$

Take an arbitrary point of  $|d_0, \dots, d_{p+1}|$ , say

$$x = \sum_{j=0}^{p+1} \mu_j d_j \quad \mu_j \geq 0, \quad \sum_{j=0}^{p+1} \mu_j = 1.$$

Then via the lemma of § 2.7 we have

$$\alpha(x) = \sum_{j=0}^{p+1} a_j d_j \quad \text{with } a_j \geq 0, \quad \sum_{j=0}^{p+1} a_j = 1, \quad a_{u_0} \geq \dots \geq a_{u_p},$$

and

$$\alpha'(x) = \sum_{j=0}^{p+1} a'_j d_j \quad \text{with } a'_j \geq 0, \quad \sum_{j=0}^{p+1} a'_j = 1, \quad a'_{u'_0} \geq \dots \geq a'_{u'_p},$$

with

$$a_{u_0} = a'_{u'_0}, \dots, a_{u_p} = a'_{u'_p} \quad \text{and} \quad a_{p+1} = a'_{p+1}.$$

Now

$$\gamma = [d_{j_0}, \dots, d_{j_k}, b(d_{j_0}, \dots, d_{j_k}), \dots, b(d_{j_0}, \dots, d_{j_p})].$$

Hence



$$\begin{aligned}
 \gamma \alpha(x) &= a_0 d_{j_0} + \dots + a_k d_{j_k} + a_{k+1} b(d_{j_0}, \dots, d_{j_p}) + \dots + \\
 & \qquad \qquad \qquad a_{p+1} b(d_{j_0}, \dots, d_{j_p}) \\
 &= a_{m_0} d_{j_{m_0}} + \dots + a_{m_k} d_{j_{m_k}} + a_{k+1} b(d_{j_0}, \dots, d_{j_k}) + \dots + \\
 & \qquad \qquad \qquad a_{p+1} b(d_{j_0}, \dots, d_{j_p}) \\
 &= a_{m_0} d_{j_{m_0}} + \dots + a_{m_k} d_{j_{m_k}} + a_{k+1} b(d_{j_{m_0}}, \dots, d_{j_{m_k}}) + \dots + \\
 & \qquad \qquad \qquad a_{p+1} b(d_{j_{m_0}}, \dots, d_{j_{m_k}}, d_{j_{k+1}}, \dots, d_{j_p}) \\
 &= a_{m_0} d_{i_0} + \dots + a_{m_k} d_{i_k} + a_{k+1} b(d_{i_0}, \dots, d_{i_k}) + \dots \\
 & \qquad \qquad \qquad + a_{p+1} b(d_{i_0}, \dots, d_{i_p}).
 \end{aligned}$$

Now take integers  $l_0, \dots, l_k$ ,  $0 \leq l_0 < \dots < l_k \leq p$ , such that  $(u_{l_0}, \dots, u_{l_k})$  is a permutation of  $0, \dots, k$ . Since  $\pi_p \in A(\tau_p)$ , we have  $m_0 = u_{l_0}, \dots, m_k = u_{l_k}$ . Hence  $a_{m_0} \geq \dots \geq a_{m_k}$ .

In a similar fashion we obtain

$$\begin{aligned}
 \gamma' \alpha'(x) &= a'_{n_0} d_{i'_0} + \dots + a'_{n_k} d_{i'_k} + a'_{k+1} b(d_{i'_0}, \dots, d_{i'_k}) + \dots \\
 & \qquad \qquad \qquad + a'_{p+1} b(d_{i'_0}, \dots, d_{i'_p}),
 \end{aligned}$$

with  $a'_{n_0} \geq \dots \geq a'_{n_k}$ ; and if  $l'_0, \dots, l'_k$ ,  $0 \leq l'_0 < \dots < l'_k \leq p$ , are integers such that  $(u'_{l'_0}, \dots, u'_{l'_k})$  is a permutation of  $0, \dots, k$ , we have

$$n_0 = u'_{l'_0}, \dots, n_k = u'_{l'_k}.$$

Applying § 1.3, we get

$$a_{m_0} d_{i_0} + \dots + a_{m_k} d_{i_k} = \sum_{l=0}^k \gamma_l b(d_{i_0}, \dots, d_{i_l})$$

with

$$\gamma_l = (l + 1)(a_{m_l} - a_{m_{l+1}}) \text{ for } l = 0, \dots, k - 1,$$

$$\gamma_k = (k + 1) a_{m_k},$$

and

$$\sum_{l=0}^k \gamma_l = \sum_{l=0}^k a_{m_l}.$$

Similarly,

$$a'_{n_0} d_{i'_0} + \cdots + a'_{n_k} d_{i'_k} = \sum_{l=0}^k \gamma'_l b(d_{i'_0}, \dots, d_{i'_l})$$

with

$$\gamma'_l = (l + 1) (a'_{n_l} - a'_{n_{l+1}}) \text{ for } l = 0, \dots, k - 1,$$

$$\gamma'_k = (k + 1) a'_{n_k}$$

and

$$\sum_{l=0}^k \gamma'_l = \sum_{l=0}^k a'_{n_l}.$$

However, since

$$\pi_p = f_{\tau_p} \pi_p^*, \quad \pi'_p = f_{\tau'_p} \pi_p^*,$$

we have

$$l_0 = l'_0, \dots, l_k = l'_k \text{ and } u_r = u'_r \text{ for } r \neq l_0, \dots, l_k.$$

Therefore,  $a_{u_{l_0}} = a'_{u'_{l'_0}}, \dots, a_{u_{l_k}} = a'_{u'_{l'_k}}$ , and hence

$$a_{m_0} = a'_{n_0}, \dots, a_{m_k} = a'_{n_k}.$$

Thus

$$\gamma_r = \gamma'_r \text{ for } r = 0, \dots, k.$$

Furthermore,

$$a_{u_r} = a'_{u'_r} \text{ for } r \neq l_0, \dots, l_k, \text{ and } a_{p+1} = a'_{p+1}.$$

Therefore,

$$\gamma \alpha(x) = \sum_{l=0}^k \gamma_l b(d_{i_0}, \dots, d_{i_l}) + \sum_{l=k}^p a_{l+1} b(d_{i_0}, \dots, d_{i_l}),$$

$$\gamma' \alpha'(x) = \sum_{l=0}^k \gamma_l b(d_{i'_0}, \dots, d_{i'_l}) + \sum_{l=k}^p a_{l+1} b(d_{i'_0}, \dots, d_{i'_l}),$$

with

$$\sum_{l=0}^k \gamma_l + \sum_{l=k}^p a_{l+1} = \sum_{l=0}^{p+1} a_l = 1.$$

Let

$$y = \sum_{j=0}^p h_j d_j$$

with

$$h_j = \gamma_j \text{ for } j = 0, \dots, k-1,$$

$$h_k = \gamma_k + a_{k+1},$$

$$h_j = a_{j+1} \text{ for } j = k+1, \dots, p.$$

Clearly,

$$h_j \geq 0 \quad (j = 0, \dots, p), \text{ and } \sum_{j=0}^p h_j = 1.$$

Then

$$\gamma \alpha(x) = \sum_{l=0}^p h_l b(d_{i_0}, \dots, d_{i_l}) = [0_{p+1} \ b_{p0} \ \tau_p](y)$$

and

$$\gamma' \alpha'(x) = \sum_{l=0}^p h_l b(d_{i_0}', \dots, d_{i_l}') = [0_{p+1} b_{p0} \tau_p'](y).$$

Therefore, since

$$T_j [0_{p+1} b_{p0} \tau_p'](y) = T_j' [0_{p+1} b_{p0} \tau_p'](y),$$

we have

$$T_j \gamma \alpha(x) = T_j' \gamma' \alpha'(x).$$

Since  $x$  is arbitrary in  $|d_0, \dots, d_{p+1}|$ , our lemma follows.

2.9. LEMMA. For any  $s, 1 \leq s \leq t$ , and  $\pi_p^* \in A$ ,

$$\sum_{(j, \tau_p) \in E_s} \lambda_j \operatorname{sgn} \bar{\tau}_p \operatorname{sgn} f_{\tau_p} \pi_p^* = 0.$$

*Proof.* Since

$$\operatorname{sgn} \bar{\tau}_p \operatorname{sgn} f_{\tau_p} \pi_p^* = \operatorname{sgn} \tau_p \operatorname{sgn} \pi_p^*,$$

we have

$$\sum_{(j, \tau_p) \in E_s} \lambda_j \operatorname{sgn} \bar{\tau}_p \operatorname{sgn} f_{\tau_p} \pi_p^* = \operatorname{sgn} \pi_p^* \sum_{(j, \tau_p) \in E_s} \lambda_j \operatorname{sgn} \tau_p = 0$$

via (5) of §2.2.

2.10. Employing §§2.8, 2.9, and (11) of §2.6, we see that  $Y_{kq} = 0$ , and hence  $P_1$  follows. Let us note also that since  $P_1 \equiv P_2, P_2$  also is valid.

### III. RESULTS

3.1. In [1, §4.2], Rado' has established a lemma, which we state here for the barycentric homotopy operator  $\rho_{*p}^R$ .

LEMMA. Let  $\{G_p\}$  be an identifier for  $R$ , such that the following conditions hold:

- (i)  $G_p \supset A_p^R$  (see [1, §3.4]),

(ii)  $c_p^R \in G_p$  implies that  $\sigma_p \beta_p^R c_p^R = 0$ ,

(iii)  $c_p^R \in G_p$  implies that  $\rho_{*p}^R c_p^R \in G_{p+1}$ .

Then  $\{G_p\}$  is an unessential identifier for  $R$ .

The proof of this lemma is identical with the proof of the corresponding lemma as given by Rado' with  $\rho_p^R$  (classical homotopy operator) replacing  $\rho_{*p}^R$ .

Since

$$\sigma_p \beta_p^R : C_p^R \longrightarrow C_p^S$$

is a chain mapping, the system  $\{N(\sigma_p \beta_p^R)\}$  of nuclei of the homomorphisms  $\sigma_p \beta_p^R$  is an identifier for  $R$  (see [1, §1.2]). Furthermore,

$$N(\sigma_p \beta_p^R) \supset A_p^R \text{ since } \sigma_p \beta_p^R = \beta_p^S \sigma_p$$

(see §1.11). Applying  $P_2$  directly, we see that  $N(\sigma_p \beta_p^R)$  satisfies (iii) of the foregoing lemma. Therefore, since  $N(\sigma_p \beta_p^R)$  is the largest identifier, satisfying (ii), we have the following maximum result yielded by the same lemma:

**THEOREM.** *The system  $\{N(\sigma_p \beta_p^R)\}$  is an unessential identifier for  $R$ .*

3.2. In order to compare our results with those of Radó [1] and Reichelderfer [3] let us first note that

$$\hat{N}(\sigma_p \beta_p^R) = N(\sigma_p \beta_p^R),$$

where  $\hat{N}(\sigma_p \beta_p^R)$  is the division hull of  $N(\sigma_p \beta_p^R)$ , since  $C_p^R$  is a free Abelian group. Then since

$$N(\sigma_p \beta_p^R) \supset \Delta_p^R = N(\beta_p^R) + A_p^R$$

(see [3, §3.6]) we have

$$N(\sigma_p \beta_p^R) \supset \hat{\Delta}_p^R \supset \hat{\Gamma}_p^R$$

(see [1, §4.7]).

The writer has been unable to determine as yet whether  $N(\sigma_p \beta_p^R)$  is effectively larger than either  $\hat{\Delta}_p^R$  or  $\hat{\Gamma}_p^R$ .

3.3. The following lemma (see [1, §4.1]) is immediate from the fact that  $\rho_{*p}^S$  satisfies the well-known "homotopy identity,"

$$\partial_{p+1}^S \rho_{*p}^S + \rho_{*p-1}^S \partial_p^S = \beta_p^S - 1.$$

LEMMA. Let  $\{G_p\}$  be an identifier for  $S$  such that the following conditions hold:

- (i)  $c_p^S \in G_p$  implies that  $\beta_p^S c_p^S = 0$ ,
- (ii)  $c_p^S \in G_p$  implies that  $\rho_{*p}^S c_p^S \in G_{p+1}$ .

Then  $\{G_p\}$  is an unessential identifier for  $S$ .

The system of nuclei  $\{N(\beta_p^S)\}$  clearly is an identifier for  $S$  since  $\beta_p^S$  is a chain mapping. Therefore, applying  $P_1$  we obtain the maximum result of the foregoing lemma.

THEOREM. The system  $\{N(\beta_p^S)\}$  is an unessential identifier for  $S$ .

#### REFERENCES

1. T. Radó, *An approach to singular homology theory*, Pacific J. Math. **1** (1951), 265-290.
2. ———, *On identifications in singular homology theory*, Rivista Mat. Univ. Parma, **2** (1951), 3-18.
3. P. V. Reichelderfer, *On the barycentric homomorphism in a singular complex*, Pacific J. Math. **2** (1952), 73-97.
4. S. Eilenberg and N. E. Steenrod, *Foundations of algebraic topology*, Princeton, 1952.



# PACIFIC JOURNAL OF MATHEMATICS

## EDITORS

R. M. ROBINSON  
University of California  
Berkeley 4, California

E. HEWITT  
University of Washington  
Seattle 5, Washington

R. P. DILWORTH  
California Institute of Technology  
Pasadena 4, California

E. F. BECKENBACH  
University of California  
Los Angeles 24, California

## ASSOCIATE EDITORS

H. BUSEMANN      P. R. HALMOS  
HERBERT FEDERER      HEINZ HOPF  
MARSHALL HALL      R. D. JAMES

BØRGE JESSEN      J. J. STOKER  
PAUL LÉVY      E. G. STRAUS  
GEORGE PÓLYA      KÔSAKU YOSIDA

## SPONSORS

UNIVERSITY OF BRITISH COLUMBIA  
CALIFORNIA INSTITUTE OF TECHNOLOGY  
UNIVERSITY OF CALIFORNIA, BERKELEY  
UNIVERSITY OF CALIFORNIA, DAVIS  
UNIVERSITY OF CALIFORNIA, LOS ANGELES  
UNIVERSITY OF CALIFORNIA, SANTA BARBARA  
UNIVERSITY OF NEVADA  
OREGON STATE COLLEGE  
UNIVERSITY OF OREGON

UNIVERSITY OF SOUTHERN CALIFORNIA  
STANFORD RESEARCH INSTITUTE  
STANFORD UNIVERSITY  
WASHINGTON STATE COLLEGE  
UNIVERSITY OF WASHINGTON  
\* \* \*  
AMERICAN MATHEMATICAL SOCIETY  
NATIONAL BUREAU OF STANDARDS,  
INSTITUTE FOR NUMERICAL ANALYSIS

---

Mathematical papers intended for publication in the *Pacific Journal of Mathematics* should be typewritten (double spaced), and the author should keep a complete copy. Manuscripts may be sent to any of the editors except Robinson, whose term expires with the completion of the present volume; they might also be sent to M. M. Schiffer, Stanford University, Stanford, California, who is succeeding Robinson. All other communications to the editors should be addressed to the managing editor, E. F. Beckenbach, at the address given above.

Authors are entitled to receive 100 free reprints of their published papers and may obtain additional copies at cost.

---

The *Pacific Journal of Mathematics* is published quarterly, in March, June, September, and December. The price per volume (4 numbers) is \$8.00; single issues, \$2.50. Special price to individual faculty members of supporting institutions and to individual members of the American Mathematical Society: \$4.00 per volume; single issues, \$1.25.

Subscriptions, orders for back numbers, and changes of address should be sent to the publishers, University of California Press, Berkeley 4, California.

Printed at Ann Arbor, Michigan. Entered as second class matter at the Post Office, Berkeley, California.

UNIVERSITY OF CALIFORNIA PRESS • BERKELEY AND LOS ANGELES

COPYRIGHT 1953 BY PACIFIC JOURNAL OF MATHEMATICS



L. Carlitz, <i>Some theorems on generalized Dedekind sums</i> .....	513
L. Carlitz, <i>The reciprocity theorem for Dedekind sums</i> .....	523
Edward Richard Fadell, <i>Identifications in singular homology theory</i> .....	529
Harley M. Flanders, <i>A method of general linear frames in Riemannian geometry. I</i> .....	551
Watson Bryan Fulks, <i>The Neumann problem for the heat equation</i> .....	567
Paul R. Garabedian, <i>Orthogonal harmonic polynomials</i> .....	585
R. E. Greenwood and Andrew Mattei Gleason, <i>Distribution of round-off errors for running averages</i> .....	605
Arthur Eugene Livingston, <i>The space <math>H^p</math>, <math>0 &lt; p &lt; 1</math>, is not normable</i> .....	613
M. N. Mikhail, <i>On the order of the reciprocal set of a basic set of polynomials</i> .....	617
Louis Joel Mordell, <i>On the linear independence of algebraic numbers</i> .....	625
Leo Sario, <i>Alternating method on arbitrary Riemann surfaces</i> .....	631
Harold Nathaniel Shapiro, <i>Iterates of arithmetic functions and a property of the sequence of primes</i> .....	647
H. Shniad, <i>Convexity properties of integral means of analytic functions</i> .....	657
Marlow C. Sholander, <i>Plane geometries from convex plates</i> .....	667