

# Pacific Journal of Mathematics

**COMPLETELY CONTINUOUS NORMAL OPERATORS WITH  
PROPERTY  $L$**

IRVING KAPLANSKY

# COMPLETELY CONTINUOUS NORMAL OPERATORS WITH PROPERTY $L$

IRVING KAPLANSKY

**1. Introduction.** Two matrices  $A$  and  $B$  are said to have property  $L$  if it is possible to arrange their characteristic roots

$$A: \lambda_1, \lambda_2, \dots, \lambda_n$$

$$B: \mu_1, \mu_2, \dots, \mu_n$$

in such a way that for every  $\alpha$ , the characteristic roots of  $\alpha A + B$  are given by  $\alpha \lambda_i + \mu_i$ . In [1] this property is investigated, and among other things a conjecture of Kac is confirmed by showing that if  $A$  and  $B$  are hermitian, then they commute. In [2] this is generalized by replacing "hermitian" by "normal".

In this note we launch the project of generalizing such results to (complex) Hilbert space. However, since it is not clear how to formulate the problem for general operators (especially in the presence of a continuous spectrum), we shall content ourselves with the completely continuous case. For self-adjoint operators we obtain a fully satisfactory generalization (Theorem 1). For the more general case of normal operators we find ourselves obliged to make an extra assumption roughly to the effect that nonzero characteristic roots are paired only to nonzero roots. In the finite-dimensional case such an assumption would be harmless; indeed, by adding suitable constants to  $A$  and  $B$ , we could even arrange to have all the characteristic roots of  $A$  and  $B$  nonzero. It would nevertheless be of interest to determine whether this blemish can be removed from Theorem 2.

**2. Remarks.** Before we state the results, some remarks are in order. The number  $\lambda$  is a characteristic root of  $A$  if  $A - \lambda I$  has a nonzero null space. If  $A$  is a completely continuous normal operator, its characteristic roots are either finite in number or form a sequence approaching zero. We have an orthogonal decomposition of the Hilbert space:

---

Received August 23, 1952. The preparation of this paper was sponsored (in part) by the Office of Naval Research.

$$H = H(0) \oplus H(\lambda_1) \oplus H(\lambda_2) \oplus \dots,$$

where  $A$  acts on  $H(\lambda_i)$  as a multiplication by  $\lambda_i$ . The dimension of  $H(\lambda_i)$  is called the multiplicity of the characteristic root  $\lambda_i$ ; it is finite except possibly for the characteristic root 0.

Now even though  $A$  and  $B$  are both to be normal,  $\alpha A + B = C$  need not (a priori) be normal. We must accordingly give further attention to the meaning of the multiplicity of a characteristic root  $\nu$  of  $C$ . For our purposes virtually any reasonable definition would do; we select the following one. We note that the null spaces of the operators  $C - \nu I$ ,  $(C - \nu I)^2$ ,  $\dots$  form an ascending chain, and we form their union; the dimension of this union is the multiplicity of  $\nu$ . Note that this agrees with customary usage in the finite-dimensional case.

We shall need the (easily proved) additivity of the multiplicity. In detail: suppose  $H$  is an orthogonal direct sum of two closed subspaces both invariant under  $C$ ; then the multiplicity of  $\nu$  in the whole space is the sum of its multiplicities in the two subspaces.

**3. Results.** We are now ready to define property  $L$ . We do this in a way that is adequate for the proof, although it does not treat  $A$  and  $B$  symmetrically.

Let  $A$  and  $B$  be completely continuous normal operators. Let there be given two sequences  $\lambda_i$ ,  $\mu_i$  of complex numbers. We say that  $A$  and  $B$  have *property  $L$*  (relative to the two sequences) provided:

- (1) The  $\lambda$ 's constitute precisely the nonzero characteristic roots of  $A$ , each counted as often as its multiplicity.
- (2) If, for a certain  $\alpha$  and  $\nu$ , there are  $k$  values of  $i$  such that  $\nu = \alpha \lambda_i + \mu_i$ , then  $\alpha A + B$  has  $\nu$  as a characteristic root at least of multiplicity  $k$ .

**THEOREM 1.** *Let  $A$  and  $B$  be completely continuous self-adjoint operators with property  $L$ . Then  $A$  and  $B$  commute.*

**THEOREM 2.** *Let  $A$  and  $B$  be completely continuous normal operators with property  $L$ , relative to the sequences  $\lambda_i$  and  $\mu_i$ . Suppose further that the  $\mu$ 's are all nonzero. Then  $A$  and  $B$  commute.*

**4. Proof.** The two theorems can conveniently be proved simultaneously. We can suppose that  $\lambda_1$  is a characteristic root of maximum absolute value, that is,  $|\lambda_1| = \|A\|$ . For brevity write  $\lambda = \lambda_1$ ,  $\mu = \mu_1$ . By an application of the definition of property  $L$ , with  $\alpha = 0$ , we see that  $\mu$  is a characteristic root of  $B$ . We are going to prove that there exists a nonzero vector  $x$  with

$$Ax = \lambda x, Bx = \mu x.$$

If  $\mu \neq 0$ , we are ready to proceed. If  $\mu = 0$ , then by hypothesis both  $A$  and  $B$  are self-adjoint. We replace  $B$  by  $A + B$  which is again self-adjoint; this replaces  $\mu$  by

$$\lambda + \mu = \lambda \neq 0.$$

So in any event we are entitled to assume that  $\mu$  is nonzero.

Let  $H(\mu)$  be the (finite-dimensional) characteristic subspace of  $B$  for the characteristic root  $\mu$ , and  $K$  the orthogonal complement; let  $E$  and  $F$  be the projections on  $H(\mu)$  and  $K$ . We note that  $B - \mu I$  is nonsingular on  $K$ ; let  $S$  be defined as its inverse on  $K$  and as 0 on  $H(\mu)$ . Thus we have

$$(1) \quad SF(B - \mu I) = F.$$

Next we consider  $E(A - \lambda I)E$  as an operator on  $H(\mu)$ , and we are going to prove that it is singular. Suppose the contrary and define  $R$  to be its inverse on  $H(\mu)$ , 0 on  $K$ . Then  $R$  will satisfy

$$(2) \quad R(B - \mu I) = 0, R(A - \lambda I)E = E, RF = 0.$$

Choose  $\alpha \neq 0$  so that

$$(3) \quad \|\alpha SF(A - \lambda I)(F - RAF)\| < 1.$$

By hypothesis, the operator  $\alpha A + B$  has  $\alpha\lambda + \mu$  as a characteristic root, say with characteristic vector  $y \neq 0$ . We have

$$(4) \quad \alpha(A - \lambda I)y + (B - \mu I)y = 0.$$

Write  $y = Ey + Fy$  in (4), apply  $R$ , and then use (2); we find that  $Ey = -RAFy$ , and so

$$(5) \quad y = Ey + Fy = (F - RAF)y.$$

Next apply  $SF$  to (4), and use (1) and (5):

$$(6) \quad Fy = -\alpha SF(A - \lambda I)(F - RAF)y.$$

On contemplating (6) in conjunction with (3) we see that  $Fy$  must be 0. But then  $y = 0$  by (5). This contradiction shows that we were in error in supposing  $E(A - \lambda I)E$  to be nonsingular on  $H(\mu)$ . Consequently we can find in  $H(\mu)$

a nonzero vector  $x$  annihilated by  $E(A - \lambda I)E$ . Then since  $Ex = x$ , we have  $EAx = \lambda x$ . Form the orthogonal decomposition

$$(7) \quad Ax = EAx + FAx = \lambda x + FAx.$$

But

$$\|Ax\| \leq |\lambda| \|x\|,$$

since the norm of  $A$  is  $|\lambda|$ . Hence in (7) we must actually have  $Ax = \lambda x$ . Also  $Bx = \mu x$  since  $x$  is in  $H(\mu)$ , and we have fulfilled our initial objective.

Let  $M$  be the orthogonal complement of  $x$ . It follows from the additivity of multiplicity (see above) that when the operator  $\alpha A + B$  is confined to  $M$ , the multiplicity of its characteristic root  $\alpha\lambda + \mu$  is diminished by precisely 1, while all other characteristic roots have unchanged multiplicity. Thus  $A$  and  $B$ , confined to  $M$ , satisfy property  $L$  relative to the sequences  $\lambda_i$  and  $\mu_i$  for  $i \geq 2$ . The procedure may now be repeated to get within  $M$  another joint characteristic vector for  $A$  and  $B$ . In this way we proceed down the nonzero characteristic roots of  $A$ . Finally we are left with the null space of  $A$ , which of course commutes with whatever is left of  $B$ . Hence  $A$  and  $B$  commute.

**5. Remark.** As soon as we know that  $A$  and  $B$  commute (and hence can be simultaneously put in diagonal form), we can assert that they satisfy property  $L$  symmetrically, and indeed various stronger statements are obvious consequences of simultaneous diagonal form.

#### REFERENCES

1. T. S. Motzkin and O. Taussky, *Pairs of matrices with property L*, Trans. Amer. Math. Soc. 73 (1952), 108-114.
2. N. A. Wiegmann, *A note on pairs of normal matrices with property L*, Proc. Amer. Math. Soc. 4 (1953), 35-36.

UNIVERSITY OF CHICAGO AND  
NATIONAL BUREAU OF STANDARDS, LOS ANGELES

# PACIFIC JOURNAL OF MATHEMATICS

## EDITORS

R. M. ROBINSON  
University of California  
Berkeley 4, California

R. P. DILWORTH  
California Institute of Technology  
Pasadena 4, California

E. HEWITT  
University of Washington  
Seattle 5, Washington

E. F. BECKENBACH  
University of California  
Los Angeles 24, California

## ASSOCIATE EDITORS

H. BUSEMANN      P. R. HALMOS  
HERBERT FEDERER      HEINZ HOPF  
MARSHALL HALL      R. D. JAMES

BØRGE JESSEN      J. J. STOKER  
PAUL LÉVY      E. G. STRAUS  
GEORGE PÓLYA      KÔSAKU YOSIDA

## SPONSORS

UNIVERSITY OF BRITISH COLUMBIA  
CALIFORNIA INSTITUTE OF TECHNOLOGY  
UNIVERSITY OF CALIFORNIA, BERKELEY  
UNIVERSITY OF CALIFORNIA, DAVIS  
UNIVERSITY OF CALIFORNIA, LOS ANGELES  
UNIVERSITY OF CALIFORNIA, SANTA BARBARA  
UNIVERSITY OF NEVADA  
OREGON STATE COLLEGE  
UNIVERSITY OF OREGON

UNIVERSITY OF SOUTHERN CALIFORNIA  
STANFORD RESEARCH INSTITUTE  
STANFORD UNIVERSITY  
WASHINGTON STATE COLLEGE  
UNIVERSITY OF WASHINGTON  
\* \* \*  
AMERICAN MATHEMATICAL SOCIETY  
NATIONAL BUREAU OF STANDARDS,  
INSTITUTE FOR NUMERICAL ANALYSIS

---

Mathematical papers intended for publication in the *Pacific Journal of Mathematics* should be typewritten (double spaced), and the author should keep a complete copy. Manuscripts may be sent to any of the editors except Robinson, whose term expires with the completion of the present volume; they might also be sent to M. M. Schiffer, Stanford University, Stanford, California, who is succeeding Robinson. All other communications to the editors should be addressed to the managing editor, E. F. Beckenbach, at the address given above.

Authors are entitled to receive 100 free reprints of their published papers and may obtain additional copies at cost.

---

The *Pacific Journal of Mathematics* is published quarterly, in March, June, September, and December. The price per volume (4 numbers) is \$8.00; single issues, \$2.50. Special price to individual faculty members of supporting institutions and to individual members of the American Mathematical Society: \$4.00 per volume; single issues, \$1.25.

Subscriptions, orders for back numbers, and changes of address should be sent to the publishers, University of California Press, Berkeley 4, California.

Printed at Ann Arbor, Michigan. Entered as second class matter at the Post Office, Berkeley, California.

UNIVERSITY OF CALIFORNIA PRESS · BERKELEY AND LOS ANGELES

COPYRIGHT 1953 BY PACIFIC JOURNAL OF MATHEMATICS

|   |     |
|---|-----|
| Paul Erdős and Gilbert Agnew Hunt, Jr., <i>Changes of sign of sums of random variables</i> .....      | 673 |
| Paul Erdős and Ernst Gabor Straus, <i>On linear independence of sequences in a Banach space</i> ..... | 689 |
| Haim Hanani, <i>On sums of series of complex numbers</i> .....  | 695 |
| Melvin Henriksen, <i>On the prime ideals of the ring of entire functions</i> .....                    | 711 |
| Irving Kaplansky, <i>Completely continuous normal operators with property L</i> .....                 | 721 |
| Samuel Karlin, <i>Some random walks arising in learning models. I</i> .....                           | 725 |
| William Judson LeVeque, <i>On uniform distribution modulo a subdivision</i> .....                     | 757 |
| Lee Lorch, <i>Derivatives of infinite order</i> .....   | 773 |
| Ernest A. Michael, <i>Some extension theorems for continuous functions</i> .....                      | 789 |
| Tyre Alexander Newton, <i>A note on the Hölder mean</i> .....   | 807 |
| Raymond Moos Redheffer, <i>On a theorem of Plancherel and Pólya</i> .....                             | 823 |
| Choy-Tak Taam, <i>On the complex zeros of functions of Sturm-Liouville type</i> .....                 | 837 |