

# Pacific Journal of Mathematics

**ON THE NUMBER OF PRIMITIVE PYTHAGOREAN  
TRIANGLES WITH AREA LESS THAN  $n$**

ROY EDWIN WILD

ON THE NUMBER OF PRIMITIVE PYTHAGOREAN TRIANGLES  
WITH AREA LESS THAN  $n$

ROY E. WILD

**1. Introduction.** In the preceding paper Lambek and Moser have shown that if  $P_a(n)$  is the number of primitive Pythagorean triangles with area less than  $n$  then

$$(1) \quad P_a(n) = cn^{1/2} + O(n^{1/3}),$$

where

$$c = \frac{\Gamma^2(1/4)}{2^{1/2} \pi^{5/2}}.$$

They conjecture that

$$(2) \quad P_a(n) = cn^{1/2} - c'n^{1/3} + o(n^{1/3}),$$

and on the basis of a table due to Miksa they find

$$(3) \quad c' \approx .295.$$

Our purpose here is to show that

$$(4) \quad P_a(n) = cn^{1/2} - c'n^{1/3} + O(n^{1/4} \ln n),$$

where

$$(5) \quad c' = -\frac{\zeta(1/3)(1+2^{-1/3})}{\zeta(4/3)(1+4^{-1/3})} \approx .297.$$

In the paper by Lambek and Moser, the problem of calculating  $P_a(n)$  has been reduced to that of counting the number of lattice points  $L(n)$  in the region  $R_1$  defined by

---

Received December 3, 1953. The author is grateful to Professor E. G. Straus for calling his attention to the problem treated in this paper, and for making valuable suggestions and criticisms.

*Pacific J. Math.* 5 (1955), 85-91

$$(6) \quad xy(y^2 - x^2) < n, \quad y > x > 0.$$

They obtain


$$(7) \quad L(n) = \frac{\Gamma^2(1/4)}{2^{5/2} \pi^{1/2}} n^{1/2} + O(n^{1/3}).$$

We shall obtain, in place of (7),

$$(8) \quad L(n) = \frac{\Gamma^2(1/4)}{2^{5/2} \pi^{1/2}} n^{1/2} + (1 + 2^{-1/3}) \zeta(1/3) n^{1/3} + O(n^{1/4}).$$

**2. Proof of (8).** Following is the graph of

$$(9) \quad K_1 : xy(y^2 - x^2) = n.$$



From  $K_1$  we obtain the curve  $K_2$  by replacing  $y$  in  $K_1$  by  $y + x$  to get

$$(10) \quad K_2 : xy(x + y)(2x + y) = n.$$

This transformation preserves the area and number of lattice points in  $R_1$ . So we count the lattice points in  $R_2$  defined by

$$(11) \quad xy(x + y)(2x + y) < n, \quad x > 0, \quad y > 0.$$

By Cardan's formulas, we obtain, from (10),

$$(12) \quad x = \left(\frac{n}{4y}\right)^{1/3} \left\{ \left[ 1 + \left(1 - \frac{y^8}{108n^2}\right)^{1/2} \right]^{1/3} + \left[ 1 - \left(1 - \frac{y^8}{108n^2}\right)^{1/2} \right]^{1/3} \right\} - \frac{y}{2}$$

and

$$(13) \quad y = \left(\frac{n}{2x}\right)^{1/3} \left\{ \left[ 1 + \left(1 - \frac{4x^8}{27n^2}\right)^{1/2} \right]^{1/3} + \left[ 1 - \left(1 - \frac{4x^8}{27n^2}\right)^{1/2} \right]^{1/3} \right\} - x.$$

In (13) take

$$x = x_1 = \left(\frac{27}{4}\right)^{1/8} n^{1/4},$$

say, so that

$$y = y_1 = \left( \left(\frac{64}{3}\right)^{1/8} - \left(\frac{27}{4}\right)^{1/8} \right) n^{1/4},$$

thus determining the point  $p_1 : (x_1, y_1)$  on  $K_2$ .

Let square brackets denote the greatest integer function. Then from the figure we have

$$(14) \quad L(n) = \sum_{x=1}^{[x_1]} [y] + \sum_{y=1}^{[y_1]} [x] - [x_1][y_1] - l(n),$$

where  $l(n)$  is the number of lattice points on  $K_2$ . Now  $l(n)$  is zero unless  $n$  is an integer  $N$ . For nonintegral  $n$  we can prove (8). For small positive  $\epsilon$  we obtain (8) for, say,  $N + \epsilon$  and  $N - \epsilon$ , so that trivially

$$(15) \quad l(n) = O(n^{1/4}) \text{ for all real } n.$$

By definition,

$$(16) \quad [x_1] = O(n^{1/4}), \quad [y_1] = O(n^{1/4}),$$

so that we may drop the brackets in (14) with an error  $O(n^{1/4})$ . Then, by use of (15) and (16), (14) becomes

$$(17) \quad L(n) = \sum_{x=1}^{[x_1]} y + \sum_{y=1}^{[y_1]} x - x_1 y_1 + O(n^{1/4}).$$

We shall estimate the above sums by the Euler-Maclaurin summation formula in the form:

$$(18) \quad \sum_{k=a}^b f(k) = \int_a^b f(x) dx + \frac{1}{2} f(b) + \frac{1}{2} f(a) + \int_a^b \left( x - [x] - \frac{1}{2} \right) f'(x) dx.$$

We obtain from (17):

$$(19) \quad L(n) = \int_1^{[x_1]} y dx + \int_1^{[y_1]} x dy - x_1 y_1 + O(n^{1/4}) \\ + \frac{1}{2} y([x_1]) + \frac{1}{2} y(1) + \frac{1}{2} x([y_1]) + \frac{1}{2} x(1) \\ + \int_1^{[x_1]} \left( x - [x] - \frac{1}{2} \right) \frac{dy}{dx} dx + \int_1^{[y_1]} \left( y - [y] - \frac{1}{2} \right) \frac{dx}{dy} dy.$$

In the first two terms of (19), we may drop brackets with an error of  $O(n^{1/4})$ , so that, if  $A$  represents the entire area of  $R_2$ , we may replace the first three terms of (19) by

$$(20) \quad A - \int_0^1 y dx - \int_0^1 x dy + O(n^{1/4}).$$

Now from (12) and (13) we have

$$(21) \quad x = (n/4y)^{1/3} (2^{1/3} + O(y^{8/3}/n^{2/3})) + O(y) \\ = (n/2y)^{1/3} + O(y^{7/3}/n^{1/3}) + O(y),$$

and similarly

$$(22) \quad y = (n/x)^{1/3} + O(x^{7/3}/n^{1/3}) + O(x).$$

Substituting in (20), we obtain

$$(23) \quad A - 3n^{1/3}/2 + O(n^{-1/3}) + O(1) \\ - 3n^{1/3}/2^{4/3} + O(n^{-1/3}) + O(1) + O(n^{1/4}) \\ = A - \frac{3}{2} (1 + 2^{-1/3}) n^{1/3} + O(n^{1/4}).$$

The fifth and seventh terms of (19) are  $O(n^{1/4})$ . Also

$$\frac{1}{2}y(1) = \frac{1}{2}n^{1/3} + O(1) \quad \text{and} \quad \frac{1}{2}x(1) = n^{1/3}/2^{4/3} + O(1).$$

Differentiating the expansions of  $x$  and  $y$  in (21) and (22) we obtain

$$(24) \quad dx/dy = -n^{1/3}y^{-4/3}/3 \cdot 2^{1/3} + O(y^{4/3}/n^{1/3}) + O(1),$$

$$(25) \quad dy/dx = -n^{1/3}x^{-4/3}/3 + O(x^{4/3}/n^{1/3}) + O(1).$$

We now rewrite the last two terms of (19) as

$$(26) \quad \int_1^{n^{1/8}} \left( x - [x] - \frac{1}{2} \right) \frac{dy}{dx} dx + \int_{n^{1/8}}^{[x_1]} \left( x - [x] - \frac{1}{2} \right) \frac{dy}{dx} dx \\ + \int_1^{n^{1/8}} \left( y - [y] - \frac{1}{2} \right) \frac{dx}{dy} dy + \int_{n^{1/8}}^{[y_1]} \left( y - [y] - \frac{1}{2} \right) \frac{dx}{dy} dy.$$

Since  $|dy/dx|$  is monotonic decreasing, we have, by the second mean value theorem for integrals, and (25),

$$(27) \quad \int_{n^{1/8}}^{[x_1]} \left( x - [x] - \frac{1}{2} \right) \frac{dy}{dx} dx = \frac{dy}{dx} \Big|_{x=n^{1/8}} \int_{n^{1/8}}^h \left( x - [x] - \frac{1}{2} \right) dx \\ = O(n^{1/6}) O(1) = O(n^{1/6}).$$

Similarly

$$(28) \quad \int_{n^{1/8}}^{[y_1]} \left( y - [y] - \frac{1}{2} \right) \frac{dx}{dy} dy = O(n^{1/6}).$$

Substituting (24) and (25) in the remaining terms of (26) yields

$$(29) \quad \int_1^{n^{1/8}} \left( x - [x] - \frac{1}{2} \right) \frac{dy}{dx} dx = -\frac{n^{1/3}}{3} \int_1^{n^{1/8}} \left( x - [x] - \frac{1}{2} \right) x^{-4/3} dx + O(1) \\ = -\frac{n^{1/3}}{3} \int_1^\infty \left( x - [x] - \frac{1}{2} \right) x^{-4/3} dx + \frac{n^{1/3}}{3} \int_{n^{1/8}}^\infty \left( x - [x] - \frac{1}{2} \right) x^{-4/3} dx + O(1) \\ = -\frac{n^{1/3}}{3} \int_1^\infty \left( x - [x] - \frac{1}{2} \right) x^{-4/3} dx + O(n^{1/6}),$$

and similarly

$$(30) \quad \int_1^{n^{1/8}} \left( y - [y] - \frac{1}{2} \right) \frac{dx}{dy} dy = -\frac{n^{1/3}}{3 \cdot 2^{1/3}} \int_1^\infty \left( y - [y] - \frac{1}{2} \right) y^{-4/3} dy \\ + O(n^{1/6}).$$

Collecting the preceding results, we have

$$(31) \quad L(n) = A - \frac{3}{2} (1 + 2^{-1/3}) n^{1/3} + O(n^{1/4}) + O(n^{1/4}) + n^{1/3}/2 + O(1) \\ + n^{1/3}/2^{4/3} + O(1) + (1 + 2^{-1/3}) c_1 n^{1/3} + O(n^{1/6}) \\ = A - (1 + 2^{-1/3}) (1 - c_1) n^{1/3} + O(n^{1/4}),$$

where

$$(32) \quad c_1 = \int_1^\infty \left( x - [x] - \frac{1}{2} \right) dx^{-1/3} = \zeta(1/3) + 1.$$

Now  $A$  is the area of  $R_2$  and therefore the area of  $R_1$ . Its value as calculated by Lambek and Moser is

$$(33) \quad A = c_2 n^{1/2}, \quad c_2 = \frac{\Gamma^2(1/4)}{2^{5/2} \pi^{1/2}}.$$

Substitution from (32) and (33) in (31) yields (8).

**3. Derivation of (4).** Let  $c_3 = -(1 + 2^{-1/3}) \zeta(1/3)$ , so that (31) becomes

$$(34) \quad L(n) = c_2 n^{1/2} - c_3 n^{1/3} + O(n^{1/4}).$$

Following the notation of Lambek and Moser, we can write (34) as

$$(35) \quad L(Rt) = c_2 t^2 - c_3 t^{4/3} + O(t).$$

From their equation (14) we have

$$(36) \quad L^*(Rt) = \sum_{i \geq 1} \mu(i) \left\{ c_2 \frac{t^2}{i^2} - c_3 \frac{t^{4/3}}{i^{4/3}} + O(t/i) \right\}$$

$$\begin{aligned}
&= \{6c_2/\pi^2 + O(1/t)\}t^2 - \{c_3/\zeta(4/3) + O(1/t^{1/3})\}t^{4/3} + O(t \ln t) \\
&= \frac{6c_2}{\pi^2} t^2 - \frac{c_3}{\zeta(4/3)} t^{4/3} + O(t \ln t).
\end{aligned}$$

Then from their equations (6), (15), and our (36), we have

$$\begin{aligned}
(37) \quad P_a(n) &= \sum_{i \geq 0} (-1)^i \left\{ \frac{6c_2 n^{1/2}}{\pi^2 2^i} - \frac{c_3 n^{1/3}}{\zeta(4/3) 4^{i/3}} \right. \\
&\quad \left. + O\left(\frac{n^{1/4}}{4^{i/4}} \ln \frac{n}{4^i}\right) \right\} \\
&= 4c_2 n^{1/2}/\pi^2 - \frac{c_3 n^{1/3}}{\zeta(4/3) (1 + 4^{-1/3})} + O(n^{1/4} \ln n) \\
&= c n^{1/2} - c' n^{1/3} + O(n^{1/4} \ln n).
\end{aligned}$$

This is (4).

UNIVERSITY OF IDAHO







# Pacific Journal of Mathematics

Vol. 5, No. 1

September, 1955

Frank Herbert Brownell, III, <i>Flows and noncommuting projections on Hilbert space</i> .....	1
H. E. Chrestenson, <i>A class of generalized Walsh functions</i> .....	17
Jean Bronfenbrenner Crockett and Herman Chernoff, <i>Gradient methods of maximization</i> .....	33
Nathan Jacob Fine, <i>On groups of orthonormal functions. I</i> .....	51
Nathan Jacob Fine, <i>On groups of orthonormal functions. II</i> .....	61
Frederick William Gehring, <i>A note on a paper by L. C. Young</i> .....	67
Joachim Lambek and Leo Moser, <i>On the distribution of Pythagorean triangles</i> .....	73
Roy Edwin Wild, <i>On the number of primitive Pythagorean triangles with area less than <math>n</math></i> .....	85
R. Sherman Lehman, <i>Approximation of improper integrals by sums over multiples of irrational numbers</i> .....	93
Emma Lehmer, <i>On the number of solutions of <math>u^k + D \equiv w^2 \pmod{p}</math></i> .....	103
Robert Delmer Stalley, <i>A modified Schnirelmann density</i> .....	119
Richard Allan Moore, <i>The behavior of solutions of a linear differential equation of second order</i> .....	125
William M. Whyburn, <i>A nonlinear boundary value problem for second order differential systems</i> .....	147