

Pacific Journal of Mathematics

**NOTE ON THE MULTIPLICATION FORMULAS FOR THE
JACOBI ELLIPTIC FUNCTIONS**

L. CARLITZ

NOTE ON THE MULTIPLICATION FORMULAS FOR THE JACOBI ELLIPTIC FUNCTIONS

L. CARLITZ

1. Introduction. For t an odd integer it is well known [4, vol. 2, p. 197] that

$$(1.1) \quad sn\,tx = \frac{sn\,x \cdot G_1^{(t)}(z)}{G_0^{(t)}(z)} \quad (z = sn^2x),$$

where

$$(1.2) \quad \begin{aligned} G_0^{(t)} &= 1 + a_{01}z + a_{02}z^2 + \cdots + a_{0t'}z^{t'}, \\ G_1^{(t)} &= t + a_{11}z + a_{12}z^2 + \cdots + a_{1t'}z^{t'} \end{aligned} \quad (t' = (t^2 - 1)/2),$$

and the a_{ij} are polynomials in $u = k^2$ with rational integral coefficients. If we define

$$\beta_m(t) = \beta_m(t, u)$$

by means of

$$(1.3) \quad \frac{sn\,tx}{t\,sn\,x} = \sum_{m=0}^{\infty} \beta_{2m}(t) \frac{x^{2m}}{(2m)!} \quad (\beta_{2m+1}(t) = 0),$$

it follows from (1.1) and (1.2) that $t\beta_{2m}(t)$ is a polynomial in u with integral coefficients for all m and all odd t . We shall show that

$$(1.4) \quad \beta_{2m}(t) = H_m(t) - \sum_{\substack{p-1 \mid 2m \\ p \mid t}} \frac{1}{p} A_p^{2m/(p-1)}(u),$$

where $H_m(t) = H_m(t, u)$ denotes a polynomial in u with integral coefficients,

Received August 8, 1953.

Pacific J. Math. 5 (1955), 169-176

the summation in the right member is over all (odd) primes p such that $(p-1) \mid 2m$ and $p \mid t$; finally $A_p(u)$ is defined [4, vol. 1, p. 399] by means of

$$(1.5) \quad sn x = sn(x, u) = \sum_{m=0}^{\infty} A_{2m+1}(u) \frac{x^{2m+1}}{(2m+1)!}.$$

so that $A_{2m+1}(u)$ is a polynomial in u with integral coefficients. We show also that

$$(1.6) \quad t \sum_{s=0}^r (-1)^{r-s} \binom{r}{s} \beta_{m+s(p-1)}(t) A_p^{r-s}(u) \equiv 0 \pmod{(p^m, p^r)},$$

where p is an arbitrary odd prime and $r \geq 1$; by (1.6) we understand that the left member is a polynomial in u every coefficient of which is divisible by the indicated power of p .

The proof of these formulas depends upon the results of [2]; for a theorem analogous to (1.4), see [1].

2. Proof of (1.4). Put

$$(2.1) \quad \frac{x}{sn x} = \sum_{m=0}^{\infty} \beta_{2m} \frac{x^{2m}}{(2m)!}.$$

Then β_{2m} is a polynomial in u with rational coefficients; indeed [2, Theorem 2],

$$(2.2) \quad p\beta_{2m} \equiv \begin{cases} -A_p^{2m/(p-1)}(u) & ((p-1) \mid 2m) \\ 0 & ((p-1) \nmid 2m). \end{cases} \pmod{p}$$

In the next place, if we write

$$\frac{sn tx}{t sn x} = \frac{sn tx}{tx} \frac{x}{sn x},$$

and make use of (1.3), (1.5), and (2.1), it follows that

$$(2.3) \quad \beta_{2m}(t) = \sum_{s=0}^m \binom{2m}{2s} \beta_{2m-2s} A_{2s+1}(u) \frac{t^{2s}}{2s+1}.$$

As already observed, $t\beta_{2m}(t)$ has integral coefficients; thus the denominator of $\beta_{2m}(t)$ is a divisor of t . Now let p denote a prime divisor of t , and assume $p^e \mid (2s + 1)$, $e \geq 1$. Then

$$2s + 1 \geq p^e \geq 3^e \geq e + 2, \quad 2s \geq e + 1.$$

Thus not only is $t^{2s}/(2s + 1)$ integral (mod p) but it is divisible by p . Since by (2.2) the denominator of β_{2m} contains p to at most the first power it therefore follows that the product

$$(2.4) \quad \beta_{2m-2s} t^{2s}/(2s + 1)$$

is integral (mod p) when $p \mid (2s + 1)$.

Suppose next that $p \nmid (2s + 1)$, where $s \geq 1$. It is again clear that (2.4) is integral (mod p) since p occurs in the denominator of β_{2m-2s} at most once while it occurs in t^{2s} at least twice. Thus as a matter of fact (2.4) is divisible by p in this case.

It remains to consider the term $s = 0$ in (2.3). Clearly we have proved that

$$(2.5) \quad p\beta_{2m}(t) \equiv p\beta_{2m} \pmod{p}.$$

Comparing (2.5) with (2.2) we may state:

THEOREM 1. *If t is an arbitrary odd integer then (1.4) holds.*

We remark that the residue of $A_p(u)$ is determined [2, § 6] by

$$(2.6) \quad \begin{aligned} A_p(u) &\equiv (-1)^{\frac{1}{2}(p-1)} F\left(\frac{1}{2}, \frac{1}{2}; 1; u\right) \\ &\equiv (-1)^{\frac{1}{2}(p-1)} \sum_{j=0}^{\frac{1}{2}(p-1)} \binom{\frac{1}{2}(p-1)}{j} u^j \pmod{p}. \end{aligned}$$

Here F denotes the hypergeometric function.

3. Some corollaries. By means of Theorem 1 a number of further results are readily obtained. By H_{2m} will be understood an unspecified polynomial in u with integral coefficients.

Since β_{2m} , as defined by (2.1), is integral (mod 2) we have first:

THEOREM 2. *If t is divisible by the denominator of β_{2m} , then*

$$(3.1) \quad \beta_{2m}(t) = H_{2m} + \beta_{2m}.$$

If t is prime to the denominator of β_{2m} , then $\beta_{2m}(t)$ has integral coefficients.

THEOREM 3. *If t_1, t_2 are relatively prime and odd, then*

$$(3.2) \quad \beta_{2m}(t_1 t_2) = H_{2m} + \beta_{2m}(t_1) + \beta_{2m}(t_2).$$

If t is a power of a prime we get:

THEOREM 4. *If p is an odd prime and $r \geq 1$ we have*

$$(3.3) \quad \beta_{2m}(p^r) = H_{2m} + \beta_{2m}(p).$$

Using (3.2) and (3.3) we get also:

THEOREM 5. *The following identity holds:*

$$(3.4) \quad \beta_{2m}(t) = H_{2m} + \sum_{p|t} \beta_{2m}(p),$$

where the summation is over all prime divisors of t .

We have also:

THEOREM 6. *If a is an arbitrary integer, then the product*

$$(3.5) \quad a(a^m - 1) \beta_{2m}(t)$$

has integral coefficients.

4. A related result. It follows from (1.1) and (1.2) that, for t odd,

$$(4.1) \quad sn \, tx = \sum_{r=0}^{\infty} C_{2r+1} sn^{2r+1} x,$$

where the C_{2r+1} are polynomials in u with integral coefficients. Clearly we have

$$(4.2) \quad \beta_{2m}(t) = \frac{1}{t} \sum_{r=0}^m A_{2m}^{(2r)} C_{2r+1},$$

where the $A_{2m}^{(2r)}$ are defined by

$$(4.3) \quad sn^{2r} x = \sum_{m=0}^{\infty} A_{2m}^{(2r)} \frac{x^{2m}}{(2m)!},$$

and like the C 's are polynomials with integral coefficients.

We shall now prove the following property of the C 's.

THEOREM 7. *For t odd we have*

$$(4.4) \quad (2m + 1)C_{2m+1} = 0 \pmod{t} \quad (m = 0, 1, 2, \dots),$$

where (4.4) indicates that every coefficient in $(2m + 1)C_{2m+1}$ is divisible by t .

Proof. Differentiating (4.1) with respect to x , we get

$$(4.5) \quad t \frac{cn \, tx \, dn \, tx}{cn \, x \, dn \, x} = \sum_{m=0}^{\infty} (2m + 1)C_{2m+1} sn^{2m} x.$$

Now we have, in addition to (1.1),

$$(4.6) \quad \frac{cn \, tx}{cn \, t} = \frac{G_2^{(t)}(z)}{G_0^{(t)}(z)}, \quad \frac{dn \, tx}{dn \, x} = \frac{G_3^{(t)}(z)}{G_0^{(t)}(z)} \quad (z = sn^2 x),$$

where G_2 and G_3 are polynomials in z of the same form as G_0 . By means of (1.1) and (4.6) it is evident that (4.5) implies

$$(4.7) \quad t \sum_{m=0}^{\infty} H_m^{(t)} z^m = \sum_{m=0}^{\infty} (2m + 1)C_{2m+1} z^m,$$

where the H_m are polynomials in u with integral coefficients. Clearly (4.4) is an immediate consequence of (4.7).

Kronecker [5, p. 439] has proved a similar result in connection with the transformation of prime order of $sn \, x$. For a result like Theorem 7 for the Weierstrass \wp -function, see [3].

Returning to (4.2) we recall [2, § 2] that

$$(4.8) \quad A_{2m}^{(2r)} \equiv 0 \pmod{(2r)!} \quad (m = 0, 1, 2, \dots).$$

We rewrite (4.2) in the form

$$(4.9) \quad \beta_{2m}(t) = \sum_{r=0}^m \frac{(2r)!}{2r+1} \frac{A_{2m}^{(2r)}}{(2r)!} \frac{(2r+1)C_{2r+1}}{t}.$$

By (4.4) and (4.8) the last two fractions in the right member of (4.9) have integral coefficients; also $(2r)!/(2r+1)$ is integral unless $2r+1$ is prime. Consequently (4.9) becomes

$$(4.10) \quad \beta_{2m}(t) = H_{2m} - \sum_{\substack{p-1 \mid 2m \\ p \mid t}} \frac{1}{p} A_{2m}^{(p-1)} \frac{{}^p C_p}{t}.$$

Comparing (4.10) with (1.4) we get:

THEOREM 8. *If the prime p divides t , then*

$$(4.11) \quad \frac{{}^p C_p}{t} \equiv 1 \pmod{p}.$$

Hence if $p^e \mid t, p^{e+1} \nmid t$ it follows that

$$(4.12) \quad C_p \equiv \frac{t}{p} \pmod{p^e}.$$

5. Proof of (1.6). Again using (5.1) we have

$$(5.1) \quad \frac{sn \, tx}{sn \, x} = \sum_{i=0}^{\infty} C_{2i+1} sn^{2i} x.$$

Now it is proved in [2, Theorem 4] that the coefficients $A_{2m}^{(2i)}$ defined by (4.3) satisfy

$$(5.2) \quad \sum_{s=0}^r (-1)^{r-s} \binom{r}{s} A_p^{(r-s)b/(p-1)} A_{2m+s}^{(2i)} \equiv 0 \pmod{(p^{2m}, p^{er})},$$

where $p^{e-1}(p-1) \mid b$. Hence using (1.3) and (5.1) we get:

THEOREM 9. *If $p^{e-1}(p-1) \mid b$, then*

$$(5.3) \quad t \sum_{s=0}^r (-1)^{r-s} \binom{r}{s} A_p^{(r-s)b/(p-1)} \beta_{2m+sb}(t) \equiv 0 \pmod{(p^{2m}, p^{er})},$$

For $b = p - 1$, (5.3) evidently reduces to (1.6).

It is of some interest to compare Theorem 9 with the results of [2, § 7].

If we take $r = 1$, (5.3) becomes

$$t \{ \beta_{2m+b}(t) - A_p^{b/(p-1)} \beta_{2m}(t) \} \equiv 0 \pmod{(p^{2m}, p^e)}.$$

If we put

$$\beta_{2m}(t) = \sum_i \beta_{2m,i} u^i$$

and recall that, by (2.6),

$$A_p(0) \equiv (-1)^{\frac{1}{2}(p-1)} \pmod{p}$$

we get exactly as in the proof of [2, Theorem 6].

THEOREM 10. *Let $p^{e-1}(p-1) \mid b$ and $p^{j-1} \leq i < p^j$. Then*

$$(5.4) \quad \beta_{2m+b,i} \equiv (-1)^{\frac{1}{2}b} \beta_{2m,i} \pmod{(p^{2m}, p^{e-j})}.$$

6. An elementary analogue of $\beta_{2m}(t)$. It may be of interest to say a word about the numbers $\phi_m(t)$ defined by

$$(6.1) \quad \frac{e^{tx} - 1}{t(e^x - 1)} = \sum_{m=0}^{\infty} \phi_m(t) \frac{x^m}{m!},$$

where t is now an arbitrary integer. Clearly (6.1) implies that

$$t\phi_m(t) = S_m(t) = \sum_{s=0}^{t-1} s^m.$$

By a theorem of Staudt (see for example [6, p. 143]),

$$(6.2) \quad \phi_m(t) = G + \sum_{p|t} \phi_m(p),$$

where G is an integer. Moreover,

$$(6.3) \quad p\phi_m(p) = \begin{cases} -1 & (p-1|m) \\ 0 & (p-1 \nmid m). \end{cases} \pmod{p}$$

It follows [6, p. 153] that

$$(6.4) \quad \phi_{2m}(t) = G - \sum_{\substack{p-1|2m \\ p|t}} \frac{1}{p}.$$

Thus Staudt's theorems (6.2) and (6.4) may be viewed as elementary analogues of (3.4) and (1.4).

Formulas like (6.2) and (6.4) hold also for the numbers $\psi_{2m}(t)$ occurring in

$$\frac{\sin tx}{t \sin x} = \sum_{m=0}^{\infty} \psi_{2m}(t) \frac{x^{2m}}{(2m)!}.$$

REFERENCES

1. L. Carlitz, *An analogue of the Bernoulli polynomials*, Duke Math. J. **8** (1941), 405-412.
2. ———, *Congruences connected with the power series expansions of the Jacobi elliptic functions*, Duke Math. J. **20** (1953), 1-12.
3. J. W. S. Cassels, *A note on the division values of $\wp(u)$* , Proc. Cambridge Philos. Soc. **45** (1949), 167-172.
4. R. Fricke, *Die elliptischen Funktionen und ihre Anwendungen*, Leipzig and Berlin, vol. 1, 1916, and vol. 2, 1922.
5. L. Kronecker, *Zur Theorie der elliptischen Funktionen*, Werke, **4** (1929), 345-495.
6. J. V. Uspensky and M. A. Heaslet, *Elementary number theory*, New York, 1939.

DUKE UNIVERSITY

PACIFIC JOURNAL OF MATHEMATICS

EDITORS

H. L. ROYDEN
Stanford University
Stanford, California

E. HEWITT
University of Washington
Seattle 5, Washington

R. P. DILWORTH
California Institute of Technology
Pasadena 4, California

A. HORN*
University of California
Los Angeles 24, California

ASSOCIATE EDITORS

H. BUSEMANN	P. R. HALMOS	R. D. JAMES	GEORGE PÓLYA
HERBERT FEDERER	HEINZ HOPF	BORGE JESSEN	J. J. STOKER
MARSHALL HALL	ALFRED HORN	PAUL LÉVY	KOSAKU YOSIDA

SPONSORS

UNIVERSITY OF BRITISH COLUMBIA
CALIFORNIA INSTITUTE OF TECHNOLOGY
UNIVERSITY OF CALIFORNIA, BERKELEY
UNIVERSITY OF CALIFORNIA, DAVIS
UNIVERSITY OF CALIFORNIA, LOS ANGELES
UNIVERSITY OF CALIFORNIA, SANTA BARBARA
MONTANA STATE UNIVERSITY
UNIVERSITY OF NEVADA
OREGON STATE COLLEGE
UNIVERSITY OF OREGON
UNIVERSITY OF SOUTHERN CALIFORNIA

STANFORD RESEARCH INSTITUTE
STANFORD UNIVERSITY
UNIVERSITY OF UTAH
WASHINGTON STATE COLLEGE
UNIVERSITY OF WASHINGTON

* * *

AMERICAN MATHEMATICAL SOCIETY
HUGHES AIRCRAFT COMPANY
SHELL DEVELOPMENT COMPANY

Mathematical papers intended for publication in the *Pacific Journal of Mathematics* should be typewritten (double spaced), and the author should keep a complete copy. Manuscripts may be sent to any of the editors. Manuscripts intended for the outgoing editors should be sent to their successors. All other communications to the editors should be addressed to the managing editor, Alfred Horn at the University of California, Los Angeles 24, California.

50 reprints of each article are furnished free of charge; additional copies may be obtained at cost in multiples of 50.

The *Pacific Journal of Mathematics* is published quarterly, in March, June, September, and December. The price per volume (4 numbers) is \$12.00; single issues, \$3.50. Back numbers are available. Special price to individual faculty members of supporting institutions and to individual members of the American Mathematical Society: \$4.00 per volume; single issues, \$1.25.

Subscriptions, orders for back numbers, and changes of address should be sent to Pacific Journal of Mathematics, c/o University of California Press, Berkeley 4, California.

Printed at Kokusai Bunken Insatsusha (International Academic Printing Co., Ltd.), No. 10, 1-chome, Fujimi-cho, Chiyoda-ku, Tokyo, Japan.

* During the absence of E. G. Straus.

PUBLISHED BY PACIFIC JOURNAL OF MATHEMATICS, A NON-PROFIT CORPORATION
COPYRIGHT 1955 BY PACIFIC JOURNAL OF MATHEMATICS

Pacific Journal of Mathematics

Vol. 5, No. 2

October, 1955

Leonard M. Blumenthal, <i>An extension of a theorem of Jordan and von Neumann</i>	161
L. Carlitz, <i>Note on the multiplication formulas for the Jacobi elliptic functions</i>	169
L. Carlitz, <i>The number of solutions of certain types of equations in a finite field</i>	177
George Bernard Dantzig, Alexander Orden and Philip Wolfe, <i>The generalized simplex method for minimizing a linear form under linear inequality restraints</i>	183
Arthur Pentland Dempster and Seymour Schuster, <i>Constructions for poles and polars in n-dimensions</i>	197
Franklin Haimo, <i>Power-type endomorphisms of some class 2 groups</i>	201
Lloyd Kenneth Jackson, <i>On generalized subharmonic functions</i>	215
Samuel Karlin, <i>On the renewal equation</i>	229
Frank R. Olson, <i>Some determinants involving Bernoulli and Euler numbers of higher order</i>	259
R. S. Phillips, <i>The adjoint semi-group</i>	269
Alfred Tarski, <i>A lattice-theoretical fixpoint theorem and its applications</i>	285
Anne C. Davis, <i>A characterization of complete lattices</i>	311