

Pacific Journal of Mathematics

**ON THE HOMOMORPHISMS OF AN ALGEBRA ONTO
FROBENIUS ALGEBRAS**

WILBUR EUGENE DESKINS

ON THE HOMOMORPHISMS OF AN ALGEBRA ONTO FROBENIUS ALGEBRAS

W. E. DESKINS

1. Introduction. A linear associative algebra possessing a nonsingular parastrophic matrix is known as a Frobenius algebra after the mathematician who first investigated the properties of such an algebra [3]. In more recent years the properties of this class of algebras have been studied in papers by a number of mathematicians, notably R. Brauer, C. C. MacDuffee, T. Nakayama, and C. Nesbitt (see References).

Since Frobenius algebras are defined in terms of the parastrophic matrices, a natural question to ask is the following: Does a parastrophic matrix of rank m of an algebra \mathcal{A} of order n determine in some manner a homomorphism of \mathcal{A} onto a Frobenius algebra of order m ? As the answer to this query is, in general, negative, it is the purpose of this paper to investigate the question: When does a parastrophic matrix of rank m determine in some manner a homomorphism of \mathcal{A} onto a Frobenius algebra of order m ? First a "manner of determination" is selected. Since the parastrophic matrices of \mathcal{A} form a double \mathcal{A} -module, various ideals of \mathcal{A} of annihilating elements correspond to each parastrophic matrix. These are studied and conditions are developed (Theorem 9) which insure the determination from these annihilators an ideal \mathcal{B} such that the difference algebra $\mathcal{A} - \mathcal{B}$ is a Frobenius algebra of order m . These requirements are shown to be necessary, also, in the sense that any homomorphism of \mathcal{A} onto a Frobenius algebra of order m implies the existence of a parastrophic matrix Q of rank m which satisfies these conditions. Furthermore, the kernel of the homomorphism will be the ideal \mathcal{B} determined from among those elements which annihilate Q as an element of a double \mathcal{A} -module.

Basic terminology is introduced in § 2, parastrophic modules are defined, and the order of such a module is discussed. In § 3 one-sided ideals determined by the parastrophic matrices are considered, while § 4 is devoted to a study of two-sided ideals determined by certain parastrophic matrices and of the homomorphisms of an algebra onto Frobenius algebras. Certain of the ideals

Received February 10, 1954. Presented to the American Mathematical Society May 1, 1954.

Pacific J. Math. 5 (1955), 501-511

introduced in § 4 have radical-like properties, and these ideals are considered in § 5. A supplementary remark on the order of the radical of a Frobenius algebra is given in § 6.

The author wishes to express his gratitude to Professor C. C. MacDuffee for his counsel during the preparation of this paper.

2. Preliminary remarks. Let \mathcal{A} be a linear associative algebra of order n over the field \mathfrak{F} , and let e_1, \dots, e_n be an \mathfrak{F} -basis for \mathcal{A} . Multiplication in \mathcal{A} follows from the multiplication of the basis elements,

$$e_i e_j = \sum_k c_{ijk} e_k \quad i, j = 1, \dots, n,$$

where the c_{ijk} are elements of \mathfrak{F} , the constants of multiplication.

The associativity condition, written in terms of these constants of multiplication, is equivalent to each of the following sets of n^2 matrix equations:

$$(1) \quad Q_j R_i = \sum_k c_{ikj} Q_k,$$

$$(2) \quad S_i Q_j = \sum_k c_{kij} Q_k \quad i, j = 1, \dots, n,$$

where the matrices R_i , S_i , and Q_i are defined as (c_{isr}) , (c_{ris}) and (c_{sri}) , respectively, where r denotes the row and s the column index.

Let $a \in \mathcal{A}$; then

$$a = a_1 e_1 + \dots + a_n e_n,$$

where the a_i are field elements. Let

$$R(a) = a_1 R_1 + \dots + a_n R_n,$$

$$S(a) = a_1 S_1 + \dots + a_n S_n,$$

$$Q(a) = a_1 Q_1 + \dots + a_n Q_n.$$

$R(a)$ is called the *first matrix*, $S(a)$ the *second matrix*, and $Q(a)$ the *parastrophic matrix*, of a . (Note that $Q(a)$ as defined here is the transpose of the parastrophic matrix as defined by other authors). The set $\mathfrak{R}(\mathcal{A})$ of all the first (second) matrices of \mathcal{A} form an algebra which is a homomorphic image of \mathcal{A} . The

set \mathcal{Q} of all the parastrophic matrices of \mathcal{A} does not in general have this property, but if the following definitions are made,

$$e_i * Q_j = S_i Q_j, \quad Q_j * e_i = Q_j R_i \qquad i, j = 1, \dots, n,$$

then \mathcal{Q} is a double \mathcal{A} -module, the *parastrophic module* of \mathcal{A} .

If a change of basis is made for \mathcal{A} , the elements of \mathcal{R} and \mathcal{S} undergo similar transformations, while the elements of \mathcal{Q} undergo congruency transformations [8]. Hence rank and symmetry are invariant set properties of \mathcal{Q} .

MacDuffee has obtained [7] necessary and sufficient conditions that \mathcal{R} and \mathcal{S} be algebras isomorphic with \mathcal{A} . A corresponding result for \mathcal{Q} is given by

THEOREM 1. *\mathcal{Q} is of order m (as an \mathcal{A} -module) if and only if the following conditions are satisfied:*

(i) \mathcal{A} contains an ideal \mathcal{W} of order m such that the difference algebra $\mathcal{A} - \mathcal{W}$ is a zero algebra.

(ii) \mathcal{A} contains *no* ideal of lower order with this property.

The proof of this theorem is a standard reversible procedure involving a change of basis for \mathcal{A} . Let there be $n - m$ linearly independent linear relations among the Q_i ; then there exist $n - m$ linearly independent row vectors

$$T_i = (t_{i1}, \dots, t_{in}) \qquad i = m + 1, \dots, n,$$

such that

$$\sum_k t_{ik} Q_k = 0 \qquad i > m.$$

If B is a nonsingular n by n matrix with the T_i as its last $n - m$ rows, and if u_1, \dots, u_n form a new basis for \mathcal{A} ,

$$u_i = \sum_k p_{ki} e_k \qquad i = 1, \dots, n,$$

where $P = (p_{rs}) = B^{-1}$, then the Q_i are transformed into

$$\begin{aligned} Q'_i &= P^T \left(\sum_k t_{ik} Q_k \right) P \\ &= 0 \qquad i > m. \end{aligned}$$

Thus if the new constants of multiplication are c'_{ijk} , then

$$c'_{ijk} = 0 \qquad i, j = 1, \dots, n; \quad k > m,$$

and u_1, \dots, u_m span an ideal \mathfrak{W} of order m such that $\mathfrak{A} - \mathfrak{W}$ is a zero algebra. The process is clearly reversible.

COROLLARY. *If \mathfrak{A} has either a left or right identity element, then \mathfrak{Z} is of order n .*

\mathfrak{A} is said to be a *Frobenius algebra* if \mathfrak{Z} contains a nonsingular element.

THEOREM 2. *\mathfrak{A} is a Frobenius algebra if and only if \mathfrak{Z} is a cyclic module of order n .*

If \mathfrak{Z} contains a nonsingular element Q , then (1) and (2) imply that

$$\mathfrak{A} * Q = Q * \mathfrak{A} = \mathfrak{Z},$$

and \mathfrak{Z} is of order n since a Frobenius algebra possesses an identity element. Conversely, if \mathfrak{Z} is generated by an element Q and is of order n , then (1), (2), and Theorem 1 imply that Q is nonsingular.

3. Ideals of \mathfrak{A} . Let \mathfrak{B} be a right ideal of \mathfrak{A} of order $n - m$. If a basis is selected for \mathfrak{A} such that the last $n - m$ elements of the basis span \mathfrak{B} , then the m matrices Q_1, \dots, Q_m have all zeros in their last $n - m$ columns. The task of determining a right ideal by a process involving reduction of certain elements of \mathfrak{Z} through changes of bases of \mathfrak{A} seems formidable, if possible. However, a somewhat similar process is given by the following theorem.

THEOREM 3. *A parastrophic matrix Q of rank m determines a right ideal \mathfrak{B} of order greater than or equal to $n - m$.*

Let \mathfrak{B} be the set of all elements $b \in \mathfrak{A}$ such that $Q * b = 0$. Clearly \mathfrak{B} is a right ideal. That its order is at least $n - m$ will follow from the next theorem.

That \mathfrak{B} may actually be of order greater than $n - m$ is proved by the following example. Let \mathfrak{A} have basis elements e_1 and e_2 , $e_1^2 = e_1 e_2 = e_2 e_1 = 0$, $e_2^2 = e_1$. Then the corresponding Q_1 has rank 1 but $\mathfrak{B} = \mathfrak{A}$.

A more desirable result is contained in the following.

THEOREM 4. *A parastrophic matrix of rank m determines a right ideal of order $n - m$.*

Let Q be a parastrophic matrix of rank m . Then there exists a nonsingular n by n matrix P such that the elements of the last $n - m$ columns of QP are all zeros. Let P effect a change of basis for A ; that is, if $P = (p_{rs})$, let

$$u_i = \sum_k p_{ki} e_k \qquad i = 1, \dots, n$$

be a new basis for \mathcal{A} . If $Q = Q(a)$, then $Q'(a)$, with respect to the new basis, is $P^T Q P$ and hence has nothing but zeros in its last $n - m$ columns.

Now assume Q is of this form. Then

$$Q = \sum_i a_i Q_i = \left(\sum_i a_i c_{sri} \right)$$

so that

$$(3) \qquad \sum_i a_i c_{jki} = 0 \qquad j > m, k = 1, \dots, n.$$

From (1) and (3) it follows that

$$(4) \qquad QR_i = \sum_j a_j \left(\sum_k c_{ikj} Q_k \right) = \sum_k \left(\sum_j a_j c_{ikj} \right) Q_k = 0 \qquad i > m.$$

Hence $Q * e_i = 0$ for $i > m$, and $\mathcal{B} = (e_{m+1}, \dots, e_n)$ is the right ideal determined by Q .

A right ideal of \mathcal{A} which may be determined in this way will be called a *parastrophic right ideal*.

THEOREM 5. *A sufficient condition that the ideal of Theorem 3 be a parastrophic right ideal is that \mathcal{L} be of order n .*

Suppose $\mathcal{B} = (e_{m+1}, \dots, e_n)$ is determined from Q as above, and consider $e_i, i \leq m$. If $Q * e_i = 0$, then if \mathcal{L} is of order n , (4) implies

$$\sum_j a_j c_{ikj} = 0 \qquad k = 1, \dots, n,$$

which is impossible since Q is assumed to be of rank m .

Let $Q = \sum a_i Q_i$ be in the reduced form described above.

THEOREM 6. *If \mathcal{A} has a right identity element, then $a_i = 0$ for $i > m$.*

Since \mathcal{A} has a right identity element there are field elements f_i such that

$$\sum_i f_i S_i = I, \quad \sum_i f_i c_{kij} = \delta_{kj}.$$

Then (3) implies

$$\sum_k f_k \left(\sum_i a_i c_{jki} \right) = 0 = \sum_i a_i \left(\sum_k f_k c_{jki} \right) = \sum_i a_i \delta_{ji} = a_j \quad j > m.$$

The results of this section are obviously valid if the word "right" is replaced by "left".

Since the existence of ideals in an algebra \mathcal{A} has been shown to be equivalent to the existence of singular elements in \mathcal{Q} , the following theorem is immediate.

THEOREM 7. *\mathcal{A} is a division algebra if and only if \mathcal{Q} contains no singular elements.*

4. Homomorphisms of \mathcal{A} . The following result is an immediate consequence of Theorem 4 and its analogue for left ideals.

THEOREM 8. *If Q is congruent to a matrix of the form*

$$\begin{pmatrix} T & 0 \\ 0 & 0 \end{pmatrix},$$

where T is a nonsingular m by m matrix, then the right parastrophic ideal \mathfrak{B} is also a left parastrophic ideal. Conversely, if \mathfrak{B} is a right parastrophic ideal determined by Q , then if \mathfrak{B} is also a two-sided ideal, Q satisfies the above condition.

Such an ideal will be called a *parastrophic ideal*, and Q will be said to have *P-rank* m . While *P-rank* is not defined for every matrix, it is a property of every symmetric matrix. Thus, if the characteristic of \mathfrak{B} is greater than n , the radical of \mathcal{A} is a parastrophic ideal. (It will be apparent shortly that this is true regardless of the field characteristic since a semisimple algebra is a Frobenius algebra.)

It does not follow that a matrix of \mathcal{Q} of *P-rank* m determines a homomorphism

of \mathcal{A} onto a Frobenius algebra of order m , for any commutative nilpotent non-zero algebra contains proper parastrophic ideals. The following indicates a necessary criterion.

LEMMA 1. *If π is a homomorphism of \mathcal{A} onto \mathcal{C} , an algebra with an identity element $\mathbf{1}$, then \mathcal{A} contains an idempotent element e such that $\pi e = \mathbf{1}$. Furthermore, the set of left annihilators of e is contained in the kernel of the homomorphism.*

This follows simply from the structure theory for algebras.

Now suppose the last $n - m$ basis elements of \mathcal{A} form a parastrophic ideal \mathcal{B} , and suppose that \mathcal{A} has an idempotent element u such that $\mathcal{A}u \cup \mathcal{B} = \mathcal{A}$. Then \mathcal{B} will be called a *regular parastrophic ideal*.

THEOREM 9. *A homomorphism of \mathcal{A} onto a Frobenius algebra of order m has as its kernel a regular parastrophic ideal of order $n - m$, and conversely if \mathcal{B} is a regular parastrophic ideal of \mathcal{A} of order $n - m$, then $\mathcal{A} - \mathcal{B}$ is a Frobenius algebra of order m .*

Suppose $\bar{\mathcal{A}}$ is a Frobenius image of \mathcal{A} , with basis $\bar{e}_1, \dots, \bar{e}_m$ and kernel \mathcal{B} spanned by e_{m+1}, \dots, e_n .

Then $\bar{\mathcal{A}}$ possesses a nonsingular m by m parastrophic matrix \bar{Q} ,

$$\bar{Q} = \sum_i^m a_i \bar{Q}_i,$$

and

$$Q = \sum_i^m a_i Q_i$$

is an element of \mathcal{Q} with P -rank m . Hence \mathcal{B} is a parastrophic ideal. By Lemma 1, \mathcal{B} is a regular parastrophic ideal.

The converse follows from the regularity of \mathcal{B} and Theorem 6.

Thus, if Q is a parastrophic matrix of rank m , if Q can be reduced to a corner matrix by a change of basis of \mathcal{A} , and if Q is associated with a linear combination of the first m of the new basis elements, then Q determines a homomorphism of \mathcal{A} onto a Frobenius algebra. Furthermore, each homomorphism of \mathcal{A} onto a Frobenius algebra may be determined in this fashion.

5. Radical-like ideals. A function f of \mathcal{A} into the set of all ideals of \mathcal{A} is called a *radical function* of \mathcal{A} if the contraction of f to the difference algebra $\mathcal{C} = \mathcal{A} - f(\mathcal{A})$ maps \mathcal{C} onto the zero ideal. The ideal $f(\mathcal{A})$ is called a *radical-like ideal* of \mathcal{A} .

Let \mathcal{P} be the set of all regular parastrophic ideals of \mathcal{A} and let \mathcal{N} be an element of \mathcal{P} of minimal order, with the agreement that \mathcal{N} is the zero ideal if \mathcal{A} is a Frobenius algebra and \mathcal{A} if \mathcal{A} is nilpotent. Then define $f(\mathcal{A}) = \mathcal{N}$.

THEOREM 10. $f(\mathcal{A}) = \mathcal{N}$ is a nilpotent ideal of \mathcal{A} .

If \mathcal{A} is nilpotent the theorem is trivially true, so assume that \mathcal{A} has the radical $\mathcal{K} \neq \mathcal{A}$. Suppose \mathcal{K} is of order r and that \mathcal{N} is of order m . Let

$$\mathcal{N} \cap \mathcal{K} = \mathcal{C}.$$

Case 1. $\mathcal{C} = (0)$. Let \mathcal{A} have a basis such that the first m basis elements span \mathcal{N} while the last r span \mathcal{K} . By the definition of \mathcal{N} there is an element Q' of \mathcal{Z} of rank $n - m$ with its first m rows and columns composed of only zeros. Now \mathcal{N} is isomorphic to a semisimple subalgebra of $\mathcal{A} - \mathcal{K}$, so there is an element Q'' of \mathcal{Z} of rank m with only zeros in its last $n - m$ rows and columns. Then $Q' + Q''$ is nonsingular.

Case 2. $\mathcal{C} \neq (0)$. Then $\mathcal{A} - \mathcal{C}$ is a Frobenius algebra by the above work.

In either case

$$\mathcal{N} \cap \mathcal{K} = \mathcal{N}$$

so that \mathcal{N} is contained in \mathcal{K} and so is nilpotent.

One important property which \mathcal{N} may lack is uniqueness. The question of whether \mathcal{N} is unique up to an \mathcal{A} -isomorphism will now be considered and partially answered.

The following result indicates a significance of the \mathcal{A} -isomorphism of two minimal elements of \mathcal{P} .

THEOREM 11. Let \mathcal{N} and \mathcal{M} be minimal elements of \mathcal{P} . Then a necessary condition that \mathcal{N} and \mathcal{M} be \mathcal{A} -isomorphic is that $\mathcal{N} - \mathcal{C}$ and $\mathcal{M} - \mathcal{C}$ be zero algebras, where $\mathcal{C} = \mathcal{N} \cap \mathcal{M}$.

It may be assumed that $\mathcal{C} = (0)$. Then let σ be an \mathcal{A} -isomorphism from \mathcal{N} onto \mathcal{M} . If a and b are elements of \mathcal{M} , then \mathcal{N} contains an element b' such that

$$a b = a(\sigma b') = \sigma(a b') = 0.$$

The isomorphism of the minimal elements of \mathfrak{P} for certain algebras will stem from the following lemma.

LEMMA 2. *If U and V are n by n matrices of rank m with elements from a field \mathfrak{F} which contains at least $m + 1$ nonzero elements, then \mathfrak{F} contains an element $t \neq 0$ such that $U + tV$ is of rank at least m .*

It will be sufficient to prove the result for $m = n$. Let D and E be nonsingular n by n matrices such that

$$DVE = I.$$

Consider the equation

$$\det(D(U - xV)E) = \det(DUE - xI) = 0,$$

which is of degree n in the indeterminate x . Since \mathfrak{F} contains at least $n + 1$ nonzero elements, one of them does not satisfy this equation.

Let U and V be n by n matrices, and let

$$U \wedge V = 0$$

mean that the two matrices do not both have nonzero elements in the same row-column position.

THEOREM 12. *If two minimal elements \mathfrak{N}_1 and \mathfrak{N}_2 of \mathfrak{P} are determined by symmetric matrices Q_1 and Q_2 of \mathfrak{Q} of rank m , if \mathfrak{F} contains at least $m + 1$ nonzero elements, and if*

$$Q_1 \wedge Q_2 = 0,$$

then $\mathfrak{A} - \mathfrak{N}_1$ is isomorphic with $\mathfrak{A} - \mathfrak{N}_2$.

The cyclic modules

$$\mathfrak{A} * Q_1 = Q_1 * \mathfrak{A}, \quad \mathfrak{A} * Q_2 = Q_2 * \mathfrak{A}$$

are of order m , and the representations of \mathfrak{A} over these double \mathfrak{A} -modules yield Frobenius algebras of order m which are images of \mathfrak{A} , isomorphic with $\mathfrak{A} - \mathfrak{N}_1$ and $\mathfrak{A} - \mathfrak{N}_2$ respectively. Let t be a nonzero element of \mathfrak{F} such that $Q_1 + tQ_2$ is of rank m (since higher rank would contradict the minimality of the order of \mathfrak{N}_1

and \mathcal{N}_2). Since Q_1 and Q_2 are symmetric

$$\mathcal{A} * (Q_1 + tQ_2) = (Q_1 + tQ_2) * \mathcal{A}$$

is a cyclic module of order m , and since $Q_1 \wedge Q_2 = 0$, (1) and (2) imply that the mapping

$$(Q_1 + tQ_2)R_i \longrightarrow Q_1 R_i \qquad i = 1, \dots, n$$

is an \mathcal{A} -isomorphism between $Q_1 * \mathcal{A}$ and $(Q_1 + tQ_2) * \mathcal{A}$. Similarly $Q_2 * \mathcal{A}$ and $(Q_1 + tQ_2) * \mathcal{A}$ are \mathcal{A} -isomorphic. Hence $Q_1 * \mathcal{A}$ and $Q_2 * \mathcal{A}$ are \mathcal{A} -isomorphic which implies that $\mathcal{A} - \mathcal{N}_1$ and $\mathcal{A} - \mathcal{N}_2$ are isomorphic.

6. A remark concerning Frobenius algebras. While Frobenius algebras are generally regarded as algebras with radicals of sufficiently small order, the following indicates that their radicals must also be of sufficiently large order.

THEOREM 13. *Let \mathcal{A} be a Frobenius algebra bound to its radical \mathcal{K} . Then if $\mathcal{A} - \mathcal{K}$ is of order m , \mathcal{K} is of order at least m . If \mathcal{K} is a zero algebra, then \mathcal{K} is of order m .*

By the results of Nakayama [9] the set of all elements of \mathcal{A} which annihilate \mathcal{K} from the right is an ideal \mathcal{L} which also annihilates \mathcal{K} from the left and has order $n - k = m$, where k is the order of \mathcal{K} . Since \mathcal{A} is bound [4] to \mathcal{K} ,

$$\mathcal{L} \subseteq \mathcal{K},$$

hence $m \leq k$, and $m = k$ if $\mathcal{L} = \mathcal{K}$.

The consideration of bound algebras is, of course, sufficient since an algebra may be written as a direct sum of a semisimple algebra and a bound algebra. (This result is due to M. Hall [4]).

REFERENCES

1. R. Brauer, *On hypercomplex arithmetic and a theorem of Speiser*, A Speiser Festschrift, Zurich, 1945.
2. R. Brauer, and C. Nesbitt, *On the regular representations of algebras*, Proc. Nat. Acad. Sci., **23** (1937), 236-240.
3. G. Frobenius, *Theorie der hyperkomplexen Groessen*, S.-B. Preuss. Akad. Wiss., Berlin (1903), 504-537.
4. M. Hall, *The position of the radical in an algebra*, Trans. Amer. Math. Soc., **48** (1940), 391-404.

5. M. Ikeda, and T. Nakayama, *Supplementary remarks on Frobeniusean algebras*, Osaka Math. J., **2** (1950), 7-12.
6. C.C. MacDuffee, *Modules and ideals in a Frobenius algebra*, Monatsh. Math. Phys., **48** (1939), 292-313.
7. ———, *On the independence of the first and second matrices of an algebra*, Bull. Amer. Math. Soc., **35** (1929), 344-349.
8. ———, *The theory of Matrices*, Chelsea, New York, 1946.
9. T. Nakayama, *On Frobeniusean algebras*, I, Ann. Math., **40** (1939), 611-633.
10. ———, *On Frobeniusean algebras*, II, Ann. Math., **42** (1941), 1-21.
11. ———, *On Frobeniusean algebras*, III, Jap. J. Math., **18** (1942),
12. ———, *Orthogonality relations for Frobenius and quasi-Frobenius algebras*, Proc. Amer. Math. Soc., **3** (1952), 183-195.
13. C. Nesbitt, *On the regular representations of algebras*, Ann. Math. **39** (1938), 634-658.
14. M. Osima, *Some studies on Frobenius algebras*, Jap. J. Math., **21** (1951), 179-190.
15. G. Simura, *On a certain ideal of the center of a Frobeniusean algebra*, Sci. Papers Coll. Gen. Ed. Univ. Tokyo, **2** (1952), 117-124.

THE OHIO STATE UNIVERSITY

PACIFIC JOURNAL OF MATHEMATICS

EDITORS

H.L. ROYDEN
Stanford University
Stanford, California

E. HEWITT
University of Washington
Seattle 5, Washington

R. P. DILWORTH
California Institute of Technology
Pasadena 4, California

* Alfred Horn
University of California
Los Angeles 24, California

ASSOCIATE EDITORS

H. BUSEMANN	P.R. HALMOS	R.D. JAMES	GEORGE PÓLYA
HERBERT FEDERER	HEINZ HOPF	BØRGE JESSEN	J.J. STØKER
MARSHALL HALL	ALFRED HORN	PAUL LEVÝ	KOSAKU YOSIDA

SPONSORS

UNIVERSITY OF BRITISH COLUMBIA
CALIFORNIA INSTITUTE OF TECHNOLOGY
UNIVERSITY OF CALIFORNIA, BERKELEY
UNIVERSITY OF CALIFORNIA, DAVIS
UNIVERSITY OF CALIFORNIA, LOS ANGELES
UNIVERSITY OF CALIFORNIA, SANTA BARBARA
MONTANA STATE UNIVERSITY
UNIVERSITY OF NEVADA
OREGON STATE COLLEGE
UNIVERSITY OF OREGON

UNIVERSITY OF SOUTHERN CALIFORNIA
STANFORD UNIVERSITY
UNIVERSITY OF UTAH
WASHINGTON STATE COLLEGE
UNIVERSITY OF WASHINGTON
* * *
AMERICAN MATHEMATICAL SOCIETY
HUGHES AIRCRAFT COMPANY
SHELL DEVELOPMENT COMPANY

Mathematical papers intended for publication in the *Pacific Journal of Mathematics* should be typewritten (double spaced), and the author should keep a complete copy. Manuscripts may be sent to any of the editors. Manuscripts intended for the outgoing editors should be sent to their successors. All other communications to the editors should be addressed to the managing editor, Alfred Horn, at the University of California Los Angeles 24, California.

50 reprints of each article are furnished free of charge; additional copies may be obtained at cost in multiples of 50.

The *Pacific Journal of Mathematics* is published quarterly, in March, June, September, and December. The price per volume (4 numbers) is \$12.00; single issues, \$3.50; back numbers (Volumes 1, 2, 3) are available at \$2.50 per copy. Special price to individual faculty members of supporting institutions and to individual members of the American Mathematical Society: \$4.00 per volume; single issues, \$1.25.

Subscriptions, orders for back numbers, and changes of address should be sent to the publishers, University of California Press, Berkeley 4, California.

Printed at Ann Arbor, Michigan. Entered as second class matter at the Post Office, Berkeley, California.

* During the absence of E.C. Straus.

UNIVERSITY OF CALIFORNIA PRESS • BERKELEY AND LOS ANGELES

COPYRIGHT 1955 BY PACIFIC JOURNAL OF MATHEMATICS

Richard Horace Battin, <i>Note on the "Evaluation of an integral occurring in servomechanism theory"</i>	481
Frank Herbert Brownell, III, <i>An extension of Weyl's asymptotic law for eigenvalues</i>	483
Wilbur Eugene Deskins, <i>On the homomorphisms of an algebra onto Frobenius algebras</i>	501
James Michael Gardner Fell, <i>The measure ring for a cube of arbitrary dimension</i>	513
Harley M. Flanders, <i>The norm function of an algebraic field extension. II</i>	519
Dieter Gaier, <i>On the change of index for summable series</i>	529
Marshall Hall and Lowell J. Paige, <i>Complete mappings of finite groups</i>	541
Moses Richardson, <i>Relativization and extension of solutions of irreflexive relations</i>	551
Peter Scherk, <i>An inequality for sets of integers</i>	585
W. R. Scott, <i>On infinite groups</i>	589
A. Seidenberg, <i>On homogeneous linear differential equations with arbitrary constant coefficients</i>	599
Victor Lenard Shapiro, <i>Cantor-type uniqueness of multiple trigonometric integrals</i>	607
Leonard Tornheim, <i>Minimal basis and inessential discriminant divisors for a cubic field</i>	623
Helmut Wielandt, <i>On eigenvalues of sums of normal matrices</i>	633