

Pacific Journal of Mathematics

**AN ALGEBRAIC CHARACTERIZATION OF FIXED IDEALS IN
CERTAIN FUNCTION RING**

LYLE E. PURSELL

AN ALGEBRAIC CHARACTERIZATION OF FIXED IDEALS IN CERTAIN FUNCTION RINGS

LYLE E. PURSELL

1. Introduction. In this paper an algebraic characterization of the fixed ideals in a certain class of function rings is given (an ideal in a function ring is fixed if there is a point at which all functions in the ideal vanish). This class of function rings includes the rings of all real-, complex-, or quaternion-valued continuous functions on a normal Hausdorff space whose points are G -delta sets and the ring of r -fold differentiable functions on an r -differentiable manifold whose coordinate covering is neighborhood finite. For these rings of functions we construct the underlying space from the fixed ideals in the same way that Gelfand and Kolmogoroff [3] have constructed a compact space from the non-unit ideals in its ring of all real-valued continuous functions.

We also show the existence of certain homomorphisms from the automorphism groups of these function rings into the group of homeomorphisms of the underlying space onto itself. In §5 we find that an isomorphism between the rings of all r -differentiable functions on two r -differentiable manifolds can be extended to an isomorphism between the rings of all continuous functions on these manifolds and that the homeomorphism determined by this isomorphism is differentiable.

2. The general case.

(2.1) *By \mathcal{R} we mean a ring of functions from a regular Hausdorff space X to a division ring D having the following properties:*

- P_1 . If f is in \mathcal{R} , then the set of zeros of f , which we denote by $Z(f)$, is closed.*
- P_2 . If x is not in a closed set F , then there is a function f in \mathcal{R} such that $Z(f)$ contains a neighborhood of F but does not contain x .*
- P_3 . If f in \mathcal{R} does not vanish at any point of a closed set F , then there is a function g in \mathcal{R} such that fg (and also gf) has the value 1 at every point of F .*
- P_4 . For each x in X there is a function f_x in \mathcal{R} which vanishes at x and only at x .*

Received June 14, 1954. This paper includes extensions of certain results given in the author's doctoral thesis which was written at Purdue University under the direction of Prof. M. E. Shanks (1952). The author wishes to express his appreciation to Prof. Shanks, Prof. M. Henriksen, and Dr. D. W. Dubois, who made several contributions to this paper.

(2.2) By the "support of a function f " in \mathcal{R} , which we denote by $\text{Sp}(f)$, we mean the set $\text{Cl}(X-Z(f))$. From the properties of closure we have:

- (i) $\text{Sp}(f) = X - \text{Int } Z(f)$,
- (ii) $\text{Int } \text{Sp}(f) = X - \text{Cl } \text{Int } Z(f)$,
- (iii) $\text{Sp}(f) = \text{Cl } \text{Int } \text{Sp}(f)$.

By the "annihilator of a function f " in \mathcal{R} , which we denote by $A(f)$, we mean the set of all g in \mathcal{R} such that $fg=0$ (and hence $gf=0$). For any ring of functions with values in a division ring the annihilator of an element is a two-sided ideal. In addition we have

$$A(f) = \{g \in \mathcal{R} \mid Z(g) \supset \text{Sp}(f)\},$$

and $A(f) = \mathcal{R}$ if and only if $f=0$.

(2.3) LEMMA. If f and g are in \mathcal{R} and $g \neq 0$, then $Z(f)$ and $\text{Sp}(g)$ are disjoint if and only if $f - A(g)$ has an inverse in the residue class ring $\mathcal{R} - A(g)$.

Proof. Since $g \neq 0$, then $Z(g) \neq X$ and $\text{Sp}(g)$ is not empty. If f does not vanish at any point of $\text{Sp}(g)$, then there is a function h in \mathcal{R} such that fh and hf have the value 1 at every point of $\text{Sp}(g)$, that is, $fh \equiv 1 \pmod{A(g)}$ and $hf \equiv 1 \pmod{A(g)}$. Hence $f - A(g)$ has an inverse in $\mathcal{R} - A(g)$. If $f - A(g)$ has an inverse in $\mathcal{R} - A(g)$, then there is a function h in \mathcal{R} such that $(fh-1)$ is in $A(g)$, that is $(fh-1)$ vanishes at every point of $\text{Sp}(g)$. Hence f does not vanish at any point of $\text{Sp}(g)$.

(2.4) If f is in \mathcal{R} , let $H(f)$ be the set of all nonzero g in \mathcal{R} such that $f - A(g)$ has an inverse in the ring $\mathcal{R} - A(g)$. An ideal I in \mathcal{R} is "bounded" if there is a function f in \mathcal{R} without an inverse such that $H(f)$ contains $H(g)$ for every g in I . We say that " I is bounded by f ". An ideal which is maximal in the set of all bounded ideals is called a "maximal bounded ideal". The set of all maximal bounded ideals is denoted by $M[\mathcal{R}]$. We observe that an ideal contained in a bounded ideal is bounded.

(2.5) LEMMA. For f and g in \mathcal{R} , $Z(g)$ contains $Z(f)$ if and only if $H(f)$ contains $H(g)$.

Proof. From (2.3) and (2.4) $H(f)$ is the set of all functions h in \mathcal{R} such that $Z(f)$ and $\text{Sp}(h)$ are disjoint. Hence if $Z(f)$ is a subset of $Z(g)$, then $H(f)$ contains $H(g)$. Suppose there is a point x in $Z(f)$

but not in $Z(g)$, then by P_2 there is a function h in \mathcal{R} which is different from zero on a neighborhood of x but vanishes on a neighborhood of $Z(g)$. For this function h , $Z(f)$ meets $\text{Sp}(h)$ but $Z(g)$ does not. Hence if $Z(f)$ is not a subset of $Z(g)$, then $H(f)$ does not contain $H(g)$.

(2.6) THEOREM. *If \mathcal{R} is a ring of functions from a regular Hausdorff space X to a division ring D which satisfies P_1, P_2, P_3 , and P_4 of (2.1), then an ideal I in \mathcal{R} is a fixed ideal if and only if it is a bounded ideal.*

Proof. If I is a fixed ideal, then there is a point x at which all elements of I vanish. From P_4 there is a function f_x in \mathcal{R} which vanishes at x and only at x . For every g in I , $Z(g)$ contains $Z(f_x)$, that is $H(f_x)$ contains $H(g)$. Since f_x has no inverse, I is bounded. If I is bounded by a function f in \mathcal{R} without an inverse, then $Z(f)$ is a subset of $Z(g)$ for every g in I . Since $Z(f)$ is not empty, I is fixed.

(2.7) *For x in X , $I(x)$ means the fixed ideal $\{f \in \mathcal{R} \mid f(x)=0\}$. From (2.6) an ideal is a maximal bounded ideal if and only if it is of this form.*

(2.8) *Let A be a subset of $M[\mathcal{R}]$. If we define*

$$J \in \text{Cl}(A) \text{ if and only if } J \supset \bigcap_{I \in A} I$$

for A nonempty and $\text{Cl}(A)=A$ for A empty, then $M[\mathcal{R}]$ is said to have the "Stone topology". We denote the set $M[\mathcal{R}]$ with the Stone topology by X^ .*

(2.9) THEOREM. *If \mathcal{R} is a ring of functions from a regular Hausdorff space X to a division ring D satisfying P_1, P_2, P_3 , and P_4 of (2.1), then X is homeomorphic to X^* .*

Proof. From (2.7) the mapping $x \rightarrow I(x)$ is a one-to-one mapping of X onto X^* . Let $a \in \text{Cl}(A)$, $A \subset X$, and let A^* be the image of A under the mapping $x \rightarrow I(x)$, then every function in \mathcal{R} vanishing on A (that is, every function in $\bigcap_{x \in A} I(x) = \bigcap_{I \in A^*} I$) also vanishes at a (that is, is in $I(a)$) and $I(a)$ is in $\text{Cl}(A^*)$. If, however, a is not in $\text{Cl}(A)$, then there is a function f in \mathcal{R} vanishing on A but not at a . Then f is in $\bigcap_{I \in A^*} I$ but not in $I(a)$, and $I(a)$ is not in $\text{Cl}(A^*)$. Hence the correspondence $x \rightarrow I(x)$ is a homeomorphism of X onto X^* .

(2.10) COROLLARY. *If the rings \mathcal{R} and \mathcal{R}' of functions from the regular Hausdorff spaces X and X' to the division rings D and D' ,*

respectively, satisfy P_1 , P_2 , P_3 , and P_4 of (2.1) and are isomorphic, then the spaces X and X' are homeomorphic.

(2.11) Let $i: \mathcal{R} \rightarrow \mathcal{R}'$ be the isomorphism referred to in the preceding paragraph. With the point x in X we associate the x' (which exists uniquely) in X' which is the common zero of all functions in the isomorphic image of the ideal consisting of all functions in \mathcal{R} which vanish at x , that is,

$$x' = \bigcap_{f \in I(x)} Z(f).$$

One can show that the correspondence $x \rightarrow x'$ is a homeomorphism from X onto X' . We will denote this homeomorphism by $\phi(i)$ and refer to it as "the homeomorphism from X onto X' corresponding to (or determined by) the isomorphism i from \mathcal{R} onto \mathcal{R}' ".

(2.12) By $\mathcal{A}(\mathcal{R})$ we mean the automorphism group of \mathcal{R} . By $\mathcal{H}(X)$ we mean the homeomorphism group of X , that is the group of all homeomorphisms of X onto itself. If i_1 and i_2 are in $\mathcal{A}(\mathcal{R})$, then it follows from (2.11) that $\phi(i_1 i_2) = \phi(i_1) \phi(i_2)$. Hence we have the theorem of the following paragraph.

(2.13) THEOREM. The mapping $\phi: \mathcal{A}(\mathcal{R}) \rightarrow \mathcal{H}(X)$ is a homomorphism from the automorphism group of \mathcal{R} into the homeomorphism group of X .

(2.14) For x in X we denote the set of values $\{f(x) | f \in \mathcal{R}\}$ by $V(x)$. From P_2 and P_3 the set $V(x)$ is a subdivision ring of D . The correspondence $f \rightarrow f(x)$ is a homomorphism from \mathcal{R} onto $V(x)$ with kernel $I(x)$, hence the correspondence $f - I(x) \rightarrow f(x)$ is an isomorphism from the residue class ring $\mathcal{R} - I(x)$ onto $V(x)$. Since $\mathcal{R} - I(x)$ is, therefore, a division ring, $I(x)$ is a maximal ideal, that is, every maximal bounded ideal is a maximal ideal.

(2.15) LEMMA. If $f \rightarrow f'$ is an isomorphism from \mathcal{R} onto \mathcal{R}' and $x \rightarrow x'$ is the corresponding homeomorphism, then the correspondence $f(x) \rightarrow f'(x')$ is an isomorphism from $V(x)$ onto $V'(x')$.

Proof. Since $I'(x')$ is the isomorphic image of $I(x)$, the correspondence $f - I(x) \rightarrow f' - I'(x')$ is an isomorphism. Since $f(x) \rightarrow f - I(x)$ and $f' - I'(x') \rightarrow f'(x')$ are isomorphisms, $f(x) \rightarrow f'(x')$ is an isomorphism from $V(x)$ onto $V'(x')$.

3. Rings of continuous functions.

(3.1) Čech [2] has shown that a subset of a normal Hausdorff

space is the zero set of some real-valued continuous function if and only if it is a closed G_δ set. Using his result and Urysohn's lemma concerning real-valued continuous functions on a normal space, one may show that the rings— $C(X, R)$ of all real-valued continuous functions on X , $C(X, K)$ of all complex-valued continuous functions on X , and $C(X, Q)$ of all quaternion-valued continuous functions on X —satisfy P_1 , P_2 , P_3 , and P_4 of (2.1) if X is a normal Hausdorff space all of whose points are G_δ sets. Hence we have the following.

(3.2) THEOREM. *Let X and X' be normal Hausdorff spaces all of whose points are G_δ sets and let F denote either the real field, the complex field, or the quaternion ring. If $C(X, F)$ and $C(X', F)$ are isomorphic, then X and X' are homeomorphic.*

(3.3) According to results obtained by Gelfand and Kolmogoroff [3], Hewitt [6], and Gillman, Henriksen, and Jerison [4], Theorem (3.2) holds for completely regular spaces satisfying the first axiom of countability. There are, however, normal spaces all of whose points are G_δ sets which do not satisfy the first axiom of countability (cf. Bing [1, p. 180, Example C]).

(3.4) For the rings $C(X, F)$ it can be established that the homomorphism $\phi : \mathcal{N}(C(X, F)) \rightarrow \mathcal{H}(X)$ of (2.11) and (2.13) is a homomorphism onto $\mathcal{H}(X)$.

4. Rings of real-valued functions.

(4.1) If \mathcal{R} is a ring of real-valued functions on X satisfying P_1 , P_2 , P_3 , and P_4 , then for each x the set of values $V(x) = \{f(x) | f \in \mathcal{R}\}$ is a subfield of the real field R . We now introduce an additional property for the ring \mathcal{R} :

P_5 . *For each x in X the set of values $V(x)$ is a subfield of the real field R which has only one isomorphism into R , the identity isomorphism.*

Property P_5 holds if $V(x) = R$; hence $C(X, R)$ satisfies P_5 . There are rings of real-valued functions satisfying P_1 , P_2 , P_3 , P_4 , and P_5 which contain discontinuous functions as is shown by the example of the following paragraph given to the author by D. W. Dubois.

(4.2) EXAMPLE. Let X be the closed interval $[0, 1]$, α be a finite subdivision $\{0 = x_0, x_1, \dots, x_{n-1}, x_n = 1\}$ of X , and A be the set of all α . Let $\theta(x) = \exp(x)$ for $x \neq 0$ and $\theta(0) = 0$. Let $B(\alpha)$ be the set of all real-valued functions f on X such that

$$f(x) = \frac{p_i(x, \theta(x))}{q_i(x, \theta(x))}, \quad x_{i-1} \leq x \leq x_i, \quad i=1, 2, \dots, n,$$

where $p_i(x, \theta)$ and $q_i(x, \theta)$ are polynomials in x and $\theta(x)$ such that $f(x)$ is continuous at x_1, x_2, \dots, x_{n-2} , and x_{n-1} and $q_i(x, \theta(x))$ does not vanish for $x_{i-1} \leq x \leq x_i$ for any i . If $\mathcal{R} = \bigcup_{\alpha \in I} B(\alpha)$, then \mathcal{R} is a ring of real-valued functions which satisfy P_1, P_2, P_3, P_4 , and P_5 but some of which are discontinuous.

(4.3) Theorem (4.4) and (4.5) may be established by using P_5 and the results of § 2.

(4.4) THEOREM. *If \mathcal{R} and \mathcal{R}' are isomorphic rings of real valued functions on regular Hausdorff spaces X and X' satisfying P_1, P_2, P_3, P_4 , and P_5 , i is the isomorphism from \mathcal{R} onto \mathcal{R}' , and h is the corresponding homeomorphism from X onto X' , then:*

- (i) $f(x) = (if)(h(x))$ for all f in R and x in X . Hence f is bounded above (below) if and only if (if) is bounded above (below); $\text{lub } f = \text{lub } f'$, $\text{glb } f = \text{glb } f'$; and the subrings of all bounded functions in R and R' are isomorphic.
- (ii) There is an isomorphism i^* from $C(X, R)$ onto $C(X', R)$ such that $i(f) = i^*(f)$ for all f in $C(X, R) \cap \mathcal{R}$.

(4.5) THEOREM. *If \mathcal{R} is a ring of real-valued functions on a regular Hausdorff space satisfying P_1, P_2, P_3, P_4 , and P_5 , then the homomorphism ϕ of (2.11) and (2.13) is an isomorphism of $\mathcal{N}(R)$ into $\mathcal{H}(X)$.*

From (3.4) and (4.5) we have the following.

(4.6) THEOREM. *The groups $\mathcal{N}(C(X, R))$ and $\mathcal{H}(X)$ are isomorphic.*

5. Rings of continuously differentiable functions.

(5.1) If $C^r(M)$ is the ring of r -fold continuously differentiable functions on an r -differentiable manifold M with a neighborhood-finite covering of coordinate neighborhoods (r may be either a positive integer or the symbol ∞), then $C^r(M)$ satisfies P_1, P_2, P_3, P_4 , and P_5 . The theorem of the following paragraph may be obtained.

(5.2) THEOREM. *If $C^r(M)$ and $C^r(M')$ are isomorphic, then M and M' are homeomorphic. The homeomorphism h determined by the isomorphism is differentiable (that is, $f(h)$ is in $C^r(M)$ if f is in $C^r(M')$) and the isomorphism can be extended to an isomorphism from $C(M, R)$ onto $C(M', R)$.*

6. Additional remarks. Since the above was written the author has observed that P_4 may be replaced by the weaker hypothesis:

P_4^* . For each x in X there is a pair of functions g and h in \mathcal{R} such that $x = Z(g) - Z(h)$.

If \mathcal{R} satisfies P_1, P_2, P_3 , and P_4^* , one can show that an ideal I is fixed if and only if there is a pair of functions g and h in \mathcal{R} such that $H(g)$ does not contain $H(h)$ but $H(gh)$ does contain $H(fh)$ for every f in the ideal I . (Lemma (2.5) holds as before.) The results of (2.9) through (2.15) may then be established if X^* is defined to be the set of maximal fixed ideals with the Stone topology.

If X is a completely regular, locally-compact space all of whose points are G_δ sets, then the rings $C_0(X, R)$, $C_0(X, K)$, and $C_0(X, Q)$ of all real-, complex-, or quaternion-valued continuous functions with compact supports satisfy P_1, P_2, P_3 , and P_4^* . Hence it follows that they determine X . (This result for $C_0(X, R)$ has already been established by Shanks [7] without assuming that points are G_δ sets). One can also show that the automorphism group $\mathcal{A}(C_0(X, R))$ is isomorphic to $\mathcal{A}(C(X, R))$ and $\mathcal{H}(X)$.

REFERENCES

1. R. H. Bing, *Metrization of topological spaces*, Canadian J. Math., **3** (1951), 175-186.
2. E. Čech, *On bicomact spaces*, Ann. Math., **38** (1937), 823-844.
3. I. Gelfand and A. N. Kolmogoroff, *On rings of continuous functions on topological spaces*, C. R. (Doklady) Acad. Sci. URSS, **22** (1939), 11-15.
4. L. Gillman, M. Henriksen, and M. Jerison, *A theorem of Gelfand and Kolmogoroff concerning maximal ideals in rings of continuous functions*, Proc. Amer. Math. Soc., **5** (1954), 447-455.
5. M. Hall, Jr., *A type of algebraic closure*, Ann. Math., **40** (1939), 360-369.
6. E. Hewitt, *Rings of real-valued continuous functions I*, Trans. Amer. Math. Soc., **64** (1948), 45-99.
7. M. E. Shanks, *Rings of functions on locally compact spaces*, (abstract no. 365), Bull. Amer. Math. Soc., **57** (1951), 295.
8. N. Steenrod, *The topology of fibre bundles*, Princeton University Press, 1951.
9. M. H. Stone, *Applications of the theory of Boolean rings to general topology*, Trans. Amer. Math. Soc., **41** (1937), 375-481.

GRINNELL COLLEGE

../../../../FrontMatter/paper.pdf

Pacific Journal of Mathematics

Vol. 5, No. 6

, 1955

Nesmith Cornett Ankeny and Theodore Joseph Rivlin, <i>On a theorem of S. Bernstei</i>	849
Louis Auslander, <i>The use of forms in variational calculation</i>	853
Paul Civin, <i>Abstract Riemann sum</i>	861
Paul Civin, <i>Some ergodic theorems involving two operator</i>	869
Eckford Cohen, <i>The number of solutions of certain cubic congruence</i>	877
Richard M. Cohn, <i>Specializations over difference field</i>	887
Jean Dieudonné, <i>Pseudo-discriminant and Dickson invariant</i>	907
Ky Fan, <i>A comparison theorem for eigenvalues of normal matrice</i>	911
Richard P. Gosselin, <i>On the convergence behaviour of trigonometric interpolating polynomial</i>	915
Peter K. Henrici, <i>On generating functions of the Jacobi polynomial</i>	923
Meyer Jerison, <i>An algebra associated with a compact grou</i>	933
Wilhelm Magnus, <i>Infinite determinants associated with Hill's equatio</i>	941
G. Power and D. L. Scott-Hutton, <i>The slow steady motion of liquid past a semi-elliptical bos</i>	953
Lyle E. Pursell, <i>An algebraic characterization of fixed ideals in certain function ring</i>	963
C. T. Rajagopal, <i>Additional note on some Tauberian theorems of O. Szás</i>	971
Louis Baker Rall, <i>Error bounds for iterative solutions of Fredholm integral equation</i>	977
Shigeo Sasaki and Kentaro Yano, <i>Pseudo-analytic vectors on pseudo-Kählerian manifold</i>	987
Eugene Schenkman, <i>On the tower theorem for finite group</i>	995
P. Stein and John E. L. Peck, <i>On the numerical solution of Poisson's equation over a rectangl</i>	999
Morgan Ward, <i>The mappings of the positive integers into themselves which preserve divisio</i>	1013
Seth Warner, <i>Weak locally multiplicatively-convex algebra</i>	1025
Louis Weisner, <i>Group-theoretic origin of certain generating function</i>	1033