

Pacific Journal of Mathematics

NOTE ON NORMAL NUMBERS

CALVIN T. LONG

NOTE ON NORMAL NUMBERS

CALVIN T. LONG

Introduction. Let α be a real number with fractional part $.a_1a_2a_3 \dots$ when written to base r . Let Y_n denote the block of the first n digits in this representation and let $N(d, Y_n)$ denote the number of occurrences of the digit d in Y_n . The number α is said to be *simply normal* to base r if

$$\lim_{n \rightarrow \infty} \frac{N(d, Y_n)}{n} = \frac{1}{r}$$

for each of the r distinct choices of d . α is said to be *normal* to base r if each of the numbers $\alpha, r\alpha, r^2\alpha, \dots$ are simply normal to each of the bases r, r^2, r^3, \dots . These definitions, due to Emile Borel [1], were introduced in 1909. In 1940 S. S. Pillai [3] showed that a necessary and sufficient condition that α be normal to base r is that it be simply normal to each of the bases r, r^2, r^3, \dots , thus considerably reducing the number of conditions needed to imply normality. The purpose of the present note is to show that α is normal to base r if and only if there exists a set of positive integers $m_1 < m_2 < m_3 < \dots$ such that α is simply normal to base r^{m_i} for each $i \geq 1$, and also to show that no finite set of m 's will suffice.

Notation. We make use of the following additional conventions.

If B_k is any block of k digits to base r , $N(B_k, Y_n)$ will denote the number of occurrences of B_k in Y_n and $N_i(B_k, Y_n)$ will denote the number of occurrences of B_k starting in positions congruent to i modulo k in Y_n .

The term "relative frequency" will denote the asymptotic frequency with which an event occurs. For example, B_k occurs in (α) , the fractional part of α , with relative frequency r^{-k} if $\lim_{n \rightarrow \infty} N(B_k, Y_n)/n = r^{-k}$.

Proof of the theorems. The following lemmas are easily proved.

LEMMA 1. *If $\lim_{n \rightarrow \infty} \sum_{i=1}^m f_i(n) = 1$ and if $\liminf_{n \rightarrow \infty} f_i(n) \geq 1/m$ for $i = 1, 2, \dots, m$; then $\lim_{n \rightarrow \infty} f_i(n) = 1/m$ for each i .*

LEMMA 2. *The real number α is simply normal to base r^k if and*

Received July 5, 1956. Results in this paper were included in a doctoral dissertation written under the direction of Professor Ivan Niven at the University of Oregon. 1955.

only if $\lim_{n \rightarrow \infty} N_1(B_k, Y_n)/n = 1/kr^k$ for every block B_k of k digits to base r .

THEOREM 1. *The real number α is normal to base r if and only if there exist positive integers $m_1 < m_2 < m_3 < \dots$ such that α is simply normal to each of the bases $r^{m_1}, r^{m_2}, r^{m_3}, \dots$.*

Proof. The necessity of the condition follows immediately from the definition of normality.

Now suppose the condition of the theorem prevails. Let ν be an arbitrary positive integer and let B_ν be an arbitrary block of ν digits to base r . Choose k so large that $m_k > \nu$. It follows from Lemma 2 that

$$\lim_{n \rightarrow \infty} \frac{N_1(A_{m_k}, Y_n)}{n} = \frac{1}{m_k r^{m_k}}$$

for each block A_{m_k} of m_k digits to base r . If B_ν occurs exactly $t = t(A_{m_k})$ times in each A_{m_k} , then it follows that

$$\liminf_{n \rightarrow \infty} \frac{N(B_\nu, Y_n)}{n} \geq \frac{T}{m_k r^{m_k}}$$

where $T = \sum t(A_{m_k})$ with the sum running over all blocks of m_k digits to base r . Now there are $r^{m_k - \nu}$ distinct blocks A_{m_k} which contain B_ν starting in position i for $i = 1, 2, \dots, m_k - \nu + 1$ so that $T = (m_k - \nu + 1)r^{m_k - \nu}$. Thus it follows that

$$\liminf_{n \rightarrow \infty} \frac{N(B_\nu, Y_n)}{n} \geq \frac{(m_k - \nu + 1)r^{m_k - \nu}}{m_k r^{m_k}} = \frac{1}{r^\nu} \frac{\nu - 1}{m_k r^\nu}.$$

But, since this argument can be made for arbitrarily large values of k and $m_k \geq k$, this implies that

$$\liminf_{n \rightarrow \infty} \frac{N(B_\nu, Y_n)}{n} \geq \frac{1}{r^\nu}.$$

With Lemma 1 this implies that

$$\lim_{n \rightarrow \infty} \frac{N(B_\nu, Y_n)}{n} = \frac{1}{r^\nu}$$

so that α is normal to base r by a result of Niven and Zuckerman [2].

The next theorem implies that no finite set of m 's will suffice in Theorem 1.

THEOREM 2. *If m_1, m_2, \dots, m_s is an arbitrary collection of distinct*

positive integers, then there exists at least one real number α simply normal to each of the bases $r^{m_1}, r^{m_2}, \dots, r^{m_s}$ but not normal to base r .

Proof. Writing to base r^m form the periodic decimal

$$\alpha = .\dot{0}12\dots(r^m - 1)$$

where m is the least common multiple of m_1, m_2, \dots, m_s . It is clear that α is simply normal to base r^m and that it is not normal to base r . To show that it is simply normal to base r^{m_i} for $i=1, 2, \dots, s$ we prove that if d divides m then α is simply normal to base r^d .

Let $m=qd$ and let B_d be an arbitrary but fixed block of d digits to base r . In view of Lemma 2 it suffices to show that

$$\lim_{n \rightarrow \infty} \frac{N_i(B_d, Y_n)}{n} = \frac{1}{dr^d}.$$

A simple counting process shows that there are precisely $\binom{q}{i}(r^d - 1)^{q-i}$ distinct blocks A_m of m digits to base r which contain B_d exactly i times starting in a position congruent to one modulo d . Therefore, since

$$\lim_{n \rightarrow \infty} \frac{N_i(A_m, Y_n)}{n} = \frac{1}{mr^m}$$

for each A_m , it follows that

$$\lim_{n \rightarrow \infty} \frac{N_i(B_d, Y_n)}{n} = \frac{1}{mr^m} \sum_{i=1}^q i \binom{q}{i} (r^d - 1)^{q-i} = \frac{1}{dr^d}$$

as required.

REFERENCES

1. Émile Borel, *Les probabilités dénombrables et leurs applications arithmétiques*, Rend. Circ. Mat. Palermo **27** (1909), 247-271.
2. Ivan Niven and H. S. Zuckerman, *On the definition of normal numbers*, Pacific J. Math., **1** (1951), 103-109.
3. S. S. Pillai, *On normal numbers*, Proc. Indian Acad. Sci., Sect. A, **12** (1940), 179-184.

PACIFIC JOURNAL OF MATHEMATICS

EDITORS

H. L. ROYDEN

Stanford University
Stanford, California

R. A. BEAUMONT

University of Washington
Seattle 5, Washington

A. L. WHITEMAN

University of Southern California
Los Angeles 7, California

E. G. STRAUS

University of California
Los Angeles 24, California

ASSOCIATE EDITORS

E. F. BECKENBACH

C. E. BURGESS

M. HALL

E. HEWITT

A. HORN

V. GANAPATHY IYER

R. D. JAMES

M. S. KNEBELMAN

L. NACHBIN

I. NIVEN

T. G. OSTROM

M. M. SCHIFFER

G. SZEKERES

F. WOLF

K. YOSIDA

SUPPORTING INSTITUTIONS

UNIVERSITY OF BRITISH COLUMBIA

CALIFORNIA INSTITUTE OF TECHNOLOGY

UNIVERSITY OF CALIFORNIA

MONTANA STATE UNIVERSITY

UNIVERSITY OF NEVADA

OREGON STATE COLLEGE

UNIVERSITY OF OREGON

UNIVERSITY OF SOUTHERN CALIFORNIA

STANFORD UNIVERSITY

UNIVERSITY OF UTAH

WASHINGTON STATE COLLEGE

UNIVERSITY OF WASHINGTON

* * *

AMERICAN MATHEMATICAL SOCIETY

CALIFORNIA RESEARCH CORPORATION

HUGHES AIRCRAFT COMPANY

THE RAMO-WOOLDRIDGE CORPORATION

Mathematical papers intended for publication in the *Pacific Journal of Mathematics* should be typewritten (double spaced), and the author should keep a complete copy. Manuscripts may be sent to any of the editors. All other communications to the editors should be addressed to the managing editor, E. G. Straus at the University of California, Los Angeles 24, California.

50 reprints per author of each article are furnished free of charge; additional copies may be obtained at cost in multiples of 50.

The *Pacific Journal of Mathematics* is published quarterly, in March, June, September, and December. The price per volume (4 numbers) is \$12.00; single issues, \$3.50. Back numbers are available. Special price to individual faculty members of supporting institutions and to individual members of the American Mathematical Society: \$4.00 per volume; single issues, \$1.25.

Subscriptions, orders for back numbers, and changes of address should be sent to Pacific Journal of Mathematics, 2120 Oxford Street, Berkeley 4, California.

Printed at Kokusai Bunken Insatsusha (International Academic Printing Co., Ltd.), No. 10, 1-chome, Fujimi-cho, Chiyoda-ku, Tokyo, Japan.

PUBLISHED BY PACIFIC JOURNAL OF MATHEMATICS, A NON-PROFIT CORPORATION

The Supporting Institutions listed above contribute to the cost of publication of this Journal, but they are not owners or publishers and have no responsibility for its content or policies.

William F. Donoghue, Jr., <i>The lattice of invariant subspaces of a completely continuous quasi-nilpotent transformation</i>	1031
Michael (Mihály) Fekete and J. L. Walsh, <i>Asymptotic behavior of restricted extremal polynomials and of their zeros</i>	1037
Shaul Foguel, <i>Biorthogonal systems in Banach spaces</i>	1065
David Gale, <i>A theorem on flows in networks</i>	1073
Ioan M. James, <i>On spaces with a multiplication</i>	1083
Richard Vincent Kadison and Isadore Manual Singer, <i>Three test problems in operator theory</i>	1101
Maurice Kennedy, <i>A convergence theorem for a certain class of Markoff processes</i>	1107
G. Kurepa, <i>On a new reciprocity, distribution and duality law</i>	1125
Richard Kenneth Lashof, <i>Lie algebras of locally compact groups</i>	1145
Calvin T. Long, <i>Note on normal numbers</i>	1163
M. Mikolás, <i>On certain sums generating the Dedekind sums and their reciprocity laws</i>	1167
Barrett O'Neill, <i>Induced homology homomorphisms for set-valued maps</i>	1179
Mary Ellen Rudin, <i>A topological characterization of sets of real numbers</i>	1185
M. Schiffer, <i>The Fredholm eigen values of plane domains</i>	1187
F. A. Valentine, <i>A three point convexity property</i>	1227
Alexander Doniphan Wallace, <i>The center of a compact lattice is totally disconnected</i>	1237
Alexander Doniphan Wallace, <i>Two theorems on topological lattices</i>	1239
G. T. Whyburn, <i>Dimension and non-density preservation of mappings</i>	1243
John Hunter Williamson, <i>On the functional representation of certain algebraic systems</i>	1251