

Pacific Journal of Mathematics

**DIMENSION AND NON-DENSITY PRESERVATION OF
MAPPINGS**

G. T. WHYBURN

DIMENSION AND NON-DENSITY PRESERVATION OF MAPPINGS

G. T. WHYBURN

1. Introduction. In this paper consideration is given to conditions under which the property of being non-dense in a space in the sense of containing no open set in that space is invariant under certain types of mappings. In some spaces and for some mapping types the issue involved is essentially equivalent to the question of dimensionality preservation. These questions are of interest and importance in numerous mathematical fields. They are especially so in the study of topological aspects of the theory of functions and it is toward this connection that the results and methods in this note will be largely directed.

A single valued continuous transformation $f(X)=Y$ will be called a *mapping*. Such a mapping is *open* if open sets in X have open images in Y and is *light* provided $f^{-1}(y)$ is totally disconnected for each $y \in Y$. Also f has *scattered point inverses* provided that for each $y \in Y$, $f^{-1}(y)$ is a *scattered set* in the sense that no point of $f^{-1}(y)$ is a limit point of $f^{-1}(y)$.

As indicated above, a set K in a space X is *non-dense in X* provided K contains no open set in X . On the other hand that K is *dense in X* means that every point of X is either a point or a limit point of K . A mapping $f(X)=Y$ is said to *preserve non-density* for compact sets provided that $f(K)$ is non-dense in Y whenever K is compact and non-dense in X . For a mapping $f(X)=Y$, a subset X_0 of X is said to be *semi-dense in X* provided X_0 is dense in some open subset of every open set U in X whose image $f(U)$ is also open in Y . Thus the property of semi-density is a property of a subset of X *relative to a mapping f on X* and not an intrinsic property of X_0 alone.

For a mapping $f(X)=Y$, the set of all $x \in X$ such that x is a component of $f^{-1}f(x)$ will be designated by the symbol D_f . Also the symbol L_f will be used for the set of all $x \in X$ such that $f^{-1}f(x)$ is totally disconnected. Thus L_f is the maximum inverse set in X on which the mapping f is light, where by an *inverse set I* we mean a set which is the inverse of its transform under f , that is, one satisfying the relation

$$I=f^{-1}f(I).$$

Accordingly L_f may be thought of as the *lightness kernel* or *0-dimensional kernel* of the mapping f . Obviously we have $L_f \subset D_f$.

2. General setting. We begin with a theorem which was suggested by the theorem of Alexandroff's [1] on invariance of dimension under countable-fold open mappings. Our proof closely parallels that of Alexandroff for his theorem.

(2.1) **THEOREM.** *Let $f(A)=B$ be open and have scattered point inverses, where A and B are locally compact separable and metric. Then A is the union $A=\sum A_n$ of a sequence of compact sets such that $f|A_n$ is topological for each n .*

Proof. Let (U_n) be a countable basis of open sets in A , so chosen that \bar{U}_n is compact for each n . For each n , let F_n be the set of all $x \in U_n$ such that $g_n^{-1}g_n(x)=x$ where g_n denotes the mapping $f|U_n$. Then F_n is closed in U_n by openness of f . Accordingly each F_n is the union of a countable sequence of compact sets and thus we can write $\sum F_n = \sum A_n$ where each A_n is compact and lies in some F_m . Thus $f|A_n$ is topological for each n . Finally, $\sum A_n = A$, because if $x \in A$, there exists an m such that $x \in U_m$ and $U_m \cdot f^{-1}f(x) = x$ and hence so that $x \in F_m \subset \sum F_n = \sum A_n$.

(2.11) **COROLLARY.** *For any closed set K in A we have*

$$\dim f(K) = \dim K .$$

(2.12) **COROLLARY.** *If K is any closed set in A and V is any open subset of $f(K)$, then V contains an open subset U which is homeomorphic with a subset of K .*

For let K_n denote the set $K \cdot A_n$ for each n . Then since $V \subset \sum f(K_n)$ and each $f(K_n)$ is compact, some $f(K_n)$ contains an open subset U of V . Then $K_n \cdot f^{-1}(U)$ maps topologically onto U under f .

(2.13) **COROLLARY.** *If A and B are 2-manifolds, (or n -manifolds), then if K is non-dense in A , $f(K)$ is non-dense in B .*

(2.2) **THEOREM.** *Let A and B be locally compact separable metric spaces and let $f(A)=B$ be a mapping preserving compact non-dense sets. Then for some $y \in B$, $f^{-1}(y)$ is totally disconnected.*

Proof. For each integer $n > 0$ and each $x \in A$, let U_x^n be an open set of diameter $< 1/n$ containing x and having a compact boundary F_x^n . Let $U_{x_1}^n, U_{x_2}^n, \dots$ be a countable collection of these sets U_x^n whose union covers A and set

$$F^n = \sum_i F_{x_i}^n, \quad F = \sum_n F^n = \sum_n \sum_i F_{x_i}^n .$$

Now $f(F) \neq B$. For if $f(F) = B$, then for some n and i the set $f(F_{x_i}^n)$ must contain an open set in B , as B is locally compact; and this is impossible by hypothesis because $F_{x_i}^n$ is compact and non-dense for each n and i . Accordingly there exists a $y \in B - f(F)$. Clearly $f^{-1}(y)$ is totally disconnected, because if it had a non-degenerate component C_y , then for $1/n <$ the diameter of C_y , we would have $C_y \cdot F_{x_i}^n \neq \emptyset$ where $C_y \cdot U_{x_i}^n \neq \emptyset$.

(2.21) COROLLARY. *Under the same hypothesis, the set Y of all $y \in B$ with $f^{-1}(y)$ totally disconnected is dense in B and $L_f = f^{-1}(Y)$ is semi-dense in A .*

For if B_0 is any open set in B , we have only to set $A_0 = f^{-1}(B_0)$ and apply the theorem to the mapping $f|_{A_0}$ to obtain the first conclusion that Y is dense in B . To prove the second conclusion suppose on the contrary that for an open set U which has an open image and which we first suppose conditionally compact, $L_f \cdot U$ is dense in no open subset of U . Then $\overline{L_f \cdot U}$ is compact and non-dense, whereas $f(\overline{L_f \cdot U})$ must contain $f(U)$ since Y is dense in $f(U)$ by openness of $f(U)$. This is a contradiction.

Finally, to see that U need not be conditionally compact, we need only show that any open set V in A with an open image contains a conditionally compact open subset U with an open image. To do this, set $V = \sum V_n$ where each V_n is open and conditionally compact and $\overline{V_n} \subset U$. Then since $f(V) = \sum f(\overline{V_n})$, some $f(\overline{V_n})$ contains an open set G . Since $f[\text{Fr}(V_n)]$ is non-dense, $Q = G - f[\text{Fr}(V_n)]$ is open and non-empty. Then $U = V_n \cdot f^{-1}(Q)$ meets our condition.

(2.3) THEOREM. *Let A and B be locally compact separable metric spaces with $\dim A = k < \infty$ and let $f(A) = B$ be a mapping such that the image of every compact non-dense set K with $\dim K < k$ is non-dense. Then the set Y of all $y \in B$ with $f^{-1}(y)$ totally disconnected is dense in B .*

For, in the preceding proofs the sets F_x^n could now be taken of dimension $\leq k - 1$.

3. Quasi-open mappings. Region on a sphere. A mapping $f(X) = Y$ is quasi-open provided that if $y \in Y$ and K is a compact component of $f^{-1}(y)$, then for any open set U in X containing K , y is interior to $f(U)$ rel. Y , and is strongly quasi-open provided y is interior to $f(U)$ relative to a larger space $Y_0 \supset Y$. A mapping $f(X) = Y$ is monotone provided $f^{-1}(y)$ is a continuum (compact and connected set) for each $y \in Y$; and f is compact provided $f^{-1}(K)$ is compact for every compact

set $K \subset Y$ or, equivalently, provided f is closed and has compact point inverses. For compact mappings, quasi-openness is equivalent to quasi-monotoneity as defined originally by Wallace [3].

(3.1) THEOREM. *Let $f(X)=Y$ be a compact and quasi-open mapping where X is a region on a sphere S , Y is a metric space and where no component of a point inverse separates X . In order that the image of every compact 1-dimensional set in X be of dimension ≤ 1 it is necessary and sufficient that the set D_f be semi-dense in X .*

Proof. Let $f=lm$, $m(X)=X'$, $l(X')=Y$ be the monotone-light factorization of f . Let the mapping m be extended to the whole sphere S by decomposing S into the sets $m^{-1}(x')$, $x' \in X'$ together with the components of $S-X$ so that we obtain a monotone mapping $\phi(S)=S'$ of S onto a sphere S' containing X' (ϕ is the natural mapping of the decomposition) which is identical with m on X . That S' is a topological sphere follows from the readily verified facts that the described decomposition of S is upper semi-continuous and no element of this decomposition separates S , together with the classical theorem of R. L. Moore [2] that the hyperspace of any such decomposition of a sphere into continua is itself a topological sphere. Then $l(X')=Y$ is a light open mapping which is compact; and since X' is a region on S' , Y is a 2-manifold by the invariance of the 2-manifold property under such mappings [4].

Now to prove the sufficiency of the condition let K be a compact 1-dimensional set in X . Then $\dim m(K) \leq 1$. For, if not, then $m(K)$ contains an open set U in X' . Then $l(U)$ is open in Y and thus $m^{-1}(U)$ is an open set in X whose image under f is open in Y . Accordingly D_f is dense in an open subset Q of $m^{-1}(U)$. Since Q cannot lie wholly in K , $Q-Q \cdot K$ contains a point x of D_f . But then since $x=m^{-1}m(x)$, $m(x)$ cannot lie in $m(K)$, contrary to the supposition that $m(x) \in U \subset m(K)$. Thus $\dim m(K) \leq 1$.

It remains to show that $\dim lm(K) \leq 1$. Since l is compact, open and light and X' is a 2-manifold, l is finite to one [4]. Hence by (2.11) we have $\dim lm(K) = \dim m(K) \leq 1$.

To prove the necessity of the condition we note first that it follows from our hypothesis that f preserves non-density for compact sets. For if K is a compact, non-dense set in X we have $\dim K \leq 1$. Whence $\dim f(K) \leq 1$; and since as shown above Y also is a 2-manifold, it follows from this that $f(K)$ is non-dense. Accordingly, by (2.21) not only D_f but also L_f must be semi-dense in X .

Clearly we have the following alternative form of (3.1) which we state as

(3.2) THEOREM. *Let f , X and Y be as described in the first sentence of (3.1). In order that f preserve non-density for compact sets it is necessary and sufficient that L_f be semi-dense in X .*

4. **Quasi-open mappings on the general 2-manifold.** We now show that the case of a mapping of this same type operating on an arbitrary 2-manifold can be reduced essentially to the case of a region on a sphere so that similar conclusions hold.

(4.1) LEMMA. *Let $f(X)=Y$ be quasi-open where X is a 2-manifold without edges and Y is a locally connected generalized continuum and suppose that $f(L_f)$ is dense in Y . If there exists in X a compact set K of dimension ≤ 1 whose image contains an open set in Y , then there exists a region R in X contained in a 2-cell of X such that $Q=f(R)$ is open in Y , the mapping $f(R)=Q$ is compact and quasi-open and for some compact subset K_1 of $K \cdot R$, $f(K_1)$ contains an open set.*

Proof. Let V be an open set in $f(K)$. Then there is a point $y \in V$ such that $f^{-1}(y)$ is totally disconnected. Now for each $x \in K \cdot f^{-1}(y)$ there exists a 2-cell E_x on X with edge C_x and interior I_x such that $f(E_x) \subset V$, $C_x \cdot f^{-1}(y) = 0$. Thus if Q_x is the component of $Y - f(C_x)$ containing y and R_x is the component of $f^{-1}(Q_x)$ containing x we have $R_x \subset I_x$ because $R_x \cdot C_x = 0$. Accordingly, R_x being conditionally compact [5], $f(R_x) = Q_x \subset V$ and the mapping $f(R_x) = Q_x$ is compact and quasi-open.

Now since $K \cdot f^{-1}(y)$ is covered by a finite union U of the sets R_x and $f(K \cdot U)$ contains an open set $V - f(K - K \cdot U)$ in V about y , some one of the sets R_x , say R , is such that $f(K \cdot R)$ contains an open set. Since $K \cdot R$ is closed in R , for some compact set $K_1 \subset K \cdot R$, $f(K_1)$ must likewise contain an open set in $Q = f(R)$. Thus the lemma is proven.

Since a region on a 2-cell may be considered as a region on a sphere (by mapping the 2-cell topologically onto a 2-cell on a sphere), this lemma together with the theorems in § 3 yield at once

(4.2) THEOREM. *Given a quasi-open mapping $f(X)=Y$ where X is a 2-manifold without edges and Y is a locally connected generalized continuum such that no component of a point inverse lying inside a closed 2-cell on X separates X , in order that f preserve non-density for compact sets it is necessary and sufficient that L_f be semi-dense in X .*

Note. Most of the results in this paper were stated without proof, or with only brief indications of proof in some cases, by the author in his Presidential Address before the American Mathematical Society [6].

For further discussion of these results, in particular for cases in which alternative dimension preserving forms of (4.2) above are possible, see [6].

5. Differentiable functions. We now show that a mapping from a region of the z -plane Z into the w -plane W generated by a function $w=f(z)$ satisfying certain differentiability conditions will satisfy the requirements needed in the preceding sections to insure the preservation of non-density for compact sets.

(5.1) **THEOREM.** *Let $w=f(z)$ be continuous in a region X of Z and differentiable at all points of a dense set $f^{-1}(Y_0)$ in X which is the inverse of an open subset Y_0 of $Y=f(X)$. Then f is strongly quasi-open, no component of a point inverse lying inside a closed 2-cell on X separates X and the set $f(L_f)$ is dense in Y . Further, L_f is semi-dense in X .*

Proof. (Note. In the proof of all but the final statement use is made of only easily established topological properties of functions meeting minimum differentiability requirements. In proving the last one, however, we use the property, rather more difficult to establish topologically, that a non-constant function everywhere differentiable in a region R cannot be constant on any open set in R .)

To prove f strongly quasi-open it suffices (see § 7 of [6]) to show that for any elementary region R in X with boundary C in X ,

$$(*) \quad f(R+C) = f(C) + \text{the union of bounded components of } W - f(C),$$

where "elementary" means that R is bounded and C consists of a finite number of disjoint simple closed curves. To accomplish this, let S be a component of $W - f(C)$ such that the set $S_0 = S \cdot f(R)$ is not empty. Since $R \cdot f^{-1}(S_0)$ is open and nonempty, it therefore intersects $f^{-1}(Y_0)$. Thus $S_0 \cdot Y_0$ is not empty. Let Q be a component of $S_0 \cdot Y_0$. Since $R \cdot f^{-1}(Q)$ is open and thus has only a countable number of components, there exists a component T of $R \cdot f^{-1}(Q)$ on which f is not constant. As f is differentiable on T by hypothesis [because $T \subset f^{-1}(Y_0)$] there exists a point $z_0 \in T$ where $f'(z_0) \neq 0$. Now using properties of the circulation index, it readily follows that Q contains the interior of a square and thus contains a point q such that $f'(z) \neq 0$ for all $z \in f^{-1}(q)$. Since this makes the circulation index equal $2\pi i$ times a positive integer when taken around any sufficiently small circle enclosing a point of $f^{-1}(q)$, it results at once that the circulation index taken over all of C of f about q must be $\neq 0$. Further, since this latter index is constant throughout S , that is, it has the same value when any $p \in S$ is substituted for q , it follows that every point p of S must belong to $f(R)$. For details of the argument needed here using the circulation index the reader is referred to the last paragraph of § 5 of [7].

Hence we have $S \subset f(R)$. This gives (*), however, because $f(R + C)$ obviously cannot contain the whole unbounded component of $W - f(C)$. Thus any component of $W - f(C)$ intersecting $f(R)$ must be bounded and must lie wholly in $f(R)$. Accordingly f is strongly quasi-open.

Suppose, contrary to the second assertion, that some component K of $f^{-1}(w_0)$, for some $w_0 \in Y$, lies inside a closed 2-cell A on X and separates X . Then one component Q of $X - K$ must lie wholly inside A since only one component of $X - K$ intersects the edge of A . Let y be a point of $f(Q + K)$ such that $|y - w_0| = \max |f(z) - w_0|$ for $z \in Q + K$. Then Q contains a component H of $f^{-1}(y)$ and H is compact. Accordingly, by the strong quasi-openness of f , y must be interior to $f(Q)$ contrary to $|f(z) - w_0| \leq |y - w_0|$ for all $z \in Q$.

That $f(L_f)$ is dense in Y is an immediate consequence of the fact that Y_0 is dense in Y and the quasi-openness of f already established. For any open set in Y thus contains the interior I of a square such that f is differentiable everywhere on $f^{-1}(I)$. Thus for some $q \in I$, $f'(z) \neq 0$ for all $z \in f^{-1}(q)$. This makes $f^{-1}(q)$ a scattered set which therefore surely lies in L_f .

Finally, to prove L_f semi-dense in X we note first that if T is any region in X on which f is non-constant and everywhere differentiable, then as shown above in the second paragraph of this proof, T contains a point z_0 where $f'(z_0) \neq 0$ and indeed $f(T)$ contains points q such that $f'(z)$ does not vanish on $f^{-1}(q)$, so that $f^{-1}(q) \subset L_f$. Accordingly any such region T intersects L_f . Now if U is any open set in X with an open image, $f(U) \cdot Y_0$ contains a region Q and if T is any component of $f^{-1}(Q) \cdot U$ on which f is not constant (and there are such components T because the collection of all components of $f^{-1}(Q) \cdot U$ is countable), then T intersects L_f as shown above. However, the same holds for an arbitrary subregion T_0 of T , because f is likewise non-constant and everywhere differentiable on T_0 . Thus L_f is dense in T .

REFERENCES

1. P. S. Alexandroff, Dokl. (C. R.) Akad. Nauk SSSR. IV (XIII), No. 7 (111), 295-299.
2. R. L. Moore, *Concerning upper semi-continuous collections of continua*, Trans. Amer. Math. Soc., **27** (1925), 416-428.
3. A. D. Wallace, *Quasi-monotone transformations*, Duke Math. J., **7** (1940), 136-145.
4. G. T. Whyburn, *Interior transformations on surfaces*, Amer. J. Math., **60** (1938), 477-490.
5. ———, *An open mapping approach to Hurwitz's theorem*, Trans. Amer. Math. Soc., **71** (1951), 114.
6. ———, *Topological analysis*, Bull. Amer. Math. Soc., **62** (1956), 204-218.
7. ———, *Lectures on functions of a complex variable*, Univ. of Michigan Press (1955), 1-14.

PACIFIC JOURNAL OF MATHEMATICS

EDITORS

H. L. ROYDEN
Stanford University
Stanford, California

R. A. BEAUMONT
University of Washington
Seattle 5, Washington

A. L. WHITEMAN
University of Southern California
Los Angeles 7, California

E. G. STRAUS
University of California
Los Angeles 24, California

ASSOCIATE EDITORS

E. F. BECKENBACH
C. E. BURGESS
M. HALL
E. HEWITT

A. HORN
V. GANAPATHY IYER
R. D. JAMES
M. S. KNEBELMAN

L. NACHBIN
I. NIVEN
T. G. OSTROM
M. M. SCHIFFER

G. SZEKERES
F. WOLF
K. YOSIDA

SUPPORTING INSTITUTIONS

UNIVERSITY OF BRITISH COLUMBIA
CALIFORNIA INSTITUTE OF TECHNOLOGY
UNIVERSITY OF CALIFORNIA
MONTANA STATE UNIVERSITY
UNIVERSITY OF NEVADA
OREGON STATE COLLEGE
UNIVERSITY OF OREGON
UNIVERSITY OF SOUTHERN CALIFORNIA

STANFORD UNIVERSITY
UNIVERSITY OF UTAH
WASHINGTON STATE COLLEGE
UNIVERSITY OF WASHINGTON

* * *

AMERICAN MATHEMATICAL SOCIETY
CALIFORNIA RESEARCH CORPORATION
HUGHES AIRCRAFT COMPANY
THE RAMO-WOOLDRIDGE CORPORATION

Mathematical papers intended for publication in the *Pacific Journal of Mathematics* should be typewritten (double spaced), and the author should keep a complete copy. Manuscripts may be sent to any of the editors. All other communications to the editors should be addressed to the managing editor, E. G. Straus at the University of California, Los Angeles 24, California.

50 reprints per author of each article are furnished free of charge; additional copies may be obtained at cost in multiples of 50.

The *Pacific Journal of Mathematics* is published quarterly, in March, June, September, and December. The price per volume (4 numbers) is \$12.00; single issues, \$3.50. Back numbers are available. Special price to individual faculty members of supporting institutions and to individual members of the American Mathematical Society: \$4.00 per volume; single issues, \$1.25.

Subscriptions, orders for back numbers, and changes of address should be sent to Pacific Journal of Mathematics, 2120 Oxford Street, Berkeley 4, California.

Printed at Kokusai Bunken Insatsusha (International Academic Printing Co., Ltd.), No. 10, 1-chome, Fujimi-cho, Chiyoda-ku, Tokyo, Japan.

PUBLISHED BY PACIFIC JOURNAL OF MATHEMATICS, A NON-PROFIT CORPORATION

The Supporting Institutions listed above contribute to the cost of publication of this Journal, but they are not owners or publishers and have no responsibility for its content or policies.

Pacific Journal of Mathematics

Vol. 7, No. 2

February, 1957

| | |
|---|------|
| William F. Donoghue, Jr., <i>The lattice of invariant subspaces of a completely continuous quasi-nilpotent transformation</i> | 1031 |
| Michael (Mihály) Fekete and J. L. Walsh, <i>Asymptotic behavior of restricted extremal polynomials and of their zeros</i> | 1037 |
| Shaul Foguel, <i>Biorthogonal systems in Banach spaces</i> | 1065 |
| David Gale, <i>A theorem on flows in networks</i> | 1073 |
| Ioan M. James, <i>On spaces with a multiplication</i> | 1083 |
| Richard Vincent Kadison and Isadore Manual Singer, <i>Three test problems in operator theory</i> | 1101 |
| Maurice Kennedy, <i>A convergence theorem for a certain class of Markoff processes</i> | 1107 |
| G. Kurepa, <i>On a new reciprocity, distribution and duality law</i> | 1125 |
| Richard Kenneth Lashof, <i>Lie algebras of locally compact groups</i> | 1145 |
| Calvin T. Long, <i>Note on normal numbers</i> | 1163 |
| M. Mikolás, <i>On certain sums generating the Dedekind sums and their reciprocity laws</i> | 1167 |
| Barrett O'Neill, <i>Induced homology homomorphisms for set-valued maps</i> | 1179 |
| Mary Ellen Rudin, <i>A topological characterization of sets of real numbers</i> | 1185 |
| M. Schiffer, <i>The Fredholm eigen values of plane domains</i> | 1187 |
| F. A. Valentine, <i>A three point convexity property</i> | 1227 |
| Alexander Doniphan Wallace, <i>The center of a compact lattice is totally disconnected</i> | 1237 |
| Alexander Doniphan Wallace, <i>Two theorems on topological lattices</i> | 1239 |
| G. T. Whyburn, <i>Dimension and non-density preservation of mappings</i> | 1243 |
| John Hunter Williamson, <i>On the functional representation of certain algebraic systems</i> | 1251 |