

Pacific Journal of Mathematics

**A SUBSTITUTION THEOREM FOR THE LAPLACE
TRANSFORMATION AND ITS GENERALIZATION TO
TRANSFORMATIONS WITH SYMMETRIC KERNEL**

ROBERT GEROGE BUSCHMAN

A SUBSTITUTION THEOREM FOR THE LAPLACE TRANSFORMATION AND ITS GENERALIZATION TO TRANSFORMATIONS WITH SYMMETRIC KERNEL

R. G. BUSCHMAN

In the problem of the derivation of images of functions under the Laplace transformation, the question arises as to the type of image produced if t is replaced by $g(t)$ in the original. Specific examples have been given by Erdélyi [3, vol. I §§ 4.1, 5.1, 6.1], Doetsch [1, 75–80], McLachlan, Humbert, and Poli [6, pp. 11–13] and [7, pp. 15–20], and Labin [5, p. 41] and a general formula is also listed by Doetsch [1, 75–80].

The Laplace transformation will be taken as

$$f(s) = \int_0^{\infty} e^{-st} F(t) dt$$

in which the integral is taken in the Lebesgue sense and which, as suggested by Doetsch [2, vol. I, p. 44], will be denoted by

$$F(t) \underset{t}{\overset{\mathcal{L}}{\circ}} \underset{s}{\bullet} f(s).$$

(The symbol will be read “ $F(t)$ has a Laplace transform $f(s)$ ”.)

THEOREM 1. *If*

(i) k , g , and the inverse function $h = g^{-1}$ are single-valued analytic functions, real on $(0, \infty)$, and such that $g(0) = 0$ and $g(\infty) = \infty$ (or $g(0) = \infty$ and $g(\infty) = 0$);

(ii) $F(t) \underset{t}{\overset{\mathcal{L}}{\circ}} \underset{s}{\bullet} f(s)$ and this Laplace integral converges for $0 < \Re s$;

(iii) there exists a function $\Phi(s, u)$, $\Phi(s, u) \underset{u}{\overset{\mathcal{L}}{\circ}} \underset{p}{\bullet} \phi(s, p)$ and this Laplace integral converges for $0 < \Re p$, and $\phi(s, p) = e^{-sh(p)} k[h(p)] |h'(p)|$; and

(iv) $\int_0^{\infty} \left[\int_0^{\infty} |e^{-up} \Phi(s, u) F(p)| du \right] dp$ converges for $a < \Re s$;

then

$$k(t) F[g(t)] \underset{t}{\overset{\mathcal{L}}{\circ}} \underset{s}{\bullet} \int_0^{\infty} \Phi(s, u) f(u) du$$

Received June 25, 1957. This paper is a portion of a thesis submitted to the Graduate School of the University of Colorado.

and this Laplace integral converges for $a < \Re s$.

Proof. From (iii) and (iv) it follows that

$$\int_0^\infty e^{-sh(p)} k[h(p)] |h'(p)| F(p) dp$$

is absolutely convergent for $a < \Re s$. There are two cases to be considered. Since from (i) both g and h are single valued, h is monotonic.

Case 1. If $g(0) = 0$ and $g(\infty) = \infty$, then $0 \leq h'(p)$.

Case 2. If $g(0) = \infty$ and $g(\infty) = 0$, then $h'(p) \leq 0$.

In either case, therefore, if the substitution $t = h(p)$ is made in the integral

$$\int_0^\infty e^{-st} k(t) F[g(t)] dt,$$

then

$$k(t) F[g(t)] \overset{\mathcal{L}}{\underset{t}{\circ-\bullet}} \int_0^\infty e^{-sh(p)} k[h(p)] |h'(p)| F(p) dp.$$

From (iii) $\Phi(s, u)$ can be introduced and by (iv) the order of integration changed so that

$$k(t) F[g(t)] \overset{\mathcal{L}}{\underset{t}{\circ-\bullet}} \int_0^\infty \left[\int_0^\infty e^{-up} F(p) dp \right] \Phi(s, u) du.$$

Finally, from (ii)

$$k(t) F[g(t)] \overset{\mathcal{L}}{\underset{t}{\circ-\bullet}} \int_0^\infty \Phi(s, u) f(u) du.$$

To show that there are functions $\phi(s, p)$ as assumed in (iii), let, for example, $g(t) = t^2$ and $k(t) = 1$ so that

$$\Phi(s, p) = (4p)^{-1/2} e^{-sp^{1/2}}$$

and

$$\phi(s, u) = (4\pi u)^{-1/2} e^{-s^2/4u}.$$

From this the known relation

$$F(t^2) \overset{\mathcal{L}}{\underset{t}{\circ-\bullet}} \int_0^\infty (4\pi u)^{-1/2} e^{-s^2/4u} f(u) du$$

is obtained.

Special cases of $k(t)$ will sometimes simplify the image of $\Phi(s, u)$.
 If $k(t)=|g'(t)|K[g(t)]$, then

$$\Phi(s, u) \underset{u \quad p}{\overset{\mathcal{L}}{\circ-\bullet}} K(p) e^{-s h(p)} .$$

If $k(t)=|g'(t)|[g(t)]^c$, then

$$\Phi(s, u) \underset{u \quad p}{\overset{\mathcal{L}}{\circ-\bullet}} p^c e^{-s h(p)} .$$

In the proof of Theorem 1 it is noted that the only important property required of the kernel is that it be symmetric. Therefore consider the transformation

$$f(s)=\int_a^b K(s, t) F(t) dt$$

in which the integral is taken in the Lebesgue sense and in which the interval (a, b) may be unbounded. This transformation will be called the \mathcal{F} -transform and denoted by

$$F(t) \underset{t \quad s}{\overset{\mathcal{F}}{\circ-\bullet}} f(s) ,$$

The following theorem is for this transformation with symmetric kernel.

THEOREM 2. *If*

(i) k, g , and $h=g^{-1}$ are single-valued analytic functions, real on (a, b) , and such that $g(a)=a$ and $g(b)=b$ (or $g(a)=b$ and $g(b)=a$);

(ii) $F(t) \underset{t \quad s}{\overset{\mathcal{F}}{\circ-\bullet}} f(s)$ and this transformation integral converges for $a < s < b$;

(iii) there exists a function $\Phi(s, u)$, $\Phi(s, u) \underset{u \quad p}{\overset{\mathcal{F}}{\circ-\bullet}} \phi(s, p)$, this transformation integral converges for $a < s < b$, and

$$\phi(s, p)=K[s, h(p)] k[h(p)] |h'(p)| ;$$

(iv)

$$\int_a^b \left[\int_a^b |K(u, p) \Phi(s, u) F(p)| du \right] dp$$

converges for $s=s_0$; and

(v) $K(u, p) \equiv K(p, u)$; then $k(t) F[g(t)] \underset{t \quad s}{\overset{\mathcal{F}}{\circ-\bullet}} \int_a^b \Phi(s, u) f(u) du$ and this

transformation integral converges for $s=s_0$.

The proof follows in a manner similar to that of Theorem 1.

Formulas which hold provided $F(t)$ satisfies (ii) or (iv) of the theorem can be obtained for various transforms for specific $k(s)$ and $g(s)$ with the aid of tables [3, formulas 14.1(6), 8.12(10), 5.5(6)].

Formula 1. For the Stieltjes transformation $K(s, t)=(s+t)^{-1}$.

$$t^{b+1}F(at^2) \underset{t}{\circ} \underset{s}{\bullet} \int_0^\infty \frac{(u/a)^{b/2}}{2\pi a} \left[\frac{(u/a) \cos b\pi/2 - s \sin b\pi/2}{s^2 + u/a} \right] f(u) du$$

for a positive.

Formula 2. For the Hankel transformation $K(s, t)=J_\nu(st)(st)^{1/2}$

$$t^{-2}F(a/t) \underset{t}{\circ} \underset{s}{\bullet} a^{-1} \int_0^\infty \sqrt{aus} J_\nu(2\sqrt{aus}) f(u) du$$

for $-1/2 < \nu$ and a positive.

The Laplace transformation will be considered in the next two formulas.

Formula 3.

$$(t+b/a)^a F(at^2 + 2bt)$$

$$\underset{t}{\circ} \underset{s}{\bullet} (1/2\pi) e^{bs/a} \int_0^\infty e^{-b^2s/a} e^{-s^2/8au} (\sqrt{2au})^{-a-1} D_a(s/\sqrt{2au}) f(u) du$$

for a and b positive and in which $D_d(z)$ is the parabolic cylinder function. The range of permissible values of d will depend, according to (iv), on the particular function $F(u)$.

Formula 4.

$$t^{a-1}F(at^{-b}) \underset{t}{\circ} \underset{s}{\bullet} b^{-1} \int_0^\infty (au)^{a/b} \phi[1/b, (d+b)/b; -s(au)^{1/b}] f(u) du$$

for a and b positive and in which $\phi(A, B; Z)$ is Wrights' function [4, vol. 3, § 18.1]. The range of permissible values of d will depend, according to (iv), on the particular function $F(u)$. In the special case $b=1$ the formula becomes

$$t^{a-1}F(a/t) \underset{t}{\circ} \underset{s}{\bullet} \int_0^\infty (\sqrt{au/s})^a J_a(2\sqrt{aus}) f(u) du .$$

REFERENCES

1. Gustav Doetsch, *Tabellen zur Laplace-transformation*, Berlin, 1947.
2. ———, *Handbuch der Laplace-Transformation*, Basel, 1950.
3. Arthur Erdélyi and others, editors, *Tables of integral transforms*, New York, 1954.
4. ———, *Higher transcendental functions*, New York, 1955.
5. Édouard Labin, *Calcul opérationnel*, Paris, 1949.
6. N. W. McLachlan and Pierre Humbert, *Formulaire pour le calcul symbolique*, Paris, 1941.
7. ———, ——— and Louis Poli, *Supplément du formulaire pour le calcul symbolique*, Paris, 1950.

UNIVERSITY OF COLORADO

PACIFIC JOURNAL OF MATHEMATICS

EDITORS

H. L. ROYDEN
Stanford University
Stanford, California

R. A. BEAUMONT
University of Washington
Seattle 5, Washington

A. L. WHITEMAN
University of Southern California
Los Angeles 7, California

E. G. STRAUS
University of California
Los Angeles 24, California

ASSOCIATE EDITORS

E. F. BECKENBACH
C. E. BURGESS
M. HALL
E. HEWITT

A. HORN
V. GANAPATHY IYER
R. D. JAMES
M. S. KNEBELMAN

L. NACHBIN
I. NIVEN
T. G. OSTROM
M. M. SCHIFFER

G. SZEKERES
F. WOLF
K. YOSIDA

SUPPORTING INSTITUTIONS

UNIVERSITY OF BRITISH COLUMBIA
CALIFORNIA INSTITUTE OF TECHNOLOGY
UNIVERSITY OF CALIFORNIA
MONTANA STATE UNIVERSITY
UNIVERSITY OF NEVADA
OREGON STATE COLLEGE
UNIVERSITY OF OREGON
UNIVERSITY OF SOUTHERN CALIFORNIA

STANFORD UNIVERSITY
UNIVERSITY OF UTAH
WASHINGTON STATE COLLEGE
UNIVERSITY OF WASHINGTON
* * *

AMERICAN MATHEMATICAL SOCIETY
CALIFORNIA RESEARCH CORPORATION
HUGHES AIRCRAFT COMPANY
THE RAMO-WOOLDRIDGE CORPORATION

Mathematical papers intended for publication in the *Pacific Journal of Mathematics* should be typewritten (double spaced), and the author should keep a complete copy. Manuscripts may be sent to any of the editors. All other communications to the editors should be addressed to the managing editor, E. G. Straus at the University of California, Los Angeles 24, California.

50 reprints per author of each article are furnished free of charge; additional copies may be obtained at cost in multiples of 50.

The *Pacific Journal of Mathematics* is published quarterly, in March, June, September, and December. The price per volume (4 numbers) is \$12.00; single issues, \$3.50. Back numbers are available. Special price to individual faculty members of supporting institutions and to individual members of the American Mathematical Society: \$4.00 per volume; single issues, \$1.25.

Subscriptions, orders for back numbers, and changes of address should be sent to Pacific Journal of Mathematics, 2120 Oxford Street, Berkeley 4, California.

Printed at Kokusai Bunken Insatsusha (International Academic Printing Co., Ltd.), No. 10, 1-chome, Fujimi-cho, Chiyoda-ku, Tokyo, Japan.

PUBLISHED BY PACIFIC JOURNAL OF MATHEMATICS, A NON-PROFIT CORPORATION

The Supporting Institutions listed above contribute to the cost of publication of this Journal, but they are not owners or publishers and have no responsibility for its content or policies.

Pacific Journal of Mathematics

Vol. 7, No. 4

April, 1957

| | |
|--|------|
| Robert Geroge Buschman, <i>A substitution theorem for the Laplace transformation and its generalization to transformations with symmetric kernel</i> | 1529 |
| S. D. Conte, <i>Numerical solution of vibration problems in two space variables</i> | 1535 |
| Paul Dedecker, <i>A property of differential forms in the calculus of variations</i> | 1545 |
| H. Delange and Heini Halberstam, <i>A note on additive functions</i> | 1551 |
| Jerald L. Ericksen, <i>Characteristic direction for equations of motion of non-Newtonian fluids</i> | 1557 |
| Avner Friedman, <i>On two theorems of Phragmén-Lindelöf for linear elliptic and parabolic differential equations of the second order</i> | 1563 |
| Ronald Kay Gettoor, <i>Additive functionals of a Markov process</i> | 1577 |
| U. C. Guha, <i>(γ, k)-summability of series</i> | 1593 |
| Alvin Hausner, <i>The tauberian theorem for group algebras of vector-valued functions</i> | 1603 |
| Lester J. Heider, <i>T-sets and abstract (L)-spaces</i> | 1611 |
| Melvin Henriksen, <i>Some remarks on a paper of Aronszajn and Panitchpakdi</i> | 1619 |
| H. M. Lieberstein, <i>On the generalized radiation problem of A. Weinstein</i> | 1623 |
| Robert Osserman, <i>On the inequality $\Delta u \geq f(u)$</i> | 1641 |
| Calvin R. Putnam, <i>On semi-normal operators</i> | 1649 |
| Binyamin Schwarz, <i>Bounds for the principal frequency of the non-homogeneous membrane and for the generalized Dirichlet integral</i> | 1653 |
| Edward Silverman, <i>Morrey's representation theorem for surfaces in metric spaces</i> | 1677 |
| V. N. Singh, <i>Certain generalized hypergeometric identities of the Rogers-Ramanujan type. II</i> | 1691 |
| R. J. Smith, <i>A determinant in continuous rings</i> | 1701 |
| Drury William Wall, <i>Sub-quasigroups of finite quasigroups</i> | 1711 |
| Sadayuki Yamamuro, <i>Monotone completeness of normed semi-ordered linear spaces</i> | 1715 |
| C. T. Rajagopal, <i>Simplified proofs of "Some Tauberian theorems" of Jakimovski: Addendum and corrigendum</i> | 1727 |
| N. Aronszajn and Prom Panitchpakdi, <i>Correction to: "Extension of uniformly continuous transformations in hyperconvex metric spaces"</i> | 1729 |
| Alfred Huber, <i>Correction to: "The reflection principle for polyharmonic functions"</i> | 1731 |