

Pacific Journal of Mathematics

ON SEMI-NORMAL OPERATORS

CALVIN R. PUTNAM

ON SEMI-NORMAL OPERATORS

C. R. PUTNAM

1. A bounded linear operator A in a Hilbert space will be called semi-normal if

$$(1) \quad H = AA^* - A^*A \geq 0 \quad (\text{or } \leq 0).$$

If A is a finite matrix, for instance, then relation (1) implies $H=0$, so that A is even normal; cf., e.g., [4]. That (1) may hold with $H \neq 0$ is seen if one chooses, for instance, A to be the isometric matrix defined by $A=D=(d_{ij})$ where $d_{i+1,i}=1$ and $d_{ij}=0$ otherwise. The purpose of this note is to investigate the spectrum of the semi-normal operator A and of the associated self-adjoint operators J_θ defined by

$$(2) \quad J_\theta = \frac{A_\theta + A_\theta^*}{2}, \quad A_\theta = Ae^{-i\theta} \quad (\theta \text{ real}).$$

It is seen that, in particular, J_θ becomes the real or the imaginary part of A according as $\theta=0$ or $\theta=\pi/2$.

A number λ belonging to the spectrum of A ($\text{sp}(A)$) will be called accessible if there exists a sequence of numbers λ_n not belonging to $\text{sp}(A)$ for which $\lambda_n \rightarrow \lambda$ as $n \rightarrow \infty$. If M is any self-adjoint operator, $\max M$ and $\min M$ will denote the greatest and the least points respectively of the set $\text{sp}(M)$.

The following theorems will be proved:

THEOREM 1. *Let A be semi-normal with $H \geq 0$ and let $\lambda = re^{i\theta}$ (r real, ≥ 0) be an accessible point of the spectrum of A . Then*

$$(3) \quad (\max J_\theta)^2 \geq \min AA^*$$

and

$$(4) \quad |r - \max J_\theta| \leq ((\max J_\theta)^2 - \min AA^*)^{1/2},$$

where J_θ is defined by (2).

THEOREM 2. *Let A be semi-normal and let $J=J_\theta$ have the spectral resolution $J = \int \lambda dE$. Then, if $S=S_\theta$ is any measurable set for which*

$$(5) \quad \int_S dE = I,$$

Received March 6, 1957. This research was supported in part by the United States Air Force under Contract No. AF 18 (603)-139.

there holds the inequality

$$(6) \quad \|H\| \leq 4\|A\| \text{ meas } S.$$

The proof of Theorem 1 will be given in § 2 below. The assertion of Theorem 2 can be considered as a supplement to Corollary 3 of [5]. The proof follows readily from the Lemma, *loc. cit.*, p. 1027 if one notes that $H/2 = J_\theta A_\theta^* - A_\theta^* J_\theta$ and that $\|A_\theta\| = \|A\|$.

Various corollaries can be obtained from the two theorems. For instance, as a consequence of Theorem 1, one has the

COROLLARY 1. *If V is isometric and not unitary, then its spectrum is the disk $|\lambda| \leq 1$ of the complex plane.*

Actually it is possible to deduce this result from a normal form for such operators; cf., e.g., [8, p. 351 ff]. It should be noted that the spectrum of the isometric matrix D defined earlier in this paper, and which occurs in the normal form, is the disk $|\lambda| \leq 1$; cf. [9, p. 279].

The proof of the corollary as a consequence of Theorem 1 however is as follows. Put $A^* = V$ so that $AA^* = I$; clearly V is semi-normal and $H \geq 0$. Let $\lambda = re^{i\theta}$ ($r \geq 0$) be an accessible point in the spectrum of A (that is, of A^* or V). Then, by (3), $|\max J_\theta| \geq 1$. On the other hand, $\|A\| = 1$, and hence $|\max J_\theta| \leq 1$. Thus $|\max J_\theta| = 1$ and (4) implies $r = 1$; consequently, the only possible accessible points of the spectrum of an isometric operator lie on the circle $|\lambda| = 1$. However, if the operator is not unitary, then $\lambda = 0$ lies in its spectrum. Hence, the entire disk $|\lambda| \leq 1$ is in the spectrum and the proof is complete.

Another consequence of Theorem 1 is

COROLLARY 2. *If A is semi-normal, if 0 lies in the spectrum of A , and if $\min AA^* > 0$, then for any θ the circular disk*

$$|\lambda| \leq \max J_\theta - ((\max J_\theta)^2 - \min AA^*)^{1/2}$$

lies in the spectrum of A (where, of course, $\max J_\theta > 0$).

The proof follows from the observation that $\lambda = 0$ is in $\text{sp}(A)$ but no accessible points of the spectrum can lie in the disk in question.

It can be remarked that if A is an arbitrary bounded linear operator (not necessarily semi-normal), and if the conditions that 0 be in $\text{sp}(A)$ and $\min AA^* > 0$ are fulfilled, then there surely exists *some* circular disk $|\lambda| \leq \text{const.}$ in the spectrum of A ; see, e.g., [7, pp. 76-78]. If however A is semi-normal, the radius of the corollary can even be specified.

An immediate consequence of Theorem 2 is the

COROLLARY 3. *If A is semi-normal but not normal, then the spectrum*

of J_θ (in particular, of the real or imaginary part of A) has a positive measure not less than $\|H\|/4\|A\|$.

It should be noted that (5) surely holds if S is the spectrum of J although it may hold for a set of measure less than that of the spectrum (but whose closure would, of course, contain the spectrum).

It seems natural to conjecture that the spectrum of (say) the real part, $J=(A+A^*)/2$ of any semi-normal, but not normal, operator A must be an interval. Evidence to support the conjecture is furnished by the isometric, but not unitary, operators V , in which case the spectrum of $(V+V^*)/2$ is the interval $-1 \leq \lambda \leq 1$. This fact also follows from the normal form for isometric operators referred to above and from the fact that the spectrum of $(D+D^*)/2$ is the interval $-1 \leq \lambda \leq 1$ (cf., e.g., [3, p. 155]). Further evidence is furnished by the (bounded) matrices $A=(c_{j-i})$, where $c_n=0$ if $n<0$, for which the spectra of the associated Toeplitz matrices $J=(A+A^*)/2$ are intervals, provided J is not a multiple of the unit matrix (in which case A is also); see [1, p. 361] and [2, p. 868]. It was shown in [6] that the matrices A are semi-normal.

The conjecture will remain unsettled. In fact, it will remain undecided whether or not the spectrum of the real part J of a semi-normal, but not normal, operator must even contain some interval. The assertion of Corollary 3 does not seem to preclude the possibility of, for instance, a nowhere dense spectrum (of positive measure).

2. Proof of Theorem 1. Let $\lambda_n=r_n e^{i\theta_n}$ be chosen so that λ_n is not in $\text{sp}(A)$ and $\lambda_n \rightarrow \lambda$ as $n \rightarrow \infty$. Put $A_n=A-\lambda_n I$. Then $A_n A_n^*=A_n A_n^* A_n A_n^{-1}$, so that the spectra of $A_n A_n^*$ and $A_n^* A_n$, hence the spectra of $AA^*-2r_n J_{\theta_n}$, and $A^*A-2r_n J_{\theta_n}$, are (respectively) identical. Since $\lambda=re^{i\theta}$ is in the spectrum of A , then either $(A-\lambda)x_m \rightarrow 0$ or $(A-\lambda)^*x_m \rightarrow 0$ for some sequence of unit vectors x_m . In either case, it follows from (1) that $\limsup(x_m, A^*Ax_m) \leq r^2$ as $m \rightarrow \infty$ and that $(x_m, J_{\theta_n}x_m) \rightarrow r$ as $m, n \rightarrow \infty$. Consequently, $\min(AA^*-2rJ_\theta) \leq -r^2$ and hence $\min AA^*-2r \max J_\theta + r^2 \leq 0$. The desired relations (3) and (4) follow and the proof of Theorem 1 is complete.

REFERENCES

1. P. Hartman and A. Wintner, *On the spectra of Toeplitz's matrices*, Amer. J. Math., **72** (1950), 359-366.
2. ———, *The spectra of Toeplitz's matrices*, Amer. J. Math., **76** (1954), 867-882.
3. D. Hilbert, *Grundzüge einer allgemeinen Theorie der linearen Integralgleichungen*, Leipzig, 1912.
4. C. R. Putnam, *On commutators of bounded matrices*, Amer. J. Math., **73** (1951), 127-131.

5. ———, *On commutators and Jacobi matrices*, Proc. Amer. Math. Soc., **7** (1956), 1026-1030.
6. ———, *Commutators and absolutely continuous operators*, Trans. Amer. Math. Soc.
7. C. R. Putnam and Winter, *The orthogonal group in Hilbert space*, Amer. J. Math., **74** (1952), 52-78.
8. M. H. Stone, *Linear Transformations in Hilbert Space*, New York, 1932.
9. A. Wintner, *Zur Theorie der beschränkten Bilinearformen*, Math. Z. **30** (1929), 228-284.

PURDUE UNIVERSITY

PACIFIC JOURNAL OF MATHEMATICS

EDITORS

H. L. ROYDEN
Stanford University
Stanford, California

R. A. BEAUMONT
University of Washington
Seattle 5, Washington

A. L. WHITEMAN
University of Southern California
Los Angeles 7, California

E. G. STRAUS
University of California
Los Angeles 24, California

ASSOCIATE EDITORS

E. F. BECKENBACH
C. E. BURGESS
M. HALL
E. HEWITT

A. HORN
V. GANAPATHY IYER
R. D. JAMES
M. S. KNEBELMAN

L. NACHBIN
I. NIVEN
T. G. OSTROM
M. M. SCHIFFER

G. SZEKERES
F. WOLF
K. YOSIDA

SUPPORTING INSTITUTIONS

UNIVERSITY OF BRITISH COLUMBIA
CALIFORNIA INSTITUTE OF TECHNOLOGY
UNIVERSITY OF CALIFORNIA
MONTANA STATE UNIVERSITY
UNIVERSITY OF NEVADA
OREGON STATE COLLEGE
UNIVERSITY OF OREGON
UNIVERSITY OF SOUTHERN CALIFORNIA

STANFORD UNIVERSITY
UNIVERSITY OF UTAH
WASHINGTON STATE COLLEGE
UNIVERSITY OF WASHINGTON
* * *

AMERICAN MATHEMATICAL SOCIETY
CALIFORNIA RESEARCH CORPORATION
HUGHES AIRCRAFT COMPANY
THE RAMO-WOOLDRIDGE CORPORATION

Mathematical papers intended for publication in the *Pacific Journal of Mathematics* should be typewritten (double spaced), and the author should keep a complete copy. Manuscripts may be sent to any of the editors. All other communications to the editors should be addressed to the managing editor, E. G. Straus at the University of California, Los Angeles 24, California.

50 reprints per author of each article are furnished free of charge; additional copies may be obtained at cost in multiples of 50.

The *Pacific Journal of Mathematics* is published quarterly, in March, June, September, and December. The price per volume (4 numbers) is \$12.00; single issues, \$3.50. Back numbers are available. Special price to individual faculty members of supporting institutions and to individual members of the American Mathematical Society: \$4.00 per volume; single issues, \$1.25.

Subscriptions, orders for back numbers, and changes of address should be sent to Pacific Journal of Mathematics, 2120 Oxford Street, Berkeley 4, California.

Printed at Kokusai Bunken Insatsusha (International Academic Printing Co., Ltd.), No. 10, 1-chome, Fujimi-cho, Chiyoda-ku, Tokyo, Japan.

PUBLISHED BY PACIFIC JOURNAL OF MATHEMATICS, A NON-PROFIT CORPORATION

The Supporting Institutions listed above contribute to the cost of publication of this Journal, but they are not owners or publishers and have no responsibility for its content or policies.

Robert Geroge Buschman, <i>A substitution theorem for the Laplace transformation and its generalization to transformations with symmetric kernel</i>	1529
S. D. Conte, <i>Numerical solution of vibration problems in two space variables</i>	1535
Paul Dedecker, <i>A property of differential forms in the calculus of variations</i>	1545
H. Delange and Heini Halberstam, <i>A note on additive functions</i>	1551
Jerald L. Ericksen, <i>Characteristic direction for equations of motion of non-Newtonian fluids</i>	1557
Avner Friedman, <i>On two theorems of Phragmén-Lindelöf for linear elliptic and parabolic differential equations of the second order</i>	1563
Ronald Kay Gettoor, <i>Additive functionals of a Markov process</i>	1577
U. C. Guha, <i>(γ, k)-summability of series</i>	1593
Alvin Hausner, <i>The tauberian theorem for group algebras of vector-valued functions</i>	1603
Lester J. Heider, <i>T-sets and abstract (L)-spaces</i>	1611
Melvin Henriksen, <i>Some remarks on a paper of Aronszajn and Panitchpakdi</i>	1619
H. M. Lieberstein, <i>On the generalized radiation problem of A. Weinstein</i>	1623
Robert Osserman, <i>On the inequality $\Delta u \geq f(u)$</i>	1641
Calvin R. Putnam, <i>On semi-normal operators</i>	1649
Binyamin Schwarz, <i>Bounds for the principal frequency of the non-homogeneous membrane and for the generalized Dirichlet integral</i>	1653
Edward Silverman, <i>Morrey's representation theorem for surfaces in metric spaces</i>	1677
V. N. Singh, <i>Certain generalized hypergeometric identities of the Rogers-Ramanujan type. II</i>	1691
R. J. Smith, <i>A determinant in continuous rings</i>	1701
Drury William Wall, <i>Sub-quasigroups of finite quasigroups</i>	1711
Sadayuki Yamamuro, <i>Monotone completeness of normed semi-ordered linear spaces</i>	1715
C. T. Rajagopal, <i>Simplified proofs of "Some Tauberian theorems" of Jakimovski: Addendum and corrigendum</i>	1727
N. Aronszajn and Prom Panitchpakdi, <i>Correction to: "Extension of uniformly continuous transformations in hyperconvex metric spaces"</i>	1729
Alfred Huber, <i>Correction to: "The reflection principle for polyharmonic functions"</i>	1731