

Pacific Journal of Mathematics

ASYMPTOTIC EXPRESSIONS FOR $\sum n^a f(n) \log^r n$

ROBERT GEROGE BUSCHMAN

ASYMPTOTIC EXPRESSIONS FOR $\sum n^a f(n) \log^r n$

R. G. BUSCHMAN

In this paper some asymptotic expressions for sums of the type

$$\sum n^a f(n) \log^r n ,$$

where $f(n)$ is a number theoretic function, are presented. (The summations extend over $1 \leq n \leq x$ unless otherwise noted.) The method applied is to obtain the Laplace transformation,

$$\mathcal{L}\{F(t)\} = \int_0^\infty e^{-st} F(t) dt = f(s)$$

of the sum and then use a Tauberian theorem either from Doetsch [2] or its modification for a pole at points other than the origin, or from Delange [1] to obtain the asymptotic relation. If $f(n)$ is non-negative, then $F(t)$ is a non-negative, non-decreasing function and hence satisfies the conditions for the Tauberian theorems. In many cases the closed form of a Dirichlet series involving the functions are known, and in this case the relation

$$\mathcal{L}\left\{\sum_{1 \leq n \leq e^t} n^a f(n) \log^r n\right\} = (-1)^r s^{-1} (d/ds)^r \sum_1^\infty n^{a-s} f(n)$$

can be used. The functions chosen for discussion and the Dirichlet series involving them can be found in Hardy and Wright [3], Landau [4], [5], or Titchmarsh [7]. We present first a few illustrations of the method and then a more extensive collection of results is presented at the end in a table.

First we choose $\sigma_k(n)$ as an example of a simpler type. Since

$$\sum_1^\infty n^{-s} \sigma_k(n) = \zeta(s) \zeta(s-k) ,$$

we have

$$\mathcal{L}\left\{\sum_{1 \leq n \leq e^t} n^{b-1-k} \sigma_k(n) \log^r n\right\} = f(s) = (-1)^r s^{-1} (d/ds)^r \{\zeta(s+1-b) \zeta(s+1-b+k)\} .$$

For $k > 0$ the pole where $\Re s$ is greatest is at $s = b$ if $b \geq 0$. At that pole, since

$$\zeta^{(m)}(s+1-b) \sim (-1)^m m! (s-b)^{-m-1} ,$$

the Laplace transformation of the sum has the form

Received October 10, 1958.

$$f(s) \sim b^{-1}\zeta(1+k)r!(s-b)^{-r-1}.$$

Now if $b > 0$, then by modifying Doetsch [2, p. 517] for poles not at the origin or from Delange [1, p. 235] we obtain

$$\sum_{1 \leq n \leq e^t} n^{b-1-k} \sigma_k(n) \log^r n \sim b^{-1}\zeta(1+k)e^{bt}r,$$

or, if $x = e^t$

$$\sum n^{b-1-k} \sigma_k(n) \log^r n \sim b^{-1}\zeta(1+k)x^b \log^r x.$$

If $b = 0$, then

$$f(s) \sim \zeta(1+k)r!s^{-r-2},$$

so that from Doetsch [2, p. 517] after substituting $x = e^t$ we obtain

$$\sum n^{-1-k} \sigma_k(n) \log^r n \sim (r+1)^{-1}\zeta(1+k) \log^{r+1} x.$$

The expressions for $\sigma(n)$ can be obtained by setting $k = 1$.

For $k = 0$, $\sigma_k(n)$ becomes $d(n)$ which will be covered as a special case of $d_k(n)$.

For $k < 0$ the pole where $\Re s$ is greatest is at $s = b - k$ so that for $b > k$

$$f(s) \sim (b-k)^{-1}\zeta(1-k)r!(s-b+k)^{-r-1}.$$

Hence

$$\begin{aligned} \sum n^{b-1-k} \sigma_k(n) \log^r n &\sim (b-k)^{-1}\zeta(1-k)x^{b-k} \log^r x, & \text{for } b > k; \\ \sum n^{-1} \sigma_k(n) \log^r n &\sim (r+1)^{-1}\zeta(1-k) \log^{r+1} x, & \text{for } b = k. \end{aligned}$$

By analogy, since

$$\sum_1^{\infty} n^{-s} \phi(n) = \zeta(s-1)/\zeta(s),$$

then

$$\begin{aligned} \sum n^{b-2} \phi(n) \log^r n &\sim \{b\zeta(2)\}^{-1} x^b \log^r x, & \text{for } b > 0; \\ \sum n^{-2} \phi(n) \log^r n &\sim \{(r+1)\zeta(2)\}^{-1} \log^{r+1} x, & \text{for } b = 0. \end{aligned}$$

If $\chi_k(n)$ represents a character, mod k , then the Dirichlet series can be represented by

$$\sum_1^{\infty} n^{-s} \chi_k(n) = L_k(s)$$

so that if χ_k is a principal character then $L_k(s)$ has a pole at $s = 1$ and

$$\begin{aligned} \sum n^{b-1} \chi_k(n) \log^r n &\sim \phi(k)(kb)^{-1} x^b \log^r x, & \text{for } b > 0; \\ \sum n^{-1} \chi_k(n) \log^r n &\sim \phi(k)\{(r+1)b\}^{-1} \log^{r+1} x, & \text{for } b = 0. \end{aligned}$$

The Dirichlet series involving $d_k(n)$ yields a power of the ζ -function, i.e.

$$\sum_1^{\infty} n^{-s} d_k(n) = \zeta^k(s),$$

so that for $k > 0$

$$\mathcal{L} \left\{ \sum_{1 \leq n \leq t} n^{b-1} d_k(n) \log^r n \right\} = (-1)^r s^{-1} (d/ds)^r \zeta^k(s+1-b).$$

Now the Laplace transform can be written to show the behavior at the pole at $s = b$,

$$f(s) \sim (r+k-1)! \{b(k-1)!\}^{-1} (s-b)^{-r-k}.$$

Thus

$$\begin{aligned} \sum n^{b-1} d_k(n) \log^r n &\sim \{b(k-1)!\}^{-1} x^b \log^{r+k-1} x, & \text{for } b > 0; \\ \sum n^{-1} d_k(n) \log^r n &\sim \{(r+k)(k-1)!\}^{-1} \log^{r+k} x, & \text{for } b = 0. \end{aligned}$$

Special cases can be obtained for $k = 1, 2$, since $d_1(n) = 1$ and $d_2(n) = \sigma_0(n) = d(n)$.

In an analogous manner we can obtain from

$$\sum_1^{\infty} n^{-s} d(n^2) = \zeta^3(s)/\zeta(2s)$$

the expressions

$$\begin{aligned} \sum n^{b-1} d(n^2) \log^r n &\sim \{2b\zeta(2)\}^{-1} x^b \log^{r+2} x, & \text{for } b > 0; \\ \sum n^{-1} d(n^2) \log^r n &\sim \{2(r+1)\zeta(2)\}^{-1} \log^{r+3} x, & \text{for } b = 0. \end{aligned}$$

Certain of the common number-theoretic functions have not been considered and do not appear in the table (in particular $\mu(n)$, $\lambda(n)$, and $\chi_k(n)$ for non-principal characters) because the sum $F(t)$ fails to satisfy the non-decreasing hypothesis for the Tauberian theorems. $\lambda(n)$ has the additional bad characteristic as shown by the poles of the closed form of the Dirichlet series

$$\sum_1^{\infty} n^{-s} \lambda(n) = \zeta(2s)/\zeta(s)$$

in that the pole of the numerator is on the line $\Re s = 1/2$ which is critical for the denominator, and thus this is not the pole where $\Re s$ is greatest as required by the theorem from Delange.

Results which he has obtained for the case $r = 0$ and the functions $\sigma(n)$, $\sigma_k(n)$, $d(n)$, and $\phi(n)$, treated by a different method, have been communicated to me in advance of their publication by Mr. Swetharanyam [6].

Table
Asymptotic expressions for $\sum n^a f(n) \log^r n$

General term of the sum	Asymptotic Expressions	
	$b > 0$	$b = 0$
$n^{b-1-k}\sigma_k(n) \log^r n$ ($k > 0$)	$b^{-1}\zeta(1+k)x^b \log^r x$	$(r+1)^{-1}\zeta(1+k) \log^{r+1} x$
$n^{b-1}\sigma_k(n) \log^r n$ ($k < 0$)	$(b-k)^{-1}\zeta(1-k)x^{b-k} \log^r x$ ($b > k$)	$(r+1)^{-1}\zeta(1-k) \log^{r+1} x$ ($b = k$)
$n^{b-2}\sigma(n) \log^r n$	$b^{-1}\zeta(2)x^b \log^r x$	$(r+1)^{-1}\zeta(2) \log^{r+1} x$
$n^{b-1}d_k(n) \log^r n$	$\{b(k-1)!\}^{-1}x^b \log^{r+k-1} x$	$\{(r+k)(k-1)!\}^{-1} \log^{r+k} x$
$n^{b-1}d(n) \log^r n$	$b^{-1}x^b \log^{r+1} x$	$(r+2)^{-1} \log^{r+2} x$
$n^{b-1} \log^r n$	$b^{-1}x^b \log^r x$	$(r+1)^{-1} \log^{r+1} x$
$n^{b-1}\wedge(n) \log^r n$	$b^{-1}x^b \log^r x$	$(r+1)^{-1} \log^{r+1} x$
$n^{b-2}\phi(n) \log^r n$	$\{b\zeta(2)\}^{-1}x^b \log^r x$	$\{(r+1)\zeta(2)\}^{-1} \log^{r+1} x$
$n^{b-1}q_k(n) \log^r n$	$\{b\zeta(k)\}^{-1}x^b \log^r x$	$\{(r+1)\zeta(k)\}^{-1} \log^{r+1} x$
$n^{b-1} \mu(n) \log^r n$	$\{b\zeta(2)\}^{-1}x^b \log^r x$	$\{(r+1)\zeta(2)\}^{-1} \log^{r+1} x$
$n^{b-1}2^{\omega(n)} \log^r n$	$\{b\zeta(2)\}^{-1}x^b \log^{r+1} x$	$\{(r+2)\zeta(2)\}^{-1} \log^{r+2} x$
$n^{b-1}d(n^2) \log^r n$	$\{2b\zeta(2)\}^{-1}x^b \log^{r+2} x$	$\{2(r+3)\zeta(2)\}^{-1} \log^{r+3} x$
$n^{b-1}d^2(n) \log^r n$	$\{6b\zeta(2)\}^{-1}x^b \log^{r+3} x$	$\{6(r+4)\zeta(2)\}^{-1} \log^{r+4} x$
$\frac{\sigma_a(n)\sigma_d(n) \log^r n}{n^{1+a+d-b}}$ ($a > 0$) ($d > 0$)	$\frac{\zeta(1+a+d)\zeta(1+a)\zeta(1+d)}{b\zeta(2+a+d)} x^b \log^r x$	$\frac{\zeta(1+a+d)\zeta(1+a)\zeta(1+d)}{(r+1)\zeta(2+a+d)} \log^{r+1} x$
$\frac{\sigma_a(n)d(n) \log^r n}{n^{1+a-b}}$ ($a > 0$)	$\frac{\zeta^2(1+a)}{b\zeta(2+a)} x^b \log^{r+1} x$	$\frac{\zeta^2(1+a)}{(r+2)\zeta(2+a)} \log^{r+2} x$
$n^{b-2}a(n) \log^r n$	$2(3b)^{-1}x^b \log^r x$	$2\{3(r+1)\}^{-1} \log^{r+1} x$
$n^{b-1}\chi_k(n) \log^r n$	$\phi(k)(kb)^{-1}x^b \log^r x$	$\phi(k)\{k(r+1)\}^{-1} \log^{r+1} x$
$n^{b-1}r(n) \log^r n$	$4b^{-1}L_4(1)x^b \log^r x$	$4(r+1)^{-1}L_4(1) \log^{r+1} x$
$n^{b-1}\wedge(n)\chi_k(n) \log^r n$	$b^{-1}x^b \log^r x$	$(r+1)^{-1} \log^{r+1} x$
$n^{b-2}\phi(n)\chi_k(n) \log^r n$	$\phi(k)\{kbL_k(2)\}^{-1}x^b \log^r x$	$\phi(k)\{(r+1)kL_k(2)\}^{-1} \log^{r+1} x$
$n^{b-1}2^{\omega(n)}\chi_k(n) \log^r n$	$4\phi(k)\{3kb\zeta(2)\}^{-1}x^b \log^{r+1} x$	$4\phi(k)\{3k(r+2)\zeta(2)\}^{-1} \log^{r+2} x$
$n^{b-1}\{\pi(n)-\pi(n-1)\} \log^r n$ ($r > 0$)	$b^{-1}x^b \log^{r-1} x$	$r^{-1} \log^r x$

REFERENCES

1. Hubert Delange, *Généralisation du théorème de Ikehara*, Ann. Sci. l'Ecole Norm. Sup. (3) **71** (1954), 213-242.
2. Gustav Doetsch, *Handbuch der Laplace-Transformation*, Basel, 1950.
3. G. H. Hardy and E. M. Wright, *An introduction to the theory of numbers*, Oxford, 1954.
4. Ddmund Landau, *Handbuch der Lehre von der Verteilung der Primzahlen*, 2nd ed., New York, 1953.
5. ———, *Vorlesungen über Zahlentheorie*, reprint, New York, 1947.
6. S. Swetharanyam, *Asymptotic expressions for certain type of sums involving the arithmetic functions of number theory*, J. Ind. Math. Soc. (to be published), Abstract: Math. Student, **25** (1957), p. 81.
7. E. C. Titchmarsh, *The theory of the Riemann Zeta-function*, Oxford, 1951.

PACIFIC JOURNAL OF MATHEMATICS

EDITORS

DAVID GILBARG

Stanford University
Stanford, California

R. A. BEAUMONT

University of Washington
Seattle 5, Washington

A. L. WHITEMAN

University of Southern California
Los Angeles 7, California

L. J. PAIGE

University of California
Los Angeles 24, California

ASSOCIATE EDITORS

E. F. BECKENBACH

C. E. BURGESS

E. HEWITT

A. HORN

V. GANAPATHY IYER

R. D. JAMES

M. S. KNEBELMAN

L. NACHBIN

I. NIVEN

T. G. OSTROM

H. L. ROYDEN

M. M. SCHIFFER

E. G. STRAUS

G. SZEKERES

F. WOLF

K. YOSIDA

SUPPORTING INSTITUTIONS

UNIVERSITY OF BRITISH COLUMBIA

CALIFORNIA INSTITUTE OF TECHNOLOGY

UNIVERSITY OF CALIFORNIA

MONTANA STATE UNIVERSITY

UNIVERSITY OF NEVADA

OREGON STATE COLLEGE

UNIVERSITY OF OREGON

OSAKA UNIVERSITY

UNIVERSITY OF SOUTHERN CALIFORNIA

STANFORD UNIVERSITY

UNIVERSITY OF TOKYO

UNIVERSITY OF UTAH

WASHINGTON STATE COLLEGE

UNIVERSITY OF WASHINGTON

* * *

AMERICAN MATHEMATICAL SOCIETY

CALIFORNIA RESEARCH CORPORATION

HUGHES AIRCRAFT COMPANY

SPACE TECHNOLOGY LABORATORIES

Printed in Japan by Kokusai Bunken Insatsusha
(International Academic Printing Co., Ltd.), Tokyo, Japan

Pacific Journal of Mathematics

Vol. 9, No. 1

May, 1959

Julius Rubin Blum and Murray Rosenblatt, <i>On the structure of infinitely divisible distributions</i>	1
Robert Geroge Buschman, <i>Asymptotic expressions for</i> $\sum n^a f(n) \log^r n$	9
Eckford Cohen, <i>A class of residue systems (mod r) and related arithmetical functions. I. A generalization of Möbius inversion</i>	13
Paul F. Conrad, <i>Non-abelian ordered groups</i>	25
Richard Henry Crowell, <i>On the van Kampen theorem</i>	43
Irving Leonard Glicksberg, <i>Convolution semigroups of measures</i>	51
Seymour Goldberg, <i>Linear operators and their conjugates</i>	69
Olof Hanner, <i>Mean play of sums of positional games</i>	81
Erhard Heinz, <i>On one-to-one harmonic mappings</i>	101
John Rolfe Isbell, <i>On finite-dimensional uniform spaces</i>	107
Erwin Kreyszig and John Todd, <i>On the radius of univalence of the function</i> $\exp z^2 \int_0^z \exp(-t^2) dt$	123
Roger Conant Lyndon, <i>An interpolation theorem in the predicate calculus</i>	129
Roger Conant Lyndon, <i>Properties preserved under homomorphism</i>	143
Roger Conant Lyndon, <i>Properties preserved in subdirect products</i>	155
Robert Osserman, <i>A lemma on analytic curves</i>	165
R. S. Phillips, <i>On a theorem due to Sz.-Nagy</i>	169
Richard Scott Pierce, <i>A generalization of atomic Boolean algebras</i>	175
J. B. Roberts, <i>Analytic continuation of meromorphic functions in valued fields</i>	183
Walter Rudin, <i>Idempotent measures on Abelian groups</i>	195
M. Schiffer, <i>Fredholm eigen values of multiply-connected domains</i>	211
V. N. Singh, <i>A note on the computation of Alder's polynomials</i>	271
Maurice Sion, <i>On integration of 1-forms</i>	277
Elbert A. Walker, <i>Subdirect sums and infinite Abelian groups</i>	287
John W. Woll, <i>Homogeneous stochastic processes</i>	293