

Pacific Journal of Mathematics

**ON THE BREADTH AND CO-DIMENSION OF A
TOPOLOGICAL LATTICE**

LEE WILLIAM ANDERSON

ON THE BREADTH AND CO-DIMENSION OF A TOPOLOGICAL LATTICE

LEE W. ANDERSON

Consider the following two conjectures:

Conjecture 1. (E. Dyer and A. Shields [7]) If L is a compact, connected, metrizable, distributive topological lattice then $\dim(L) = \text{breadth of } L$.

Conjecture 2. (A. D. Wallace [10]) If L is a compact, connected topological lattice and if $\dim(L) = n$ then the center of L contains at most $2^n - 2$ elements.

The purpose of this note is to prove the following results:

(1) If L is a locally compact distributive topological lattice and if each pair of comparable points is contained in a closed connected chain then the breadth of $L \leq \text{codim}(L)$.

(2) If L is a compact, connected, distributive topological lattice and if $\text{codim}(L) \leq n$ then the center of L contains at most $2^n - 2$ elements.

1. NOTATION. The terminology and notation used in this paper is the same as in [1] [2] and [3]. If L is a lattice, then the *breadth* of L [4], hereafter denoted by $Br(L)$, is the smallest integer n such that any finite subset, F , of L has a subset F' of at most n elements such that $\inf(F) = \inf(F')$.

If A is a subset of a lattice, let $\bigwedge A^n$ denote the set of all elements of the form $x_1 \wedge x_2 \wedge \cdots \wedge x_n$ where $x_i \in A$.

2. $Br(L) \leq cd(L)$. The proof of the following lemma is quite straight forward and will be omitted.

LEMMA 1. *If L is a lattice then the following are equivalent:*

(i) $Br(L) \leq n$

(ii) *If A is an $n + 1$ - element subset of L then A contains an n -element subset B , such that $\inf(A) = \inf(B)$.*

(iii) *If A is a subset of L and if $m, p \geq n$ then $\bigwedge A^m = \bigwedge A^p$.*

If L is a topological lattice, then L is *chain-wise connected* if for each pair of elements, x and y , in L with $x \leq y$ there is a closed connected chain from x to y . Clearly a compact connected topological lattice is chainwise connected.

Problem. Is a locally compact (or locally connected), connected topological lattice chain-wise connected?

THEOREM 1. *If L is a distributive (chain-wise connected) topological lattice then $Br(L) \leq n$ if, and only if, L does not contain a sublattice topologically isomorphic with a Cartesian product of $n + 1$ nondegenerate (closed and connected) chains.*

Proof. If $Br(L) \not\leq n$ then L contains an $n + 1$ element subset, A , such that if B is any proper subset of A then $\inf(A) \neq \inf(B)$. Let x_1, \dots, x_{n+1} be an enumeration of A . Let $b_i = \inf(A \setminus x_i)$, $i = 1, 2, \dots, n + 1$ and let $a = \inf(A)$. Then $b_i \neq a$, $i = 1, 2, \dots, n + 1$ and $b_i \neq b_j$ if $i \neq j$. Let C_i , $i = 1, 2, \dots, n + 1$ be a chain from a to b_i . If L is chain-wise connected we can choose C_i closed and connected. Let $C = C_1 \times C_2 \times \dots \times C_{n+1}$ and define $f: C \rightarrow L$ by $f(x_1, x_2, \dots, x_{n+1}) = x_1 \vee x_2 \vee \dots \vee x_{n+1}$. It is shown in [3] that f is a topological isomorphism, hence the result follows.

If L contains a sublattice, L' , isomorphic with a product of $n + 1$ nondegenerate chains then $Br(L) \not\leq n$ since $Br(L) \geq Br(L') \geq n + 1$.

COROLLARY 1. *If L is a locally compact, chain-wise connected, distributive topological lattice then $Br(L) \leq cd(L)$.*

Proof. Suppose $cd(L) \leq n$ and $Br(L) \not\leq n$. Since L is locally compact and connected it follows that L is also locally convex [1]. Since L is locally convex, the chains C_1, \dots, C_{n+1} chosen in the proof of Theorem 1 can be taken to be compact [2], hence L contains a sublattice topologically isomorphic with a Cartesian product of $n + 1$ nondegenerate compact connected chains. It follows from a result of Cohen [6] that the Cartesian product of $n + 1$ nondegenerate compact connected chains has codimension $n + 1$. Thus it follows that $cd(L) \geq n + 1$ which is a contradiction.

If X is a compact metric space, we denote by 2^X the set of all closed nonvoid subsets of X with the usual Hausdorff metric.

LEMMA 2. *If L is a compact, connected, metrizable topological lattice and if $f: 2^L \rightarrow L$ defined by $f(A) = \inf(A)$ is continuous then L is an absolute retract.*

Proof. If L is a compact topological lattice and if A is a nonvoid subset of L then $\inf(A)$ exists, hence $f(A)$ is defined. If we embed L in 2^L in the usual way and if f is continuous then L is a retract of 2^L . Since L is compact, connected and metrizable, it follows that L is a

Peano continuum [2]. Therefore 2^L is an absolute retract [9] and so L is also an absolute retract.

COROLLARY 2. (Dyer and Shields [7]) *If L is a compact, metrizable, distributive topological lattice and if $cd(L)$ is finite then L is an absolute retract.*

Proof. If $cd(L) = n$ then $Br(L) \leq n$ and so $\bigwedge A^n = \bigwedge A^{n+1} = \dots$ for all $A \subset L$. Let \mathcal{A} denote the set of $A \in 2^L$ such that $\inf(A) \in A$. It is known [5] that $f: \mathcal{A} \rightarrow L$ defined by $f(A) = \inf(A)$ is continuous. Define $g: 2^L \rightarrow \mathcal{A}$ by $g(A) = \bigwedge A^n$ then clearly g is continuous and so $F: 2^L \rightarrow L$ defined by $F(A) = f(g(A)) = \inf(A)$ is continuous. Thus it follows from Lemma 2 that L is an absolute retract.

Problem. Is $A \rightarrow \inf(A)$ continuous if L is not distributive and not finite dimensional?

3. *On the set $\mathcal{B}(x)$.* If L is a lattice and $a \in L$, let $\mathcal{M}(a)$ denote the set of all subsets, M , of L that satisfy

- (i) $M \wedge M \subset M$
- (ii) $a \notin M$
- (iii) M is maximal with respect to (i) and (ii).

Let $\mathcal{B}(a)$ denote the set of all complements of elements in $\mathcal{M}(a)$.

LEMMA 3. *If L is a lattice and $a \in L$ then $\bigcap \{B: B \in \mathcal{B}(a)\} = \{a\}$.*

Proof. If $x \in \bigcap \{B: B \in \mathcal{B}(a)\}$ and if $x \neq a$ then by the Hausdorff Maximality Principle, there is a maximal \wedge -closed set, M , containing x but not containing a . But then $M \in \mathcal{M}(a)$ and so $x \notin L \setminus M \in \mathcal{B}(a)$. It is clear that $a \in \bigcap \{B: B \in \mathcal{B}(a)\}$, hence the result is established.

LEMMA 4. *If L is a lattice and if $a \in L$, $B \in \mathcal{B}(a)$ then $a \vee L \subset B$ if, and only if, $a = 1$.*

Proof. If $a \vee L \subset B$ then $a \vee L \subset \bigcap \{B: B \in \mathcal{B}(a)\} = \{a\}$ and so $a = 1$. If $a = 1$ then $a \vee L = \{1\} = B$.

LEMMA 5. *If L is a lattice and if $a \in L$, $a \neq L$, $B \in \mathcal{B}(a)$, $M = L \setminus B \in \mathcal{M}(a)$ then $x \in B$ if, and only if, $a \in x \wedge M$.*

Proof. If $a \in x \wedge M$ and if $x \notin B$ then $x \in M$ and so $a \in x \wedge M \subset M$ which is a contradiction. If $x \in B$ and $x = a$ then, since $a \neq 1$, by

Lemma 4 we have $M \cap (a \vee L) \neq \varnothing$ and hence $a \in a \wedge M$. If $x \in B$ and if $x \neq a$ and if $a \notin x \wedge M$ then, since

$$(\{x\} \cup (x \wedge M)) \wedge (\{x\} \cup (x \wedge M)) \subset \{x\} \cup (x \wedge M),$$

we have $\{x\} \cup (x \wedge M) \subset M$. This, however, is a contradiction since $x \in B = L \setminus M$.

LEMMA 6. *If L is a lattice and if $a \in L$, $b \in B \in \mathcal{B}(a)$ and if $y \geq a$ then $y \wedge b \in B$.*

Proof. If $b \in B \in \mathcal{B}(a)$, there is an $x \in M = L \setminus B$ such that $b \wedge x = a$. Now $x \wedge (b \wedge y) = (x \wedge b) \wedge y = a \wedge y = a$ and so, by Lemma 5, $b \wedge y \in B$.

LEMMA 7. *If L is a lattice and if $a \in L$, $b \in B_0$, $b \neq a$ and $b \notin \cup \{B: B \in \mathcal{B}(a), B \neq B_0\}$ then*

$$\{y \in L: y \wedge b = a, y \neq a\} \subset \cap \{B: B \in \mathcal{B}(a), B \neq B_0\} \cap M_0$$

where $M_0 = L \setminus B_0$. Moreover if $y \wedge b = a$ and $y \neq a$ then

$$B_0 = \{x \in L: x \wedge y = a\}.$$

Proof. Let $y \in L$ such that $y \wedge b = a$ and let $B \in \mathcal{B}(a)$ be distinct from B_0 . Now if $y \notin B$ then $y, b \notin B$ and so $y, b \in L \setminus B \in \mathcal{M}(a)$. But $y \wedge b = a$ which is a contradiction and so $y \in \cap \{B: B \in \mathcal{B}(a), B \neq B_0\}$. Now $y \neq a$ and $y \wedge y = y$, thus there is an $M \in \mathcal{M}(a)$ with $y \in M$. However $y \in \cap \{B: B \in \mathcal{B}(a), B \neq B_0\}$ and therefore $M = M_0$. Now if $y \wedge b = a$ and $x \in B_0$ then $y \wedge x \in \cap \{B: B \in \mathcal{B}(a)\} = \{a\}$ and so $y \wedge x = a$. Also if $y \wedge b = a$ and $y \neq a$ then $y \in M_0$, and so if $y \wedge x = a$ then $x \in B_0$.

LEMMA 8. *If L is a distributive lattice and if $Br(L) = n$ then $\sup \{\text{card}(\mathcal{B}(x)): x \in L\} = n$.*

Proof. Suppose that for some $a \in L$, $\text{card}(\mathcal{B}(a)) \geq n + 1$. Pick $n + 1$ distinct members of $\mathcal{B}(a)$, say B_1, \dots, B_{n+1} . Since L is distributive, we can pick, for each $i = 1, 2, \dots, n + 1$, an $x_i \in B_i$ such that $x_i \notin B_j$ if $i \neq j$. Thus it follows that

$$\inf \{x_i: i = 1, 2, \dots, n + 1\} \in B_1 \cap B_2 \cap \dots \cap B_{n+1}$$

but $\inf \{x_i: i \neq j \text{ and } i = 1, 2, \dots, n + 1\} \notin B_j$ and so $Br(L) \geq n + 1$. Therefore $\text{card}(B(x)) \leq n$ for all $x \in L$.

Now $Br(L) = n$ and so there is an n -element set, say A , such that

$\inf(A) \neq \inf(A')$ for all proper subsets A' of A . Thus for each $a \in A$ we can find a $B \in \mathcal{B}(\inf(A))$ such that $a \in B$ and $A \setminus \{a\} \subset L \setminus B$ and so $\text{card}(\mathcal{B}(\inf(A))) \geq n$.

LEMMA 9. *If L is a distributive topological lattice and if $a \in L$ and if $\text{card}(\mathcal{B}(a))$ is finite then each $B \in \mathcal{B}(a)$ is a closed sublattice of L .*

Proof. Let B_1, B_2, \dots, B_n be an enumeration of $\mathcal{B}(a)$. We will show that B_1 is a closed sublattice of L . Since L is distributive, we can pick $b \in B_1$ so that $b \notin B_i$ if $i \neq 1$. Thus there is a $y \in B_2 \cap \dots \cap B_n$ such that $y \neq a$ and $y \wedge b = a$. By Lemma 7, $B_1 = \{x \in L: x \wedge y = a\}$ and so B_1 is closed. Since L is distributive, B_1 is clearly a sublattice of L .

Problem. If L is a topological lattice and if $a \in L$, and $B \in \mathcal{B}(a)$ is B closed?

THEOREM 2. *If L is a compact, connected, distributive topological lattice and if $cd(L) \leq n$ and if $a \in L$ and $B \in \mathcal{B}(a)$ then $cd(B) \leq n - 1$.*

Proof. We first prove the theorem for the case $n > 1$. By way of a contradiction let us assume that $cd(L) \leq n$ and $cd(B) > n - 1$. Then for some closed set $A \subset B$ we have $H^n(B, A) \neq 0$. Since B is a closed sublattice of L we have, letting $b = \sup(B), b \in B$. To simplify our notation, we let $C = \{x \in L: x \wedge b = a\}, c = \sup(C), D = c \vee L, E = C \vee A$ and $F = B \cup E \cup D$. It follows that $B \cap C = \{a\}$ and $B \cap (E \cup D) = A$, and that C, D, E , and F are closed. We will now show that if $p > 0, H^p(E \cup D) = 0$. Define $f: (E \cup D) \times C \rightarrow E \cup D$ by $f(x, y) = x \vee y$. Clearly f is defined and continuous. For each $y \in C$ define $F_y: E \cup D \rightarrow E \cup D$ by $F_y(x) = f(x, y)$ then, since $E \cup D$ is compact and C is connected, it follows from the Generalized Homotopy lemma that $F_a^* = F_c^*$.

Now F_c retracts $E \cup D$ onto D and, since $H^n(D) = 0$, it follows that $F_c^* = 0$. Also F_a is the identity function and therefore $H^n(E \cup D) = 0$. Now consider the following Mayer-Victoris exact sequence [8]:

$$H^{n-1}(E \cup D) \times H^{n-1}(B) \xrightarrow{I^*} H^{n-1}(A) \xrightarrow{\Delta^*} H^n(F) \xrightarrow{J^*} H^n(E \cup D) \times H^n(B).$$

Now $H^{n-1}(E \cup D) = H^{n-1}(B) = H^n(E \cup D) = H^n(B) = 0$, and so Δ^* is an isomorphism onto. It therefore follows that $H^n(F) \neq 0$ which contradicts the fact that $cd(L) \leq n$ and $H^n(L) = 0$.

In the case $n = 0, L$ is a single point and therefore the result is trivial. If $n = 1$ then L is a chain [1] and so B is at most a single point which implies that $cd(B) \leq 0$.

We recall (see e.g. [3] or [4]) that if L is a lattice with 0 and 1 then the center of L , denoted by $\text{Cen}(L)$, is the set of all $x \in L$ other than 0 and 1 such that for some $y \in L$, $x \wedge y = 0$ and $x \vee y = 1$. If L is distributive and if $x \in \text{Cen}(L)$ then there is a unique element, denoted by $c(x)$, such that $x \wedge c(x) = 0$ and $x \vee c(x) = 1$.

COROLLARY. *If L is a compact, connected, distributive topological lattice and if $cd(L) \leq n$ then $\text{card}(\text{Cen}(L)) \leq 2^n - 2$.*

Proof. We proceed by finite induction. If $cd(L) \leq 1$ then L is a chain and so $\text{card}(\text{Cen}(L)) \leq 0$. Suppose the theorem is true for all $n < k$ and suppose $cd(L) \leq k$. If $a \in \text{Cen}(L)$, choose $M \in \mathcal{N}(0)$ such that $a \in M$ so that $B = L \setminus M \in \mathcal{B}(0)$. Thus if $\mathcal{B}(0)$ is empty then $\text{Cen}(L)$ is also empty and the result is established. If $\mathcal{B}(0)$ is not empty, let $B \in \mathcal{B}(0)$. It follows from lemma [9] that B is a closed sublattice of L . Letting $b = \sup(B)$ we have that $b \in B$. We will now show that if $a \in \text{Cen}(L)$ then either $b \wedge a = 0$, $b \wedge a = b$, $a \in \text{Cen}(B)$ or $c(a) \in \text{Cen}(B)$. If $a \wedge b \neq 0$, b and if $a \notin B$ and if $c(a) \notin B$ then $a, c(a) \in L \setminus B$ and so $a \wedge c(a) \neq 0$ which is a contradiction. Therefore $a \in B$ or $c(a) \in B$. Now if $a \in B$ then $a \wedge (c(a) \wedge b) = 0$ and

$$a \vee (c(a) \vee b) = 1 \vee (a \vee b) = 1 \wedge b = b$$

and so $a \in \text{Cen}(B)$. Similarly if $c(a) \in B$ then $c(a) \in \text{Cen}(B)$. If $a, c(a) \in B$ then $a \vee c(a) = 1 \in B$ which is a contradiction. If $a \wedge b = 0$ then $a \notin B$ and since $b \notin \cup \{A \in \mathcal{B}(0): A \neq B\}$ we have, by Lemma 7, that $B = \{x \in L; x \wedge a = 0\}$. Thus it follows that $c(a) \in B$. Therefore $1 = a \vee c(a) \leq a \vee b$ which implies that $c(a) = b$ and $a = c(b)$. If $a \wedge b = b$ then $c(a) \wedge b = 0$ and so $c(a) = c(b)$ which implies that $a = b$. It follows, therefore, that $\text{card}(\text{Cen}(L)) \leq 2 \text{card}(\text{Cen}(B)) + 2$. Now $cd(B) \leq k - 1$ and so $\text{card}(\text{Cen}(B)) \leq 2^{k-1} - 2$ and so

$$\text{card}(\text{Cen}(L)) \leq 2(2^{k-1} - 2) + 2 = 2^k - 2.$$

REFERENCES

1. L. W. Anderson, *On one dimensional topological lattices*, Proc. Amer. Math. Soc., to appear.
2. ———, *On the distributivity and simple connectivity of plane topological lattices*, Trans. Amer. Math. Soc., to appear.
3. ———, *Topological lattices and n -cells*, Duke Math. Jour., **25** (1958), 205-208.
4. G. Birkhoff, *Lattice theory*, Amer. Math. Soc., Colloquium Pub, Vol. 25, New York, 1948.
5. C. Capel and W. Strother, *Multi-valued functions and Partial order*, Port. Math., **17** (1958), 41-47.
6. H. Cohen, *A cohomological definition of dimension for locally compact Hausdorff*

spaces, Duke Math. Jour., **21** (1954), 209-224.

7. E. Dyer and A. Shields, *On a conjecture of A. D. Wallace*, Mich. Math. Jour., to appear.

8. S. Eilenberg and N. Steenrod, *Foundations of algebraic topology*, Princeton University Press, 1952.

9. J. L. Kelley, *Hyperspaces of a continuum*, Trans. Amer. Math. Soc., **52** (1942), 23-36.

10. A. D. Wallace, *Factoring a lattice*, Proc. Amer. Math. Soc., **9** (1958), 250-252.

THE UNIVERSITY OF OREGON, EUGENE, OREGON

PACIFIC JOURNAL OF MATHEMATICS

EDITORS

DAVID GILBARG
Stanford University
Stanford, California

R. A. BEAUMONT
University of Washington
Seattle 5, Washington

A. L. WHITEMAN
University of Southern California
Los Angeles 7, California

L. J. PAIGE
University of California
Los Angeles 24, California

ASSOCIATE EDITORS

E. F. BECKENBACH
C. E. BURGESS
E. HEWITT
A. HORN

V. GANAPATHY IYER
R. D. JAMES
M. S. KNEBELMAN
L. NACHBIN

I. NIVEN
T. G. OSTROM
H. L. ROYDEN
M. M. SCHIFFER

E. G. STRAUS
G. SZEKERES
F. WOLF
K. YOSIDA

SUPPORTING INSTITUTIONS

UNIVERSITY OF BRITISH COLUMBIA
CALIFORNIA INSTITUTE OF TECHNOLOGY
UNIVERSITY OF CALIFORNIA
MONTANA STATE UNIVERSITY
UNIVERSITY OF NEVADA
OREGON STATE COLLEGE
UNIVERSITY OF OREGON
OSAKA UNIVERSITY
UNIVERSITY OF SOUTHERN CALIFORNIA

STANFORD UNIVERSITY
UNIVERSITY OF TOKYO
UNIVERSITY OF UTAH
WASHINGTON STATE COLLEGE
UNIVERSITY OF WASHINGTON
* * *
AMERICAN MATHEMATICAL SOCIETY
CALIFORNIA RESEARCH CORPORATION
HUGHES AIRCRAFT COMPANY
SPACE TECHNOLOGY LABORATORIES

Mathematical papers intended for publication in the *Pacific Journal of Mathematics* should be typewritten (double spaced), and the author should keep a complete copy. Manuscripts may be sent to any one of the four editors. All other communications to the editors should be addressed to the managing editor, L. J. Paige at the University of California, Los Angeles 24, California.

50 reprints per author of each article are furnished free of charge; additional copies may be obtained at cost in multiples of 50.

The *Pacific Journal of Mathematics* is published quarterly, in March, June, September, and December. The price per volume (4 numbers) is \$12.00; single issues, \$3.50. Back numbers are available. Special price to individual faculty members of supporting institutions and to individual members of the American Mathematical Society: \$4.00 per volume; single issues, \$1.25.

Subscriptions, orders for back numbers, and changes of address should be sent to Pacific Journal of Mathematics, 2120 Oxford Street, Berkeley 4, California.

Printed at Kokusai Bunken Insatsusha (International Academic Printing Co., Ltd.), No. 6, 2-chome, Fujimi-cho, Chiyoda-ku, Tokyo, Japan.

PUBLISHED BY PACIFIC JOURNAL OF MATHEMATICS, A NON-PROFIT CORPORATION

The Supporting Institutions listed above contribute to the cost of publication of this Journal, but they are not owners or publishers and have no responsibility for its content or policies.

Pacific Journal of Mathematics

Vol. 9, No. 2

June, 1959

Lee William Anderson, <i>On the breadth and co-dimension of a topological lattice</i>	327
Frank W. Anderson and Robert L. Blair, <i>Characterizations of certain lattices of functions</i>	335
Donald Charles Benson, <i>Extensions of a theorem of Loewner on integral operators</i>	365
Errett Albert Bishop, <i>A duality theorem for an arbitrary operator</i>	379
Robert McCallum Blumenthal and Ronald Kay Getoor, <i>The asymptotic distribution of the eigenvalues for a class of Markov operators</i>	399
Delmar L. Boyer and Elbert A. Walker, <i>Almost locally pure Abelian groups</i>	409
Paul Civin and Bertram Yood, <i>Involutions on Banach algebras</i>	415
Lincoln Kearney Durst, <i>Exceptional real Lehmer sequences</i>	437
Eldon Dyer and Allen Lowell Shields, <i>Connectivity of topological lattices</i>	443
Ronald Kay Getoor, <i>Markov operators and their associated semi-groups</i>	449
Bernard Greenspan, <i>A bound for the orders of the components of a system of algebraic difference equations</i>	473
Branko Grünbaum, <i>On some covering and intersection properties in Minkowski spaces</i>	487
Bruno Harris, <i>Derivations of Jordan algebras</i>	495
Henry Berge Helson, <i>Conjugate series in several variables</i>	513
Isidore Isaac Hirschman, Jr., <i>A maximal problem in harmonic analysis. II</i>	525
Alfred Horn and Robert Steinberg, <i>Eigenvalues of the unitary part of a matrix</i>	541
Edith Hirsch Luchins, <i>On strictly semi-simple Banach algebras</i>	551
William D. Munro, <i>Some iterative methods for determining zeros of functions of a complex variable</i>	555
John Rainwater, <i>Spaces whose finest uniformity is metric</i>	567
William T. Reid, <i>Variational aspects of generalized convex functions</i>	571
A. Sade, <i>Isomorphisme d'hypergroupoï des isotopes</i>	583
Isadore Manual Singer, <i>The geometric interpretation of a special connection</i>	585
Charles Andrew Swanson, <i>Asymptotic perturbation series for characteristic value problems</i>	591
Jack Phillip Tull, <i>Dirichlet multiplication in lattice point problems. II</i>	609
Richard Steven Varga, <i>p-cyclic matrices: A generalization of the Young-Frankel successive overrelaxation scheme</i>	617